



22400025944

Med
K4542

to John Drummond Hay K.
with the kind remembrances of
the Author.

SPICILEGIUM
FLORÆ MAROCCANÆ.

BY

JOHN BALL, F.R.S., F.L.S.

[*Extracted from the* LINNEAN SOCIETY'S JOURNAL—BOTANY,
vol. xvi.]

LONDON:

PRINTED BY TAYLOR AND FRANCIS, RED LION COURT, FLEET STREET.

1878.

867-



17822 736

WELLCOME INSTITUTE LIBRARY	
Coll.	weIMOmec
Call	
No.	GK

EXPLANATION OF THE PLATES.

PLATE

- I. MARATTIA FRAXINEA.—Views of the rootstock and of its microscopical structure, to illustrate Mr. J. Buchanan's paper thereon.
- II. FUNGI, viz. *Uromyces deformans* and *Puccinia corticioides*, and magnified details of same, illustrating the Rev. M. J. Berkeley's description of remarkable Fungi among the 'Challenger' Expedition collection.
- III. DISARTICULATION OF BRANCHES.—Sketches showing mode of, according to the observations of Mr. R. Irwin Lynch.
- IV. } HEPATICÆ.—Structural peculiarities of new species from the Cape of
V. } Good Hope described by Mr. W. Mitten.
- VI. ALGÆ.—New freshwater forms from the Cape of Good Hope, referred to by P. F. Reinsch in his paper on the Rev. A. Eaton's 'Transit-of-Venus' collection.
- VII. } RUBIACÆ.—New species of African forms, viz. *Pentas parvifolia*
VIII. } and *Trichostachys vaginalis*, characterized by Mr. W. P. Hiern.
- IX. NASTURTIUM ATLANTICUM.
- X. ARABIS DECUMBENS.
- XI. ARABIS CONRINGIODES.
- XII. BRASSICA ELATA.
- XIII. BRASSICA RERAYENSIS.
- XIV. RESEDA ATTENUATA.
- XV. LOTONONIS MAROCCANA.
- XVI. ONONIS POLYPHYLLA.
- XVII. ONONIS ATLANTICA.
- XVIII. TRIFOLIUM ATLANTICUM.
- XIX. RUBUS DEBILIS.
- XX. SEDUM MODESTUM.
- XXI. GALIUM ACUMINATUM.
- XXII. GALIUM BOURGÆANUM, var. MAROCCANUM.
- XXIII. GNAPHALIUM HELICHRYSOIDES.
- XXIV. ANACYCLUS DEPRESSUS.
- XXV. ANTHEMIS TENUISECTA.
- XXVI. CHRYSANTHEMUM ATLANTICUM.
- XXVII. THYMUS MAROCCANUS.
- XXVIII. SALVIA MAURORUM.

New species of Plants,
with structural details,
described by
Mr. J. Ball in his
"Speilegium Floræ
Maroccanæ."

CORRIGENDA ET ADDENDA.

- Page 2, line 15 from bottom, for "*glaucomeria*" read *glaucomaria*.
 — 6, — 8 from bottom, for "*Chroospora*" read *Chnoospora*.
 — 9, — 12 from bottom, for "*Psiacelia*" read *Psiadia*.
 — 11, — 19 from bottom, for "*Hyprestes*" read *Hypoëstes*.
 — 11, — 10 from bottom, for "*rodriguesii*" read *rodriguesiana*.
 — 282, — 4 from bottom. "*Broussonnet*" here and elsewhere throughout the paper is, according to the common usage of botanists, spelt with a double n; but that author himself, in the "*Opuseules*," Paris, 1785, spells his name with but a single n, *i. e.* Broussonet.
 — 325, — 15 from top, for "*D. auriculata*" read *D. tenuisiliqua*.
 — 325, — 9 from bottom, for "*Sinapis hispida*" read *Sinapis radicata*.
 — 332, after line 15 from top *insert*:—*Ar. Geog. Regio Mediterranea occidentalis. Proxime affinis B. Columnæ, Ten., in Oriente occurrit.*
 — 332, before line 18 from bottom *insert*:—*Ar. Geog. Regio Mediterranea occidentalis.*
 — 352, line 3 from bottom, at end *insert*:—*In hortis urbis Maroece*
 — 361, — 20 from bottom, for "*quæ*" read *quam*.
 — 401, — 16 from top, for "*florentem*" read *florens*.
 — 406, — 16 from top, for "*Mogadorensis*" read *Boreali-Africana*.
 — 408, — 3 from bottom, for "*O. Schounii*" read *O. Schouwii*.
 — 421, — 4 from top, after "*raya!*" *insert*:—*Vidimus in horto Ben Dreis urbis Maroece!*
 — 488, — 19 from bottom, for "*Aparini*" read *Aparinæ*.
 — 499, — 20 from top, for "*ebracteolati*" read *braeteolati*.
 — 534, — 3 from top, for "*phaiolepis*" read *phialolepis*.
 — 564, after line 20 from top *insert*:—*Ar. Geog. Algeria interior. Marocco meridionalis.*
 — 585, line 21 from top, for "*Reuta*" read *Reuter*.
 — 626, — 6 from bottom, for "*insula*" read *insulam*.
 — 628, — 20 from top, for "*nemini*" read *nullo*.
 — 630, — 18 from top, for "*nemini*" read *nullo*.
 — 631, — 17 from bottom, transpose "*et*" after "*T. Scorodoniæ*."
 — 633, — 15 from top, for "*aut*" read *vel*.
 — 675, — 17 from bottom, for "*variatates*" read *varietates*.
 — 694, — 17 from bottom, for "*faice*" read *faeie*.
 — 696, — 7 from top, for "*nostra*" read *nostro*.
 — 702, — 8 from bottom, for "*australi*" read *australis*.
 — 717, — 16 from top, for "*Lim*" read *Limnetis*.
 — 719, — 3 from top, for "*multis*" read *ineultis*.
 — 720, — at bottom for "*nex*" read *nee*.
 — 726, — 4 from bottom, *strike out* "*cum*" before "*typica*."

Spicilegium Floræ Maroccanæ.
By JOHN BALL, Esq., F.R.S., F.L.S.

[Read March 1, 1877.]

(PLATES IX.-XXVIII.)

INTRODUCTORY OBSERVATIONS.

THE territory of Marocco extends from the Straits of Gibraltar, with the neighbouring part of the Mediterranean at its northern extremity, to the great desert on the southern side of the Great Atlas, and from the frontier of Algeria to the Atlantic coast of Northern Africa. Of this extensive region, about equal in extent to Spain, it may be truly said that none other so easy of access is so little known to geographers.

Although the chief ports are within a few days' journey from London or Marseilles, and are freely open to Europeans, so little is known of the interior that the maps, all founded on native information, which profess to represent the direction of the mountain-ranges and the course of the chief rivers, are hopelessly at variance, and of the best of them it may be said that the little positive knowledge we possess shows that even in its main features it is very wide of the truth.

The causes of our ignorance of the country are easily stated. The traditional policy of the government has been hostile to the admission of strangers into its territory; and as regards the natives of Christian States the difficulties have been aggravated by the fanatical character of a great part of the native population. But a still more serious, and probably a more enduring, obstacle to exploration arises from the fact that fully two thirds of the entire country is inhabited by independent tribes, who recognize no external authority, even when they accord a nominal supremacy to the Sultan of Marocco. These tribes, descendants of the original Berber population of Northern Africa, which has never been subjugated by any of the foreign rulers who have held the coast and the open country, occupy nearly all the mountain region. Possessing, it would seem, many of the rude virtues common to such populations, they are constantly engaged in internal petty warfare, and always disposed to regard strangers not of their own tribe as lawful prey.

Even in the parts of Marocco where the natives have approached the limits of European civilization, access to mountain districts is

all but impossible to strangers. The fine peaks of the Beni Hassan, within sight of the rock of Gibraltar and easy reach of Tangier and Tetuan, have never been approached by a traveller; and Sir John Drummond Hay, whose knowledge of the people and influence amongst them surpass that of any other European, pronounced against the feasibility of the excursion which we earnestly desired to make in that direction.

As a necessary consequence of what has been stated, our knowledge of the Flora of Marocco has been extremely limited, and, in spite of what has been done during the last few years, is still most imperfect; and the time is yet distant when it will be possible to give more than a rude outline of the relations between the vegetable population and those of the surrounding regions.

The earliest notice that I have found of botanical research into the Marocco Flora is in the 'Istoria Botanica' of Zanoni (Curator of the Botanic Garden at Bologna), published in 1675. He had received plants and seeds from Tangier from a Mr. Alexander Balaam, who appears to have been a merchant trading with Tangier during some part of the time when that town was held by the Portuguese and the English. The next contribution, and a more considerable one, is contained in a list published (1696) in the 19th vol. of the 'Philosophical Transactions' (p. 239 *et seq.*), but drawn up in 1673 by Mr. Spotswood, a surgeon who lived there during the English occupation, and gathered plants which he gave to the same Mr. Balaam and to Dr. Morison at Oxford. There is some difficulty in identifying many of the species named; and I have not thought it worth while to cite the plants contained in Mr. Spotswood's list.

After the evacuation of Tangier by the English, under circumstances little creditable to our national honour, communication between Europe and Marocco became more unfrequent than ever, and, excepting some occasional negotiation for the release of Christian captives, seems to have almost ceased. At the latter end of the last century relations between some European powers and the Marocco Government became more frequent and almost amicable, so that travelling in the country seems to have been easier than it has been in more recent times. A French botanist, M. Broussonnet, who also visited the Canary Islands and some parts of Spain, spent some time in Marocco during the last ten years of the century. He explored the neighbourhood of Mogador and Tangier, visited several of the ports on the Atlantic coast,

and made at least one journey into the interior, in which he reached Fez and Mequinez. Broussonnet sent specimens to the chief botanists of his day—to Willdenow at Berlin, to Desfontaines at Paris, to Gouan at Montpellier, and to Cavanilles at Madrid. Most of the latter were published by that excellent botanist in the '*Anales de Ciencias Naturales*,' a now scarce periodical which secured for Spain a temporary place in the republic of science, which she did not long retain. A few of Broussonnet's plants were also published by Willdenow in his edition of the '*Species Plantarum*;' and a few others are referred to in Desfontaines's '*Flora Atlantica*;' but the greater part of his collections, now in the museum at Montpellier, remained unexamined until M. Cosson visited that city. The results of the latter's study of that collection form a portion of the materials, which we hope soon to see published. Unfortunately Broussonnet seems to have been somewhat careless respecting the localities whence his specimens came, and to have sometimes intermixed those from the Canary Islands, South Morocco, Tangier, and Spain. Several undoubted errors that have been detected throw doubt on the authenticity of his localities when unconfirmed by other authorities.

About the time of Broussonnet's visits to Morocco Mr. Schousboe was appointed as Danish consular agent at Mogador, and subsequently at Tangier. He was an active and intelligent observer, who studied carefully the vegetation of both the districts above named. Besides this he collected a few plants in the course of a journey in which he reached the city of Morocco, and another which led him to Fez and Mequinez. He was the author of the first attempt at a Flora of Morocco, the first portion of which, including the Linnean classes from Monandria to Enneandria, with a few new species of other classes, was published in Danish and in German in 1801. Unfortunately, the work remained unfinished. Although Schousboe lived for many years afterwards chiefly at Tangier, and added very largely to his collections, I am not aware that he worked at the continuation of the work, which, in the German edition, bears the title "*Beobachtungen über das Gewächsreich in Marokko*." Schousboe's herbarium is preserved at Copenhagen; but the large collection of his duplicates passed not long ago into the possession of M. Cosson, who has liberally given sets of them to the Kew Herbarium and to myself.

Early in this century a French ecclesiastic, the Abbé Philippe

Durand, visited Morocco, reached Fez and Mequinez, and, so far as I know, alone of European travellers traversed some part of the great forest of Mamora, respecting which many a strange story circulates among the natives of Morocco.

In 1809 was published the first edition of the work which gives by far the fullest account of Morocco that has yet appeared, 'An Account of the Empire of Morocco,' by James Grey Jackson. Mr. Jackson lived for the greater part of sixteen years in South Morocco; he became familiar with the language and manners of the people, and must have acquired more of their confidence than any previous European visitor, as he not only travelled by several routes through the lower country near the coast, but was allowed to join a military force proceeding over the Atlas from Morocco to Tarudant.

Mr. Jackson does not, however, seem to have ventured on entering the city of Tarudant; nor, though he approached Wadnoon, did he actually visit that place.

Though not a naturalist, Jackson was a very intelligent observer; and on some points, especially regarding the curious cactoid Euphorbias of South Morocco, he has given information interesting to botanists.

The next contributor to the knowledge of the Morocco flora was M. Salzmann, an active collector who, after previous visits to other interesting spots in the Mediterranean, spent a considerable time at Tangier in 1825, and seems to have explored very carefully the district within convenient reach of that city. Salzmann does not appear to have had much scientific knowledge; but he was a sharp-eyed collector, and often detected differences between plants found by him and allied species, with which they had been confounded by other botanists. Several of these were published by the late M. A. P. Decandolle, then at the zenith of his scientific fame and authority, who had shortly before commenced the publication of that colossal work, the 'Prodromus Systematis Naturalis.' Others remained unpublished, except for the manuscript name attached to them by Salzmann. As, however, his collections were sold, and were sufficiently numerous to supply the demands of the chief botanists of his time, this has been, I think, justly regarded as equivalent to publication, and as constituting a title to priority over names subsequently given by other botanists.

Very complete sets of Salzmann's Tangier plants were con-

tained in the herbarium of Mr. Bentham (presented by him to the national collection at Kew), and in that of my late lamented friend M. Jacques Gay (purchased by Dr. Hooker, and also given to the Kew Herbarium); so that I have had the opportunity of examining original specimens of nearly all Salzmann's plants.

In 1827 my excellent friend the late Philip Barker Webb spent three weeks at Tangier, and then visited Tetuan, where he was able to ascend two of the neighbouring mountains, and to discover amongst other novelties a new genus of Cruciferae, for which, up to the present time, no other locality is known. His collections, preserved along with the remainder of his great herbarium at Florence, have been examined by M. Cosson and by myself.

During the years between 1840 and 1870 many botanists touched at Tangier, including my friend M. Boissier, with the late M. Renter, Dr. Lagrange, M. Blanche, M. Marés, and M. Jourdan. Several of the plants collected by MM. Boissier and Reuter were described by them in a little volume styled 'Pugillus Plantarum Novarum Africæ borealis Hispaniæque australis.' In that work the criteria for the admission of new species, so justly maintained in the great works on which M. Boissier's fame depends, were somewhat unduly relaxed.

In 1859 the late Rev. R. T. Lowe contributed to the 'Proceedings' of this Society a list of plants observed by him in the immediate neighbourhood of Mogador; and most of these are preserved in the herbarium bequeathed by him to the Royal Gardens at Kew.

In June 1851 I made an attempt to reach the higher summits of the Lesser Atlas in the neighbourhood of Tetuan; but, owing to the disturbed condition of the district, I was limited to the immediate neighbourhood of the city, which, however, offers many species of interest not found about Tangier.

In 1869 my indefatigable friend, Mr. George Maw, was somewhat more successful than myself, and was permitted to ascend part of the Beni Hosmar, previously visited by Webb.

In 1867, chiefly through the influence and with the aid of M. Cosson, whose intimate acquaintance with the flora of Northern Africa is unrivalled by any competitor, the successful botanical traveller Mr. Balansa was despatched to Mogador, with the object of attempting to investigate the flora of the Great Atlas range as far as circumstances would permit. Although receiving every

assistance from the French diplomatic agents in Morocco, his progress was impeded at every step; and after about a fortnight he was forced to abandon the attempt and return to Mogador. He nevertheless was able to collect a large number of the new and remarkable species which characterize the skirts of the great mountain-range and the lower mountains that diverge from the main chain.

The short time at his disposal enabled M. Balansa to collect so few duplicates that many of the new species subsequently described by M. Cosson are not to be found in the principal herbaria of Europe, and several of these, as well as the other plants of his journey, are preserved only in the magnificent herbarium of M. Cosson.

Along with other materials scarcely to be found outside that herbarium, I should mention a collection made by M. Warion, in the medical service of the French army, a good observer and collector, who gathered many plants on the eastern frontier of Morocco, adjoining the French province of Oran.

Early in the year 1871 Dr. Hooker, who had long felt a desire to explore the range of the Great Atlas, applied to Earl Granville, then Secretary of State for Foreign Affairs, to use the influence of the English Government at the court of Morocco to obtain the necessary permission of the Sultan. The information received from Sir John Drummond Hay, our minister at the court of Morocco, was so far favourable that Dr. Hooker determined to leave England in the spring of that year; and I was fortunate enough to be able to accompany him, along with our mutual friend Mr. George Maw, who, as well as myself, had already made an attempt to visit the extremity of the chain of the Lesser Atlas of Morocco in the neighbourhood of Tetuan.

We received before starting valuable information and aid from our excellent friend M. Cosson, who was kind enough to prepare a manuscript list of all the plants then known or believed to inhabit the Morocco territory, as far as his unequalled means of information would go.

In our subsequent course we were under deep obligations to our active and energetic representative in Morocco already named, and to all the British consular agents on the coast, and especially to the late Mr. Carstensen, then Vice-consul at Mogador, through whose hospitality and active assistance our stay was rendered agreeable, and our journey into the interior materially expedited.

After three days passed at Tangier, during which we made an excursion to Cape Spartel, and thence along the Atlantic coast to the so-called Cave of Hercules, we started for Tetuan, which we reached early on the 12th of April. On the following day we ascended the highest of the picturesque mountains that enclose the verdant valley of Tetuan, called Beni Hosmar, the same which had been visited more than forty years before by the late Mr. Webb. Although the season was not yet sufficiently advanced, we collected nearly all the species found by our predecessor, with several others of much interest. After another day well spent on the rocky ground in the immediate neighbourhood of the city, we started on the 15th to ride along the coast to Ceuta, thereby adding some additional species to our rapidly growing collections. The botanical exploration of the coast between Ceuta and Tangier, on the seaward slopes of the Angera Mountains, would be of great interest; and until this has been effected it will not be possible to say whether *Rhododendron ponticum*, *Helianthemum lasianthum*, and several other remarkable species now known on the northern side of the Straits of Gibraltar are really absent from the southern shores. The country, however, is considered too unsafe, especially for travellers starting from Ceuta, and we traversed the Straits in the felucca, by which the garrison and population of Ceuta keep up communication with Algeciras. Two days' unexpected detention on the Spanish side was not ill employed in a survey of the spring vegetation of the hills above Algeciras and the rock of Gibraltar. Returning to Tangier we were detained two days more, which were employed in putting in order and further increasing our collections. Mr. Maw, who had returned direct from Tetuan to Tangier, made in the interval two fruitful excursions to some swampy brackish lakes about 14 miles S.S.W. of the latter city.

Having at length received the necessary letter from the Sultan to the Governor of Mogador, we left Tangier on the 20th of April in the French steamer 'Vérité,' bound for the Atlantic ports of Morocco and the Canary Islands. On the following day we lay off Rabat, but, owing to the heavy swell, were unable to land. On the following day, however, we landed at Casa Blanea, or Dar el Beida, a place not mentioned by any of the botanists who have visited the coast. Having made a tolerably large collection on the bare hills near Casa Blanea, and been hospitably received by Mr. Dupuis, the British vice-consul, we continued our voyage to

Mogador, not having any favourable opportunity for landing at Mazagan or Saffi. We reached Mogador on the 25th of April; and after attending to the requisite arrangements for our journey we visited the curious neighbourhood of that town. The vegetation of the neighbourhood, and especially that of the low rocky island which forms the port, had suffered severely from the periodical visitation of locusts which had lately passed over the district; but we were able to secure most of the interesting species found by our predecessors; and on the 29th we started for the interior, making a slight detour by keeping southward near the coast for some five or six miles, and thereby making fuller acquaintance with the Argan forest, and then across the country till we rejoined our tents and baggage outside the walls of the extensive castle of the Governor of Shedma. During most of the day we had traversed the northern part of Haha, a very large province that extends along the coast from Mogador to Agadir and to the skirts of the Atlas chain. It is needless to say that on this, as on each succeeding day of our ride to Marocco, we encountered many interesting plants, some altogether new to us, some known only from dried specimens, and several hitherto undescribed. On the evening of the 30th we reached Aïn Oumast, a spring where travellers necessarily encamp, as for many miles eastward the country is an arid stony desert, closely resembling in aspect, as well as in its vegetation, some parts of the Sahara.

On the 1st of May we encamped at Sheshaoua, a veritable oasis, where in the shade of fruit-trees we were surprised to find many common European species. The soil surrounding it appears to contain much gypsum, and produced an unusual proportion of *Chenopodiaceæ*.

A long ride on the following day took us to Misra ben Kara, a place near the banks of a considerable stream that flows northward from the Great Atlas; and on the following day a short but hot ride across the glowing plain took us to the ancient capital of Marocco.

While detained in Marocco Mr. Maw made an interesting excursion to the nearest of some rocky hills of metamorphic rock that rise a few miles N.W. of the city. He brought back a new species of *Boerhavia*, *Forsākhlia tenacissima*, *Andropogon laniger*, and several other species not hitherto seen by us.

On the 8th of May, after encountering various tokens of unfriendly disposition on the part of some of the native authorities,

we turned our faces towards the chain of the Great Atlas, which rises in full view of the city. Early on the following day we reached Tasseremont, and there began to make acquaintance with the special flora of the Great Atlas, gathering several of the species first seen by M. Balansa, as well as some others which he had not met. Being unable to continue our course towards the high range immediately south of Tasseremont, we turned S.W., and encamped on the same evening near a place called Ourika, at the opening of a valley to which we have given the same name. This valley appears to penetrate deeply the main chain of the Atlas, which rose patched with snow in the background. We naturally desired to ascend its course, and started on the 10th of May for that purpose. The scenery and the vegetation were full of interest; and it was with much annoyance that, at the pressing instance of our escort, who alleged an insurrection among the people of the upper valley, we were forced to return to a village near the opening. On the following day we skirted the base of the mountains through a district called Reraya, and encamped near to a stream descending from the main chain. It now became clear that there existed an organized plan on the part of our escort, in concert with the native chiefs, to prevent us from accomplishing our design of penetrating to the interior of the Great Atlas chain. The firmness of Dr. Hooker prevailed, however, over the reluctance of the escort; and the circumstance that we were able to secure the cooperation of a native Berber Sheik, finally enabled us to do a good deal more than the latter originally contemplated. On the following day we crossed a low range dividing the stream last mentioned from a parallel one which flows through the valley, which, from the name of the tribe inhabiting it, we called Aït Mesan*.

On the following day the friendly Sheik undertook to conduct us *to the snow*; and, as we had seen no snow save in patches on

* To the best of my belief the Aït Mesan valley, as well as the parallel and nameless one in which we camped on the 11th of May, is included within the district known as Reraya; and accordingly the labels for all Dr. Hooker's plants collected in both valleys bear the latter name. As there appeared to be no doubt that the valley wherein we passed five days is correctly named Aït Mesan, I have adopted that name in my labels, and have reserved the name Reraya for the valley first reached by us and the ridges which we traversed in approaching and departing from it. I have mentioned this, as without such explanation botanists might hereafter suppose that plants gathered at the same time and place by Dr. Hooker and myself came from two distinct localities.

the highest part of the chain, this was taken to mean the ascent of at least one of the higher summits. On the 13th we followed a steep and difficult track, amid a vegetation mainly new to us, —where it was tantalizing to be content with a specimen or two that could be snatched while pressing onward—and some time after midday reached the last village, called Arround, standing at nearly 7000 feet above the sea-level, a short distance beyond a barrier partly composed of rock and partly of ancient moraine, the only undoubted moraine seen by us in the Atlas. Here the scenery assumed a singularly stern character. The floor of the valley extends for some way nearly level, and then slopes gradually upward, enclosed on every side but one by steep and rugged peaks, nearly bare of snow. But in the hollows and recesses of the valley, at no great height above the village, snow lay in various directions; and towards one snow-bed our guide, in accordance with his promise, led us. Although many plants were not yet in flower, we found at every step something of engrossing interest—among the novelties that which most struck us being the curious *Chrysanthemum* with scarious involucre, which I have called *Chrysanthemum Catananche*. Our return to our camp in the lower valley in great part by night, and in almost pitch darkness, was difficult and even somewhat dangerous.

Our object now was to persuade the Sheik to let us sleep at the village of Arround, whence we felt that we could in one way or other attain the higher region, and thus accomplish in some measure the main object of our journey. After some negotiation this plan was arranged; and on the 15th of May, before the sun had well risen, we started from Arround with a very threatening sky, determined to ascend as far as might be possible. Fortunately the Sheik did not insist on accompanying us, but appointed two guides with injunctions not to let us go beyond a small stone hut at the base of the ascent to the pass of Tagherot. It should have been stated that on our first visit two days before, we ascertained that a somewhat frequented track leads from the head of Aït Mesan to the upper part of the Sous valley on the opposite side of the main chain. The weather had become cold and rainy when we reached the hut above referred to. Here our guides endeavoured to make us understand that it was too dangerous to go further; and we made no show of resistance, but encouraged them to make a fire in the hut while we collected plants round about. As soon as they were thus engaged we started up the track, which

throughout a steep ascent of over 3000 feet is well engineered, sometimes in regular zigzags, such as are seen in Switzerland or Tyrol. It is needless to say that every step in our ascent was of engrossing interest; but the weather grew steadily worse, as the rain became sleet, and then snow. We were overtaken by our guides, who used every means of threat and entreaty to induce us to return: for a time we silenced them by some silver coins; but they were in a pitiable state as we approached the summit. Except on some ledges of rock it was no longer possible to get at plants; and the snow driven by a violent wind made it impossible to see more than a few feet. Mr. Maw alone reached the actual crest of the ridge, but could see nothing. I, as well as Dr. Hooker, was a little lower down; and when Mr. Maw returned we readily consented to forego the barren honour of touching the summit, which is, by our observations, about 11,500 feet above the sea.

On the following day we returned to our camp, the valley above Arrond being deep in snow, and no more botanizing being there practicable. The following day was needed to arrange and pack our collections; and on the 18th we reached a place called Sektana, near the base of the main range, of which it commands a very fine view. On the next morning Mr. Maw left us, to return to England, taking with him two of our escort; while we proceeded westward by a place called Gurguri, and, after crossing the wide stony bed of a torrent called Oued en Fisk, reached Amsmiz, the largest place we had yet seen on the skirts of the Atlas. Here, after some trouble, we succeeded in making another excursion into the interior of the range; and from a village near the head of the valley, where we slept, we were able on the 23rd of May to ascend the highest summit near its head, called Djebel Tezah. The chain has here diminished considerably in height; for while the Tagherot pass is as nearly as possible 11,500 feet in height, the summit of Djebel Tezah only attains to about 11,000 feet. Thence we were able to see across the great valley of Sous, and for the first time beheld the parallel range, or Anti-Atlas, some 60 miles distant, which, at a rough estimate, can scarcely attain the height of 10,000 feet.

Returning to Amsmiz, our course lay through the sloping plain at the foot of the mountains to Mzouda; and on the 26th we passed Keira (where M. Balansa had found a friendly reception), and encamped near the village of Seksaoua. Two days, on one of which we ascended a neighbouring mountain to the height of some

4600 feet above the sea, were very productive in species not before seen, and one or two of them altogether new. We next went to Milhaïn, passing on the way Imintenout, where the main track from Morocco to Tarudant enters the mountains. Our next camp was outside the Kasbah or Castle of the Governor of Mzouda, then engaged in local war with the Governor of Haha. This affair caused us, in compliance with an urgent letter forwarded by courier from Mr. Carstensen, to alter our route and proceed northward, to a place called Mskala, on the border of the provinces of Haha and Shedma. Here the Governor of the latter province was encamped with a considerable force of his retainers, watching the progress of the petty war pending between the neighbouring tribes. On the next evening we met Mr. Carstensen by appointment at the Kasbah of the Governor of Shedma; and instead of returning direct to Mogador, we proceeded together on the 1st of June through a pleasant country to Aïn el Hadjar, a place where numerous springs break out from the foot of the Djebel Hadid, or Iron Mountain, a long, flat-topped ridge that rises near the coast some 15 miles north of Mogador. Although the heat was by no means oppressive, the season was rather far advanced for seeing the vegetation to full advantage; but we passed a not unprofitable day upon the mountain, the chief drawback on our enjoyment being the extraordinary number and variety of spiny and prickly bushes which cover its slope. One of these, which escaped the vigilant eyes of M. Balansa when he visited the mountain, is the widely spread tropical *Celastrus*, which, when found in the south of Spain, was taken to be a new species and called *C. europæus*, but which Professor Oliver has shown to be the *C. senegalensis* of Lamarck. We had already gathered this plant near the foot of the Great Atlas; but until then it was not known to grow anywhere between Senegal and the south of Spain.

On June 3rd we returned to Mogador, and were again hospitably entertained by Mr. and Mrs. Carstensen, until the arrival of a British steamer enabled us to take our passage homeward. We were able to spend a great part of one day at Saffi, where we embarked cargo, and, among other plants of interest, were enabled to add to the short list of Canary-Island plants that extend to the African coast the curious fleshy *Zygophyllum Fontanesii*, figured by Webb in the 'Phytographia Canariensis.'

We also landed at Mazagan, and had time enough to make a

short walk along the shore near that town, and add two or three species to our list of the coast-flora.

On reaching Tangier Dr. Hooker's engagements forced him to proceed at once to Gibraltar, and thence return to England. I was enabled to spend one day at Tangier, and after an absence of not quite two months found the aspect of the vegetation wonderfully altered, since the appearance of a large number of species not before seen gave the neighbourhood an aspect of complete novelty. The flowers of the *Cisti* had disappeared; but in their place were many *Compositæ* and *Labiatae* and other summer-flowering species. The most conspicuous was the magnificent *Salvia bicolor*, which attains a height of 7 or 8 feet, each of its numerous branches bearing the conspicuous white-and-blue flowers to which it owes its name.

During the period of our stay in the interior we had arranged, through the kind assistance of Mr. Carstensen, to send two natives to collect plants near Agadir, about 80 miles south of Mogador, near to Cape Guer, where the Great Atlas range finally subsides into the Atlantic Ocean. The result, which seemed to show that the character of the vegetation does not vary much along this part of the coast, yet sufficed to prove that with due perseverance something might be accomplished towards extending our knowledge of the Marocco flora through native collectors.

M. Cosson, having succeeded in securing the active assistance of the late M. Beaumier, French Consul at Mogador, and sparing on his own side neither trouble nor expense to effect his object, has caused two native collectors to travel into the interior for several successive seasons. One of these, a very intelligent Jew, a native of Akka, a place on the southern side of the Great Atlas between the rivers Sous and Noun, has been chiefly employed in the region, utterly inaccessible to European travellers, between the oasis of Akka and the neighbourhood of the Atlantic coast. The other, a Schleuh by birth, and a native of the mountains, has been mainly engaged in making collections in a district nearly adjoining the portion of the Great Atlas which we were able to explore. The collections first sent home were in indifferent condition, and the specimens incomplete; but M. Cosson's perseverance overcame all difficulties, and the collections lately received would do credit to a professed naturalist.

When I consulted my excellent friend M. Cosson on the subject of the present publication, he was good enough to suggest that we should jointly bring out a work which might bear to the

future Flora of Marocco the same relation that Desfontaine's 'Flora Atlantica' will hold to the, I trust, not distant Flora of Algeria.

I may at once say that I have no claim whatever to approach M. Cosson's extensive and accurate knowledge of the flora of North Africa and South-western Europe, and I feel that it would be a high honour to be associated with him in such a work as that suggested; but seeing that circumstances have placed me in possession of a considerable mass of materials that must go towards its production, I have thought it better without further delay to present it to this Society under the title "*Spicilegium Floræ Maroccanæ*," accompanied by lithographed plates illustrating some of the new species described, presented to the Society jointly by Dr. Hooker and myself.

I trust that M. Cosson will carry out his intention of speedily publishing in a compendious form the rich materials for the Flora of Marocco, most of which are in his exclusive possession, and thus preparing the way for a more complete work.

In the following descriptive catalogue I have, as a rule, avoided referring to plants not yet published by M. Cosson, excepting where this was necessary to name or illustrate specimens collected by us or other recent travellers, although, through his liberality, I am in possession of many species not enumerated in the following pages.

Although I feel that the time has not yet come for attempting a general survey of the Marocco flora, and the materials for discussing the subject must remain further incomplete until the publication of the long expected Flora of Algeria, it would be inexpedient to close these introductory remarks without a brief view of the geographical relations of the Marocco species to those of the surrounding regions. For this purpose I have enumerated in tabular form the Natural Orders indigenous to Marocco under headings that show the distribution of the species hereafter enumerated.

The following is a tabular view of the Marocco flora, showing:—

1. The whole number of species belonging to each Natural Order found in Marocco;
2. The number of these which are cosmopolitan or extend in the Old World beyond the Mediterranean region;
3. Species of the Mediterranean region including Macaronesia;
4. Mediterranean excluding Macaronesia;
5. Confined to Western Mediterranean region excluding Macaronesia;
6. Species

exclusively confined to Algeria and Morocco; 7. Species confined to the Iberian peninsula and Morocco; 8. Species known only in Macaronesia and Morocco; 9. Species peculiar to Morocco:—

Natural Order.	Total of Morocco species.	Cosmopolitan or Temperate Old World.	Mediterranean region including Macaronesia.	Mediterranean region excluding Macaronesia.	Western Mediterr. region excluding Macaronesia.	Algeria and Morocco.	Iberian peninsula and Morocco.	Macaronesia and Morocco.	Peculiar to Morocco.
Ranunculaceæ	33	11	5	9	4	1	1	..	2
Berberideæ	1	1					
Papaveraceæ	21	10	...	4	2	1	2	...	2
Cruciferae	73	25	12	16	2	5	3	...	10
Capparideæ	1	1					
Resedaceæ	13	3	...	2	1	1	2	...	4
Cistineæ	28	3	2	13	2	1	6	1	
Violariææ	3	1	1	1
Polygaleæ	5	3	2
Frankeniaceæ	2	2							
Caryophylleæ	69	29	9	13	6	2	5	1	4
Portulacaceæ	2	2							
Tamariscineæ	4	1	...	2	1
Hypericineæ	7	1	...	1	5				
Malvaceæ	12	3	1	5	2	...	1		
Lineæ	10	3	2	...	3	...	1	...	1
Zygophylleæ	3	...	1	1	1	
Geraniaceæ	21	9	4	4	3	1
Rutaceæ	4	1	...	2	1
Celastrineæ	1	1							
Rhamneæ	4	3	1				
Ampelideæ	1	1					
Acerineæ	1	1					
Anacardiaceæ	4	...	1	2	...	1			
Coriariææ	1	1				
Leguminosæ	189	27	46	37	24	11	16	3	25
Rosaceæ	16	7	1	5	1	...	1	...	1
Saxifrageæ	5	3	...	1	1
Crassulaceæ	18	7	...	1	5	...	1	...	4
Droseraceæ	1	1		
Haloragææ	2	2							
Myrtaceæ	1	...	1						
Lythrarieæ	6	2	2	2					
Onagrarieæ	3	3							
Cucurbitaceæ	4	1	1	2					
Ficoideæ	4	2	1	1					
Umbelliferae	86	17	11	18	20	5	4	...	11
Araliaceæ	1	1							
Caprifoliaceæ	7	3	2	1	1				
Rubiaceæ	30	8	3	9	4	3	1	...	2
Valerianeæ	8	2	1	5					
Dipsaceæ	6	1	1	...	3	1
Totals	711	191	107	166	91	31	45	6	74

Natural Order.	Total of Morocco species.	Cosmopolitan or Temperate Old World.	Mediterranean region including Macaronesia.	Mediterranean region excluding Macaronesia.	Western Mediterr. region excluding Macaronesia.	Algeria and Morocco.	Iberian peninsula and Morocco.	Macaronesia and Morocco.	Peculiar to Morocco.
Brought forward	711	191	107	166	91	31	45	6	74
Compositæ :									
Corymbiferae } 208 ..	83	10	19	17	11	9	1	1	15
Cynaraceae }	58	6	8	12	13	4	5	...	10
Cichoriaceae }	67	13	14	14	10	6	5	1	4
Lobeliaceae	1	1
Campanulaceae	13	1	3	...	2	...	3	...	4
Ericaceae	9	4	1	1	2	...	1
Sapotaceae	1	1
Plumbagineae	18	3	...	6	4	...	2	...	3
Primulaceae	7	3	...	1	2	...	1
Oleaceae	7	1	1	3	1	1
Apocynaceae	2	1	1
Asclepiadeae	3	1	1	1
Gentianeae	8	5	3
Boragineae	35	9	5	8	7	1	1	1	3
Convolvulaceae	10	4	3	1	2
Solanaceae	11	4	4	3
Scrophularineae	61	16	5	9	9	4	7	1	10
Orobanchae	12	1	1	2	3	...	3	...	2
Lentibulariæ	1	1
Acanthaceae	1	...	1
Verbenaceae	3	1	1	1
Selagineae	1	1
Labiata	81	21	6	10	14	2	5	...	23
Plantagineae	11	2	5	3	...	1
Nyctagineae	2	1	1
Illecebreæ	10	4	2	2	2
Amaranthaceae	6	4	...	1	...	1
Chenopodiaceae	20	13	4	1	2	...
Polygonæ	17	9	3	2	2	1
Thymeleæ	7	2	1	1	1	...	2
Laurineæ	1	1
Santalaceæ	3	1	2
Cytinæ	1	...	1
Aristolochiæ	3	...	1	...	2
Euphorbiaceæ	24	7	3	4	3	7
Urticaceæ	7	4	1	1	1
Ulmaceæ	2	2
Cupuliferæ	4	3	1
Salicinæ	6	4	...	1	1	...
Ceratophylleæ	1	1
Gnetaceæ	2	...	1	1
Coniferae	7	1	2	2	1	1
Orchideæ	18	7	2	9
Irideæ	14	3	2	4	2	...	1	1	1
Amaryllideæ	11	4	2	...	4	...	1
Totals	1381	359	212	297	190	61	87	14	161

Natural Order.	Total of Morocco species.	Cosmopolitan or Temperate Old World.	Mediterranean region including Macaronesia.	Mediterranean region excluding Macaronesia.	Western Mediterr. region excluding Macaronesia.	Algeria and Morocco.	Iberian peninsula and Morocco.	Macaronesia and Morocco.	Peculiar to Morocco.
Brought forward ...	1381	359	212	297	190	61	87	14	161
Dioscoreæ	1	1							
Alismaceæ	3	3							
Juncagineæ	1	1					
Potameæ	3	3							
Palmeæ	2	1	...	1					
Typhaceæ	1	1							
Aroideæ	3	1	1	1					
Lemnaceæ	1	1							
Liliaceæ	43	7	11	12	7	2	4		
Melanthaceæ	4	2	...	1	1		
Smilacere	3	1	1	1					
Asparageæ	7	...	3	1	1	1	1
Juncææ	15	12	...	1	1	1
Cyperaceæ	25	18	1	2	2	...	1	...	1
Gramineæ	134	57	25	32	15	1	3	...	1
	1627	467	254	350	216	64	96	15	165
Dicotyledones	1338	349	208	280	186	61	82	13	159
Monocotyledones	289	118	46	70	30	3	14	2	6

It is obvious to all who have considered the subject, that a mere enumeration of the species known to exist in a given region gives of itself but little information as to the character of its flora ; nevertheless the statistical method, to which somewhat too much importance has, in my opinion, been attributed, does lead towards some probable conclusions, and gives indications which may not be unimportant.

Taken as a whole, the figures in the preceding Table show very distinctly that the Morocco flora is altogether a portion of that great Mediterranean flora which, with local peculiarities, extends from the Indus to the Atlantic Islands. Excluding all the species widely spread beyond those limits either through the tropics or the temperate zone, we have remaining 1160 species, of which 820 are common to some considerable portions of the Mediterranean region. Of the remaining 340 species about one half, or 165, are peculiar to Morocco. Next come 96 common to Spain and Morocco, nearly all of these being, as far as we know, confined to the neighbourhood of Tangier and Tetuan. A much smaller proportion, only 64, are common to Algeria and Morocco,

and only 15 species are confined to the Macaronesian Islands, under which name I include Madeira along with the Canary Islands. This proportion will doubtless be somewhat increased when we know more of the coast vegetation between Agadir and Wadnoun; but the figures tend to show that the separation of those islands must date from a period even geologically speaking remote.

With a view to exhibit further the characteristics of the Marocco flora it may be convenient to give a summary of the results of the foregoing Table, showing the absolute number of species or subspecies of each of the principal Natural Orders, and the percentage which they bear to the total number of Phanerogamic species, this being 1627.

	Total number of species.	Percentage proportion.
Dicotyledones	1338	82·2
Monocotyledones	289	17·8
Compositæ	208	12·8
Leguminosæ	189	11·6
Gramineæ	134	8·2
Umbelliferæ	86	5·3
Labiatae	81	5·0
Cruciferae	73	4·5
Caryophyllæ	69	4·2
Scrophularinæ	61	3·7
Liliaceæ	43	2·6
Boraginæ	35	2·2
Ranunculaceæ	33	2·0
Rubiaceæ	30	1·8
Cistineæ	28	1·7
Cyperaceæ	25	1·5
Euphorbiaceæ	24	1·5

It will be seen that the proportion of Compositæ, Leguminosæ, and Liliaceæ is unusually large, whilst that of Gramineæ and Ranunculaceæ is exceptionally small. Still more characteristic of the Marocco flora is the small proportion borne by some natural orders that usually take a prominent place among the vegetable population in mountain countries in the north temperate zone. Thus we have of Rosaceæ but 16 species, of Saxifrageæ 5 species, of Primulaceæ 7 species, of Gentianeæ 8 species, and of

Cyperaceæ only 25 species; and very few of these belong to sub-alpine or alpine types.

In my opinion, however, the general numerical results obtained from tabular statements, such as those I have already given, are very likely to mislead unless critically controlled. The boundaries of states are determined by historical accidents, and, to a slight extent only, by physical geography; and it often happens that in general summaries the characteristics of what form natural divisions of the earth's surface are masked by these being thrown in along with others wherein different conditions predominate.

Bearing in mind that the entire of Marocco in a botanical sense forms part of the Mediterranean region, it includes within its boundaries several provinces, partly corresponding to geographical limits, which exhibit well-marked features, and are distinguished by a considerable number of species not common to other provinces.

Speaking broadly, and excluding the wide-spread species, which belong to two types—1st, those widely spread through the cooler temperate zone, either Old World or extending to the New World, and, 2nd, tropical species which extend more or less widely into the warmer temperate regions—there are within the Marocco territory five elements, of which four correspond more or less closely to geographical limits.

The Mediterranean flora, including within that term species confined to that region and widely spread throughout it, forms what may be considered as the substratum extending throughout the whole territory, excepting only the higher range of the Great Atlas. Overlying, so to say, this substratum, we find in the northern extremity of Marocco a strong infusion of another element, which is now generally denominated the Peninsular flora. This is recognizable, even to the unscientific observer, by the number of conspicuous flowering shrubs, including, along with more wide-spread species of those genera, several showy species of *Cistus* and *Erica*, with numerous herbaceous plants, such as the curious *Drosophyllum lusitanicum*. We do not know the limits of this province along the coast between Tetuan and the Algerian frontier; but it does not seem to extend more than 30 or 40 miles southward from Tangier and Cape Spartel, and in point of fact nearly all the 96 species set down as common exclusively to Spain and Marocco have been found in the neigh-

bourhood of Tangier and Tetuan, and are not seen in the central or southern provinces of Marocco.

The Algerian flora, so far as it offers special characteristics to distinguish it from the general Mediterranean type, owes its individuality mainly to species endemic in the mountain region of the Lesser Atlas, or on the high plateaux that present such a peculiar feature in the geography of Southern Algeria.

It is highly probable that most of these extend into Eastern and Central Marocco; but in the limited region known to us these characteristic species of the Algerian flora are for the most part wanting, and even in the Great Atlas but a small number of them have yet been seen.

Although several of the species characteristic of the Desert flora extend beyond their original boundary, and a few of them are found even in the south-east of Spain, there are few better-marked botanical provinces than that of the desert regions of Western Asia and Northern Africa. Considering the wide portion of the earth's surface occupied by the hot stony or sandy plains that extend with unimportant interruptions nearly from the banks of the Lower Indus to the Atlantic coast of Southern Marocco, the general uniformity of its scanty vegetable population is a fact very remarkable in botanical geography. Unfortunately our knowledge of the true desert-region of Southern Marocco, namely that which extends along the Southern side of the Great Atlas chain, is extremely limited; but when the collections recently received by M. Cosson from his collector are fully enumerated, it will be seen that many species of this peculiar type extend to within a few leagues of the Atlantic coast. But even in the low country, on the northern side of the Great Atlas, traversed by us there are some considerable tracts closely approximating in their physical conditions to the northern skirts of the Sahara. Many of the characteristic desert-species there reappear, although they are separated by lofty mountain-ranges from what may be considered their natural home.

The least-important element that goes to make up the Marocco flora, if we measure it by the number of representative species, but in some respects one of the most interesting, is that revealed by the presence of a small number of species common to the Maroccan and the Macaronesian* floras. Of these all but one are

* I have ventured to use the name Macaronesia in a rather wider sense than some preceding writers, including under it Madeira with its dependent islets,

Canary-Island species, the remaining one being common only to Madeira and Western Morocco. It is true that the short list of fifteen species might be somewhat extended if we added certain species peculiar to Morocco but closely allied to Canary-Island endemic species. Such a list would include three cactoid Euphorbias, a *Sonchus* (*S. acidus*, nearly allied to *S. pinnatus*), a Senecio of the *Kleinia* group, and *Monanthes atlantica*, nearly allied to its Canary-Island congeners.

It must be remarked, however, that the types to which these last-mentioned species belong are rather generally West-African than specially Macaronesian, as all are common to the Cape de Verde Islands, if not to a wider region. Of the fifteen Macaronesian species found in the coast region of South Morocco, I think it is safe to say that the facts rather tend to show the accidental diffusion of a few Macaronesian species on the adjacent coast of Africa than to indicate the existence of a direct connexion between the continent and those islands within a geological period at all recent.

There remains to be considered the flora of the Great Atlas, the only one of the constituent portions of the general Morocco flora that seems to be confined within the boundaries of the empire. I am tempted to enter into some detail in discussing this part of my subject; but when I recollect what a large mass of additional unpublished matter is already in the possession of my friend M. Cosson, I feel that it would be unsatisfactory to attempt any such detailed discussion at the present time; and I hope to be able to resume the subject to greater advantage on a future occasion.

In the mean time it is allowable to point out some characteristic features of the Great-Atlas flora, as far as this is known to me from our collections. Dividing the mountain region into two zones, an upper and a lower one, and fixing the limit between them at about 1500 metres above the sea-level, I find in each of these a considerable number of endemic species, amounting in regard to the upper zone to about one fourth of the whole number of species. But there is little indication of that multiplication of specific forms that is so characteristic of the mountain-floras of

along with the Canary-Island archipelago. Though the flora of Madeira be more limited than that of the Canaries, it is impossible to regard it as other than a detached member of that group.

Southern Spain, Greece, and Asia Minor. On the arenaceous and porphyritic rocks of the Aït Mesan valley we found to a great extent the same endemic species afterwards gathered on the schistose and calcareous rocks of Djebel Tezah, at least 30 miles distant.

Another fact worth noticing is, that, among a considerable number of new specific forms, no new generic type has yet been found in the Great Atlas.

But in my opinion the most singular point about the Great-Atlas flora is the presence of a large number of the common species of Central Europe, both annual and perennial, coupled with the fact that many of these are not species that have proved their power of adapting themselves to different climatal conditions, and that many of them have no nearer habitat than the mountains of Central or Southern Spain, while others must be sought at a still greater distance.

The conclusion to which I am led by a consideration of the facts is the same that I have derived from rather long observation of the mountain-vegetation of Central and Southern Europe—namely, that it is impossible to explain the facts without admitting that a large portion of our present flora is relatively very ancient, and that the species had assumed their present characters at least as early as the Miocene epoch.

I ought not to omit mentioning that two years after our visit two German naturalists, MM. Rein and Fritsch, were able to make a short expedition in the Great Atlas, nearly following our footsteps. They were not exclusively engaged in collecting plants; but they have deposited at Kew specimens of most, if not all, the plants found by them. Having been able to examine these, I have enumerated their specimens in the following catalogue.

In determining the specimens of our collections I have been materially assisted, in regard to the Orchideæ, by my friend Prof. H. G. Reichenbach. The friendly aid and assistance of Professor Oliver and Mr. J. G. Baker have been frequently invoked by me; and to the latter I am especially indebted for his thorough knowledge of the Liliaceæ and Amaryllideæ of the Mediterranean region, which he kindly contributed for my benefit. To Mr. Bentham, it is needless to say that in any case of doubt or difficulty I never applied in vain. The resources of his unequalled knowledge are always at the service of the younger and less-gifted workers in the field of science; and I cannot better close than by expressing the sense of affectionate regard with which I acknowledge his guidance.

ENUMERATIO GENERUM AC SPECIERUM.

PRÆMONENDA.

Ordines naturales plantarum secundum normam Candolleanam a celeb. Bentham et Hooker reformatam enumeravi; generum definitiones a Bentham et Hooker in opere classico 'Genera Plantarum' stabilitas quatenus profuerunt fere semper secutus sum.

Nomina trivialia specierum antiquiora, nisi plane erronea, semper prætuli. Auctorem speciei, qui primus eam ab affinis distinxit et publici juris fecit, esse duxi, et prius citavi. Auctores cæteros parca citavi, opus tamen cyclopædicum Candolleanum 'Prodromus Systematis Universalis,' necnon Kunthianum 'Enumeratio Plantarum,' semper adduxi.

Formas istas plantarum a speciebus affinis notis evidentibus et satis stabilibus dignoscendas cujus tamen differentiae tales sint ut una pro alterius prole habenda esse videatur, sicut subspecies enumeravi, et typis diversis eorum nomina a cæteris distinxī.

Territorium Imperii Maroccani in regiones quinque limitibus certis nondum definitas distinxī: scilicet 1. Marocco septentrionalis—ditionem Tingitanam et Tetuanensem necnon regionem litoralem inter hanc et fines Algerienses amplectens; 2. Marocco occidentalis—vel regio litoralis et sublitoralis Atlantica, ab urbi El Araïsch usque Agadir; 3. Marocco centralis, botanicis fere ignotus, inter zonam sublitoralem et convallem fluminis Moulouya; 4. Marocco orientalis, regio vix nota inter flumen Moulouya et fines Algerienses; 5. Marocco meridionalis, quo comprehenditur catena centralis Atlantis Majoris et regio inferior a montium

radicibus usque flumen Oum erbia. His addenda erit regio vasta desertorum, Sahara Maroccana dicta, ab Atlante Majore meridiem versus se protendens, quæ tamen species nullas operi nostro subministravit.

Locorum natalium plantarum quibus adest signum (!) testis ipse sum. Idem signum, dum sequitur nomina abbreviata, plantas indicat quarum specimina vidi et scrutavi. Horum sequitur brevis elenchus :

Webb. Herb.—Philippus Barker Webb. Stirpes circa Tingidem et Tetuan anno 1827 lectæ, in Herbario Florentino asservatæ.

Salzm. Exsicc.—Salzmann. Plantæ in ditione Tingitana anno 1825 lectæ quarum series venales duæ in Herbario Kewensi asservantur.

Lowe.—R. T. Lowe. Plantæ in agro Mogadorensi anno 1859 lectæ, quarum catalogum in Actis Soc. Linnæanæ Londinensis edidit. Harum exemplaria fere omnium in Herbario Kewensi exstant.

Schsb. spec.—Plantæ olim a Schousboe lectæ, ex ejus herbario duplicatæ, a cel. Cosson benevole communicatæ.

Balansa spec.—Plantæ a cl. Balansa anno 1867 lectæ, quarum perpaucæ venales in Herbario Kewensi vel in cæteris herbariis inveniuntur.

J. D. H.—Plantæ quas in itinere Maroccano amicissimus Eques Josephus Dalton Hooker solus legit.

Coll. Indig.—Plantæ mense Majo 1871 a duobus indigenis in ditione Agadir lectæ, quarum exemplaria in herbario meo ac in Kewensi extant.

R. et Fr.—Plantæ a ell. Rein et Fritsch anno 1873 lectæ, quarum specimina in Herbario Kewensi conservantur.

RANUNCULACEÆ (*Juss. Gen.* 231).CLEMATIS (*L. Gen.* 696).

C. FLAMMULA (*L., DC. Pr.* i. 2).

Mar. sept.—Tanger. (*Salzm. sp. in herb. nost.*)! Tetuan (1851)!

Mar. merid.—Ourika! Ansmiz!

Ar. Geog. Regio Mediterranea exelusa Macaronesia.

C. CIRRHOSA (*L., DC. Pr.* i. 9).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Blackmore*)!

Mar. occid.—Mogador (*Lowe*)!

Mar. merid.—Visa pr. Ourika!

Ar. Geog. Regio Mediterranea calidior (exelusa Macaronesia).

C. Balearica (*Rich, Journ. Phys.* 1779, 127; *DC. Prod.* i. 9), = *C. polymorpha* (*Viv. Fl. Cors.* p. 9), inclusa forma foliis minus profunde divisis *C. semitriloba* (*Lag. Cat. Matr.* 17).

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

Ar. Geog. Insul. Mediterraneæ (Balears, Corsica, Sicilia). Marocco occidentalis.

A plerisque botanicis cum præcedente conjuncta, sed formæ intermediæ vix inveniendæ. Confer Boissier in *Fl. Or.* i. 2. Mihi videtur melius ad subspeciem reducenda. Specimina ex monte supra Agadir a eel. Cosson benevole communicata ad var. seu lusum *C. semitrilobam* pertinent.

THALICTRUM (*L. Gen.* 697).

T. GLAUCUM (*Desf. Cat. H. Par.* ed. 2, p. 126; *DC. Pr.* i. 15).

Mar. merid.—Mesfioua!

Ar. Geog. Penins. Iberica. Africa boreali-occidentalis. In Italia erronee indicatum.

ANEMONE (*L. Gen.* 694.).

A. PALMATA (*L., DC. Pr.* i. 19).

Mar. sept.—Tanger, in monte Djebel Kebir! Tetuan, in monte Beni Hosmar, usque 400 m.!

Ar. Geog. Penins. Iberica. Africa boreali-occidentalis.

ADONIS (*L. Gen.* 698).

A. AUTUMNALIS (*L., DC. Pr.* i. 23).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Mar. merid.—Prov. Haha!

Ar. Geog. Europa media. Regio Mediterranea omnis (exelusa Macaronesia).

Specimina nostra carpellorum juniorum rostro uncinato referunt *A.*

baticam (Coss. Not. Crit. Plant. d'Esp. ii. 25), quæ mihi videtur
var. *A. autumnalis*.

ADONIS ÆSTIVALIS (*L., DC. Pr. i. 24*).

Mar. merid.—Shedma! Amsmiz! Sektana!

Ar. Geog. Europa media (vix indigena?). Regio Mediterranea (Europa, Oriens, Africa borealis). Mont. Himalaya.

Variat in reg. nostra petalis sanguineis vel citrinis.

A. MICROCARPA (*DC. Syst. i. 223, DC. Pr. i. 24*), = *A. Cupamana*,
Guss. Syn. Fl. Sic. ii. 36), = *A. intermedia* (*Webb, Phyt. Can. i. 12*).

Mar. merid.—Sektana!

Ar. Geog. Penins. Iberica; Sicilia. Afr. borealis. Macaronesia.

Petala in. spec. nostris citrina.

— —, var. *A. DENTATA* = *A. dentata* (*Del. Fl. Eg. Ill. 17; DC. Pr. i. 24*).

Mar. merid.—Sheshaoua!

Ar. Geog. Syria; Arabia; Persia. Africa borealis in reg. Saharensi.

In hoc genere species plurimæ extricatu diffieillimæ et forsitan non satis constantes.

RANUNCULUS (*L. Gen. 699*).

R. AQUATILIS (*L., DC. Pr. i. 26*).

Mar. sept.—Tanger! Inter Tetuan et Ceuta!

Mar. merid.—Amsmiz!

Ar. Geog. In reg. temperata et frigida hemisph. borealis (Europa, Asia, Africa, Macaronesia, et America). Rarius occurrit in Asia subtropica.

R. Baudotii (*Godr. Monog. p. 14, fig. 4*), = *Batrachium marinum* (*Fr. Nov. Fl. Suec. Mant. iii. 51*).

Inter specimen nostra Tingitana video exemplaria quæ verisimiliter huc spectant. Habuerim pro varietate speciei polymorphæ (*R. aquatilis*) sed auctoritate tantorum virorum devinetus hanc in numerum subspecierum enumerare malui.

Ar. Geog. Gallia; Insul. Britannicæ. Batavia. Ditio Tingitana.

R. FICARIA (*L., B. et H. Gen. Pl. i. 6.*), = *Ficaria ranunculoides* (*Mench. Meth. 215; DC. Pr. i. 44*).

Mar. sept.—Inter Tanger et Tetuan! In monte Beni Hosmar prope Tetuan!

Ar. Geog. Europa borealis et media et Mediterranea. Marocco septentrionalis.

— —, var. *INTERMEDIUS*, *nob.*

Mar. sept.—In monte Djebel Kebir prope Tanger!

R. calthæfolius (*Jordan, Obs. vi. 2*), = *Ficaria calthæfolia* (*Reich. Fl. Exc. ii. 718*), a planta Linnæana statuta et floribus duplo majoribus facile dignoscendus, foliis profunde cordatis, lobis incumbens

adeo ut primo intuitu peltata videntur, nobis non obvia fuit in itinere Maroccano. Varietas supra memorata gaudet floribus aliquid majoribus, foliis paullo profundius cordatis, sed typo Linnæano multo propior est quam *R. calthæfolio*, Jord.

Quoad sciam *R. Ficaria*, L., hucusque extra territorium Europæ non citatur.

RANUNCULUS BULLATUS (L., DC. Pr. i. 27).

Mar. sept.—Tanger! folia tantum visa.

Mar. occid.—Mogador (*Brouss. sec. Boissier*).

Ar. Geog. Regio Mediterranea australior e Creta usque in Lusitaniam.

R. CHÆROPHYLLUS (L., DC. Pr. i. 27), var. γ *FLABELLATUS*, DC. l. cit. = *R. flabellatus* (*Desf. Fl. Atl. i. 438, tab. 114*).

Mar. sept.—Valde frequens in regione collina circa Tanger! et Tetuan! et in monte Beni Hosmar, usque 800 m.! Variat caule simplici vel ramoso (usque 12-floro), foliis radicalibus integris ternatis vel biternatis.

Ar. Geog. Species per regionem Mediterraneam (exclusa Macaronesia) et montes Himalaya diffusa.

R. leucothrix (*Ball, Journ. Bot. 1873, 296*).

Inter priorem et *R. oxyspermum* (*Steph. in Willd. Sp. Pl. ii. 1328, DC.*

Pr. i. 28) collocandus; *R. chærophyлло* propior et sicut subspecies enumerandus. Differt imprimis grumis cylindricis, nec ovatis, indumento ex pilis longis villosis inferne patulis superne adpressis, nec brevi adpresso sericeo, pedicellis fructiferis rigidioribus subincrassatis, foliorum tripartitorum segmento medio longiore sæpe tripartito calyce fructifero persistente reflexo.

Mar. merid.—In regione inferiore Atlantis (*Distr. Rerava*) circ. 1000 m.! Petala subemarcida quam in typo minora; carpella immatura fere omnia abortiva.

A *R. chærophyлло* facie valde diversus, et forsán pro specie omnino distincta habuerim, si specimina meliora adfuissent. Calyx sub anthesi reflexus est in hoc genere nota specifica gravis momenti.

R. SPICATUS (*Desf. Fl. Atl. i. 438, tab. 115; DC. Pr. i. 29*).

Mar. sept.—In monte Beni Hosmar prope Tetuan circ. 900 m.!

Ar. Geog. Africa borealis. Lusitania?

In speciminibus nostris spica fructifera quam in spec. Algeriensibus paullo brevior.

— —, var. *R. BLEPHARICARPOS*, = *R. blepharicarpos* (*Boiss. El. 5*).

Mar. merid.—In regione subalpina Atlantis Majoris, Djebel Tezah circ. 2600 m.!

Ar. Geog. Peninsula Iberica. Marocco meridionalis. Speciminibus Nevadensibus a nobis lectis omnino conformis.

R. GRAMINEUS (L., DC. Pr. i. 32).

Mar. sept.—In monte Beni Hosmar prope Tetuan circ. 1000 m.!

Ar. Geog. Europa occidentalis et australis, nullibi frequens. Marocco septentrionalis. Extra Europam non prius detectus?

RANUNCULUS ATLANTICUS (*Ball, Journ. Bot.* 1873, p. 296).

Subspecies *R. acris* (*L., DC. Pr.* i. 36) sed formæ intermediæ nondum observatæ?

Inter subspecies et varietates *R. acris* distinguitur imprimis statura maxima 3-5-pedali, corolla (pro grege) maxima, rhizomate crasso horizontali fibras validas emittente, foliis magnis pentagonis profunde 3- vel 5fidis, segmentis ineisis, indumento sericeo-piloso adpresso, præsertim in foliorum pagina inferiore (folia igitur bicoloria). Carpella parva numerosa rostrata, rostro subreeto longiuseulo.

Mar. merid.—Frequens in convallibus septentrionalibus Atlantis Majoris —Urika! Reraya! Ait Mesan! Amsmiz! a 1000 m. ad 1900 m.

Huic proximus est *R. Friesianus*, *Jord. Obs.* vi. p. 17, = *R. nemorivagus*, *Jord. Diagn.* i. p. 74.

In regione Mediterranea *R. acris* est fere semper planta montana, et vix in zonam olivetorum se protendit. *R. atlanticus* noster non fugit loca calidiora.

R. PALUSTRIS (*L., DC. Pr.* i. 41), var. *MACROPHYLLUS*, = *R. macrophyllus* (*Desf. Fl. Atl.* i. 437), = *R. villosus* (*Salzm. Exsicc. non DC.*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Prope Cap. Spartel! Tetuan 1851! In monte Beni Hosmar!

Ar. Geog. Africa borealis. Formæ seu varietates arcte affines proveniunt in Lusitania (*Brot. Fl. Lus. sub R. adscendente*), in Hispania (*Boiss. Voy.*), in Sardinia (*Moris. Fl. Sard.*), in Corsica (*DC. fl. fr. sub R. corsico*) et in Oriente (*Boiss. Fl. Or.*).

Planta Fontanesiana differt ab orientali foliis minus profunde divisis, pedunculis fructiferis crassioribus. Ab his paululum differt planta Europæ australis foliorum lobo medio longiore, rostro apice subrecurvato.

R. procerus (*Moris, Fl. Sard.* i. 45; *Icon.* tab. 2)?

Mar. sept.—Tanger (*Webb Herb.*)!

Ar. Geog. Sardinia. Ager Tingitanus.

A *R. palustri*, cui simillimus, differt carpellis etiam junioribus tuberculoso-setigeris. De differentia specifica dubitavit ipse beatus Moris; nobis videtur ad subspeciem reducendus.

R. BULBOSUS (*L., DC. Prod.* i. 41), var. *NEAPOLITANUS*, = *R. Neapolitanus* (*Ten. Syll.* p. 272).

Mar. merid.—In regione media Atlantis Majoris; in convalle Amsmiz circ. 1500 m. ! in monte Djebel Tezah usque 2000 m. !

Ar. Geog. Italia; Sicilia?; Corsica; Hispania (*Bourg. Pl. d'Esp.* no. 2249). Oriens (*Boiss. Fl. Or.*). Maroeo meridionalis.

A typo facile dignoseitur rhizomate parum inerassato fibras plurimas crassas edente, sed pæce eel. Boissier carpella habet *R. bulbosi*, rostris brevissimis plus minusve arcuatis.

Confer notulam MS. beati J. Gay, et frustulum ex exemplari authenticæ Tenorcano herb. Desfontaines, in Herb. Kew conservata. Meo sensu *R. neapolitanus* est omnino forma meridionalis *R. bulbosi* et verisimiliter per totam regionem Mediterraneam diffusus.

— —, var.? GIGANTEUS, nob. 5-pedalis et ultra; rhizoma inerassatum; foliorum radicalium petioli 2-pedales; folia pinnatipartita, segmenta triseeta, varie incisa; flores *R. bulbosi*. Carpella desunt.

Mar. merid.—In regione inferiore Atlantis seeus torrentem Ourika!

R. rhæadifolius (DC. Syst. i. 284) ex descript. celeb. auctoris nostræ plantæ haud absimilis, sed ex icone in Delessert Icon. i. 40, hic mihi videtur forma *R. philonotidis*, Retz, seu *R. trilobi*, Desf.; sed ob defectum carpellorum res incerta manet. Planta nostra est fere certe *R. bulbosi* forma insignis.

RANUNCULUS PHILONOTIS (Retz, Obs. vi. 3; DC. Pr. i. 41), var.

INTERMEDIUS, nob.=*R. intermedius* (Poir. Dict. vi. 116)?

Mar. sept.—Tanger!

Ar. Geog. Hinc inde in regione Mediterranea?

R. trilobus (Desf. Fl. Atl. i. 437, tab. 113; DC. Pr. i. 42).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb. herb.*)! Tetuan seeus ripas fluminis (1851)!

Mar. merid.—Sektana!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

Pæce auctorum celeb. qui *R. trilobum* pro specie probe distincta enumerarunt hic mihi mera forma *R. philonotis* videtur, et nisi obstitisset auctoritas majorum pro varietate potius quam pro subspecie habuerim. Notæ diagnosticæ ex carpellis desumptæ non gravis momenti videntur. In *R. philonoti* vulgari carpella ostendunt tubereula circa marginem concentricæ disposita, sed hæc non raro omnino desunt. In *R. trilobo* tubereula crebriora sunt sed secundum eandem normam distribuuntur. Varietas nostra supra memorata, quæ verisimiliter est *R. intermedius*, Poir., habitu et corolla est *R. philonotis*, sed glabritie et carpellis propius accedit *R. trilobo*.

R. ARVENSIS (L., DC. Pr. i. 41).

Mar. merid.—In regione inferiore et media Atlantis Majoris; in convalle Aït Mesan circ. 1200 m.! et usque 2200 m!

Ar. Geog. Europa media, Asia borealis (*Gmel.*). Regio Mediterranea omnis. Montes Himalaya.

R. MURICATUS (L., DC. Pr. i. 42).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb. herb.*)! Prope Cap. Spartel!

Mar. merid.—In regione inferiore et media Atlantis Majoris. Reraya eire. 1100 m. ! Amsmiz ! Djebel Tezah usque 1750 m. !

Ar. Geog. Regio Mediterranea omnis. India boreali-occidentalis. America borealis et Meridionalis (vix indigenus?).

RANUNCULUS PARVIFLORUS (L., DC. Pr. i. 42).

Mar. sept.—Tanger (*Salzm. Exsicc.*) ! Vidimus in sepibus haud procul a Tanger ! et in dumosis montis Djebel Kebir !

Mar. merid.—In reg. inferiora Atlantis Majoris. distr. Reraya ! Ex prov. Mtouga specimina nana attulerunt R. et Fr.

Ar. Geog. Europa occidentalis ex Hibernia ad fretum Hereulis. Europa australis, ager Byzantinus. Africa boreali-occidentalis. Macaronesia. America borealis.

R. OPHIOGLOSSIFOLIUS (*Vill. Dauph.* iv. 732, tab. 49 ; DC. Pr. i. 43).

N.B. In exemplari nostro operis Villarsii anno 1789 impressi inventum III. 731.

Mar. sept.—In uliginosis pr. Tanger ! (*Salzm. Exsicc.*)

Ar. Geog. Europa occidentalis ex Sarnia ad Lusitaniam. Regio omnis Mediterranea. Formæ affines in America boreali (*R. pusillus*, Poir.), et in America meridionali (*R. bonariensis*, Poir.).

In regione subalpina Atlantis Majoris (in convalle Aït Mesan eire. 2550 m. loco frigido uliginoso) invenimus *Ranunculum* nondum florentem, rhizomate brevi recto, fibris validis elongatis (6–10-polliaribus) præditum. Folia radicalia sunt glabra, pinnatim triseeta, lobis cuneato-ovatis, in pagina superiore albo variegata. Eandem plantam florentem ex reg. superiore Atlantis anno 1873 attulerunt ell. Rein et Fritsch. In hae seapus solitarius superne adpresso-pilosus, folium unicum subsimplex gerens, petala lutea mediæ magnit., carpella (nimis immatura) rostrata, rostro spiraliter recurvo. Est verisimiliter nova species e grege *R. montani*, W., et proxima *R. procumbenti* (Boiss., Ann. Se. Nat., et Fl. Or. i. 51).

NIGELLA (*L. Gen.* 685).

N. ARVENSIS (L., DC. Pr. i. 49).

Mar. merid.—Aïn Tarsil, prov. Mtouga !

Ar. Geog. Europa media. Regio Mediterranea (exclusa Macaronesia).

— —, var. COSSONIANA, nob. = N. hispanica, var. parviflora (*Coss. Pl. Crit.* 49).

Mar. oecid.—Prope Mogador ! et Agadir (*Coll. Indig.* 1871) !

Mar. merid.—Prov. Shedma pluribus locis ! Prov. Mtouga (*R. et Fr. spec.*) !

Ar. Geog. Gallia australis. Hispania. Africa borealis.

Cel. Boiss. (in Fl. Or. i. 66) sub *N. arvensi* plurimas sic dict. species adunavit et sicut varietates descripsit ; seil. *N. tuberculata*, Gris. Spic., *N. aspera*, C. Koch., *N. fœniculacea*, DC., *N. divaricata*, Beaupr.

in DC. Syst., *N. cretensis*, Stev., *N. aristata*, Sibth. et Sm., sed revera vix pro varietatibus habendæ sunt. *N. arvensis*, L., est enim species summiopere ludibunda, per totam regionem Mediterraneam (exclusa Macaronesia) late diffusa, et rarius invenies specimen typo unius alteriusve formæ descriptæ omnino conveniens.

Cel. Cosson loc. cit. plantam nostram a *N. arvensi* distinxit et pro varietate *N. Hispanicæ* suæ vindicavit. Sed vir oculatissimus limites frustra quæsivit, dum natura invicta obstat. Forma et color sepalorum, ratio inter se labiorum petali, carpella extus plus minusve glanduloso-rugosa, horum rostra erecta, vel patentia, vel recurva, sunt omnes notæ variabiles, sicut jam me docuerunt specimina plurima Marocæana in vivo observata. Nee valet nota diagnostica a ell. Gr. et God. (Fl. Fr. i. 45, 46) proposita. In *N. arvensi* nervi laterales carpelli basin usque rarius producti; dum in *N. Hispanica* rostrum semper trinerve est, nervi in dorso carpelli haud raro producti.

N. Hispanica, Cosson, *l.c.*, in Hispania Gaditana haud infrequens (Bourgeau, Pl. d'Esp. 1849, No. 13, et a me olim prope Vejer et pr. Zahara lecta), statura maxima, sepalis illis *N. arvensis* 2-3ve majoribus, basi vix attenuatis, pro subspecie habenda videtur. Hæc in ditione Marocæana nondum detecta fuit. Specimina nostra ex Agadir et prope Mogador lecta gaudent floribus quam in *N. arvensi* typica paullo majoribus. Hæc forma prope Mogador a Balansa lecta sistit verisimiliter *N. Hispanicam*, var. *intermediam* (Coss., Index in Bull. Soc. Bot. Fr.).

NIGELLA DAMASCENA (L., DC. Pr. i. 49), forma spontanea floribus minoribus = *N. Bourgaei*, Jord. pug.

Mar. sept.—Tetnan (*Webb. herb.*)!

Mar. oceid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid. In prov. Shedma! et Haha (*J. D. H.*)!

Ar. Geog. Regio Mediterranea occidentalis. Europa. Afr. borealis. Macaronesia. In Oriente vix indigena?

AQUILEGIA (*L. Gen.* 684).

A. VULGARIS (L., DC. Pr. i. 50) var.?

Mar. merid.—In monte Djebel Tezah circ. 1800 m. legit *J. D. H.*

Ar. Geog. Europa borealis (excl. arctica), et media, et australis (in umbrosis montanis). Macaronesia. In Asia temperata et in mont. Himalaya. Formæ affines in America boreali.

Stirps polymorpha et niro modo ludibunda. Specimina nostra gaudent foliis radicalibus triternatis, foliolis profunde ineisis, omnibus glabrescentibus, petiolis longis, caule bipedali superne ramoso, floribus minoribus viridi-flavescentibus. Tota planta (foliis exceptis) pube brevi non viscosa prædita. Carpella desunt. Experientia docebit an varietas sit, vel subspecies nomine novo salutanda.

DELPHINIUM (*L. Gen.* 681).

D. AJACIS (*L., DC. Pr.* i. 51).

Mar. merid.—Mesfioua!

Ar. Geog. Europa meridionalis, Mæaronesia (an indigenum?).

Legi specimen unicum valde incompletum, et ideo minus certum.

D. HALTERATUM (*Sibth. et Sm. Fl. Gr.* vi. tab. 107; *Boiss. Fl. Or.* i. 86, et forsā *DC. Pr.* i. 53).

Mar. sept.—Tanger (*Salzm. Exs.* sub *D. junceo*)! (*Webb. herb.*)!

Mar. oeeid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Sat frequens in regione inferiore. Prov. Iaha! Shedma! Mtouga!

Ar. Geog. Regio Mediterranea. Europa. Africa borealis. Formæ affines in Oriente.

— —, var. *CARDIOPETALUM*, = *D. cardiopetalum* (*DC. Syst.* i. 347).

Mar. sept.—Tanger! Tetuan (anno 1851)! Forma elata, racemo laxo, foliis rigidis.

Mar. oeeid.—In monte Djebel Hadid! prope Aïn el Hadjar!

Ar. Geog. Gallia; Hispania. Africa borealis.

— —, var. *MACROPETALUM*, = *D. macropetalum* (*DC. Syst.* i. 350).

Mar. oeeid.—Frequens in arenosis prope Mogador! ubi legerunt etiam Broussonet, Sehouboe, Lowe, et Balansa.

Ar. Geog. Hucusque nonnisi pr. Mogador detectum.

Primo aspectu a *D. halterato* et formis affiniibus differt floribus duplo majoribus, pedunculis longioribus, foliorum segmentis latioribus. Characteres stabiles frustra quæsi dum hæ notæ omnes variabiles sunt. Vix aut ne vix quidem a *D. obcordato*, *DC. Syst. Nat.* i. 350, ex specimine maneo in 'Barbaria' lecto a el. Durand descripto, diversum est. Nihil dubito quin *Delphinia* omnia § *Delphinellum*, *DC.*, regionis Mediterraneæ (exclasis orientalibus et *D. Balansæ*) eum *D. peregrino*, *L.*, nexu genetico eolligari, et pro subspeciebus vel varietatibus enumeranda esse. *D. halteratum* dignoscitur petalorum lateralium lamina orbiculari basi rotundata vel leviter obeordata. *D. junceum* (*DC. Fl. Fr.* v. 364, et *Prod.* i. 52) habet laminam ellipticam in unguem attenuatam. Foliorum forma, indumentum, ratio longitudinis calcaris et sepalorum notæ sunt omnino variabiles. Formæ arete affines in tota regione Mediterranea (exclusa Mæaronesia) vulgares sunt.

DELPHINIUM, sp. nov.? Forsan subspecies *D. Balansæ* (*Boiss. et Reut. Diagn. Pl. Or.*, ser. 2, v. 12)?

Mar. merid.—In regione subalpina Atlantis Majoris. In convalle Aït Mesan supra Arround ad 2400 m.! In monte Djebel Tezali 2300–2500 m.!

Stirpem nondum florentem ineertæ sedis inter *Delphinia* (suadente el.

Baker) retuli; accuratius visa affinis *D. Balansæ* mihi videtur, sed pluribus notis differt. Radix perennis lignosa, plerumque obliqua; caules floriferi e basi procumbente adscendentes; folia radicalia et inferiora palmatum (nec pinnatum) partita, caulina magis divisa subbiternata; indumentum *D. Balansæ*.

DELPHINIUM PENTAGYNUM (*Lam. Dict.* ii. 264; *DC. Pr.* i. 54).

Mar. occid.—In monte Djebel Hadid haud procul a Mogador. *R. et Fr. spec. in Herb. Kew.*!

Ar. Geog. Peninsula Iberica. Africa borealis. Sicilia = *D. emarginatum* (Presl, Bert. Fl. It. v. 419).

Pace beati Bertoloni planta Algeriensis (Desf. Fl. Atl. i. 427, tab. 111) est certe non diversa a stirpe Calpensi quæ Maroccanis conformis. Specimina lusitanica non vidi, sed *Lam. l. c.* plantam lusitanicam sub oculis habuit, et eandem speciem descripsit (confer *Brotero, Fl. Lus.* ii. 304) quam pro *D. emarginato*, Psl. habuit auctor Fl. Italicæ. Notæ diagnosticæ mihi videntur incertæ et variabiles.

BERBERIDEÆ (*Vent. Tabl.* iii. 83).

BERBERIS (*L., Gen.* 442).

B. CRETICA (*L., DC. Pr.* i. 106), var. = *B. Hispanica* (*B. et R. Pug.* 3)?

Mar. merid.—In regione subalpina Atlantis Majoris. In monte Djebel Tezah (*J. D. H.*)!

Ar. Geog. Hispania. Africa boreali-occidentalis. Forma arete affinis in Oriente et in insulis Italicis Corsica, Sardinia, Sicilia (*B. Ætuensis*, Presl).

Specimina nostra inter *B. creticam* et *B. cratæginam*, DC., ambigua videntur sed propiora plantæ Nevadensi (olim *B. vulgaris*, L., var. *australis*, Boiss. = *B. Hispanica*, B. et R.). Folia subintegerrima in ramis fertilibus spinas subæquantia, in sterilibus (hornotinis?) duplo longiora. Baccæ nigro-cærulæ stigmatibus breviter stipitato coronatæ. *B. cretica*, L., cum *B. cratægina*, DC., et *B. densiflora*, Boiss. et Buhse, melius sicut subspecies *B. vulgaris* enumerandæ videntur.

PAPAVERACEÆ (*Juss. Gen.* 235).

PAPAVER (*L. Gen.* 648).

P. HYBRIDUM (*L., DC. Pr.* i. 118).

Mar. occid.—Casa Blanca! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Tam in planitie, prov. Shedma! et prope Marocco! quam in reg. inferiore Atlantis. Aït Mesan!

Ar. Geog. Europa media. Regio Mediterranea omnis.

P. DUBIUM (*L., DC. Pr.* i. 118), var. *Maroccanum*, nob.

Mar. sept.—Tanger! Inter Tetuan et Ceuta!

Mar. occid.—Casa Blanca!

Differt a typo foliis radicalibus pinnatis, caulinis integris basi latis subamplexicaulibus inæqualiter incisus vel crenato-incisus, interdum subtrilobis, pilosis vel subglaberrimis. Quoad foliorum formam videtur a *P. dubio* prorsus diversum et potius ad *P. gracile* (Boiss. Fl. Or. i. 115) accedens, sed formæ intermediæ non desunt.

P. dubium est species per totam Europam late diffusa, nec deest in regione Mediterranea (inclusa Macaronesia). In India boreali-occidentali adest forma glabra (*P. dubium* β . *lævigatum*, Elk. Mon.) quæ forma, monente Boissier (Fl. Or. i. 114), ab ipso Elkan cæterisque auctoribus cum *P. lævigato*, M. B., perperam confusa est. Vercor ne characteres a disci crenis incumbensibus vel inter se discretis a cel. Boissier allati, in hac grege fallaces sint. In stirpe nostra crenæ disci interdum incumbentes fere ut in *P. Rhœas*.

PAPAVER RHÆAS (L., DC. Pr. 118).

Mar. sept.—Tanger!

Mar. occid.—Mogador (Lowe); Agadir (Coll. Indig. 1871)!

Mar. merid.—Hinc inde in planitie Maroccana—Shedma! Sheshaoua!

Pr. urbem Marocco! Formam novam parvifloram legi pr. Sektana!

Ar. Geog. Europa media. Regio Mediterranea omnis.

P. TENUE (Ball, Journ. Bot. 1873, 296).

Annuum (seu bienne?); planta tota pilis longis setosis vestita (folia inferiora et sepala interdum glabrescentia); folia primaria pinnata, pinnis inciso-lobatis segmentis rotundatis, caulina circuitu late triangulari-ovata, bipinnatifida, segmentis lineari-oblongis acutis; caulis subsimplicis setæ inferiores patulæ, superiores adpressæ; petala obovato-oblonga; discus convexiusculus capsulæ glabræ obovato-turbinatæ latitudine vix æqualis, crenatus, crenis vix incumbensibus; stigmata 6-7.

Mar. merid.—In regione media et superiore Atlantis Majoris, in convalle Aït Mesan! et in jugo Tagherot! a 1800 m. ad 3000 m., in convalle Amsmiz! et in monte Djebel Tczah! a 1600 m. ad 2500 m.

Proximum *P. arenario*, M. B., at differt petalis angustis nec subrotundatis, setis caulinis adpressis nec patulis, cæterisque notis. Capsula fere *P. Decaisnei*, Hochst. et Steud., sed in hoc discus complanatus, crenæ magis discretæ.

P. SOMNIFERUM (L., DC. Pr. i. 119), var. SETIGERUM, = *P. setigerum* (DC. Fl. Fr. v. 585; DC. Pr. i. 119).

Mar. sept.—Tanger, versus Cap. Spartel (Webb. herb.)!

Mar. occid.—Mogador (Lowe Cat.).

Ar. Geog. In regione Mediterranea hinc inde, sed patria incerta.

P. somniferum colitur, et sæpe cum seminibus offertur in Imperio Maroccano, et ex cultis evadit forma sylvestris, *P. setigerum*, sed verisimiliter haud indigenum est.

PAPAVER RUPIFRAGUM (*Boiss. et Reut. Pug. p. 6*), var. ATLANTICUM (*Ball, Journ. Bot. 1873, 296*).

Percune, caespitosum; rhizoma multicaule; folia (fere omnia radicalia) setis longis albidis dense obteeta, circuitu obverse lanceolata, grosse dentata vel pinnatipartita, segmentis inæqualibus plus minusve incis; caulis scapiformis, 8-20-pollicaris, simplex vel ima basi furcatus, adpresse setosus; flores ante anthesin nutantes; petala late obovata, sordide aurantiaca; capsula clavata, glabra, ima basi (ad petalorum insertionem) annulo glandulifero instructa; discus convexus, crenatus, capsulae latitudinem superans.

Mar. merid.—In regione subalpina Atlantis Majoris; Aït Mesan! Djebel Tezali! a 2000 m. ad 2600 m.

Aspectu a planta Boissieriana diversissima; hæc enim obscure virens subglaberrima, dum varietas nostra canescens undique pilosissima, forsan melius pro subspecie habenda.

P. rupifragum est species distinctissima, hucusque tantum in Hispania Arundana detecta.

REMERIA (*Med. in Ust. Ann. 1792, 15*).

R. HYBRIDA (*DC. Syst. ii. 92, et Prod. i. 122, L. sub Chelidonio*).

Mar. merid.—In regione inferiore Atlantis Majoris. Pr. Sektana circ. 1200 m.! In convalle Amsmiz, circ. 1500 m.!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia). In Europa media vix indigena.

Siliquæ in spec. nostris aliquid attenuatæ fere ut in *R. rhæadiflora*, Boiss.

GLAUCIUM (*Scop. Fl. Carn. ed. 2, i. 369*).

G. LUTEUM (*Scop. l. c.*), = G. flavum (*Cltz. Fl. Aust. i. 141; DC. Pr. i. 122*), = G. fulvum (*Lois. Fl. Gall. i. 376; DC. Pr. i. 122*), = Chelidonium glaucium, L.

Mar. sept.—Legi nondum florens in arenosis maritimis inter Tetuan et Ceuta!

Ar. Geog. Europa borealis et media. Regio Mediterranea omnis (inclusa Macaronesia).

G. CORNICULATUM (*L., sub Chelidonio, Curt. Lond. vi. tab. 32; DC. Prod. i. 122*).

Mar. occid.—In prov. Haha prope Mogador! (*Schsb.*) et prope Agadir (*Coll. Indig. 1871*)!

Mar. merid.—Prov. Shedma! Mefsoua! Seksaoua!

Ar. Geog. Germania. Regio Mediterranea (exclusa Macaronesia).

HYPECOUM (*L. Gen. 171*).

II. PENDULUM (*L., DC. Pr. i. 124*).

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Amsmiz, circ. 1400 m.! Prope Mzonda, *J. D. H.*!

Ar. Geog. Europa media. Regio Mediterranea (exclusa Macaronesia).

CORYDALIS (DC. *Fl. Fr.* iv. 636; *Benth. et H. fil. Gen. Pl.* i. 55).

C. HETEROCARPA (*Durieu in Parl. Giorn. Bot.* i. 336, sub *Ceratocapno*),
= *Ceratocapnos umbrosa* (*Dur. in Walp. Ann.* ii. 29).

Mar. occid.—Prope Mogador (*Lowe Cat.*).

Mar. merid.—In regione inferiore Atlantis Majoris, in convalle Aït Mesan
detexit G. Maw!

Ar. Geog. Africa borealis. Hucusque nonnisi in Algeria detecta.

De genere *Ceratocapno* confer B. & H. *l. c.* p. 56. A *Corydali claviculata*, cui simillima, differt fructu dimorpho. Superiores cujusvis racemi capsulæ 1–2-spermæ, bivalves, in rostrum longum cornuto-recurvum attenuatæ; inferiores nuces indehiscentes, monospermæ, compressæ, truncatæ, fenestrato-sulcatæ. Nomen specificum a clar. inventore prius editum, aptum et nullomodo mutandum.

FUMARIA (*L. Gen.* 849).

F. AFRICANA (*Lam. Dict.* ii. 569), = *F. corymbosa* (*Desf. Act. Soc. Hist. Nat. Par.* i. 26, tab. 6; *DC. Pr.* i. 130; *Ol. Ham. Mon.* 42).

Mar. sept.—In rupibus juxta Tetuan florentem et fructiferam legi mense
Junio 1851 et iterum Aprili 1871!

Ar. Geog. Hispania Granatensis; Africa borealis. Species sat rara.

F. OFFICINALIS (*L., DC. Pr.* i. 130; *Ol. Ham. Mon.* 9).

Mar. merid.—In regione media Atlantis Majoris, in convalle Aït Mesan
a 2000 m. ad 2200 m.!

Ar. Geog. Patria verisimiliter regio Mediterranea. Nunc fere cosmopolita per Europam, Asiam, Africam borealem (inclusa Macaronesia), Americam tam borealem quam meridionalem late diffusa.

F. VAILLANTII (*Lois. Not.* p. 102; *DC. Pr.* i. 130; *Ol. Ham. Mon.* 14).

Mar. merid.—In regione inferiore Atlantis majoris pr. Tasseremout!

Ar. Geog. Europa et Asia temperata; regio Mediterranea (inclusa Macaronesia); India orientalis (*sec. Ol. Hammar*).

Notæ quibus hæc et sp. sequens dignoscuntur ex fructu desumptæ
vix stabiles sunt. Fructus *F. parvifloræ* in Marocco rarius apiculatus.

F. PARVIFLORA (*Lam. Dict.* ii. 567; *DC. Pr.* i. 130; *Ol. Ham. Mon.* 16).

Mar. occid.—Casa Blanca!

Mar. merid.—Sat frequens in plautic et in regione inferiore Atlantis
Majoris. Prov. Shedma, pluribus locis! Prope urbem Marocco!
In convalle Aït Mesan, *J. D. H.*! In convalle Amsmiz circ. 1500 m.!

Ar. Geog. Europa media (vix indigena?). Regio Mediterranea omnis,
unde in Americam subtropicam (Mexico, *sec. Parlatore*, Buenos Ayres,
herb. Hooker) et in Indiam boreali-occidentalem effugit.

In spec. Maroccanis flores sæpissime albi, rarius dilute rosei.

FUMARIA CAPREOLATA (*L., DC. Pr. i. 130; Ol. Ham. Mon. 24*), var. *CONDENSATA, nob.*

Mar. merid.—Ad radices Atlantis Majoris. Frequens circa Amsmiz! Mzouda! Keira!

Ar. Geog. *F. capreolata* per Europam fere totam et regionem Mediterraneam (inclusa Macaronesia) late diffusa est, et inde in Americam extratropicalem aufuga. De distributione formarum seu varietatum ardua sententia et forsitan inutilis est.

Var. nostra est forma parviflora, racemis densis, et forsitan eum var. γ , *albiflora*, *Ol. Ham. l. c.* conjungenda.

F. media (*Lois Not. 102; Ol. Ham. Mon. 28, non DC. Syst. nec Prod.*), var. α , *Ol. Ham. l. c.*, = *F. Boræi* (*Jord. Cat. Jard. Bot. Gren. 1849*), = *F. Munbyi* (*Bois. et Reut. Pug. p. 5*), = *F. capreolata* (*Eng. Bot. tab. 943, et auct. Brit. fere omnium*).

Mar. sept.—Circa Tanger! Prope Tetuan, anno 1851!

Ar. Geog. Europa occidentalis a Norvegia ad fretum Herculis. Africa boreali-occidentalis ab agro Tingitano ubi valde frequens usque Oran. Hue verisimiliter referenda est *F. capreolata* (*Salzm. Exsicc.*).

F. media est nobis subspecies eum *F. capreolata* pluribus formis intermediis conjuncta.

— —, var., cui proxima *F. SEPIUM* (*Boiss. Diagn. Pl. Or. ser. 2, i. 16; Ol. Ham. Mon. 27*).

Mar. merid.—In regione media Atlantis Majoris. In convalle Aït Mesan 1700–1800 m.!

Var. nostra inter formas intermedias enumeranda quibus adsocianda est *F. sepium*, *Boiss.* Habet sepala magna corollæ dimidium subæquantia, subintegra, basi leviter dentata, et bractæas pedicellis apice valde inerassatis æquilongas.

F. MACROSEPALA (*Bois. El. p. 8; Ol. Ham. Mon. 36*).

Mar. sept.—In monte Beni Hosmar legi specimen manebundum! In eodem monte olim legit beatus P. B. Webb.

Ar. Geog. Hispania Granatensis et Gaditana. Montes Tetuanenses.

Species rara, verisimiliter probe distincta.

F. AGRARIA (*Lag. El. Pl. Hort. Matrit. 1816, p. 21; Ol. Ham. Mon. 38*), = *F. media* (*DC. Syst. ii. 134; et Prod. i. 130, non Lois.*).

Mar. sept.—Tanger (*Salzm. Exsicc. sub F. media*)! Tetuan!

Mar. occid.—Casa Blanca! Mazagan! Mogador (*Lowe*). Agadir (forma fl. minoribus, *Coll. Indig. 1871*)!

Ar. Geog. Regio Mediterranea Europæ et Africæ borealis. Deest in Oriente et in Macaronesia. Ex Europa in Americam australem allata.

— —, var. *ELATA, nob.*, an *F. rupestris*, var. *laxa* (*Boiss. et Reut. Pug. p. 4*)?

Mar. sept.—In rupibus calcareis pr. Tetuan!

Mar. merid.—Prov. Haha! Shedma! Prope Marocco!

F. rupestris (B. et R. *l. c.*; Ol. Ham. Mon. 40), mihi videtur forma rupestris *F. agrariæ*, et var. *laxa* est ejusdem forma sylvestris scandens. Var. nostra differt a typo sepalis longius acuminatis apice subherbaceis, fructu insigniter compresso et carinato.

— —, var. ATLANTICA, nob.

Mar. merid.—In regione media Atlantis Majoris. Aït Mesan circ. 1600 m. ! Planta hand omnino certa—facies *F. capreolata*, sed characteres potius *F. agrariæ*; a priore differt sepalis multo latioribus. Multis notis, et foliorum segmentis subflabellatis, ad *F. flabellatam* accedit, sed bracteæ duplo longiores, ut in *F. rupestri*, et pedicelli erecto-patuli nec reflexi.

F. flabellata (Gasp., Rendie. Ac. Se. Nap. i. 51; Ol. Ham. Mon. 41) verisimiliter ut subspecies *F. agrariæ* enumeranda erit.

Fumaria tenuisecta (Ball, Journ. Bot. 1873, 297).

Affinis *F. agrariæ*, sed insigniter differt sepalis angustis minimis acutis dentatis diametrum fructus non attingentibus; a *F. judaica*, Boiss., petalis multo majoribus, et ab omnibus hujus gregis valde ludibundi foliis tripinnatisectis in lacinias lineares planas mucronatas dissectis.

Mar. merid.—In regione inferiore Atlantis majoris in convalle Aït Mesan ad 1400 m. specimen unicum legi !

Invito animo novam subspeciem proposui pro hac stirpe singulari. Folia fere *F. Kralikii*, Jord.

CRUCIFERÆ (Juss. Gen. 237).

MATTHIOLA (R. Br. in H. Kew. iv. 119).

M. TRISTIS (R. Br. *l. c.* p. 120; DC. Pr. i. 134).

Mar. occid.—Mogador (Lowe).

Ar. Geog. Europa mediterranea. Africa borealis. In Asia Minore rarior. Macaronesia (sec. Lowe).

M. TRICUSPIDATA (R. Br. *l. c.*; DC. Pr. i. 134).

Mar. sept.—In arenosis maritimis inter Tetuan et Ceuta !

Ar. Geog. Per oras maris Mediterraneæ late diffusa. In littoribus Atlanticis Europæ et Africæ hucusque non indicata.

M. PARVIFLORA (R. Br. *l. c.*; DC. Pr. i. 135).

Mar. merid.—Frequens in incultis saxosis calidioribus. Aïn Oumast ! Misra ben Kara ! Gurguri ! Mtonga ! Mskala ! Prope urbem Marocco (Schsb. spec.) !

Ar. Geog. Hispania; Marocco; Macaronesia.

In Marocco fit sæpe robustior, sesquipedalis et ultra.

CHEIRANTHUS (R. Br. in H. Kew. iv. 118).

C. SEMPERFLORENS (Schousb. Gew. in Marok. p. 181; DC. Pr. i. 136).

Mar. occid.—Frequens in arenosis circa Mogador ! ubi jam legerunt Broussonnet, Schousboe, Lowe et Balansa. Inde se protendit usque radices montes Djebel Hadid ad Aïn el Hadjar !

Ar. Geog. Species Marocana e grege Macaronesiaco *Dicroantho*, Webb, sed satis diversa. Specimen mancum sub hoc titulo, ex Mostaganem in herb. Kewensi est aliena planta. Semina immatura tenuiter marginata.

NASTURTIIUM (*R. Br. in H. Kew.* iv. 109).

N. OFFICINALE (*R. Br. l. c.* ; *DC. Pr.* i. 137).

Mar. sept.—Tanger !

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—In regione inferiore Atlantis Maj. pr. Ainsmiz, *J. D. H.* ! et in subalpinis supra Arround circ. 2400 m. !

Ar. Geog. Herba amphigæa, in temperatis, frigidioribus et calidioribus hemisphæræ borealis frequens.

N. ATLANTICUM (*Ball, Journ. Bot.* 1873, 297). (Plate IX.)

Bienné (?), cæspitosum, glaberrimum, ex collo radicis longæ verticalis caules breves debiles emittens ; folia (fere omnia basilaria) profunde pinnatipartita, lobo terminali anguloso-rotundato, lateralibus inæqualibus, spathulatis vel oblongo-ellipticis, angulatis ; pedicelli breves graciles, non incrassati ; petala parva, ochroleuca ; siliqua (immatura) lævis, subtorulosa, enervis ; stylus apice subclavatus.

Mar. merid.—In regione superiore Atlantis Majoris—Aït Mesan supra Arround ! Tagherot ! a 2200 m. usque 2800 m.

Ab affinibus (*N. asperum*, Boiss. (*Sisymbrium*, L. ; *DC. Prod.*) et *N. Boissieri*, Coss.) differt imprimis habitu et foliis diversissimis, floribus minoribus, ochroleucis (fere albidis) nec flavis, pedicellis gracilibus nec incrassatis, siliqua cuervi, nec ut in illis valvæ basi nervo notata. Stylus longitudine variabilis.

ARABIS (*L. Gen.* 818).

A. VERNA (*R. Br. in H. Kew.* iv. 105 ; *DC. Pr.* i. 142, *L.* sub *Hesperide*).

Mar. sept.—In monte Beni Hosmar ! In eodem monte et in Djebel Dersa olim legit Webb !

Ar. Geog. Regio Mediterranea, exclusa Macaronesia.

A. ALBIDA (*Stev. Cat. II. Gor.* 51 ; *DC. Pr.* i. 142).

Mar. merid.—In reg. subalpina Atlantis Majoris, in ascensu jugi Tagherot versus 2600 m. ! Fere eodem loco, anno 1873, specimen mancum legerunt R. et Fr. !

Ar. Geog. Montes regionis Mediterraneæ, sed deest in Gallia et in peninsula Iberica. In Italia media (territ. Lucensi) descendit in zonam *Oleæ Europææ*. In montibus ins. Teneriffa a 2200 ad 2700 m. se tenet.

Beatus P. B. Webb (in *Phyt. Can.* i. 63) et cel. Boissier (in *Fl. Or.* i. 174) plurimas sic dict. species orientales *A. albidæ* rite conjunxerunt,

scil. *A. caucasica* (W. Enum. Supp. 45), *A. viscosa* (DC. Syst. ii. 216), *A. thyrsoides* (Sibth. et Sm. Fl. Gr. tab. 642), *A. longifolia* (DC. Syst. ii. 219), *A. brevifolia* (DC. Syst. ii. 218), *A. Billardieri* (DC. Syst. ii. 218). Ab his omnibus specimina atlantica discrepant foliis inferioribus, et caudiculorum profunde sinuatis vel repando-dentatis, caulinis etiam magis profunde dentatis.

Procul dubio est forma *A. alpina* sed satis constans et ideo pro subspecie habenda.

ARABIS ERUBESCENS (Ball, Journ. Bot. 1873, 297).

Stirps male cognita e grege *A. alpina* sed cum ea confundi nequit. Ab hac differt insigniter sepalis basi æqualibus coloratis, petalis dimidio minoribus erubescens, corum lamina angustiore, pilis caulinis simplicibus vel apice fureatis, nec stellato-ramosis.

Mar. merid.—In regione alpina Atlantis Majoris. In jugo Tagherot, ultra 3000 m., specimen unicum immaturum et incompletum legi!

A. AURICULATA (Lam. Dict. i. 219; DC. Pr. 143).

Mar. merid.—In regione subalpina Atlantis Majoris. In convalle Aït Mesan supra Arround a 2400 m. ad 2550 m. ! In jugo Tagherot usque 2800 m. ! In monte Djebel Tezah a 2500 m. ad 2800 m. !

Ar. Geog. Europa media. Montes regionis Mediterraneæ, exclusa Macaronesia.

Specimina in monte Djebel Tezah lecta habent siliquas breviores. Nonnulla (versus 2800 m. lecta) sunt tenuiora pauciflora, fere var. β *Sinica* (Boiss. Fl. Or. i. 169).

A. PUBESCENS (Poir. Suppl. i. 413; DC. Pr. i. 145), = *Turritis pubescens* (Desf. Fl. Atl. ii. 92, tab. 163).

Mar. sept.—In monte Beni Hosmar ! et in rupibus calcareis pr. Tetuan ! ubi eam olim legit P. B. Webb.

Ar. Geog. Species propria Algeriæ et montium Tetuanensium.

Siliquæ erectæ strictæ subsessiles pubescentes, valvis dorso obsolete l-nervosis, in stylum attenuatæ. Semina uniseriata exalata. Stigma bifidum.

A. decumbens (Ball, Journ. Bot. 1873, 297). (Plate X.)

Perennis, cæspitosum, e collo radicis caules plurimos debiles decumbentes emittens; folia radicalia obverse lanceolata, acute repando-dentata, in petiolum attennata, caulina ovato-lanceolata, profunde dentata, suprema linearia subintegra, omnia ex pilis rigidis trifurcatis scabriuscula; pedicelli filiformes, patentes, siliquæ $\frac{1}{4}$ partem æquantes; petala alba seu dilute carnea; siliquæ compressæ subtorulosæ erecto-patentis valvæ margine parum incrassatæ, dorso obsolete nervosæ; stylus filiformis, glaber, pro genere longiusculus; stigma vix (aut ne vix) bifidum.

Mar. merid.—In regione suhalpina Atlantis Majoris—Aït Mesan, supra Arround ! a 2100 m. ad 2600 m. Djebel Tezah ! a 2400 m. ad 2500 m. Differt ab *A. pubescente*, cui proxima, habitu omnino diverso, pedicellis patentibus nec adpressis, siliquis magis torulosis, stylo longiore basi discreto. Tabula Fontanesiana *A. pubescentis* habet siliquas truncatas, stylo subnullo præditas ; in speciminibus nostris prope Tetuan lectis video siliquas semper in stylum attenuatas.

ARABIS CONRINGIOIDES (*Ball, Journ. Bot.* 1873, 297). (Plate XI.)

Radix lignosa, perennis, caules plurimos erectos, subsimplices, foliatis emittens ; herba tota glaberrima, glaucovirens ; folia inferiora orbiculata, in petiolum longum sensim attenuata, caulina inferiora elliptica, basi auriculata, auriculis subangulatis, superiora ovato-cordata, semi-amplexicaulia, omnia integerrima ; pedicelli numerosi, conferti, filiformes, erecto-patuli, siliquæ $\frac{1}{3}$ vel $\frac{1}{4}$ partem æquantes ; calyx basi subæqualis, sepalis purpurascentibus ; petala alba, apice rotundata in unguem sensim attenuata, suberecta ; siliqua erecta, elongata, lævis, subcylindrica, nervo dorsali subobsoleto notata, in stylum brevem truncatum attenuata ; semina 1-seriata.

Mar. merid.—In regione alpina et subalpina Atlantis Majoris—in jugo Tagherot ! a 3000 m. ad 3500 m. ; in monte Djebel Tezah ! a 2100 m. ad 2800 m.

Species distinctissima, nulli proxima, facie *Conringiæ orientalis* sive *C. clavatæ*, Boiss.

CARDAMINE (*L. Gen.* 812).

C. HIRSUTA (*L., DC. Pr.* i. 152).

Mar. sept.—Tanger (in monte Djebel Kebir) ! In umbrosis collium inter Tanger et Tetuan ! In monte Beni Hosmar pr. Tetuan !

Ar. Geog. Regiones temperatæ hemisphæræ borealis. Per regionem Mediterraneam (exclusa Macaronesia) late diffusa.

— —, var. *SYLVATICA*, = *C. sylvatica* (*Lk. in Hoffm. Phyt. Bl.* i. 50).

Mar. sept.—In monte Beni Hosmar pr. Tetuan circa 1000 m. !

Mar. merid.—In regione subalpina Atlantis Majoris—in monte Djebel Tezah circa 2200 m. !

Ar. Geog. Europa borealis et media (excl. arctica). Montes Hispaniæ et Italiæ, sed verisimiliter rara in reg. Mediterranea.

NOTOCERAS (*R. Br. II. Kew.* iv. 117).

N. CANARIENSE (*R. Br. l. c. ; DC. Pr.* i. 140).

Mar. sept.—Tetuan (*Webb Herb.*) !

Mar. merid.—Inter Misra ben Kara et Marocco !

Ar. Geog. Regio Mediterranea calidior a Belutschia usque in insulas Canarienses. India boreali-occidentalis.

ALYSSUM (*L. Gen.* 805; *H. et B. Gen. Pl.* i. 73).

A. ALPESTRE (*L., DC. Pr.* i. 161), var. *SERPYPILLIFOLIUM* = *A. serpyllifolium* (*Desf. Fl. Atl.* ii. 70).

Mar. merid.—In regione media et superiore (?) Atlantis Majoris. In convalle Aït Mesan circa 1400 m. florens et fructiferum legi 15 Majo 1871! Ejusdem stirpis specimen mancum legerunt in jngo Tagherot cl. Rein et Fritsch anno 1873! Denuo ecl. Cosson benevole communicavit specimina in monte Djebel Afougueur ab indigena Ibrahim lecta mense Junio 1874!

Ar. Geog. *A. alpestre* est species polymorpha in montibus Europæ mediæ et regionis mediterraneæ late diffusa. Var. nostra montium peninsulae Ibericæ et Africæ borealis incola est.

Forma Fontanesiana a nobis lecta vix differt ab *A. alpestre*, var. *β. suffrutescens* (*Boiss. Fl. Or.* i. 268). Injussu (monente ecl. Boissier) synonyma sunt *A. tortuosum β. orientale* (*DC., Prod.* i. 164), *A. Savranicum*, *Andr.*, et *A. serpyllifolium* (*Griseb. Spic.* i. 276).

— —, var. *MACROSEPALUM* (*Ball, Journ. Bot.* 1873, 298).

Differt a typo sepalis majoribus subpersistentibus siliculam semimaturam æquantibus, siliculis basi ellipticis, nec cuneatis, stylo siliculam subæquante.

Mar. merid.—In regione media Atlantis Majoris, in convalle Amsmiz!

A. MONTANUM (*L., DC. Pr.* i. 162), var. *ATLANTICUM*, nob., an *Boiss.?* = *A. atlanticum* (*Desf. Fl. Atl.* ii. 71, tab. 149; *DC. Pr.* i. 161).

Mar. merid.—In regione subalpina et alpina Atlantis Majoris. In monte Djebel Tezah a 2500 m. usque 3300 m.!

Ar. Geog. Varietas nostra habitat in montibus Africæ borealis et forsan Hispaniæ Granatensis. Formæ arcte affines per Europam mediam et regionem Mediterraneam (exclusa Macaronesia) diffusæ.

Specimina nostra bene convenient spec. in monte Djebel Cheliah Algeriæ lectis (*Balansa, Pl. d'Algérie*, No. 898). Ab icone Fontanesiana differre videntur caulibus subprostratis, et sepalis persistentibus. Specimina Hispanica quæ vidi et ipse legi fructifera in montibus Sierra Nevada ad aliam formam hujus speciei polymorphæ pertinere videntur. In his caules erecti rigidiusculi, planta tota ex squamis stellatis multiradiatis lepidotis dense incana. In spec. ab ipso Boissier in Sierra de Mijas anno 1837 lectis filamenta majora alata, nec dentata ut in *A. montano* typico, et in pl. nostra.

A. montanum est certe species polymorpha quibus adsociandæ sunt velut varietates vel subspecies plurimæ ab auctoribus jam descriptæ, et formæ non paucæ hucusque ineditæ. Inter varietates cum *A. atlantico*, *Desf.*, enumeranda sunt *A. diffusum*, *Ten.*, et *A. repens*, *Baumg.* Subspecies, notis stabilioribus agnoscendas, sistunt *A. flexicaule*, *Jord.*, *A. cuneifolium*, *Ten.*, et *A. Wulfenianum*, *Bernh.* Hoc novissimum, in Alpibus Carnicis et Juliis sat frequens, non habet siliculas semper glabratas.

Hæ, pace eel. Boissier, non raro sunt squamis lepidotis adpersæ, basi tamen semper cuneata, stylo longiore, ab illis *A. montani* genuini dignoscendæ.

ALYSSUM CAMPESTRE (L., DC. Pr. i. 163).

Mar. merid.—In planitie, unde se protendit usque in regionem subalpinam Atlantis Majoris. Prov. Shedma! Tasseremout! Aït Mesan circ. 1400 m. ! et iterum ad 2300 m. !

Ar. Geog. Europa media. Regio Mediterranea omnis (exclusa Macaronesia).

A. CALYGINUM (L., DC. Pr. i. 163).

Mar. merid.—Rarius occurrit in planitie et in regione subalpina Atlantis Majoris. Prov. Shedma, prope Aïn Oumast In convalle Aït Mesan ad 1200 m. et in jugo Tagherot usque 2700 m. !

Ar. Geog. Europa media. Regio Mediterranea (exclusa Macaronesia). In calidioribus fit plantula subsimplex erecta 2-3-pollicaris ; in frigidioribus e basi ramosissima subcæspitosa.

A. SPINOSUM (L., DC. Pr. i. 164 ; Boiss. Voy. Esp. ii. 45, sub *Ptilotricho*).

Mar. merid.—Crescit abunde in regione subalpina et alpina Atlantis Majoris in convalle Aït Mesan a 2200 m. usque 3300 m.

Ar. Geog. Regionis Mediterraneæ pars occidentalis, e Gallia Narbonensi ad Sierra Nevada ; et in Africa boreali-occidentali. In Macaronesia deest.

A. MARITIMUM (Lam. Dict. i. 98 ; DC. Pr. i. 164 ; Desv. Journ. Bot. iii. 162, sub *Lobularia* ; R. Br. in Clapp. Voy. App. p. 214, sub *Kæniga*).

Synonyma Linnæana confusa. Est enim *Clypeola maritima* (L. Mant. 426), et *Alyssum halimifolium* (L. Sp. Pl. ii. 207), et *A. minimum*, L., Sp. Pl. ii. 908).

Mar. sept.—Frequens pr. Tanger! (*Salzm. Exsicc.*)! prope Tetuan! Legi etiam in monte Beni Hosmar! ubi jam legerat Webb.

Mar. occid.—Djebel Hadid haud procul Mogador, *J. D. II.*!

Mar. merid.—In regione inferiore Atlantis Majoris—pr. Milhain!

Ar. Geog. Regio Mediterranea littoralis (inclusa Macaronesia).

— —, var. LEPIDOIDES, nob.

Differt a typo habitu graciliore, siliculis minimis in racemo confertis, nec ut in typo racemo basi laxo.

Mar. merid.—Ad radices Atlantis Majoris, supra Seksaoua! et in prov. Mtouga!

A. LIBYCEUM (*Viv. Pl. Lib.* p. 34, tab. 16, sub *Lunaria* ; R. Br. in Clapp. Voy. App., sub *Kæniga*).

Mar. occid.—Mogador in arenosis! (*Lowe Cat.*) et in rupibus insule! Prope Aïn el Hadjar!

Ar. Geog. Regio Mediterranea calidior imprimis occidentalis (Hispania australis, Africa boreali-occidentalis, Macaronesia). Nascitur etiam in Persia australi et in Arabia Petræa.

DRABA (*L. Gen.* 800).

D. HISPANICA (*Boiss. El.* p. 8, *et Voy. Esp.* tab. 13; *Walp. Rep.* i. 147). Mar. merid.—In regione superiore Atlantis Majoris legimus in monte Djebel Tezah a 2100 m. ad 3000 m.!

Specimina nostra ab exemplaribus Hispanicis et Algeriensibus paululum recedunt siliculis ovatis stylum versus attenuatis, minus hispidis (interdum glabrescentibus), pilis semper simplicibus, nec pilis ramosis commixtis, petalis minoribus styli longitudinem vix attingentibus.

Ar. Geog. Hispania Granatensis, Africa borealis.

Formæ arcte affines per montes Europæ meridionalis, Asiæ minoris et Syriæ diffusæ.

EROPHILA (*DC. Syst.* ii. 356).

E. VULGARIS (*DC. l. c.*; *DC. Pr.* 172), = *Draba verna*, *L.*

Mar. sept.—In monte Beni Hosmar pr. Tetuan!

Ar. Geog. Europa omnis (excl. arctica). Oriens. Africa borealis, ubi rara. In Macaronesia non indicata.

MALCOLMIA (*R. Br. in H. Kew.* iv. 121).

M. LITOREA (*R. Br. l. c.*; *DC. Pr.* i. 187).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)!

Mar. occid.—Casa Blanca (*R. et Fr. spec. in Hb. Kew.*)!

Ar. Geog. Litora Oceani a Cherbourg usque ad fretum Herculis. Secus mare Mediterraneum in Hispania et Hetruria. Africa boreali-occidentalis.

M. LACERA (*L. Sp.* 926, sub *Cheirantho*; *Lam. Dict.* iii. 332, sub *Hesperide*; *DC. Syst.* ii. 445, *et Prod.* i. 188), var. *PATULA*, = *M. patula*, (*DC. Syst.* ii. 444, *et Prod.* i. 187).

Mar. occid.—Valde frequens in arenosis prope Mogador! et in reg. litorea usque Aïn el Hadjar!

Ar. Geog. Hispania centralis. Ager Mogadorensis. Arcte affinis *M. lacera* habitat in reg. litorali Lusitaniæ et Hispaniæ meridionalis.

Huc verisimiliter specimina a Broussonnet in agro Mogadorensi lecta quæ ill. A. P. De Candolle sub *M. lacera* (*Syst. l. c.*) disposuit. Re vera *M. lacera* et *M. patula* sunt formæ ejusdem speciei. In hac folia linearia integra, in illa oblongo-linearia, pinnatifido-sinuata. Ab his differt *M. Broussonnetii* stylo basi scabrido, foliis sæpe latoribus integris, vel breviter repando-dentatis; nec video aliam notam diagnosticam. In omnibus stylus filiformis elongatus, apicem versus glaber, vel tenuissime papilloso-scaber.

— —, var. *Broussonnetii*, = *M. Broussonnetii* (*DC. Syst.* ii. 445, et *Prod.* i. 184; *Deless. Icon. Sel.* ii. tab. 69).

Mar. occid.—In arenosis pr. Mogador! (*Brouss.*) *DC. l. c.* Inter Shedma et Mogador legit G. Maw et specimina pulchra mihi benevole dedit!

Mar. merid.—Prope Mesfioua unicum specimen legi!

Ar. Geog. Nimis incerta, vix ultra territ. Maroccanum.

M. Broussonnetii est *M. patulæ* forma vegetior, sæpius statura major, ramis rigidioribus, floribus majoribus. In spec. nostris tamen flores paullo minores quam in icone Delessertiana. Quid sit *M. Broussonnetii* auct. Floræ Hispanicæ est mihi res dubia. Specimina sub hoc titulo a Boissier et Reuter prope Gibraltar anno 1837 lecta, et quæ ipse legi anno 1871, mihi videntur magis affinia *M. litoreæ*, et forte ejusdem varietas foliis profunde sinuatis. An diversa sit *M. erosa* (*Lag. Cat. Hort. Matr.* 1814, p. 20, sub *Hesperide*, *DC. Syst.* ii. 446) ex descriptione incompleta non satis apparet.

M. NANA (*DC. Syst.* ii. 486, et *Prod.* i. 195; *Coss. in Bull. Soc. Bot. Fr.* x. 397, sub *Sisymbrio*, *Boiss. Fl. Or.* i. 222), = *M. binervis* (*Boiss. Ann. Sc. Nat.* 1842, 71), = *Sisymbrium binerve* (*C. A. M. Enum.* p. 189; *Jaub. et Sp. Ill. Pl. Or.* tab. 289), = *Hesperis ramosissima* (*DC. Prod.* i. 189), = *M. confusa*, *Boiss. Fl. Or.* i. 221.

Mar. sept.—Specimina duo semipollicaria florentia detexi inter cæspites graminæ ejusdam in saxosis prope Tetuan lectæ!

Ar. Geog. Species per regionem Mediterrancam fere omnem late dispersa. In Gallia meridionali, in Italia, Græcia, et in Africa boreali hinc inde occurrit. In litore Maris Caspii legerunt Hohenacker, C. A. Meyer, et Buhse; et in maritimis Anatoliæ (suadente Boissier) olim legit Tournefort.

Pace cel. Boissier differentia inter *M. nanam* et *confusam*, Boiss., omnino instabiles sunt.

Speciminibus ad Mare Caspium prope Baku lectis plantulæ nostræ adamussim conveniunt nisi stylo aliquantulum brevior. Distributio hujus speciei, forsitan ex statura minima a viatoribus prætermisæ, admodum singularis.

SISYMBRIUM (*All. Fl. Ped.* i. 274).

S. OFFICINALE (*L.*, sub *Erysimo*, *Scop. Fl. Carn.* ii. 20; *DC. Pr.* i. 191).

Mar. sept.—In calcareis prope Tetuan! et in monte Beni Hosmar!

Ar. Geog. Europa media et borealis (excl. arctica). Regio Mediterranea (inclusa Macaronesia). In Americam advecta.

S. IRIO (*L.*, *DC. Pr.* i. 192).

Mar. sept.—Tanger (*Schsb. sec. Cosson*). Visum pr. Tetuan!

Mar. occid.—Casa Blanca!

Mar. merid.—Sheshaoua! Inter rudera urbis Maroeco!

Ar. Geog. Europa occidentalis a Britannia ad fretum Herculis. Regio Mediterranea (inclusa Macaronesia). India boreali-occidentalis.

SISYMBRIUM THALIANUM (*Gay in Gaud. Fl. Helv.* iv. 348; *L. et DC.*, sub *Arabide*).

Mar. merid.—In regione subalpina Atlantis Majoris, in monte Djebel Tezah a 2100 m. ad 2300 m.!

Ar. Geog. Europa omnis (inclusa arctica). Sibiria. Regio Mediterranea (excl. Macaronesia).

S. RUNCINATUM (*Lag. Fl. Hisp. ined.*; *DC. Pr.* i. 194).

Mar. orient.—Maharoug (*Warion*), *Coss. Index*.

Ar. Geog. Hispaniæ prov. Murcia. Africa borealis.

— — —, var. *HIRSUTUM*, *nob.*, = *S. hirsutum* (*Lag.*, *DC. Pr.* l. c.)

Mar. merid.—In regione inferiore Atlantis Majoris, in convalle Amsmiz circa 1500 m. specimen unicum legi exemplaribus Hispanicis prope Matritum lectis omnino conforme.

Ar. Geog. Hucusque nonnisi in agro Matritensi et in Atlante Majore detectum.

S. ERYSIMOIDES (*Desf. Fl. Atl.* ii. 84, tab. 158; *DC. Pr.* i. 195).

Mar. merid.—Sat frequens in ruderalis et locis saxosis reg. inferioris.

Prov. Shedma! Ain Beida! in ruderalis urbis Marocco! Seksaua! Milhain! et alibi visum.

Ar. Geog. Regio Mediterranea imprimis occidentalis, scil. Macaronesia, Hispania australis, Africa boreali-occidentalis, Sardinia; deinde occurrit in Persia australi et Arabia et in Abyssinia. Deest ut videtur in Italia continentali, Græcia, Asia Minore, Syria et Ægypto.

Specimina in solo arido creseentia graciliora sunt, sed vix pro varietate habenda.

ERYSIMUM (*L. Gen.* 814).

E. AUSTRALE (*Gay, Erysim. Diagn.* p. 6), = *F. pallens* (*Boiss. Voy.*

Esp. in Add. p. 716), var. *GRANDIFLORUM*, = *E. grandiflorum* (*Desf.*

Fl. Atl. ii. 85), = *E. longifolium* (*DC. Syst.* ii. 504, et *Prod.* i. 199).

Mar. merid.—In regione media et superiore Atlantis Majoris a 1900 m. usque 3000 m. In convalle Aït Mesau unde se protendit usque in jugo Tagherot! In monte Djebel Tezah! Specimina in monte Djebel Afougueur (*Coll. Indig. Ibrahim*) siliquis maturis (nobis prius deficientibus) mense Augusto 1874 lecta, benevole comm. cel. Cosson!

Ar. Geog. Var. nostra montium Africæ borealis et Hispaniæ incola est.

Formæ arete affines per Europam mediam et totam regionem Mediterraneam (exclusa Macaronesia) diffusæ.

Inter tot tantosque viros rei Botanices expertissimos qui genus *Erysimum* tetigerunt sententiæ omnino discordes. Invita natura characteres stabiles frustra quærun, et limites certos indicare conantur in catena formarum affinium quam maxime proximarum.

In speciminibus nostris folia sæpius anguste linearia sed interdum latiora, semper tamen integerrima. Pedicelli sæpissime calycis longitudinem dimidiam attingentes, sed interdum longiores. Siliquæ tenues ex pilis adpressis undique (etiam in angulis) canescentes. Semina exangulata. Confer Gr. et God. Fl. Fr. i. 89.

Exemplaribus permultis hujus gregis accurate observatis species s. d. Europæ videntur omnes sub *E. Cheiranthio*, Pers., velut subspecies seu varietates enumerandæ. Synonymia extricatu difficillima.

BRASSICA (*L. Gen.* 820; *B. et H. Gen. Pl.* i. 84).

[In genere a cel. Benthām et Hooker reformato cum *Brassicis* veris (Linnæano sensu) enumerantur sicut subgenera *Sinapodeudron*, Lowe, *Sinapis*, L., *Erucastrum*, Schimp. et Spenn., et *Hirschfeldia*, Moench. His forsā melius addenda erit *Diploaxis*, genus caractere nullo satis stabili fulcratum. Plurimæ species habitu proprio a *Brassicis* distinguuntur, sed *Diploaxis virgata*, *D. siifolia*, *D. unriculata* nec habitu nec caractere a *Brassica* dignoscendæ, et nisi obstitisset auctoritas gravissima *Diploaxis* cum cæteris generibus supra memoratis sub *Brassica* enumerare maluissē, duce cel. Boissier (*Voy. Esp.* ii. 32), etsi vir celeb. in Flora Orientali sententiam mutavit.]

B. TORULOSA (*Durieu in Rev. Bot.* ii. 434; *Expl. Bot. Alg.* i. 74; *Walp. Ann.* i. 48).

Mar. sept.—Tanger (*Salzm. Exsicc. sub* *Diploaxis catholica*)! (*Webb Herb.*); Tetuan (*Webb Herb.*)!

Mar. occid.—Mazagan! Mogador (*Balansa, Coss. Index*).

Ar. Geog. Africa boreali-occidentalis; Hispania?

Nimis proxima *Diploaxis siifolia*, Kze.; hanc ægre distinguo siliquæ rostro longiore tenuiore insigniter trinervi, nec ut in *D. siifolia* sæpius obsolete trinervi.

B. FRUTICULOSA (*Cyr. Pl. Var. Neap.* fasc. 2, p. 7, tab. 1; *DC. Pr.* i. 216).

Mar. occid.—Prope Mogador (legit Lowe, sed determinatio incerta)?

Cel. Cosson spec. Loweanum cum *Sinapi radicata* (Desf. Fl. Atl. ii. 98, tab. 167) comparavit et in eadem sententia facile hæreo. *Sinapis hispida*, Desf. est enim, ut mihi fidem fecerunt specimina (*Balansa, Pl. d'Alg. No. 50*) ex Mostaganem, varietas hispida *Brassicæ fruticulosæ*, Cyr. Confer tamen Boiss. Fl. Or. i. 393, de *B. fruticulosa*.

B. VARIA (*Durieu in Rev. Bot.* ii. 434; *Walp. Ann.* i. 48), = *Erucastrum varium* (*Durieu, Expl. Bot. Alg.* tab. 75).

Mar. merid.—In saxosis aridis pr. Seksaoua! Milhain!

Ar. Geog. Marocco; Algeria occidentalis.

Inter spec. duo supra Seksaoua lecta unum refert potius var. β . mon-

tanam, Dur. *l. c.* Siliquæ pubescentes in spec. nostris sunt magis patientes quam in spec. e loco classico pr. Oran receptis.

BRASSICA ELATA (*Ball, in Journ. Bot.* 1873, 298). (Plate XII.)

Perennis; caulis erectus, simplex, 3-6-pedalis, superne glaberrimus subaphyllus; folia radicalia (10-14-pollicaria) et caulina inferiora lyratopinnatifida, lobo terminali-maximo, rhomboideo, inæqualiter crenatodentato, lateralibus oppositis alternisve, ovato-cuneatis, dentatis, rachilata cum nervis lateralibus albo-rubente, omnia hispido-incana; inflorescentia pyramidalis e ramis brevibus patulis subæqualibus; pedicelli breves, stricti; sepala erecta, obtusa, pilosiuscula; petala saturate lutea, limbo subrotundo; siliqua subtorulosa, superne nervosa, in rostrum conicum 1-2-ovulatum trinerve attenuata.

Mar. merid.—In regione inferiore Atlantis Majoris—in rupibus apricis prope Seksaouna! et Milhain! circ. 1000 m. Huc verisimiliter frustulum mancum in monte Djebel Afougneur ab indigena Ibrahim lectum et a cel. Cosson benevole communicatum!

Species insignis, ab affinibus omnino dissimilis. Proxima est *B. bætica*, Boiss., sed in hac rami laterales distantes erecti, nec velut in nostra planta versus apicem caulis approximati patuli, siliquæ rostrum bi-ovulatum, pedicelli erecto-patentes. Ill. Boissier calycem *patentem* *B. bætica* tribuit, et sepala *obtusa*, dum in tabula (Boiss. Voy. en Esp. pl. 9) pictor sepalum *acutum* perperam delineavit. In speciminibus siccis mihi obviis sepala erecta nec patentia videntur. In eadem tabula siliquæ rostrum ovulis 2 abortivis ostenditur. Ubi ovulum unum alterumve fertile adest rostrum fit longius et siliquæ dimidiam longitudinem attingit. In specie nostra et in sequente siliqua quam in *B. bætica* multo minor.

B. RERAYENSIS (*Ball, Journ. Bot.* 1873, 298). (Plate XIII.)

Radix perennis; caulis e basi adscendente erectus 1-2 furcatus, glaberrimus, subaphyllus; folia radicalia parva, incano-hirsuta, runcinato-lyrata, segmentis basin versus alternis, caulina pauca demum linearia; racemus brevis, pauciflorus; pedicelli erecto-patuli, tenues, siliquam $\frac{1}{2}$ æquantes; rostrum tenue, aspermum(?), siliquæ $\frac{1}{4}$ seu $\frac{1}{5}$ partem attingens; sepala elliptico-oblonga, pilosa; petala saturate lutea.

Mar. merid.—In regione subalpina Atlantis Majoris in convalle Aït Mesan, a 2000 m. ad 2500 m.

Fructus in speciminibus nostris omnino immaturnus, melius exploratus forsitan ovulum in rostro detectum fuerit. Quocunque fit a *Brassicis* planta nostra nullomodo divellenda. *B. bætica*, Boiss., proxime affinis, differt foliis glabris vel ex setis brevibus sparsis scabrida, racemo elongato, siliqua duplo longiore, rostro crasso dispermo, nec tenui aspermo.

B. NIGRA (*L. et DC. Pr.* i. 218, sub *Sinapi*; *Koch, Deutsch. Fl.* iv. 713; *Boiss. Fl. Or.* i. 390).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Europa (exclusa arctica); regio tota Mediterranea.

Specimen immaturum et incertum a *J. D. H.* prope Tingidem lectum forsā huc spectat.

BRASSICA ARVENSIS (*L. et DC. Pr.* i. 219, sub *Sinapi*, = *B. sinapis* (*Vis. Fl. Dalm.* iii. 136).

Mar. sept.—Tanger (*Webb Herb.*)!

Mar. merid.—? Specimen mancum valde incertum legi pr. Sheshaoua!

Spec. alterum pariter incertum pr. Ourika legerunt *R. et Fr.*!

Ar. Geog. Europa et Asia temperata; regio Mediterranea.

B. ALBA (*L. et DC. Pr.* i. 220, sub *Sinapi*; *Boiss. Voy. Esp.* ii. 39).

Mar. occid.—Mogador (*Lowe Cat.*).

Forsan est potius species sequens. *B. alba* hucusque in Imp. Maroecano non alibi visa sæpe cum seminibus cœrealibus in terras longinquas aufertur.

Ar. Geog. Verisimiliter indigena in regione Mediterranea orientali, unde versus occidentem usque Insulas Canarienses, et per Europam mediam diffusa est.

B. HISPIDA (*Schsb. Gen. Marok.* p. 182, tab. 4, et *DC. Pr.* i. 220, sub *Sinapi*, non Tenore).

Mar. occid.—Verisimiliter ex agro Mogadorensi orta sunt semina a Schousboe culta.

Mar. merid.—In arvis pr. Seksaoua!

Ar. Geog. Algeria; Marocco; Macaronesia.

Tabula Schousboeana effigies plantæ cultæ ex seminibus Maroccanis—verisimiliter ex agro Mogadorensi. Planta spontanea gracilior est, et foliis multo minoribus prædita.

B. hispida, Tenore, est *Eruca sativa*, Lam.

B. ADPRESSA (*Moench, Meth.* 261, sub *Hirschfeldia*; *Bois. Voy. Esp.* ii. 38), = *Sinapis ineana* (*L., DC. Pr.* i. 220), = *Erueastrum incanum* (*Koch, Syn.* p. 56).

Mar. sept.—Tetuan (1851)!

Mar. merid.—Misra ben Kara haud procul Marocco! Keira (*Balansa*), *Coss. Index.*

Ar. Geog. Europa occidentalis; regio Mediterranea fere omnis (inclusa Macaronesia).

— —, var. *LASIOCARPA*, nob. = *Sinapis heterophylla* (*Lag. Cat. Hort. Matr.* 1816; *DC. Pr.* i. 220).

Mar. merid.—Mesfioua! In convalle Ait Mesan cire. 1300 m.!

Spec. nostra a typo longius recedunt quam spec. quæ ex Hispania vidi.

B. GENICULATA (*Desf. Fl. Atl.* ii. 98; et *DC. Pr.* i. 218, sub *Sinapi*).

Mar. merid.—Sheshaoua! In hortis et ruderatis urbis Maroeco specimina immatura legimus!

Ar. Geog. Africa borealis.

Hæc species, cum *B. nigra* ab ill. A. P. De Candolle adsociata, meliore jure in § *Hirschfeldia* enumeratur. Habet rostrum 1-2-spermum et siliquam nullomodo tetragonam, gaudet enim hæc nervo medio parum prominulo nervis lateralibus evidentibus, quibus notis *B. adpressæ* potius quam *B. nigre* propius accedit.

BRASSICA NERVOSA (*Ball, Journ. Bot.* 1873, 299).

Annua (?), ramosa; folia radicalia lyrata ineiso-dentata, caulina pauca lanceolata, omnia in petiolo et nervis hirta; petala (pro grege) magna læte lutea; siliquæ brevis insigniter nervosæ pube brevi velutina vestitæ rostrum glabrum, aspermum (?).

Mar. merid.—Specimen unicum maneum legi in prov. Mtouga!

Forsan ineapte nomen novum proposui (*loc. cit.*) pro planta male cognita, verisimiliter in § *Hirschfeldia* enumeranda. Ex structura siliquæ et cotyledonibus conduplicatis ad *Brassicas* certe referenda.

BRASSICA, spec. nov. ?

Mar. sept.—Cel. J. D. Hooker specimen legit, verisimiliter inter Tanger et Tetuan, cui desunt flores et folia inferiora. Proximum videtur *Sinapi Allionii*, Jacq., sed in spec. Hookeriano rostrum valvarum dimidio æquilongum, eximie trinerve, compressum, subaneeps. Foliolum caulinum unicum quod adest omnino *Barbareæ vulgaris*, R. Br., unde nomen triviale *barbareæfolia* aptum fuerit.

BRASSICA, spec. nov. ?

Mar. sept.—Alterum specimen ex regione collina inter Tanger et Tetuan attulit cel. Hooker, verisimiliter ad *Brassicam* referendum, sed ob fructum nimis immaturum incertum. Planta tota glabra, nisi foliis inferioribus setis nonnullis retrorsis præditis. Folia inferiora runcinato-pinnatifida. An forma singularis *B. nigre* ?

BRASSICA, spec. nov. ?

Mar. merid.—In convalle Amsmiz ad 1400 m. specimen unicum legi!

Annuum; caulis e basi ramosus, ramis subaphyllis 6-8-pollicaribus; folia radicalia et inferiora pauca obverse lanceolata in petiolum attenuata, repando-dentata, præsertim secus nervos strigulosa; racemus elongatus; pedicelli glabriuseuli, stricti, calyceem subæquantes; sepala æqualia, basi non saecata, purpurascens; petala in ungucem sensim attenuata, oehroleuca, venis saturate violaceis; siliquæ immaturæ subcylindricæ elongatæ erectæ, cum caule ex setis brevibus retrorsis seabræ; rostrum glabrum, tenue, valvarum sextæ parti æquilongum, verisimiliter aspermum. Stigma capitatum breviter bifidum. Habitu a *Brassicis* recedit; planta ulterius inquirenda.

DIPLOTAXIS (*DC. Syst.* ii. 220).

D. HARRA (*Forsk. Fl. Æg.-arab.* 118, sub *Sinapi*; *Boiss. Fl. Or.* i. 388), = *D. pendula* (*DC. Syst.* ii. 630, et *Prod.* i. 222), = *D. hispida* (*DC. l. c.*), = *D. crassifolia* (*DC. l. c.*), = *Sisymbrium pendulum* (*Desf. Fl. Atl.* tab. 156).

Mar. merid.—Prope Amsmiz circa 900 m. !

Ar. Geog. Hispania; Sicilia. Oriens. Africa borealis.

Siliquæ in spec. nostris sæpius non pendulæ.

D. ERUCOIDES (*L.*, sub *Sinapi*; *DC. Syst.* ii. 631, et *Prod.* i. 222).

Mar. sept.—Tanger !

Ar. Geog. Regio Mediterranea, sed non ubique; deest in Macaronesia, Græcia, Asia Minore et in Ægypto.

D. VIRGATA (*Cav. Præl.*; *DC. Syst.* ii. 631, et *Prod.* i. 222).

Mar. sept.—Prope Tanger !

Ar. Geog. Hispania. Africa borealis.

Specim. nostra Tingitana sistunt hanc vel varietatem seminibus sub-uniseriatis. Formam majorem foliis profundius divisis legi prope S. Roque !

D. CATHOLICA (*DC. Syst.* ii. 632, et *Prod.* i. 222; *L.* sub *Sisymbrio*).

Mar. sept.—Prope Tetuan legi anno 1851 specimina incompleta verisimiliter huc referenda !

Ar. Geog. Europa australis; Africa borealis.

D. SIIFOLIA (*Kunze in Flora*, xxix. 685; *Walp. Ann.* i. 49).

Mar. sept.—Frequens circa Tanger ! In regione collina inter Tanger et Tetuan ! In maritimis inter Tetuan et Centa !

Mar. occid.—Frustulum plantæ 4-pedalis haud procul Mogador legit J. D. Hooker verisimiliter huc referendum ! ?

Ar. Geog. In utroque litore freti Herculis, et (?) prope Mogador.

D. siifolia est *D. catholicæ* proxima species, sed forsau satis ditincta foliis et caule pilis patentibus vel subreflexis hispidulis, foliorum laciniis ovatis vel inciso-lobatis acutis, nec linearibus obtusis, petalis ochroleucis nec luteis. Ludit glabrescens sed foliorum indole scuper diversa. Confer *Brassicam torulosam*, Dur.

D. TENUIFOLIA (*L.*, sub *Sisymbrio*; *DC. Syst.* ii. 632, et *Prod.* i. 222).

Mar. sept.—Tanger (*Salzm. Exsicc.* sub *D. catholica*) !

Ar. Geog. Europa media et Mediterranea. Tauria. Asia Minor.

D. TENUISILIQUA (*Del. Ind. Sem. II. Mousp.* 1847, 7; *Walp. Ann.* i. 49), = *D. auriculata* (*Durieu in Expl. Bot. Alg.* tab. 76, sine descr.).

Mar. occid.—Casa Blanca ! Aïn el Hadjar, haud procul Mogador, J. D. H. !

Mar. merid.—Satis frequens in regione inferiore. In aridis fit tenuior,

pumila parviflora. Prov. Shedma prope Aïn Oumast ! prope Mskala ! Misra Ben Kara ! Amsmiz ! Prov. Mtouga ! et alibi visa.

Ar. Geog. Africa boreali-occidentalis.

Nomen triviale *tenuisiliqua* jure prioritatis conservatur.

Typo Algeriensi sat bene congruunt specimina nostra ; paululum variant siliquis longioribus, caule interdum pilis retrorsis hispidulo.

— —, var. RUPESTRIS, *nob.*

A typo differt habitu graciliore, caule ramosissimo cum foliis hispidulo, seminibus sæpissime uniseriatis.

Mar. merid.—In rupibus aridis supra Seksaoua ! Eandem formam legunt in prov. Mtouga el. *R. et Fr.* !

In hac specie pedicelli et siliquæ semper patentes vel erecto-patentes.

ERUCA (*DC. Syst.* ii. 636).

E. SATIVA (*Lam. Fl. Fr.* ii. 496 ; *DC. Pr.* i. 223), var. *STENOCARPA*, *Coss.*, = *E. stenocarpa* (*Boiss. et Reut. Pug.* 8).

Mar. merid.—Frequens in regione inferiore. Tasseremout ! Ourika ! Reraya ! Milhain ! Seksaoua usque 1500 m. ! Sektana (*R. et Fr. spec. in herb. Kew.*) ! Keira (*Balansa*) ; Akka (*Mard.*), *Coss. Index.*

Ar. Geog. Africa borealis. Forma typica per regionem Mediterraneam (inclusa Macaronesia) late diffusa et in cultis ac ruderalis Europæ mediæ cum seminibus adveeta.

Varietas distinguitur habitu tenuiore, siliquis angustioribus subuniseriatis. Adsunt formæ intermediæ.

CARRICHTERA (*DC. Syst.* ii. 64).

C. VELLÆ (*DC. Syst., Ic. et Prod.* i. 224), = *Vella* annua, *L.*

Mar. merid.—Sat frequens in regione inferiore. Prov. Shedma (pluribus locis) ! Sheshaona ! Misra Ben Kara !

Ar. Geog. Europa meridionalis calidior. Oriens. Africa borealis.

SUCCOWIA (*Med. in Ust. Ann.* i. 64).

S. BALEARICA (*DC. Syst.* ii. 642, et *Prod.* i. 224).

Mar. sept.—In monte Beni Hosmar spec. unicum detexit G. Maw et mihi benevole dedit !

Ar. Geog. Regio Mediterranea occidentalis a Sicilia usque Ins. Canarienses, nullibi a mare remota.

CAPSELLA (*Vent. tabl.* iii. 110).

C. BURSA-PASTORIS (*L.*, sub *Thlaspi* ; *Moench. Meth.* p. 271 ; *DC. Prod.* i. 177).

Mar. sept.—Tanger !

Mar. merid.—In regione inferiore et subalpina et alpina Atlantis Majoris.

In convalle Aït Mesan supra Arronnd (2200 m.?), *R. et Fr. spec. in herb. Kew.*! In jugo Tagherot ultra 3000 m.!

Ar. Geog. Herba amphigæa in temperatis utriusque hemisphæræ et usque in regiones arcticas diffusa, hominum vestigia secuta.

SENEBIERA (*Poir. Dict.* vii. 75).

S. CORONOPUS (*L.*, sub *Cochlearia*, *Poir. Dict. l. c.*; *DC. Prod.* i. 203).

Mar. sept.—Tetuan!

Mar. merid.—Misra Ben Kara, haud procul urbe Marocco!

Ar. Geog. Europa media; regio Mediterranea (exclusa Macaronesia).

LEPIDIUM (*L. Gen.* 801).

L. SATIVUM (*L.*, *DC. Prod.* i. 204).

Mar. merid.—In ruderatis urbis Marocco!

Ar. Geog. Regio natalis incerta. Per orbem late diffusum, sed sæpius ex hortis aufuga.

L. NEBRODENSE (*Rafin. Préc.* p. 38, sub *Nasturtio*; *Guss. Fl. Sic.* ii. 154; *Boiss. Fl. Or.* i. 355), = *L. pubescens* (*Tin. Cat. H. R. Panorm.* 150, non *Desv.*), = *L. Bonannianum* (*Guss. Prod. Fl. Sic.* p. 211), = *Lepia Bonanniana* (*Psil. Fl. Sic.* 84), var. ATLANTICUM, nob.

Folia radicalia spathulata longe petiolata, integra vel lyrata, caulina sessilia vix auriculata, omnia velutina; sepala glabra, purpurascentia; silicula glabra, matura interdum pubescens, breviter emarginata; stylus brevissimus.

Mar. merid.—In regione superiore Atlantis Majoris, vall. Aït Mesan! a 2500 m. ad 3000 m.

An diversum sit *L. stylatum* (*Lag. Ann. Cienc. Nat.* 1802) non sat liquet; confer *Guss. Fl. Sic. l. c.* Nomen infaustum, dum styli longitudo in hac grege nimis variabilis.

Ar. Geog. Formæ arete affines per regionem Mediterraneam (exclusa Macaronesia) late diffusæ, nullibi frequentes.

Plurimæ s. d. species sicut varietates sive subspecies enumerandæ, scil. *L. calycotrichum*, Kze. = *L. granatense*, Coss. = *L. Dhayense*, Munby, quocum forsân jungendum est *L. microstylum* (*Boiss. & Held. Diag. ser.* 2, fasc. vi. p. 21, et *Boiss. Fl. Or.* i. p. 355). *L. nebrodensis* forma aberrans est (ni fallor) *L. petrophilum*, Coss. Valde affinis est species Europæ occidentalis *L. hirtum*, L., quocum adnumeranda sunt *L. heterophyllum*, Benth., *L. Smithii*, Hook., et *L. Villarsii*, G. et G., dum inter has formas characteres firmos adhuc frustra quærunt botanici. *L. oxyotum*, DC. Syst. = *L. humifusum*, Req., est species insulæ Corsicæ propria, erronee in DC. Prod. vol. i. p. 204, et *Boiss. Fl. Or.* i. 356, ad Floram Syriacam relata. Specimina enim in herb. Labillardière loci natalis indicatione earent, sed spec. Corsicis omnino conformia sunt. Hoc tamen, etsi siliculæ forma sat bene definitum, est

sp. nostræ forsā nimis proxima; vidi inter specimina Sicula *L. nebrodensis* formam intermediam.

LEPIDIUM LATIFOLIUM (*L.*, *DC. Pr.* i. 207).

Mar. sept.—Legi anno 1851 prope Tetuan ad ostium fluminis!

Ar. Geog. Europa temperata; regio Mediterranea (exclusa Macaronesia); Asia media. Verisimiliter stirps orientalis in regionibus occidentali-bus advena.

BISCUTELLA (*L. Gen.* 808).

B. APULA (*L.*, *DC. Pr.* i. 182).

Mar. sept.—In monte Djebel Kebir prope Tanger! Prope Tetuan in Monte Dj. Dersa (*Webb Herb.*)!

— —, var. **MICROCARPA**, = *B. microcarpa* (*DC. Pr. l. c.*).

Mar. occid.—Aïn el Hadjar, haud procul Mogador!

Mar. merid.—In regione inferiore Atlantis Majoris. Tasseremout! Ourika! Aït Mesan usque 1700 m.!

B. lyrata (*L.*, *DC. Pr.* i. 182), = *B. raphanifolia* (*Poir.*), = *B. maritima* (*Ten. Prod. Fl. Neap.* 38).

Mar. sept.—Tanger in saxosis! In monte Beni Hosmar supra Tetuan, *J. D. H.*! Prope Tetuan ad rivos fluminis legi anno 1851!

Mar. merid.—In prov. Mtouga! In reg. subalpina Atlantis Majoris supra Arround ad 2200 m. legi specimen unicum siliculis minimis ad varietatem *B. scutulatam* (*Boiss. et Reut. Diagn. Pl. Or. ser. 1, i. 41*) accedens.

Nihil dubito quin *Biscutellæ* omnes annuæ regionis Mediterraneæ (exclusa *B. obovata*, sp. omnino dubia) pro unius speciei prole haberi debeant. Forma foliorum radicalium, qua nota *B. lyrata*, *L.*, *B. raphanifolia*, *Poir.*, et *B. maritima*, *Ten.*, a cæteris dignoscuntur, subconstans, et hæ meo sensu subspeciem sistunt. Sed notæ diagnosticæ ex caule plus minusve folioso et ramoso, ex siliculis minoribus majoribusve, disco et margine puberulis, vel scabris, vel lævibus, vel disco lævi et margine scabro, omnino fallaces sunt, et inter exemplaria eodem loco nata plane variabiles. Habeo specimen prope Tanger lectum ad *B. baticam* (*Boiss. et Reut. Diagn. Pl. Or. ser. 1, i. 42*) accedens. Specimina secus flumen pr. Tetuan lecta sunt proxima *B. maritimæ*, *Ten.* Sed in hac grege qui nolit speciem in catervam inordinatam divellere de nominibus auctorum parum curare debet.

THLASPI (*L. Gen.* 802).

T. PERFOLIATUM (*L.*, *DC. Pr.* i. 176).

Mar. sept.—In monte Beni Hosmar prope Tetuan!

Mar. merid.—In regione media Atlantis Majoris, in convalle Aït Mesan circ. 2100 m.!

Ar. Geog. Europa media (in Scandinavia advena). Asia temperata. Regio Mediterranea.

— — —, var. *ROTUNDIFOLIUM*, *nob.* = *T. rotundifolium* (*Tin. Pl. Rar. Sic.* fasc. i. 46), = *T. Tineanum* (*H. de Pav. Pl. Sic. Exsicc.*).
Mar. merid.—In regione alpina Atlantis Majoris. In jugo Taghierot usque 3200 m. ! In monte Djebel Tezali !
Spec. nostra Siculis omnino conformia. Secundum auctores præcipuos siliculæ *T. perfoliati* sunt in quovis loculo 4-ovulatæ. In spec. nostris video sæpe semina 5 in quovis loculo fructus maturi.
Clar. Rein et Fritsch ex regione alpina versus jugum Taghierot retulerunt specimen mancum nimis incompletum, verisimiliter in genere *Thlaspi* ponendum.

IBERIS (*L. Gen.* 834).

I. PARVIFLORA (*Munby, in Bull. Soc. Bot. Fr.* ii. 282), ? = *I. Numidica* (*Jord. Diagn.* 290).

“Pedicellis fructiferis corymbosis; radice annua; canle a basi ramoso parce hirtulo; foliis glabrescentibus linearibus, valde dentatis quasi pectinatis; siliculis emarginatis, sinu latiusculo, stylo dimidium sinus æquante; petalis subæqualibus vix calycem violaceum superantibus.”

See. diagnosi Munbyanum, *l. c.*

Mar. merid.—Specimina duo fructifera prope Tassercimont legit el. G. Maw; quorum unum mihi communicavit !

Ar. Geog. Africa boreali-occidentalis.

Specimina nostra diagnosi supra citatæ hand omnino conformia, sed speciminibus Bourgeanis prope Gharrouban in prov. Oran lectis sat bene conveniunt. In spec. nostro racemus fructifer corymbosus, folia margine setosa, subpinnatifida lobis paucis 1-2 utrinque instructa; pedicelli fructiferi rigidi superne papilloso-scabri, inferne læves; stylus basi alæ siliculæ vix adnatus.

Pace cel. Cosson hæc mihi potius quam *I. pectinata*, Boiss., videtur *I. odorata*, L., proxima, et forsan ejus subspecies.

I. GIBALTARICA (*L., DC. Pr.* i. 181).

Mar. sept.—In fissuris rupium montis Beni Hosmar prope Tetuan ! eodem loco quo eam olim legit fructiferam beatus P. B. Webb.

Ar. Geog. In promontorio Calpes et in montibus Tetuancensibus.

HUTCHINSIA (*R. Br. in Hort. Kew.* iv. 82).

II. PETRÆA (*L., sub Lepidio; R. Br. l. c.; DC. Pr.* i. 178).

Mar. sept.—Inter Tetuan et Ceuta ! In monte Beni Hosmar prope Tetuan !

Mar. merid.—In regione subalpina Atlantis Majoris. Aït Mesan supra Arround circ. 2300 m. ! In monte Djebel Tezali ad 2500 m. !

Ar. Geog. Europa (exclusa arctica). Montes regionis Mediterraneæ (exclusa Macaronesia).

ISATIS (*L. Gen.* 824).

I. TINCTORIA (*L., DC. Pr.* i. 211), var. *LÆTEVIRENS*, *nob.*

Differt a planta Europæa satis variabili foliis læte virentibus, sed non video notas graviores differentiales. Siliculæ obverse lanceolatæ, sæpius apice obtusangulo.

Mar. merid.—Frequens in regione media Atlantis Majoris unde secus torrentes in reg. inferiorem descendit. In convalle Aït Mesan a 1600 m. usque 2000 m. ! In convalle Amsmiz 1100–1700 m. ! Specimina nostris conformia sub “*I. virens*, Coss. MSS.” in montibus Djebel Onensa et Djebel Afougucur ab indigeno Ibrahim lecta benevole comm. cel. Cosson.

Ar. Geog. Species per Europam mediam et australem, et per Orientem diffusa. Varietas Maroccana est.

CRAMBE (*L. Gen.* 825).

C. HISPANICA (*L., DC. Pr.* 226), = *C. reniformis* (*Desf. Fl. Atl.* ii. 78, tab. 151 ; *DC. Pr. l. c.*), = *C. filiformis* (*Boiss. Voy., an Jacq. ?*).

Mar. merid.—In regione media Atlantis Majoris. In convalle Amsmiz a 1500 usque 1800 m. !

Ar. Geog. Peninsula Iberica. Africa borealis.

C. hispanicam, *L.*, et *C. reniformem*, *Desf.*, sistere unam eandemque speciem mihi haud dubium videtur. Nec in descript. auctorum, nec in exemplaribus plurimis Hispanicis et Algeriensibus a me sedule investigatis, nec in aspectu et habitu plantæ spontaneæ, quam pluries vivam observavi, notam differentialem nisi levissimam detexi. Quid sit vera *Crambe filiformis*, *Jacq.*, hic non dijudicare licet. Nomen triviale Linnæanum certe conservandum.

RAPISTRUM (*Desv. Journ. Bot.* iii. 160).

R. RUGOSUM (*L.*, sub *Myagro* ; *Berg. Phyt.* iii. 171 ; *DC. Pr.* i. 227).

Mar. sept.—Tanger ! (*Webb Herb.*) !

Ar. Geog. Europa media ; regio Mediterranea omnis. Ludit siliculis hirtis vel glabris.

CERATOCNEMUM (*Coss. et Bal. in Bull. Soc. Bot. Fr.* xx. 239).

“Sepala erecto-patula, lateralia haud saccata. Petala longe unguiculata, limbo integro. Stamina tetradynama, filamentis liberis subfiliformibus inferne complanatis edentulis. Glandulæ hypogynæ quatuor, duæ ad staminum longiorum paria, duæ ad stamina lateralia. *Siliqua*

abbreviata, *indehiscens*, coriaceo-indurata mesocarpio suberoso, *biarticulata*; articulo inferiore pedicello vix latiore, tereti-subcompresso, *uiloculari*, *monospermo*, *valvis distinctis* sed *cohærentibus in appendicem terminalem triangulari-lanceolatam articulo superiori æquilongam* et in anfractu dorsali articuli superioris receptam *productis*; articulo superiore vix secedente, *subrhombeo* apice truncato stylo cuspidato, a latere *compresso*, nervo præminente suturam valvarum articuli inferioris continuante donato, utrinque dorso ad receptionem processus valvaris excavato, *uiloculari*, *monospermo*, evalvi. Stylus siliqua longior, lanceolato-linearis, tetragonus. Stigma depressum, integrum. Semina oblonga, funiculo brevissimo, *semen articuli inferioris pendulum*, *superioris erectum*. *Cotyledones* rectæ, obovato-suborbiculatæ, retusæ, *cauliculato-complicatæ*, radiculam amplexantes.

“Planta annua, erecta, ramosa, pilis simplicibus hirta. Folia oblonga, grosse sinuata dentatave, superiora angustiora sæpe subintegra. Racemi ebracteati, virgati, demum elongati, pedicellis filiformibus demum subclavato-incrassatis. Flores lutei. Siliquæ parvulæ, erecto-adpressæ.

“*Ceratocnemum*, novum genus ex ordine Cruciferarum, juxta *Rapistrum* collocandum, sed distinctum valvis articuli inferioris siliquæ in appendicem articulo superiori æquilongam productis, non abrupte truncatis, articulo superiore subrhombeo compresso utrinque uninervio, non ovoideo vel subgloboso pluricostato.—Nomen generis ob formam insignem articuli inferioris siliquæ e verbis græcis *κέρας* (*cornu*) et *κνημῖς* (*ocrea*) conflatum.”

CERATOCNEMUM RAPISTROIDES (*Coss. et Bal. l. c.*).

“Planta annua. Caulis erectus, a parte inferiore vel a basi ramosus, teres, pilis albis rigidulis deflexis hirtus. Folia hinc inde præsertim in nervis et ad margines hirta, oblonga, in petiolum brevem angustata, grosse sinuato-subpinnatifida dentatave, superiora angustiora dentata vel subintegra. Racemi ebracteati, primum subconferti, demum elongati virgati laxiusculi. Pedicelli calyce subbreviares, suberecti, demum incrassato-subclavati, hirti. Sepala hirta. Petala lutea, calyce subtriplo longiora, limbo obovato integro venis saturatoribus picto, in unguem exsertum attenuata. Stamina longiora calycem longe excedentia. Siliquæ pilis patentibus hirtæ, 8–10 millim. longæ, erectæ, adpressæ. Semina stramineo-fuseescentia, sub lente tenuissime punctulata. Maio 1867 floriferum et jam fructiferum lectum.”

Diagnoses accuratas gen. et spec. nov. ad calcem conservare malui.

Mar. merid.—In monte Djebel Orguis supra Imintenout detexit cl. Balansa! Legimus 4–6-pedale ramosissimum in sepibus distr. Mesfionia haud procul urbem Marocco! et nanum (vix pedale) prope Tasseremout!

CAKILE (*Adans. Fam. Pl. ii. 423*).

C. MARITIMA (*Scop. Fl. Carn. No. 844; DC. Pr. i. 185*), = *Bunias Cakile, L.*

Mar. sept.—Tanger in arenosis! In arenosis inter Tetuan et Ceuta!

Ar. Geog. Litora maritima totius Europæ et Africæ borealis, in Macaronesia et Africæ litore Atlantico nondum detecta.

Forma arcte affinis in America boreali et in Antillis.

HEMICRAMBE (*Webb, in Ann. Sc. Nat. ser. 3, xvi. 246, tab. 19*).II. FRUTICULOSA (*Webb, l. c.; Walp. Ann. iv. 215*).

Descriptionem Webbianum hic referre licet.

“Suffruticulus glaberrimus caudicibus lignosis, tortuosis, nodosis, nudis ad ramorum apicem foliosis. Folia petiolata, læte viridia, lyrata, basi sæpe auriculata, lobis 2 inferioribus oppositis, 7-nervata, nervis a medio ad lobos divergentibus, venis reticulatis sub epidermide insignita, basi subcordata truncata vel cuneata, lamina 2–2½ poll. longa, 1½–2 poll. lata. Petioli 2–3 poll. longi, ascendentes, basi latiores, subsemivaginant, supra canaliculati, juniores glandulis nigris inconspicuis conspersi. Ramuli floriferi foliis subduplo longiores. Racemi laxè paniculati. Pedicelli filiformes, floribus duplo longiores, fructiferi elongati, ebracteati. Calyx basi fere æqualis, luteus, patens, mox deciduus, foliolis lanceolatis, concavis, intus nervosis, tenuibus, diaphanis. Petala lutea, calyce duplo longiora, longe unguiculata, unguiculis angustis, laminis obovato-lanceolatis, basi attenuatis, apice rotundatis, integris, ramoso-nervosis. Filamenta edentula, sub medio dilatata, apice subulata. Antheræ æquales lanceolatae, sub medio affixæ, basi sagittatae, flavæ. Glandulæ hypogynæ 4, quarum 2 post stamina longiora sitæ iisque alternantes, calcariformes, obtusæ, 2 ante stamina breviora iisque oppositæ, tenues laminiformes ovario applicatæ. Ovarium ellipticum, planum, basi attenuatum, trinerve, margine acutum, ad medium usque alatum. Stylus nullus. Stigma sessile globosum, centro depressum. Siliquæ ellipsoideæ utrinque attenuatæ, apice rostratæ, glabræ. Funiculi recurvi, usque ad medium dissepimento rudimentario immersi. Ovula globoso-pyriformia, sub apice affixa. Semina ovata, compressa, reticulata, fusca. Cotyledones rotundatæ, tenues. Radicula cylindracea cotyledonibus longior.”

Mar. sept.—In fissuris rupium calcarearum montis Beni Hosmar supra Tetuan primus detexit beatus P. B. Webb. Eodem fere loco ad 900 m. s. m. legimus vix florentem die 12 Aprilis 1871!

Ovarium (utrinque) trinerve sec. descript. Webbianum. In flore innupto video nervos duos laterales conspicuos, medium vero subobsoletum.

Planta singularis inter *Brassicæas* et *Cakiliæas* transitum præbens,

septo rudimentario, siliquæ articulo inferiore interdum aspermo vel prorsus abortivo. Habitus *Brassicæ*. Caudex basi diametro tripollicaris et ultra.

RAPHANUS (*L. Gen.* 812).

R. RAPHANISTRUM (*L., DC. Pr.* i. 229).

Mar. sept.—Tanger! (*Webb Herb.*)! In regione collina inter Tanger et Tetuan!

Mar. occid.—Casa Blanca!

Ar. Geog. Europa omnis; regio Mediterranea (inclusa Macaronesia).

CAPPARIDÆ (*Juss. Gen.* p. 242).

CAPPARIS (*L. Gen.* 643).

C. SPINOSA (*L., DC. Pr.* i. 245), var. β , CANESCENS (*Coss. Not. Pl. Crit.* i. 38), = *C. herbacea* (*Willd. Enum. Ber.* 560; *DC. Pr.* i. 245).

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Aït Mesan a 1300 ad 1600 m.!

— —, var. γ , ÆGYPTIA (*Boiss. Fl. Or.* i. 420), = *C. Ægyptia* (*Lam. Dict.* i. 605; *DC. Pr.* i. 245).

Mar. merid.—Sat frequens in apricis saxosis ad radices Atlantis Majoris. Gurguri! Seksaoua! Inter Sektana et Frouga (*G. Maw spec.*)! et alibi pluries visa!

Optimo consilio eel. Boissier loc. cit. species plurimas auctorum notis diagnosticis stabilibus non definitas sub typo Linnæano sicut varietates enumeravit.

Ar. Geog. Species per regionem omnem Mediterraneam late diffusa est, sed deest in Macaronesia.

RESEDACEÆ (*DC. Theor. Elem.* p. 214).

RESEDA (*L. Gen.* 608).

R. ALBA (*L., Müll. Arg. in DC. Pr.* xvi. sect. 2, 557).

Mar. sept.—Tetuan!

Mar. occid.—Casa Blanca! Mazagan! Mogador (*Lowe*)!

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia). Occurrit in Europa media, præsertim occidentali, forsân non indigena. Verisimiliter introducta in Indiæ orientalis montibus Nilgherry, et in Capite Bonæ Spei.

R. PROPINQUA (*R. Br. Obs. Oudn. Denh. Clapp.* p. 22; *Müll. Arg. in DC. Prod.* xvi. sect. 2, 559), = *R. eremophila* (*Boiss. Diagn. ser.* 1, viii. 88).

Mar. occid.—Secus Oued Kseb prope Mogador!

Mar. merid.—Prov. Shedma prope Mskala! Prov. Mtouga!

Ar. Geog. Regio desertorum a Syria et Arabia Petræa per Ægyptum, et

in plaga Africæ Tripolitanæ et Tunetanæ, et in Marocco meridionali.

R. propinqua est species sat variabilis, in eodem specimine video flores tetrameros et pentameros; scil. sepalis et petalis 4 vel 5. Petala in spec. Maroccanis iconi Müllerianæ (Monog. tab. vi. fig. 90) haud omnino conformia, sed differentię parvi momenti. Sepala sæpius subinæqualia, superiora paullo longiora. Capsulæ glandulosæ, sæpissime octocostatæ, costis 4 vel rarius 3 placentis oppositis elevatis subalatis. Semina fere ut in *R. alba*.

R. tricuspis (Coss. in Bull. Soc. Bot. Fr. xx. 241) cujus specimen prope urbem Marocco lectum ex herb. Schousboe debeo humanitati cæl. auct., differt a *R. propinqua* imprimis capsula trigona ore tridentato. Forsan pro subspecie habenda; video tamen capsulam in *R. propinqua* interdum, etsi rarius, tridentatam.

RESEDA ATTENUATA (Ball, in Journ. Bot. 1873, 299). (Plate XIV.)

Radix perennis; caules 2-3 exiles, 8-12-pollicares; folia inæqualiter pinnatifida, caulina abbreviata, 2-4-juga; spica gracillima, pauciflora; bracteæ lineares, pedicellos brevissimos sæpius superantes; sepala ovato-lanceolata, superiora 3 interdum coalita; petala brevissima calyceem vix superantia, staminibus breviora, 2 superiora gaudent appendicula (pro grege) magna, papilloso-ciliolata, concava, et lamina appendiculam æquante, profunde trifida, lobis æqualibus, petala media superioribus subæqualia, petalum inferius in laciniis tres æquales uninervias ultra medium divisum; capsula dentibus quatuor brevibus erectis coronata, pube rara brevissima adspersa.

Mar. merid.—In regione alpina et subalpina Atlantis Majoris, Ait Mesan! et in jugo Tagherot! a 2100 m. ad 3000 m. Ex montibus Djebel Afougueur et Djebel Ouensa (*Coll. Indig. Ibrahim*) specimina benevole comm. cæl. Cosson.

Proxima *R. Gayanæ* (Boiss. Voy. Esp. p. 76, tab. 21; Müll. Arg. Monog. p. 111). Differt præsertim petalis minimis, lamina fere ad basin trifida, omnibus appendiculatis, et capsulæ dentibus magis prominentibus. Forsan sicut subspecies enumeranda.

In DC. Prod. xvi. sect. 2, 558, cl. Müller, *R. Gayanam* sub nomine "*R. undata*, L." enumeravit. De *R. undata* inter auctores gravissimos magna est orta dissensio. Ipse Linnaeus speciem male cognovit et quoad fructum descriptio (Spec. Pl. ed. 2, p. 644) plane erronea, et auctores fere omnes ad aliam speciem nomen retulerunt. Sententiam primam cl. auctoris Monographiæ secutus sum, et nomen *R. undata* delendum esse censeo.

R. PHYTEUMA (L., Müll. Arg. in DC. Pr. xvi. sect. 2, 563).

Mar. merid.—Prov. Shedma! In convalle Amsmiz circa 1200 m.!

Ar. Geog. Europa media. Regio Mediterranea imprimis occidentalis, sed deest in Macaronesia.

Reseda media (*Lag. Nov. Gen. et Spec.* p. 17; *Müll. Arg. in DC. Pr.* xvi. sect. 2, 563), = *R. macrosperma* (*Rehb. in Flora*, 1830, i. 130; *Walp. Repert.* ii. 752; *Müll. Arg. Monog.* 133).

Mar. sept.—In monte Djebel Kebir prope Tanger! et in saxosis prope Cap Spartel!

Ar. Geog. Peninsula Iberica. Forma arcte affinis (*R. orientalis*, Boiss.) in Oriente infrequens.

Specimina nostra Hispanicis conformia, sed folia fere omnia integra, capsulae demum horizontales, nec pendulae. Praeter semina paululum majora petalorum lacinulae spathulatae, nec lineares, notam diagnosticam praebent. Adsunt tamen, sed rarius, formae intermediae, et hanc pro subspecie *R. Phyteumatis* enumerare malui.

R. diffusa (*Ball, Journ. Bot.* 1873, 299).

Radix biennis (?), caulis e basi ramosissima, diffusa; folia integra vel rarius trifida, oblongo-linearia, obtusa; bractea filiformes, (pro grege) longiuseculae; sepala ut in *R. media*, Lag.; petalorum superiorum lamina brevis, antheris vix longior, 3-partita, laciniis lateralibus profunde 3-5fidis, lacinulis apice abrupte spathulatis (fere cochleariformibus); capsula (nondum matura) insigniter 3-dentata, dentibus divergentibus.

Mar. merid.—In rupibus calcareis ab urbe Marocco septentrionem versus legit. cl. G. Maw, et mihi benevole communicavit.

A proxima *R. media*, Lag., recedit petalorum lamina multo minore, lacinulis cochleariformibus, nec obovato-oblongis apice rotundatis, capsulae dentibus divergentibus, qua nota ad *R. inodoram*, Rehb., proprius accedit. *R. media*, Lag. et *R. inodora* ad subspecies, sub typo *R. Phyteumate*, L., reducendae mihi videntur. Planta nostra longius a typo recedit.

R. LUTEA (L., *Müll. Arg. in DC. Pr.* xvi. sect. 2, 569), var. **MUCRONULATA**, = *R. mucronulata* (*Tin. Cat. H. Panorm.* 1828, p. 280).

Mar. merid.—Prov. Mtouga! Aïn Tarsil! Shedma!

Distinguitur caule stricto subsimplici elongato, capsulae papillosae, praesertim immaturae, dentibus longioribus. Individua vegetiora ad typum revocantur.

Ar. Geog. Varietas nostra (ex *Müll. Arg. l. c.* 570) nascitur in peninsula Iberica et in Algeria et in insula Zante.

Species per Europam mediam et regionem Mediterraneam late diffusa est, sed in Africa boreali infrequens. Deest in Macaronesia, ubi locum tenet species affinis *R. crystallina*, Webb.

R. neglecta (*Müll. Arg. Monog.* p. 178, et in *DC. Pr.* xvi. sect. 2, 568).

Mar. merid.—In rupibus supra Seksaoua!? Frustulum ex Oasi Akka (*Coll. Indig. Mardochaus*) benevole comm. ecl. Cosson.

Ar. Geog. Huensque nonnisi in Algeria meridionali ad fines Saharæ detecta.

Frustula spec. ab Oasi Akka, quæ humanitati eel. Cosson debeo, specimenibus Kralikianis a Müll. Arg. citatis non omnino conformia. Specimen a nobis ad Seksaoua lectum, *R. luteæ* habitu simile, ex forma capsularum propius *R. neglectæ*. Hæc mihi certe forma *R. luteæ*, et forsitan potius varietas quam subspecies, videtur. Forma petalorum inter varietates Müllerianas *R. luteæ* valde indibunda. Quoad notam diagnosticam a magnitudine capsulæ desumptam confer verba ipsius auctoris. Sub *R. neglecta* habet "capsulæ 12 mm. longæ," dum sub *R. lutea* var. α legitur "capsulæ sæpius 12 mm. longæ." Denique semina in *R. neglecta* minora et magis numerosa, sed ni fallor adsunt individua intermedia.

RESEDA LANCEOLATA (*Lag. Gen. et Spec.* p. 17; *Müll. Arg. in DC. Pr.* xvi. sect. 2, 572), var. CONSTRICTA, nob., = *R. constricta* (*Lange, Descr. Pl. Nov. præc. Hisp.* fasc. i. 6, tab. viii., *icon optima*).

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Aït Mesan a 1200 m. usque 1600 m.! Supra Seksaoua circ. 1500 m.! Djebel Sidi Fars (*Balansa, Coss. Index*).

Ar. Geog. Hucusque nonnisi in Hispania australi visa.

R. constricta sistit varietatem satis insignem, pedicellis longis patulis subhorizontalibus, capsulis longius dentatis erectis. Folia in spec. ab Aït Mesan omnia trifida segmentis latis. Capsulæ dentes divergentes.

R. STRICTA (*Pers. Ench.* fii. 10; *Müll. Arg. in DC. Pr.* xvi. sect. 2, 373), var. SUBSESSILIS (?).

Mar. occid.—Aïn el Hadjar haud procul Mogador!

Specimen unicum legi nondum fructiferum et ideo incertum; differt a spec. Hispanicis pedicellis brevissimis.

Ar. Geog. *R. stricta* est imprimis species Hispanica: varietates tres a Müller *l. c.* enumeratæ in Africa boreali sat late diffusæ.

R. ELATA (*Müll. Arg. in DC. Pr.* xvi. sect. 2, 578; *Coss. in Bull. Soc. Fr.* xx. 242).

Descriptionem Cossonianam optimam hic referre licet.—

"Planta biennis vel induratione perennans. Caules elati, indurato-frutescentes, ramosi, brevissime puberuli, ramis elongatis virgatis cortice olivaceo-viridi striis albidis. Folia sub lente brevissime puberula, in petiolum attenuata, plana, indivisa lineari-lanceolata elongata acutiuscula, vel rarius trisecta segmentis foliis indivisis conformibus. Racemi densiflori, bracteis apice subcomosi, demum elongati, capsulis inferioribus remotiusculis, superioribus approximatis. Bractæ lanceolato-lineares longe acuminatæ, late albido-marginatæ, glabræ,

flores sub anthesi longe excedentes, post anthesin cito deciduæ. Calyx 6-sepalus, sepalis oblongo-lanceolatis acutiusculis, late albo marginatis, glabris vel dorso brevissime puberulis, cito deciduis. Petala calyce subduplo longiora, omnium ungue albo suborbiculato villosa, lamina pallide lutea, superiorum trisecta lobo medio lineari-spathulato lobis lateralibus subduplo brevioribus in lacinias inæquales lineares 6-9-sectis, lateralium trisecta lobo medio oblongo inferne angustato lobis lateralibus indivisis, inferiorum lobo medio conformi lobis lateralibus sæpius obsoletis. Disci hypogyni lamina majuscula ovata acutiuscula ascendente marginibus recurvis extus concava, intrinque villosa. Stamina circiter 40, filamentis inæqualibus, glabris, sub anthera haud dilatatis, mox deciduis. Ovarium oblongo-cylindricum, papilloso-subvelutinum, apice profunde 3-dentatum dentibus glabris, placentis integris. Ovula inordinatim in placentis 3-serialia, in placenta quaque circiter 18-20. Capsula longiuscule pedicellata, suberecta, longa, trigono-subcylindrica, apice tridentata, infra dentes vix angustata, brevissime puberula, placentis integris. Semina minuta (immatura tantum nota), tuberculis minutissimis undique obsita."

Mar. merid.—In montibus ad austro-occidentem urbis Maroc, prope Keïra, circiter ad 900 m. detexit cl. Balansa. Legimus in collibus aridis supra Imintenout ad 1100 m. ! et in monte supra Seksaoua circ. 1500 m. !

A proxima *R. villosa*, Coss., differt, ut nobis docet cel. auctor, caule foliisque puberulis, non longe pubescenti-villosis, sepalis citius deciduis, petalorum superiorum lobis lateralibus haud cristæformibus, capsulis cylindræco-trigonis, nec oblongis infra dentes vix angustatis.

In spec. nostris capsula infra dentes evidenter angustata, dentes divergentes multo longiores quam in *R. villosa*, indumentum totius plantæ brevissime hirtulo-scabrum.

RESEDA LUTEOLA (L., Müll. Arg. in DC. Pr. xvi. sect. 2, 583).

Mar. sept.—Tanger (*Salzm. Exsicc.*) ! (*Webb Herb.*) ! Formam latifoliam legimus inter Tetuan et Ceuta !

— —, var. AUSTRALIS (Müll. Arg. l. c.).

Mar. merid.—In monte supra Seksaoua circa 1400 m. !

— —, var. CRISPATA, = *R. crispata* (Link, Enum. Pl. H. Ber. 1822, p. 8).

Mar. sept.—In regione collina inter Tanger et Tetuan !

Species per Europam mediam et regionem totam Mediterraneam (inclusa Maearonesia) late diffusa est. De distributione varietatum ab auctoribus non rite definitis fusius disserere inutile duco.

CISTINÆ (*Juss. Gen.* 249 ; *Spach, Ann. Sc. Nat.*
ser. 2, vi. 357).

CISTUS (*L. Gen.* 673 ; *Willk. Ic. et Desc. Pl.* tab. 75 ad 109).

C. ALBIDUS (*L., DC. Pr.* i. 264 ; *Willk. l. c.* tab. 77).

Mar. sept.—In monte Beni Hosmar supra Tetuan ! In rupibus prope Tetuan (1851) !

Ar. Geog. Regio Mediterranea occidentalis. Deest in Macaronesia. Hic et sequens *C. crispus* desunt in Græcia. Conf. *Boiss. Fl. Or.* i. 437.

C. CRISPUS (*L., DC. Pr.* i. 264 ; *Willk. l. c.* tab. 78).

Mar. sept.—Prope Tanger in monte Djebel Kebir ! et alibi visus ! (*Salzm. Exsicc*) !

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

Ludit foliis breviter pubescentibus, vel (præsertim junioribus) longius villosis.

C. POLYMORPHUS (*Willk. l. c.* tab. 79–82), = *C. vulgaris* (*Spach, l. c.* pro parte). Forma genuina = *C. villosus* (*L., Sp. Pl.* 736 ? et *Lam. Dict.* ii. 12), cui ut synonym. verisimiliter conjungendi sunt *C. pilosus*, *L.*, et *C. incanus*, *L.*, et *C. eriocephalus* (*Viv. Cors.* p. 8).

Mar. occid.—Djebel Hadid haud procul Mogador (*Rein et Fritsch, Spec. in hb. Kew.*) !

Mar. merid.—In collibus apricis sat frequens, et in regione inferiore Atlantis Majoris. Tasseremout ! Distr. Reraya ! In convalle Aït Mesan circ. 1300 m. ! In rupibus pr. Seksaoua !

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia) ; in peninsula Iberica minus frequens.

— —, var. CORSICUS (*Willk. l. c.* tab. 82).

Mar. merid.—Ad radices Atlantis Majoris prope Ourika legit G. Maw !

— —, var. CRETICUS (*Boiss. Fl. Or.* i. 437), = *C. creticus* (*L., DC. Pr.* i. 264).

Mar. occid.—In monte Djebel Hadid haud procul Mogador !

Ar. Geog. Regio Mediterranea præsertim orientalis. Hæc. var. deest, ut videtur, in peninsula Iberica.

Nomen Spachianum (*C. vulgaris*) antiquius foret, sed cl. auctor cum formis plurimis *C. polymorphi* etiam *C. albidum* et *C. crispum* in unam speciem comprehendit. Hæc novissimæ mihi videntur species probe distinctæ nec transitum unquam observavi. Nomen Willkommianum nobis ideo præferendum. Pace cel. Boissier *C. villosus*, *L.*, est nomen nimis incertum, et ab ipso Linnæo pro parte minima catervæ formarum affinium sine nota diagnostica idonea propositum, et speciem collectivam indicare nequit.

Formam in Djebel Hadid lectam non sine dubio sub. var. *cretico*, Boiss.,

enumeravi. Habet folia minora magis rotundata, valde rugosa, et nisi in calyce desunt pili longiores.

CISTUS MONSPELIENSIS (L., DC. Pr. i. 265).

Mar. sept.—Tanger (*Schsb. spec.*)! (*Webb Herb.*)! In regione collina inter Tanger et Tetuan! Inter Tetuan et Centa!

Mar. merid.—Ad radices Atlantis Majoris prope Tasseremout!

Ar. Geog. Regio Mediterranea præsertim occidentalis (inclusa Macaronesia). Nisi in Græcia et in insulis Cypro et Creta deest in Oriente.

C. SALVIÆFOLIUS (L., DC. Pr. i. 265).

Mar. sept.—Sat frequens in ditione Tingitana. Prope Tanger! Prope Tetuan! “In fruticetis totius imperii” (*Schsb. in herb.*) Schousboe tamen non nisi ditionem Tingitanam et provinciæ meridionalium partem minimam cognovit.

Mar. occid.—Mogador (*Lowe Cat.*). Prope Aïn el Hadjar haud procul Mogador!

Ar. Geog. Regio Mediterranea tota (exclusa Macaronesia).

— —, var. β , *MACROCALYX* (*Willk. Ic. et Descr. Pl. ii. 38. tab. 92*).

Mar. occid.—In monte Djebel Hadid!

Mar. merid.—In regione inferiore Atlantis Majoris in convalle Aït Mesau, usque 1200 m.!

C. POPULIFOLIUS (L., DC. Pr. i. 266).

Mar. sept.—In monte Djebel Kebir prope Tanger! (*Salzm. Exsicc.*)! (*Schsb. spec.*)!

Ar. Geog. Peninsula Iberica, adjecta Gallia Narbonensi finitima, et ditione Tingitana.

C. LADANIFERUS (L., DC. Pr. i. 266).

Mar. sept.—In monte Djebel Kebir prope Tanger! (*Salzm. Exsicc.*)!

Var. vel lusum petalis immaenlatis.

Ar. Geog. Peninsula Iberica, cum ditione Tingitana et Gallia Mediterranea finitima.

HELIANthemum (*Pers. Syn. ii. 75.*)

H. LIBANOTIS (L., sub *Cisto*; *Willd. Enum. 570*; DC. Pr. i. 267, non *Cistus Libanotis*, *Lam., nec Dsf.*), = *Halimium rosmarinifolium* (*Spach, Ann. Sc. Nat. ser. 2, vi.*).

Mar. sept.—Prope Cap Spartel!

Ar. Geog. Peninsula Iberica et ditio Tingitana. Forsan etiam in Sicilia.

Olim inter auctores sententiæ diversæ de vero *C. Libanotide*, L.; hodie tamen omnibus notum est hoc esse cum stirpe Linnaëana gemina identicum. Nomen triviale igitur nullomodo mutandum, et nomen *rosmarinifolium* (hodie pro sp. Americana Purshiana vindicatum)

verisimiliter ad *Cistum fastigiatum* (Guss. Pr. Fl. Sic. ii. 13) meliore jure spectat. An stirps Gussoniana a nostro *II. Libanotide* probe diversa sit mihi adhuc incertum est.

HELIANTHEMUM UMBELLATUM (*Mill. Dict.* No. 5 ; *DC. Pr.* i. 267),
var. *VERTICILLATUM*, = *Cistus verticillatus*, *Brot.*

Mar. sept.—In collibus saxosis prope Tetuan !

Ar. Geog. Gallia media et meridionalis. Græcia. Peninsula Iberica. Marocco Tingitanus. Legi formam arete affinem in Libano ; conf. *Boiss. Fl. Or.* i. 439.

Specimina Gallica plurima quæ vidi inflorescentiam terminalem nec verticillatam ostendunt.

II. HALIMIFOLIUM (*L.*, sub *Cisto* ; *Willd. Enum.* 569 ; *DC. Pr.* i. 268),
= *Halimium lepidotum* (*Spach, l. c.*).

Mar. sept.—Tanger (*Webb Herb.*) ! In saxosis prope Tetuan !

Mar. occid.—In parte superiore montis Djebel Hadid ! In sylva *Callitris quadrivalvis* et alibi prope Mogador, et inde versus Aïn el Hadjar !

Ar. Geog. Europa Mediterranea occidentalis, ex Italia usque Lusitaniam. Africa borealis. Deest in Macaronesia.

— —, var. *SEPALIS STELLATO-TOMENTOSIS*, = *II. multiflorum* (*Salzm. Exsicc.* ; *Willk. Ic. et Descr.* ii. 67, tab. 108).

Mar. sept.—Ex fruticetis et aridis regionis Tingitanæ (*Schsb. spec.*) benevole comm. cel. Cosson. (*Salzm. Exsicc.* !) In monte Djebel Kebir ! (*Webb Herb.*) !

— —, var. *LASIO-CALYGINUM*, = *II. lasio-calycinum* (*Boiss. et Reut., Diagn. Pl. Or. ser. 2, i. 50*), = *II. hirsutissimum* (*Willk. l. c. tab. 106*).

Mar. sept.—In monte Djebel Kebir prope Tanger ! (*Boiss. et Reut. l. c.*).

II. multiflorum et *H. lasiocalycinum* certo certius ad varietates *H. halimifolii* reducenda, notis nullis stabilibus distinguendas. *H. multiflorum* sæpe, sed non semper, foliis minoribus præditum, gaudet sepalis ex pilis longiuscule stellatim ramosis dense obteectis. Folia in hoc uninervia, in *H. halimifolio* obsolete trinervia dicuntur, sed discrimen omnino fallax et minime constans. In hac specie, ut fit in *Cistis* plurimis, indumentum variabile. In *H. lasiocalycino* folia et sepala pilis longis subvillosis plus minusve vestita, et indumentum stellatum fit rarius. Specimina nostra fere intermedia sunt sed propiora *H. lasiocalycino*. Nomen hujus formæ, vix pro varietate enumerandæ, el. Willk. perperam mutavit.

II. TUBERARIA (*L.*, sub *Cisto* ; *Mill. Dict.* No. 10 ; *DC. Pr.* i. 270).

Mar. sept.—Tanger (*Salzm. Exsicc.*) ! (*Webb Herb.*) ! Djebel Kebir prope Tanger ! Cap Spartel ! Prope Tetuan !

Ar. Geog. Per regionem Mediterraneam occidentalem ex Dalmatia usque Macaronesiam diffusum. In Græcia verisimiliter errore indicatum ex *Boiss. Fl. Or.* i. 440.

HELIANTHEMUM GUTTATUM (L., sub *Cisto*; *Mill. Dict.* No. 18; *DC. Pr.* i. 270).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Djebel Kebir prope Tanger! Cap. Spartel!

Mar. occid.—Casa Blanca!

Ar. Geog. Regio Mediterranea a Syria usque Macaronesiam. Gallia occidentalis. Forma arcte affinis in Anglia.

— —, var. *MACROSEPALUM* (*Coss. Pl. Crit.* 29), = *II. macrosepalum* (*Salzm. Exsicc.*), = *Tuberaria macrosepala* (*Willk., l. c.* tab. 117).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)! Spec. ex herb. Schousboe (sub *Cisto glanduloso*, Schsb.) “in fruticetis pr. Tanger,” lectum benevole comm. cel. Cosson. Djebel Kebir! (*Rent.*). Tetuan (1851)!

Legi etiam prope San Roque haud procul Gibraltar!

— —, var. *INCONSPICUUM*, = *II. inconspicuum* (*Thib. in Pers. Syn.* ii. 77; *DC. Pr.* i. 271).

Mar. sept.—Prope Tanger legit *J. D. H.* spec. Hispanicis omnino conforme!

Ar. Geog. Tam hæc quam var. *macrosepalum* in peninsula Iberica et in ditioe Tingitana obviæ sunt.

Hæc est inter typum et formam pro specie nova in Boiss. et Rent. Pug. p. 13, propositam, sub nom. *II. brevipes*, varietas intermedia. Characteres in hoc grege ex præsentia vel defectu stipularum, ex pilis longis plus minusve frequentibus, ex longitudine pedicellorum et magnitudine petalorum et sepalorum valde ludibundi et ideo fallaces.

II. NILOTICUM (L., sub *Cisto*; *Pers. Syn.* ii. 78; *DC. Pr.* i. 272).

Mar. sept.—Tanger (*Webb Herb.*)! Inter Tanger et Tetuan!

Mar. occid.—Casa Bianca! Prov. Haha prope Mogador!

Mar. merid.—Prov. Shedma! Mesfioua! Tasseremout! Milhain!

Ar. Geog. Regio Mediterranea tota (inclusa Macaronesia).

— —, var. *MICROCARPUM* (*Coss. in Bourg. Pl. Hisp.*; *Boiss. Fl. Or.* i. 441).

Mar. merid.—In saxosis aridis regionis inferioris. Prov. Shedma prope Aïn Oumast! Mskala! et alibi.

Ar. Geog. Varietas per reg. omnem Mediterraneam diffusa.

— —, var. ? *PUMILUM*.

Mar. merid.—In reg. subalpina Atlantis Majoris supra Arround legi exemplaria duo 2-3-pollicaria deflorata, foliis superioribus et sepalis tomentoso-hirtis, racemo contracto, capsula fere typica sed multo minore, seminibus angularibus omnino lævibus. Verisimiliter est hujus speciei polymorphæ forma nana.

Utrum *Cistus ledifolius*, L., Sp. 742, foret synonymon *C. nilotici*, sive melius ad var. *microcarpum* referri debet non satis eminet. Nomen speciei ab auctoribus posterioribus confusum hodie negligendum.

Helianthemum papillare (*Boiss. Voy. Esp.* ii. 63, tab. 14).

Mar. merid.—In saxosis aridis supra Seksaoua !

Ar. Geog. Peninsula Iberica ; Africa boreali-occidentalis.

II. ÆGYPTIACUM (*L.*, sub *Cisto* ; *Mill. Dict.* No. 23 ; *DC. Pr.* i. 273).

Mar. occid.—Legi prope Casa Blanca specimen exsiccatum sed satis certum !

Ar. Geog. Per regionem Mediterraneam calidiorem (exclusa Macaronesia) diffusa. Deest in Græcia, Dalmatia et in Gallia Mediterranea.

II. LIPPII (*L.*, sub *Cisto* ; *Pers.* ii. 78 ; *DC. Pr.* i. 273). Var *a* (*Boiss.*

Fl. Or. i. 443), = *Cistus ellipticus* (*Desf. Fl. Atl.* i. 418, tab. 107), = II. *rosmarinifolium*, *Psl. Symb.* tab. 21.

Mar. occid.—In monte Djebel Hadid specimen unicum legi ! Ex spec. in Oasi Akka lecto (*Coll. Indig. Mardochæus*) frustulum benevole comm. Cosson.

In herb. Kew. conservatur specimen a ell. Rein et Fritsch in Marocco (verisimiliter in ditione Mogadorensi) lectum ; a nostro parum differt floribus et capsulis aliquid majoribus et sepalis longius villosis.

Ar. Geog. Oriens. Africa borealis. Sicilia.

H. CANARIENSE (*Jacq. Ic. Rar.* i. tab. 97, sub *Cisto* ; *Pers. Syn.* ii. 78, *DC. Pr.* i. 274).

Mar. occid.—Legimus prope Mogador ! et Aïn el Hadjar ! et in monte Djebel Hadid ! et iterum legi in litore Atlantico ab urbi Saffi meridiem versus !

Ar. Geog. Species propria insularum Canariensium et terræ continentalis proximæ.

Nisi herba densius ineana spec. nostra Canariensibus omnino conformia.

H. RUBELLUM (*Psl. Del. Prag.* 25 ; *Boiss. Voy. Esp.*), = II. *rotundifolium* (*Dun. in DC. Pr.* i. 277), = *Cistus nummularius* (*Cav. Ic.* tab. 142 ; *Desf. Fl. Atl.* i. 423, non *L.*).

Mar. merid.—In regione media Atlantis Majoris. In convalle Amsmiz 1200–1600 m. !

Ar. Geog. Hispania ; Africa boreali-occidentalis ; Sicilia, et forma arcte affinis (*H. Hymettium*, *Boiss.* et *Heldr.*) in Græcia.

Specimina nostra gaudent foliis oblongo-ellipticis acutiusculis, et sistunt transitum in *H. Hymettium*, quod nobis mera varietas videtur.

II. LAVANDULÆFOLIUM (*Lam. Dict.* ii. 25, sub *Cisto* ; *DC. Fl. Fr.* iv. 820, et *Prod.* i. 278).

Mar. occid.—Ab urbe Mogador septentrionem versus ! In monte Djebel Hadid ! Inter Mogador et Shedma (*G. Maw spec.*) ! Agadir (*Coll. Indig.* 1871) !

Ar. Geog. Peninsula Iberica. Gallia Mediterranea et Corsica. Oriens. Africa boreali-occidentalis. Deest in Italia, Dalmatia, Africa boreali-orientali et in Macaronesia. Fere semper in regione litorali occurrit.

H. Broussonnetii, species Canariensis ex scheda in herb. Bouchet prope Mogador indicata, verisimiliter ad floram Maroccanam non pertinet. Broussonnet plantas patriæ diversæ in herbario sæpe miscuit, et hæc pulchra stirps in agro Mogadorensi nemini recentiorum obvia fuit.

HELIANTHEMUM GLAUCUM (*Cav. Ic.* iii. 31, tab. 261, non *Desf.*, sub *Cisto*; *Pers. Syn.* ii. 78; *DC. Pr.* i. 279).

Mar. merid.—In convalle Amsmiz circa 1400 m.!

Specimina nostra Hispaniis omnino conformia nisi statura minore.

Ludit petalis albis ut fit in spec. Bourgaenis, No. 1074, Pl. d'Espagne.

— —, var. *STÆCHADIFOLIUM*, = *H. stæchadifolium* (*Brot. Fl. Lus.* ii. 270, sub *Cisto*; *DC. Pr.* i. 279).

Mar. merid.—In rupibus supra Seksaoua!

Distinguitur foliis superioribus insigniter revolutis et ramis abortivis in axillis foliorum ad fasciculos foliorum juniorum rednetis, unde affinitas aliqua, ut jam innotuit Brotero *l. c.*, eum *H. lavandulæfolio*.

Specimina Maroccanis simillima legi olim prope Malaga. Petala vidi semper lutea.

Ar. Geog. H. glaucum, per regionem Mediterraneam occidentalem late diffusum, in Macaronesia deest. Est stirps valde ludibunda, ejus limites ægre definiendi.

H. VIRGATUM (*Desf. Fl. Atl.* i. 432, sub *Cisto*; *Pers. Syn.* ii. 79; *DC. Pr.* i. 282. Inelusis *H. racemosum* (*Lam. Diet.* ii. 25, sub *Cisto*; *Dun.* in *DC. Pr.* i. 281), et *H. pilosum* (*Pers. Syn.* ii. 79; *DC. Pr.* i. 282, an *L.*, sub *Cisto*).

Mar. merid.—Frequens ad radices Atlantis Majoris et in convallibus calidioribus. Tasseremout! Ourika! Reraya! Aït Mesan usque 1400 m.! Sektana! Amsmiz! Milhain! Ex Prov. Mtuga (*R. et Fr. spec. in herb. Kew.*)!

Notæ diagnosticæ quibus *H. racemosum* et *H. pilosum* ab *H. virgato* separantur sunt adeo instabiles et ludibundæ ut ne quidem varietates sistunt. Si verbis auctorum hæremus, calyx in *H. virgato* cano-pulverulentus pubescens, in *H. piloso* nervoso-striatus pilosiusculus, in *H. racemoso* nervoso-suleatus violaceo-rufescens. In ditione nostra calyces sæpius subglaberrimi, rarius tenuiter incano-pubescentes, vel pilosiusculi, folia angusta, plus minusve margine revoluta, in pagina superiore viridia vel subincana, subtus stellato-tomentosa. In speciminibus Hispanicis et Algeriensibus *H. pilosi* video calyces valde variabiles ut in spec. nostris. Petala ludunt alba vel pallide rosea.

— —, var. *STRICTUM*, = *H. strictum* (*Cav. Ic.* iii. 32, tab. 263, sub *Cisto*; *Pers. Syn.* ii. 79; *DC. Pr.* i. 281), = *H. virgatum*, var. *Maroccanum*, nob. *MSS.*

Mar. merid.—Valde frequens in saxosis aridis regionis inferioris et rarius in convallibus Atlantis Majoris. Legimus in Prov. Shedma

prope Aïn Oumast! in glareosis torrentis Ourika! prope Seksaoua! et in convalle Aït Mesan! In convalle Amsmiz legi varietatem seu lusum calyce piloso, sepalis exterioribus longioribus et acutioribus, stipulis pariter longioribus foliorum dimidiam longitudinem subæquantibus. Petala ex albo pallide rosea.

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

Affinitas inter *H. virgatum* et *H. apenninum* (quocum velint synonyma enumeranda sunt *H. pulverulentum*, DC., *H. polifolium*, DC., et plurimæ species ab auctoribus enumeratæ) tam areta est ut vix pro typis specificis probe distinctis haberentur.

FUMANA (*Spach, Ann. Sc. Nat.* ser. 2, vi. 359).

Genus *Fumana* staminibus exterioribus sterilibus moniliformibus, seminibus anatropis raphe præditis, facile recognoscendum, et præterea habitu proprio (in *F. glutinoso* minus eminente) gaudens, a celeb. Benth. et H. f. in Gen. Pl. 114, ad subgenus *Helianthemum* reductum est. Pace tantorum virorum vestigia auctorum recentiorum qui hoc (eodem jure quo *Callianthemum* et plurima genera subnaturalia) nomine proprio salutarunt secuti sumus.

F. GLUTINOSA (L., sub *Cisto*; *Pers. Syn.* ii. 79, et DC. *Pr.* i. 276, sub *Helianthemum*; *Boiss. Fl. Or.* i. 449), = *F. viscida* (*Spach, loc. cit.*).

Mar. sept.—Tanger (*Schsb. spec. sub C. Fumana*)! (*Webb Herb.*)! In rupibus prope Tetuan 1851, et iterum 1871!

Mar. occid.—Mogador (*Lowe*)! Ain el Hadjar! Djebel Hadid!

Mar. merid.—Sat frequens ad radices Atlantis Majoris et in convallibus calidioribus. Mesfioua! Ourika! Aït Mesan usque 1400 m.! Seksaoua! Amsmiz!

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia).

— —, var. *VIRIDIS*, = *Helianthemum viride* (*Ten. Pr. Fl. Neap.* p. 31).

Mar. sept.—In rupibus prope Cap. Spartel! Inter Tanger et Tetuan! In monte Beni Hosmar supra Tetuan!

Optimo consilio cl. Spach plurimas s. d. species nullis notis stabilibus dignoscendas in unam reduxit, sed minus bene nomen triviale Linæum antiquius et satis aptum mutavit. Verum est in var. *viridi* folia sæpius subglaberrima, laxiora et longiora quam in typo; sed pedicelli semper plus minusve glanduloso-viscidi, et formæ intermediæ non desunt. Specimina Schousbœana Tingitana quæ benevolentia eel. Cosson debeo inter typum et genuinum *H. viride*, Ten. transitum præbent.

FUMANA LÆVIPES (L., sub *Cisto*; *Pers. Syn.* ii. 76, et DC. *Pr.* i. 275, sub *Helianthemum*; *Spach, loc. cit.*).

Mar. sept.—Tetuan! (*Webb Herb.*)! In monte Beni Hosmar!

Mar. merid.—In rupibus supra Seksaona !

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

F. CALYCINA, *Clauson*, = *F. arbuscula* (*Ball, Journ. Bot.* 1873, 300) = *Fumana Spachii*, var. *calycina* (*Coss. Index Pl. Mar.* 14).

Frutex humilis, ramosissimus, intricatus, trunco demum diametro pollicari; ramuli recti, rigidi, foliacei; folia alterna, coriacea, infima ovalia, vix lineam longa, superiora ovali-oblonga circa 3 lineas attingentia, suprema brevissima, cum ramulis junioribus puberula; flores in quovis ramulo pauci (2–4); pedicelli subpollicares, demum recurvati; sepala exteriora (epicalyx) brevina, linearia, scabrida, interiora membranacea, carnea, ex costis 3 viridibus hirtulis et nervo 1 utrinque non costato lineata; petala (nondum expansa) calyci æquilonga, saturate flava; semina maxima, in quavis capsula sex, pallida, lævia, oculo armato minute granulata.

Mar. merid.—In regione inferiore Atlantis Majoris admodum rara. Exemplaria duo, jam deflorata, legimus in glaucosis ad torrentem Ourika (circ. 1000 m.)! et alterum, nondum florens, in convalle Aït Mesan (circ. 1300 m.)!

Semina illis *F. vulgaris* ter quaterve longiora et latiora. Planta Atlantica a nobis visa multo major est quam spec. quæ postea vidi in herb. Cosson, et mihi a *F. Spachii* Europæa omnino diversa visa est. Melius edoctus ex spec. intermediis Algeriensibus nuper scrutatis hodie hæc stirps insignis sicut subspecies *F. Spachii* enumeranda mihi videtur. Monente cel. Cosson *F. calycinam* legit Balansa in sylva *Calitris* haud procul Mogador. Nascitur etiam in Algeria.

VIOLARIÆ (*DC. Fl. Fr.* iv. 801).

VIOLA (*L. Gen.* 1007).

V. ODORATA (*L., DC. Pr.* i. 296).

Mar. sept.—Tanger (*Webb Herb.*)! Tetuan (*G. Maw*)!

Mar. merid.—In umbrosis irrigatis urbis Marocco!

Ar. Geog. Europa media. Regio Mediterranea tota (inclusa Macaronesia). Sibiria. Forma orientalis montana *V. suavis*, M. B.; a typo aliquid recedit, sed sicut varietas a cel. Boissier (*Fl. Or.* i. 458) enumeratur. Specimina nostra ex urbe Marocco carent flore et fructu; forsitan referunt varietatem *maderensis* (Lowe, *Man. Fl. Mad.*).

V. ARBORESCENS (*L., DC. Pr.* i. 299), = *V. suberosa* (*Desf. Fl. Atl.* ii. 313).

Mar. sept.—Tanger (*Schsb. spec.*)! In rupibus maritimis haud procul Cap. Spartel! In rupibus calcareis prope Tetuan! In monte Beni Hosmar supra Tetuan (exemplaria pulcherrima sesquipedalia et ultra)!

Mar. oeeid.—Seeus rivulum Oued Kseb prope Mogador! ubi eam jam legerat Balansa. In monte Djebel Hadid!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia); prædiligat rupes zonæ sublitoralis.

Ludit foliis latioribus angustioribusve, margine integerrimis vel denticulatis vel serratis, sed hæ notæ nimis inconstantes et veras varietates non sistunt. Confer DC. Prod. l. c.

VIOLA TEZENSIS, nob., = *V. modesta* (*Ball in Journ. Bot.* 1873, 300; non *Fzl. Ill. Pl. Taur.* p. 41).

Egrege *V. tricoloris*, L., et verisimiliter juxta *V. Heldreichianam*, Boiss. colloecanda, sed notis plurimis distincta. Folia inferiora ovata, superiora lanceolata in petiolum attenuata, profunde erenata vel erenato-dentata, præsertim margine et nervo medio hirtula. Stipulæ *V. Heldreichianæ*, oblongo-lineares, integræ vel basi externa lacinula auctæ, foliis breviores. Pedicelli glaberrimi, apicem versus bibracteolati. Sepala viridia, oblonga, margine ciliolata, cæterum glabra. Petala calyceem vix æquantia. Calcar brevissimum.

Mar. merid.—Exemplar unicum in monte Djebel Tezah alt. circiter 2400 m. detexi!

POLYGALÆ (*Juss. Ann. Mus.* xiv. 386).

POLYGALA (*L. Gen.* 851).

P. RUPESTRIS (*Pourr. Act. Toul.* iii. 325), = *P. saxatilis* (*Desf. Fl. Atl.* ii. 128, tab. 175; *DC. Pr.* i. 324).

Mar. oeeid.—Aïn el Hadjar!

Mar. merid.—In regione inferiore et media Atlantis Majoris, et rarius in planitie. In convalle Aït Mesan 1400–1800 m.! In convalle Amsmiz circ. 1500 m.! In monte Djebel Tezah circ. 1800 m.! Ex prov. Shedma specimen retulerunt ell. Rein et Fritsch!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

P. ROSEA (*Desf. Fl. Atl.* ii. 128, tab. 176; *DC. Pr.* i. 324), = *P. Preslii* (*Spreng. Syst. Veg.* v. 531) = *P. niæensis* (*Risso ex Koch, Syn.* i. 98).

Mar. sept.—Pr. Tanger in monte Djebel Kebir versus Cap. Spartel! In rupibus prope Tetuan! In monte Beni Hosmar! ubi eam jam legerat P. B. Webb.

Ar. Geog. Per regionem Mediterraneam, imprimis occidentalem, late diffusa. Deest in Macaronesia, sed insulas Azoricas incolit.

Icon Fontanesiana supra citata minus bona, sed de identitate stirpis vix dubitandum. Confer tamen Boissier, *Fl. Or.* i. 475.

P. MONSPELIACA (*L., DC. Pr.* i. 325).

Mar. sept.—In regione collina prope Fondak, inter Tanger et Tetuan!

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia), sed potius infrequens.

POLYGALA WEBBIANA (Coss. in Bull. Soc. Bot. Fr. xx. 240).

Descriptionem eel. auctoris præferre iuvat.

“Planta perennis, ut videtur a basi ramosa, ramis diffusis, teretiuseulis, inferne denudatis pulvinis foliorum delapsorum tuberculatis, superne dense foliosis, pilis arcuato-ascendentibus dense pubescentibus. Folia eodem modo ac rami sed parcius pubescentia, oblonga, plana, in petiolum contracta, acutiuscula vel inferiora obtusa haud mucronata. Flores majusculi, in axillis foliorum superiorum in racemos brevissimos cymiformes 2-3-flosos vel sæpissime abortu unifloros dispositi, bracteolis 3 membranaceis obovatis amplis deciduis ad basin pedicelli involnerum efforinantibus suffulti. Sepala omnia citius decidua, glabrescentia, 3 exteriora albida membranacea vix herbacea quorum superius majus fornicatum subsaccatum et duo inferiora plauiuscula ovata, 2 interiora petaloidea alba vel pallide purpurascencia corollam subæquantia obovata obtusa inferne angustata. Petala 3, inter se et cum vagina staminali ad tertiam partem superiorem in tubum superne fissum coalita, duorum lateraliu limbo inæquilatero concavo-subcarinato pallide purpurascence apice obtuso, inferioris limbo æquilatero concavo-carinato genitalia amplectente apice truncato in carina procul ab apice crista purpurascence in lobos nonnullos oblongos latiusculos haud incrassatos divisa donato superne pallide lutescente. Ovarium superne basi glandula stipatum. Capsula (immatra tantum visa) membranaceo-carnosula, a latere compressa, obovata, apice marginata.”

Mar. sept.—Florentem legimus in monte Beni Hosmar supra Tetnan ubi eam olim legit P. B. Webb!; et iterum spec. 2 detexi in rupibus juxta urbem Tetuan! In monte Djebel Moussa supra fretum Herculis eminente legit el. Blackmore.

Monente eel. Cosson:—“*P. Webbiana*, floribus magnis, in racemos axillares 1-3-flosos dispositis, sepalis deciduis, superiore fornicato subsaccato, crista carinali petali inferioris parce et grosse lobata, ad sectionem *Chamæbuxus* pertinet.—A *P. Munbyana* (Boiss. et Reut. Diagn. Pl. Or. ser. 2, v. 50) differt ramis teretiuseulis, non acute angulatis, dense pubescentibus, non glabrescentibus, foliis haud mucronatis, petalis lateralibus haud truncato-obtusis.—A *P. Chamæbuxo*, L., cui valde affinis, distincta indumento ramorum densiore, foliis minus coriaceis haud mucronatis, petalis lateralibus haud truncato-obtusis, lobis cristæ tennibus plauiusculis, non incrassatis subeoreulatis.”

Facie simillima est *P. Chamæbuxo*, var. *rhodoptera*, stirpi Alpij Vene-tarum solenni, sed in hac, præter notas supra indicatas. sepala interiora saturate purpureo-rosea, petala lateralia apice lutescentia; dum in *P. Webbiana*, sepala interiora in viva pallide purpurascencia, petala lateralia læte purpureo-rosea. Folia in *P. Chamæbuxo* longiora, angustiora, nec sensim in petiolum attenuata.

P. Chamæbuxus est Alpij Europæ (latiore sensu) incola, unde in eri-

cetis Baviaræ et per Pannoniam usque Transsylvaniam, necnon per Apenninum Liguriæ in Alpes Apuanas se protendit. Deest in peninsula Iberica et in Pyrenæis et in Europa austro-orientali. Localitas in ducatu Lucemburgensi verisimiliter spuria.

POLYGALA BALANSÆ (Coss. in *Bal. Pl. Maroc.* 1867, et in *Bull. Soc. Bot. Fr.* xx. 240).

Sequitur descriptio Cossoniana:—“*Frutex* erectus, multicaulis, dumosus, divaricatum ramosus, ramis rigidis *spinescentibus*, pulvinis foliorum delapsorum tuberculatus, sub lente punctato-puberulus. *Folia parva*, oblongo-linearia, concavo, cito decidua. *Racemi* breves, 3–4-flori, sæpius abortu 1–2-flori. *Sepala* omnia citius decidua, glabrescentia, 3 exteriora subherbacea margine purpurea, quorum superius fornicatum subsaccatum et duo inferiora planiuscula ovata; 2 interiora petaloidea purpurea corollam subæquantia ovata, inferne angustata, superne margine involuto apiculata. *Petala* 3, inter se et cum vagina staminali ad quartam partem superiorem in tubum superne fissum coalita, duorum lateralium limbo inæquilatero concavo-carinato pallide purpureo; inferioris limbo latiore æquilatero concavo-carinato genitalia amplectente, apice emarginato, in carina infra apicem crista indivisa irregulariter plieata donato, superne luteo. Ovarium superne basi glandula stipatum. Capsula membranaceo-carnosula, a latere compressa, suborbiculata, apice vix retusa, angusta, marginata, venis concentricis parce ramosis donata. Semen nigrescens, pilosum, caruncula triloba lobis lateralibus seminis quartam partem inferiorem attinentibus.”

Mar. merid.—Species per clivum septentrionalem Atlantis Majoris hucusque exploratum in saxosis aridis et rupestribus usque 1800 m. s. m. frequentissima. Primus eam detexit el. Balansa anno 1867. Legimus pr. Tasseremont! Ourika! Seksaoua! In convallibus Aït Mesan! Amsmiz! et alibi vidimus.

Stirps insignis sæpe 6–8-pedalis. Ramis teretibus apice spinescentibus *Spartium scoparium* interdum refert.

Individua minora habitu et facie stirpem Sinaiticam *P. spinescens*, Decaisne, miro modo simulant etsi characteres valde diversi. Species nostra structura floris et fructus ad sectionem *Chamæbuxus* pertinet.

FRANKENIACEÆ (*St.-Hil. Mém. Plac. Cent.* p. 39).

FRANKENIA (*L. Gen.* 445).

F. PULVERULENTA (*L., DC. Pr.* i. 349).

Mar. occid.—Casa Blauca! Mogador!

Mar. merid.—In saxosis aridis Seksaoua!

Ar. Geog. Regio Mediterranea omnis inclusa Macaronesia. Senegalia. Songaria. Olim in Anglia meridionali.

FRANKENIA HIRSUTA (L., *Boiss. Fl. Or.* i. 780; *Schousb. Gew. Marok.* 168). Includis *F. lævis* (L., DC. *Pr.* i. 349; *Schousb. Gew. Marok.* 167) et *F. intermedia* (DC. eodem loco).

Mar. sept.—Tanger specimen (cui scheda *F. lævis*, L.) ex herb. Schousboe benevole communicavit cel. Cosson! et eandem formam, foliis basi ciliolatis cæterum glaberrimis, legimus prope Cap. Spartel! et prope Tetuan legit Webb! Formam foliis tenuiter tomentellis legi prope Tetuan anno 1851!

Mar. occid.—Casa Blanca in litore Atlantico, formam foliis subtoментellis! et eandem legimus prope Mazagan! Mogador (*Schsb. l. c.*).

Optimo consilio cel. Boissier, *l. c.*, *Frankenias* fere omnes perennes regionis Mediterraneæ sub *F. hirsuta*, L., reduxit. Pace tanti viri varietates α et β sola præsentia vel absentia tomenti tenuissimi innixas separare nequeo, dum in eodem loco specimina utriusque formæ semel crescunt. Formas sequentes a typo remotiores pro speciebus rite distinctis habuerim, sed characteres nimis instabiles sunt et formæ intermediæ non desunt.

Ar. Geog. *F. hirsuta*, L., sensu latiore est species late diffusa per oras Atlanticas Europæ et Africæ, ex Britannia ad Cap. Bon. Spei, et per totam regionem Mediterraneam. In Europa media deest, sed terras continentales Asiæ temperatæ non fugit.

— —, var. VELUTINA, nob. = *F. velutina* (DC. *Pr.* i. 350).

Stirps insignis differt ab affinibus statura majore, floribus majoribus pulchre roseis, herba tota velutino-canesciente; sed specimina intermedia in litore Atlantico lecta hanc cum typo conjungunt.

Mar. occid.—Legimus in rupibus insulæ Mogador! verisimiliter eodem loco ubi eam primus detexit Broussonnet et postea legerunt Schousboe et Balansa. Specimina fere intermedia legimus inter hanc et *F. intermedium*, DC., prope Saffi! et pr. Casa Blanca!

— —, var. δ , REVOLUTA, *Boiss. l. c.*, = *F. revoluta* (*Forsk. Fl. Æg.-arab.* p. 75; DC. *Pr.* i. 350), = *F. corymbosa* (*Desf. Fl. Atl.* i. 315, tab. 93; DC. *Pr.* i. 350).

Mar. merid.—In planitie aridissima inter Aïn Oumast et Sheshaoua! et iterum prope Misra ben Kara!

Specimina nostra ludibunda. Nonnulla omnino similia spec. ex Ajanque prov. Murcia (*Bourgeau, Pl. d'Esp.* no. 2315), quæ *F. Webbii* (*Boiss. et Reut. Pug.* 16), et aliis a me olim lectis pr. Almeria et prope Chiclana. Alia propiora sunt *F. pallidæ* (*Boiss. et Reut. Diagn. Pl. Or. ser.* 2, i. 61), quæ mihi hucusque levis nec vera varietas videtur.

F. thymifolia (*Desf. Fl. Atl.* i. 316; DC. *Pr.* i. 350) a Schousboe indicatur in arenosis maritimis prope Mogador (*Schousb. Gew. Marok.* 167); nos non vidimus et floræ nostræ videtur dubia civis. Hæc forsitan velut subspecies enumeranda est. Facies sat singularis et

charaeteres minus variabiles quam in affnibus. Hujus var. levis mihi videtur *F. Reuteri* (Boiss. Diagn. Pl. Or. ser. 2, i. 60).

CARYOPHYLLÆ (*Juss. Gen.* 299).

VELEZIA (*L. Gen.* 447).

V. RIGIDA (*L., DC. Pr.* i. 387).

Mar. oecid.—Agadir (*Coll. Indig.* 1871)!

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia).

DIANTHUS (*L. Gen.* 565).

D. ATTENUATUS (*Sm. Trans. Linn. Soc.* ii. 301; *DC. Pr.* i. 358), = *D. lusitanus* (*Brot. Fl. Lusit.* ii. 177).

Mar. merid.—In regione media et superiore Atlantis Majoris. In convalle Aït Mesan usque in jugo Tagherot a 1800 m. ad 3000 m.! Specimina ex montibus Djebel Ouensa et Djebel Afougneur (*Coll. Indig. Ibrahim*) benevole comm. eel. Cosson sub "*D. lusitanicus*, Brot."

Ar. Geog. Peninsula Iberica. Pyrenæi Orientales. Atlas Marocceanus. *D. lusitanus*, a Brotero *loc. cit.* bene descriptus, est sæpius forma multicaulis, caulibus pluries fureatis, petalis profunde inciso-serratis basi pilosiusculis, dum stirps Pyrenaica (*D. attenuatus* typicus) habet ramos pauciores breviores 1-2-floros, petala erenata seu leviter dentata imberbia. Sed hæ notæ a solo et temperie pendent. Vidi ex Pyrenæis orientalibus specimina nullomodo a *D. lusitano* distinguenda. Nostra specimina ex convalle Aït Mesan potius ad formam vulgarem Pyrenaicam; contra quæ debeo humanitati eel. Cosson ad formam Lusitanicam propius accedunt.

D. VIRGINEUS (*L. Sp. Pl.* 590; *Gr. et God. Fl. Fr.* i. 238, non *DC. Pr.* i. 361).

Mar. sept.—? Specimen nondum florens legi in rupibus pr. Tetuan verisimiliter hue referendum!

Mar. merid.—In regione media Atlantis Majoris. In convalle Aït Mesan a 1600 m. ad 2000 m.! In monte Djebel Tezali spec. legit J. D. H.!

Ar. Geog. Europa media. Regio Mediterranea occidentalis (exclusa Macaronesia).

D. siculus (*Psl. Del. Prag.* p. 59; *Guss. Fl. Sic.* i. 479).

Mar. oecid.—Inter Mogador et Aïn el Hadjar! In monte Djebel Hadid! (forma calycibus coloratis). Inter Mogador et Shedma, *G. Maw, sp.*! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In rupestribus supra Seksaoua! Ourika (*R. et Fr. spec. in herb. Kew.*)!

Ar. Geog. Regio Mediterranea occidentalis, imprimis in calidioribus.

Genus omne deest in Macaronesia. Specimina classica Panormitana et

quæ legi in Hispania sunt omnia valde variabilia quoad calyceem et squamas externas plus minusve elongatas, chartaceas vel herbaceas, et specimina ex ditone Mogadorensi non minus ludibunda sunt. Quæ legimus pr. Seksaoua propiora *D. virgineo*. An *D. sculus* pro varietate vel subspecie meliore jure enumerandus sit judicet monographus in hoc genere valde desideratus.

DIANTHUS GADITANUS (*Boiss. Diagn. Pl. Or. ser. 2, i. 67*), = *D. Horne-
manni* (*Salzm. Exsic., non Ser. in DC. Pr. i. 362*), = *D. Schousbœi*
(*Coss. MSS.*).

Mar. sept.—Tanger (*Schsb. herb. sub "D. glaucus"*), *Coss. spec.!*
(*Salzm. Exsic. in herb. Kew.*)! In monte Beni Hosmar supra Tetuan
nondum florentia, sed verisimiliter huc referenda, specim. 2 legimus!

Ar. Geog. Hispania Gaditana et Marocco Tingitanus.

Affinis ut videtur *D. Liboschitziano*, Ser.

TUNICA (*Scop. Fl. Carn. i. 300*).

T. COMPRESSA (*Desf. Fl. Atl. i. 343. tab. 97, et DC. Pr. i. 354, sub Gyp-
sophila; Fisch. et Mey. Ind. Sem.*).

Mar. merid.—Sat frequens in planitie et in calidioribus reg. inferioris
Atlantis Majoris. Aïn Oumast! Mskala! Seksaoua! In convalle
Amsmiz, circ. 1400 m.!

Ar. Geog. Africa boreali-occidentalis.

T. PROLIFERA (*L., et DC. Pr. i. 355, sub Diantho; Scop. Fl. Carn. i.
300*).

Mar. merid.—In regione media Atlantis Majoris. In convalle Aït Mesan
circ. 1500 m.! In monte Djebel Tezah usque 2100 m.!

Ar. Geog. Europa media. Regio Mediterranea (inclusa Macaronesia),
sed rara in calidioribus. Orientem versus vix ultra Byzantium se
protendit.

SAPONARIA (*L. Gen. 584*).

S. VACCARIA (*L., DC. Pr. i. 365*).

Mar. merid.—In distr. Mesfioua hand procul urbem Marocco spec. uni-
cum legit J. D. H.!

Ar. Geog. Europa media. Regio Mediterranea omnis (inclusa Macaro-
nesia). In regionibus alienis, e. g. Sibiria Altaica et India orientali,
verisimiliter eum seminibus introducta.

SILENE (*L. Gen. 772; Rohrbach, Monog. Sil.*).

S. INFLATA (*Sm. Fl. Brit. ii. 292; DC. Pr. i. 368*), = *S. cucubalus*
(*Wib. Prim. Werth, 241; Rohrb. Sil. 84*).

Mar. sept.—Tanger! In monte Beni Hosmar pr. Tetuan! Prope Tetuan
(1851)!

Mar. occid.—Casa Blanca! Agadir (*Coll. Indig. 1871*)!

Mar. merid.—Milhain! formam foliis angustioribus legimus in convalle Aït Mesan, alt. 1200–1600 m.!

Ar. Geog. Europa media et borealis. Regio Mediterranea omnis (inclusa Macaronesia). Asia temperata. India orientalis. In America boreali introducta.

— —, var. *RUBRIFLORA*, nob. Petalis brevissimis calycem vix excedentibus saturate rubellis.

Mar. sept.—In sepibus prope Tanger!

Nomen triviale ab omnibus fere botanicis receptum cl. Rohrbach infanste mutavit. Si solo antiquitatis jure legamur, *S. vulgaris* erit. Nomen *Behen vulgaris* (Mœnch. Meth.) est cuim Wibeliano pluribus annis antiquius.

S. DISTICHA (Willd. Enum. 476; Rohrb. Sil. 96), = *S. parvifolia* (Otth in DC. Pr. i. 376, non Zea, in Poir. Dict. Suppl.).

Mar. sept.—Specimen ex herb. Schousboc in monte Djebel Kebir pr. Tanger lectum sub "*S. hirsuta*" Schsb. herb., benevole comm. Cosson.

Ar. Geog. Algeria. Ins. Balcares. Ditio Tingitana.

S. GALLICA (L., DC. Pr. i. 371; Rohrb. Sil. 96).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Cap. Spartel! Inter Tanger et Tctuan!

Mar. occid.—Casa Blanca!

— —, var. *MINOR*, magis pilosa, = *S. sardoa*, *Moris.*?

Mar. merid.—Mesfioua haud procul urbem Marocco!

Hæc est verisimiliter *S. insitanica* (L. Sp. Pl. i. 416).

Ar. Geog. Species verisimiliter Europæ mediæ et australis indigena, per totam fere orbem introducta et facta spontanea; ex cœli et soli diversitate formæ plurimæ enatæ pro speciebus distinctis ab auctoribus descriptæ.

S. CERASTOIDES (L., DC. Pr. i. 372; Rohrb. Sil. 38), var. *ANOMALA* (Ball, Journ. Bot. 1873, 301).

Habitu et seminum structura typo omnino conformis, sed insigniter differt calyce subconvenio, nervis in parte calycis superiore obscure anastomosantibus, et filamentis basi puberulis.

Mar. merid.—Prope Mzouda!

Forsan melius cognita pro specie diversa habenda.

Ar. Geog. Species per regionem Mediterraneam (exclusa Macaronesia) dispersa.

S. Giraldii (Guss. Fl. Inar. 36), quæ gaudet seminibus et calyce *S. gallicæ* et filamentis glabris *S. cerastoidis*, mihi videtur varietas prioris, nec cum *S. cerastoide* adsocianda.

S. TRIDENTATA (Desf. Fl. Atl. i. 349; DC. Pr. i. 371; Rohrb. Sil. 99).

Mar. merid.—Amsmiz ad radices Atlantis Majoris!

Ar. Geog. Africa boreali-occidentalis. Peninsula Iberica. Macaronesia. Priori similis sed ut videtur bene distincta species, floribus subsessilibus, calyce ex nervis eminentibus eximie costato, fructifero ore contracto dentibus longis divergentibus, capsula subsessili globosa longe rostrata, nec, ut in *S. cerastoidi*, ovoidea carpophoro brevi falcata et rostro brevi prædita.

SILENE NOCTURNA (L., DC. *Pr.* i. 372; *Rohrb. Sil.* 100).

Mar. merid.—Hinc inde in regione inferiore et in convallibus calidioribus Atlantis Majoris. In convalle Aït Mesan circ. 1300 m.! Prope Mzouda! Ourika (*R. et Ft. spec. in herb. Kew.*)! Seksaoua (*J. D. II.*)!

— —, β , PARVIFLORA (*Otth in DC. Pr. l. c.*), = var. micrantha (*Willk. Ic. et Desc.* i. 69, tab. 50), = var. γ , permixta (*Rohrb. l. c.*), = *S. permixta* (*Jord. Pug. Pl. Nov.* 32), = *S. mutabilis* (L. *Sp. Pl.* 596)?

Mar. merid.—Tasseremout!

Ar. Geog. Species per regionem Mediterraneam (inclusa Macaronesia) late diffusa. In America boreali advena.

Var. nostra e tvar. *brachypetala* (*Rohrb. l. c.*) verisimiliter per totam aream inveniendæ.

S. decipiens (*Ball in Journ. Bot.* 1873, 300).

Proxima *S. nocturnæ*, cui facie similis; differt tamen pedicellis parce glandulosis vix viscidulis, calyce fructifero ex nervis fusco-viridibus costato, dentibus erecto-patentibus capsulam superantibus, ejus diametrum æquantibus, petalis capsulam vix superantibus, lamina brevi atrorubente, seminibus minimis dorso vix canaliculatis.

Mar. occid.—Legimus in arenosis prope Casa Blanca!

Mar. merid.—Specimen incompletum sed cæteris plane conforme legi prope Amsuiz ad radices Atlantis, alt. circ. 1100 m.!

Stirps singularis calyce et seminibus a *S. nocturna* nimis diversa ut pro varietate haberetur.

S. obtusifolia (*Willd. Enum.* p. 473; *DC. Pr.* i. 374; *Rohrb. Sil.* 102), = *S. colorata* (*Schousb. ex Horn. H. Hafn.* i. 412; *DC. Pr.* i. 374, non *Poir.*).

Mar. sept.—In arenosis pr. Tanger! In arenosis maritimis inter Tetuan et Ceuta!

Ar. Geog. Peninsula Iberica. Macaronesia. Africa boreali-occidentalis. Habitat etiam in Ægypto inferiore (*ex Boiss. Fl. Or.* i. 593).

S. hirsuta (*Lag. Var. de Cienc.* 1805, 212; *Rohrb. Sil.* 102, non *Poir. Voy.*), = *S. hirsutissima* (*Otth in DC. Pr.* i. 372), var. *TUBERCULATA*, nob.

Mar. sept.—In monte Djebel Kebir pr. Tanger die 13 Jun. 1871, florentem et fructiferam legi! Ex eodem monte specimina seminibus

nondum maturis a b. Schousboc lecta (*S. pilosa*, Schsb. herb.) benevole communicavit cel. Cosson. Eandem varietatem, nisi fallor, sistit *S. hispida*, Salzm. Exsicc. Tingitana.

Mar. occid.—Specimina jam deflorata legi prope Casa Blanca (Apr. 22, 1871) seminibus spec. nostris Tingitanis omnino conformibus.

Hæc varietas speciminibus typicis *S. hirsutæ* quæ in Hispania legi et pluries habui omnino conformis est, nisi seminibus dorso et præsertim margine tuberculatis, facie paululum depressis nullomodo excavatis. Speciminibus plurimis Hispanieis accurate scrutatis transitum nullum detexi; demum specimen *S. hirsutæ* prope Malaga a Rambur lectum semina nostris valde accedentia mihi præbuit. Semina in genere *Silene* notas optimas diagnosticas inter species affines præbuerunt, sed *vis variativa* naturæ, vitæ socia inseparabilis, etiam in seminum structura se palam facit.

Utrum *S. hirsuta*, Lag., a *S. vespertina* (Retz. Obs. Bot. iii. 31; Rohrb. Sil. 95, non auctorum plurimorum) probe distincta sit est mihi res valde dubia. Pace cl. Rohrbach inter semina utriusque speciei discrimen vix ullum invenies, nec notæ differentiales ex inflorescentia allatæ satis constantes. Calyces dentes in *S. vespertina* supra capsulam contracti forsitan ad speciem dignoscendam satis valent. *S. vespertina*, Retz, ab auctoribus fere omnibus cum specie omnino diversa (*S. colorata*, Poir.) confusa, a Rohrbach *loc. cit.* suo loco restituta fuit. Hujus synonyma sunt *S. hirsuta* (Poir. Voy. ii. 163), *S. hispida* (Desf. Fl. Atl. i. 348), *S. bellidifolia* (Jacq. Hort. Vind., non Pourr.), et *S. sabuletorum*, Dub., non Lk.

S. hirsuta genuina est peninsulæ Ibericæ civis, formæ affines in Italia calidiore et Græcia et in Africa boreali obviæ sunt.

SILENE PSAMMITIS (Link in Spreng. Nov. Prov. p. 39), = *S. villosa* (Boiss. Voy. Esp. ii. 90, non aliorum), = *S. lasiostyla* (Boiss. Diagn. Pl. Or. ser. 1, viii. 79).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Species propria peninsulæ Ibericæ et ditionis Tingitanæ.

S. COLORATA (Poir. Voy. ii. 163; Rohrb. Sil. 114, non W., nec Schousb., nec DC. Pr. i. 374), = *S. bipartita* (Desf. Fl. Atl. i. 352, et Willk. Ic. et Descr. i. 45, tab. 30), = *S. scircea* (Guss. et auct. plurim., non All.), = *S. vespertina* (Rchb. Ic. Fl. Germ. tab. 279, et auct. mult., non Retz., nec Boiss. Voy.).

Mar. sept.—Tanger! (*Salzm. Exsicc.* sub *S. vespertina*)! Inter Tetuan et Tanger! Prope Tetuan (1851 et iterum 1871)!

Mar. occid.—Prope Mogador! Aïn el Hadjar!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia).

— — —, var. *LASIOCALYX* (Soy. W. et God. Sil. Alg. p. 26, sub *S. bipartita*).

Mar. occid.—In arenosis maritimis prope Mazagan!

SILENE SETACEA (*Viv. Fl. Lib.* 23; *Rohrb. Sil.* 105, non *Oth*), = *S. colorata*, var. *angustifolia*, *nob. olim*.

Mar. merid.—In regione inferiore pr. Mesfioua! Ourika! Sektana (*R. et Fr. spec. in herb. Kew.*)!

Accedit ad *S. coloratam* var. β , *decumbens*, *Rohrb. l. c.*, = *S. decumbens* (*Biv. Cent.* i. 75); sed in *S. setacea* folia omnia tenuia linearia, vel infima lineari-oblonga, caulis tenuis e basi ramosus, tota planta pube tenui adspersa. *S. coloratae* affinis et post eam enumeranda.

Ar. Geog. In regione desertorum ex Mesopotamia et Palæstina per Africam borealem interiorem diffusa.

S. GLAUCA (*Pourr. El. H. Matr.* 1803; *Lag. Gen. et Sp.* p. 15; *Rohrb. Sil.* 117), = *S. secundiflora* (*Oth in DC. Pr.* i. 375), = *S. ambigua*, *Cambess. in Soy. W. et God. Sil. Alg.* 24).

Synonyma alia plurima in *Rohrb. l. c.*

Mar. sept.—Tanger (*Salzm. Exsicc.*, sub *S. decumbens*, *fide Willk. Ic. et Descr.*, et *Rohrbach, l. c.*).

Ar. Geog. Peninsula Iberica. Algeria et forsan ager Tingitanus.

Floræ Maroccanæ dubia civis. Salzmann hanc speciem legit prope Malaga et forsan inter plantas suas Tingitanas specimina nonnulla commisevit. Qui series Salzmannianas habuerunt plurimi botanici hanc a Tingide nunquam viderunt, et ager Tingitanus a multis botanophilis perlustratus speciem hanc nemini post Salzmann obtulit.

S. APETALA (*Willd. Sp. Pl.* ii. 307; *DC. Pr.* i. 369; *Rohrb. Sil.* 118).

Mar. sept.—Prope Tetuan!

Mar. merid.—Prov. Shedma! Sheshaoua! Mesfioua! Seksaoua!

Ar. Geog. In regione Mediterranea calidiore ex Affghanistan usque Macaronesiam late diffusa. Deest in Europa in reg. inter Atticam et Hispaniam Muricam, et in Africa boreali inter Mare Rubrum et Mositanem.

S. CORRUGATA (*Ball in Journ. Bot.* 1873, 301), = *S. canariensis* (*Willd. in Spreng.*)? Nomen verisimiliter falsum.

E collo radiceis ramosissima, diffusa, glanduloso-pilosa; folia inferiora spathulato-acuminata, superiora lanceolata; pedicelli calycem demum æquantes; bractee 2 inæquales, major herbaacea, minor subsetacea; flores distantes, ante anthesin nutantes; calyx membranaceus, viridistriatus, nervis parallelis, basi truncatus, vix umbilicatus; petala carnea, biloba, ungue quam calyx longiori; carpophorum capsulae æquilongum; semina fusca, sat profunde canaliculata, dorso corrugata.

Mar. merid.—In regione media Atlantis Majoris; Ait Mesan! circa 1750 m.!

Affinis *Silene pseudo-Atocion*, Dsf., a pl. nostra differt petalis integris, seminibus multo majoribus vix canaliculatis, calycis tubo longiore. Habitu propior est *S. pendula*, structura omnino diversa.

Monente cel. Cosson spec. nostra ad *S. Canariensem* (Willd. in Spreng. Neue Entd. iii. 60; Rohrb. Sil. 194, non Otth in DC. Pr. i. 372, referenda sunt, et eandem plantam legit Balansa, anno 1867, in monte Djebel Haid, et secus rivulum Oued Kseb prope Mogador. Vix dubito specimen in herb. Willd. (No. 8633) a Broussonnet communicatum ex ditioe Mogadorensi ortum esse, et errore schedæ ex insula Teneriffa prodisse fertur. Dum nomen *Canariensis* incertum et falsum videtur, plantam enumeravi sub nomine triviali a me proposito. Iudicent magistri scientiæ nostræ. Icon Webbiana (Phyt. Can. tab. 18) minime exacta vix ad plantam nostram referenda est.

— —, var. *ADUSTA*, = *S. adusta* (Ball in Journ. Bot. l. c.). Proxima præcedenti et pro subspecie a me olim edita.

Planta pumila, erecta, pauciflora; differt insuper a *S. corrugata* foliis angustioribus, bracteis subæqualibus setaceis, petalis albidis, magis profunde bilobis, seminibus (non omnino maturis) pallide luteo-fuscis minus profunde canaliculatis, in fossa dorsali tuberculosus.

Mar. merid.—Prope Seksaoua! in rupibus calidissimis.

S. RUBELLA (L., DC. Pr. i. 369; Rohrb. Sil. 155).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Mar. merid.—In planitie Maroecana pr. Sheshaoua!

Ar. Geog. Regio Mediterranea, imprimis occidentalis. In Oriente, admodum rara, deest in Macaronesia.

S. INAPERTA (L., DC. Pr. i. 376; Rohrb. Sil. 165).

Mar. merid.—Ad radices Atlantis Majoris pr. Amsmiz!

Ar. Geog. Regio Mediterranea occidentalis a Corsica et Galloprovincia usque Macaronesiam.

S. MUSCIPULA (L., Rohrb. Sil. 170, non DC. Prod.), = *S. stricta* (DC. Pr. i. 379, non L.).

Mar. merid.—In regione inferiore Atlantis Majoris. Distr. Reraya! In convalle Aït Mesan! Amsmiz! Mzouda! Oued Ghagaia (*Balansa*), *Coss. Index*.

Ar. Geog. Regio Mediterranea, imprimis occidentalis. Deest in Macaronesia, et rarius in Oriente occurrat, forsan cum seminibus introducta.

Silene stricta (L., Rohrb. Sil. 170), = *S. pteropleura* (Boiss. et Rent. Pug. 18), species affinis sed bene distincta, nascitur in Hispania australi proxima (e. g. pr. San Roque!); hactenus in Imp. Maroecano desideratur.

S. ITALICA (L., sub *Cucubalo*, Pers. Syn. i. 498; DC. Pr. i. 381; Rohrb. Sil. 218).

Mar. sept.—In monte Beni Hosmar supra Tetnan legit P. B. Webb. Specimen habui ex herb. Webb sub "*S. viscosa*"!

Mar. merid.—In regione inferiore et media Atlantis Majoris. In convalle Aït Mesan (forma typica) circ. 1400 m.! In monte Djebel Tezah (forma depauperata) usque 1900 m.!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia). Forma arete affinis (*S. nemoralis*, W. K.) in Germania orientali, Hungaria, et Rossia meridionali.

SILENE MOLLISSIMA (L., sub *Cucubulo*, *Sibth. et Sm. Pr. Fl. Gr.* i. 298; *Rohrb. Sil.* 226, non DC. *Pr.* i. 382), = *S. velutina* (*Pourr. in Herb. Dsf.*; DC. *Pr.* i. 373; *Willk. Ic. et Desc.* tab. 43), = *S. Salzmanni Otth in DC. Pr.* i. 381, non Bad.). Var. *rosulata*, = *S. rosulata* (*Soy. W. et God. Sil. Alg.* 50, *Exp. bot. Alg.* tab. 82).

Mar. sept.—Tanger!

Ar. Geog. Peninsula Iberica. Africa borealis. Corsica. Scriptores Floræ Orientalis, et imprimis cel. Boissier, siluerunt de hae specie, a Sibthorp, *l. c.* indicata inter Smyrnam et Bursam.

Notæ diagnosticæ quibus *S. rosulata* a *S. mollissima* distinguitur mihi videntur nimis leves et instabiles. Specimen unicum nostrum Tingitanum simillimum est spec. *S. gibraltariæ* in monte Djebel Santo prope Oran lectis. Quæ vidi et ipse legi in rupibus Gibraltar orientem spectantibus paululum diversa sunt et aliam varietatem sistunt.

——, var. *GIBALTARICA*, = *S. tomentosa* (*Otth in DC. Pr.* i. 383), = *S. gibraltaria* (*Boiss. El.* p. 29, et *Voy. Esp.* tab. 266; *Rohrb. Sil.* 227).

Mar. sept.—Prope Tetuan legit Webb (spec. in herb. nostro sub "*S. viscoso*")!

Ar. Geog. Nascitur in fissuris rupium Gibraltar et in rupestribus Tetuanensibus.

Pace cl. Rohrbach, ex descriptione incompleta, sed omnino conformi, et loco natali plane apparet *S. tomentosa* esse stirps ipsissima quæ postea cel. Boissier melius descripsit et illustravit. Hanc ego olim legi in rupibus Gibraltar orientem spectantibus iconi Boissierianæ simillimam; sed, ut mihi videtur, est nil nisi var. *S. mollissimæ* quæ fatente cel. Boissier in rupibus septentrionem versus ejusdam montis nascitur.

S. LÆTA (*Ait. H. Kew.* ii. 118, sub *Lychnide*, *Al. Br. in Flora*, 1843, 373; *Gren. et God. Fl. Fr.* i. 220).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Ar. Geog. Peninsula Iberica, Gallia austro-occidentalis. Ditio Tingitana.

LYCHNIS (*L. Gen.* 584; *DC. Fl. Fr.* iv. 761).

L. CÆLI ROSA (L., sub *Agrostemma*; *Desr. in Lam. Dict.* iii. 644; *DC. Pr.* i. 386).

Mar. sept.—Prope Tetuan legit P. B. Webb! flores, see. MSS. Webb., albi.

Ar. Geog. Peninsula Iberica. Gallia meridionalis. Italia. Dalmatia. Ditio Tingitana.

——, var. *ASPERA* (*DC. Pr. l. c.*).

Mar. sept.—Tanger (*Webb Herb.*)! (*Schousb. spec.*)!

LYCHNIS MACROCARPA (*Boiss. et Rent. Diagn. Pl. Hisp.* p. 8; *Expl. Scient. Alg.* tab. 80).

Mar. merid.—Ad radices Atlantis Majoris prope Ainsmiz!

Ar. Geog. Peninsula Iberica. Africa borealis.

L. DIURNA (*Sibth. Fl. Ox.* 145; *Koch, Syn. Fl. Germ.* i. 146), = L. dioica, var *a*, L., = L. sylvestris (*DC. Pr.* i. 386), = *Melandrium pratense* (*Rochl. Deutsch. Fl.* i. 254; *Boiss. Fl. Or.* i. 660).

Mar. sept.—In itinere inter Tanger et Tetuan specimen unicum legit J. D. H.!

Ar. Geog. Europa fere omnis, exclusa arctica. Africa boreali-occidentalis.

HOLOSTEUM (*L. Gen.* 333).

II. UMBELLATUM (*L., DC. Pr.* i. 393).

Mar. merid.—In regione media Atlantis Majoris. Supra Arround circ. 2300 m. ! In monte Djebel Tezah circ. 2100 m. !

Ar. Geog. Europa media. Regio Mediterranea omnis (exclusa Macaronesia).

CERASTIUM (*L. Gen.* 585).

C. GLAUCUM (*Gren. Mon. Cer.* 47), var. OCTANDRUM (*Gren. l. c.*), = C. tenue (*Viv. Tent. Fl. Cors.* 7), = *Moenchia octandra* (*Gay in Perr. Cat. Fréjus*, 55).

Mar. sept.—Tanger! (*Schsb. spec.*)! (*Salzm. Exsicc.* sub *Sagina octandra*)! (*Webb Herb.*)! Prope Tetuan!

Ar. Geog. Species a Greniero *l. c.* definita per Europam mediam et regionem Mediterraneam (exclusa Macaronesia) diffusa, fit rarior in Asia Mediterranea.

C. GLOMERATUM (*Thuill. Par.* 225; *Koch, Syn. Fl. Germ.* i. 132), = C. vulgatum (*Sm. Fl. Brit.* ii. 496; *DC. Pr.* i. 415), = C. viscosum (*Gren. Mon.* p. 25; *Boiss. Fl. Or.* i. 722).

Mar. sept.—Tanger! Cap. Spartel! Prope Tetuan!

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—In planitie et in Atlante Majore usque in regionem subalpinam. Mesfiona! In convalle Aït Mesan usque 2400 m. ! In monte Djebel Tezah circ. 2200 m. !

Ar. Geog. Species per orbem terrarum fere totum diffusa, vestigia hominum et animalium secuta.

Monentibus celeb. Mertens et Koch, Bentham, aliisque magistris, nomina Linnæana *C. vulgatum* et *C. viscosum* diu vexata et ab ipso auctore in Herbario confusa hodie negligenda videntur.

C. PUMILUM (*Curt. Fl. Lond.* i. fasc. 6, tab. 30; *Gren. Mon. Cer.* 33), = C. pentandrum (*Moris Fl. Sard.* i. 266, an *L.?* an *DC. Pr.* i. 416).

Mar. sept.—Prope Tanger et inter Tanger et Tetuan legit Lagrange sec. Cosson. In monte Beni Hosmar supra Tetuan specimen nannum inter radices gramineæ cujusdam inveni, nimis incompletum et ideo incertum.

Ar. Geog. Per Europam totam (exclusa aretica) et verisimiliter per omnem regionem Mediterraneam diffusum. Synonymia auctorum incerta. In Macaronesia hæc species et affines hucusque non detectæ.

CERASTIUM BRACHYPETALUM (*Desp. in Pers. Syn.* i. 520; *DC. Pr.* i. 416; *Gren. Mon. Cer.* 36).

Mar. sept.—In monte Beni Hosmar supra Tetuan!

Mar. merid.—In regione subalpina Atlantis Majoris. In convalle Aït Mesan supra Arround, circ. 2400 m.! In monte Djebel Tezah usque 2600 m.!

Ar. Geog. Europa media. Regio Mediterranea omnis (exclusa Macaronesia).

Facie a typo Europæo paululum differt, sed notis gravioribus, et præsertim petalorum unguibus basi barbatis, bracteis herbaceis, inflorescentiæ indole, huc referendum.

C. ARVENSE (*L., DC. Pr.* i. 419), = C. mutabile (*Gren. Mon. Cer.* 68) pro parte.

Mar. merid.—In regione media Atlantis Majoris. In convalle Aït Mesan a 1800 m. usque 2300 m.! In monte Djebel Tezah circ. 2000 m. In jugo Tagherot (*R. et Fr. in Herb. Kew.*)!

Ar. Geog. Herba amphigea. Per omnem Europam, Asiam temperatam, Americam borealem et per montes regionis Mediterraneæ (inclusa Macaronesia) diffusa; nec deest in America australi frigidior.

STELLARIA (*L. Gen.* 568).

S. MEDIA (*L., sub Alsine; DC. Pr.* i. 397).

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—Mesfoua! In convalle Aït Mesan usque 1400 m.! verisimiliter communis sed cito marcescens.

Ar. Geog. Per orbem terrarum fere totum diffusa, hominum et animalium (imprimis avium) vestigia secuta.

— —, var. MAJOR, *Koch*, = *S. latifolia* (*Pers. Syn.* i. 501; *DC. Pr.* i. 396), = *S. neglecta* (*Weihe in Bluff et Fingerh. Comp. Germ.* i. 560).

Mar. sept.—Inter Tanger et Tetuan spec. legit J. D. II.!

Ar. Geog. Varietas in umbrosis regionis Mediterraneæ et Europæ calidioris frequens.

S. ULIGINOSA (*Murr. Prod. Goett.* 55; *Rehb. Ic.* fig. 3669), = *S. aquatica* (*Poll. Pl. Pal.* No. 422; *DC. Pr.* i. 398).

Mar. merid.—In regione subalpina Atlantis Majoris. In ascensu versus jugum Tagherot ultra 2500 m.!

Ar. Geog. In temperatis frigidioribus hemisphæræ borealis. In reg. subalpina montium regionis Mediterraneæ rarius occurrit. Hucusque quoad sciam in Africa non detecta fuit.

STELLARIA ARENARIA (*Salzm. Exsicc.*; an *L. et DC. Pr. i. 398?*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Species mihi non satis cognita, facie *Arenariæ spathulatæ*, Dsf., valde similis. An propria ditionis Tingitanæ?

ARENARIA (*L. Gen. 777*).

Genus a celeb. Bentham et Hooker (*Gen. Plant. i. 149*) reformatum genera plurima auctorum notis non satis firmis defuita includit, quæ sistunt sectiones sequentes. 1. *Arenaria* (auct. recent.). 2. *Mœhringia* (*L. Gen. 494*). 3. *Rhodalsine* (Gay, in *Ann. Sc. Nat. ser. 3, iv. 25*). 4. *Alsine* (Wahl. *Fl. Lapp. 125*). 5. *Minuartia* (*L. Gen. 107*). 6. *Cherleria* (*L. Gen. 570*). 7. *Hymenella* (*DC. Pr. i. 389*). 8. *Gouffieia* (Rob. et Cast. in *DC. Fl. Fr. v. 609*). 9. *Lepyrodiclis* (Fenzl, in *Endl. Gen. Pl. p. 966*). 10. *Ammoderia* (Gincl. *Fl. Sib. iv. 160*). 11. *Merckia* (Fisch. ex Cham. et Schlecht. in *Linnæa, i. 59*).

A. PUNGENS (*Clem. in Lag. Gen. et Sp. 15*; *DC. Pr. i. 403*).

Mar. merid.—In reg. subalpina Atlantis Majoris. In monte Djebel Tezah circ. 2500 m.!

Ar. Geog. Montes Hispaniæ Granatensis. Atlas Major.

— —, var. GLABRESCENS (*Ball, in Journ. Bot. 1873, 301*).

Differt a typo Hispanico sepalis glabris, nervo medio prominulo basin versus ciliato, caulibus subglaberrimis, foliis ad basin membranaceam connatis ciliatis, antheris fusco-griseis, nec rubentibus.

Mar. merid.—In regione subalpina Atlantis Majoris, Aït Mesan! versus 2500 m.!

A. SERPYLLIFOLIA (*L., DC. Pr. i. 411*).

Mar. merid.—In planitie Marocana et in Atlante Majori. Mesfioua haud procul urbe Marocco! Ourika! Aït Mesan 2000–2200 m. In jugo Tagherot usque 2800 m. In monte Djebel Tezah circ. 2200 m.!

Ar. Geog. Species per Europam fere omnem et Asiam temperatam et regionem Mediterraneam (inclusa Macaronesia) late diffusa. In ditione Tingitana nondum detecta.

Specimina nostra sæpe inter typum et *A. leptocladon* (Rehb. Cent. xv. tab. 32, = *A. serpyllifolia*, var. *tenuoir*, Koch) media videntur.

In distr. Reraya ad radices Atlantis Majoris legi specimina duo nana nimis incompleta ad *Arenarias* e grege *A. serpyllifoliæ* referenda. In his folia in petiolum brevissimum margine ciliatum sensim attenuata, sepala acuminata pubescenti-scabra, apice setis longioribus coronata. An spec. nova?

A. SPATHULATA (*Desf. Fl. Atl. i. 358*; *DC. Pr. i. 413*), = *Stellaria arenaria* (*L., Sp. Pl. App. 1196*), = *Arenaria cerastoides* (*Poir. Dict. vi. 363*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! Inter Tanger et Tetuan!

Ar. Geog. Africa boreali-occidentalis. Hispania Granatensis.

ARENARIA EMARGINATA (*Brot. Fl. Lus.* ii. 202; *DC. Pr.* i. 406), =
A. rosea (*Salzm. Exsicc.*), = *A. bætica* (*Salzm. Exsicc.*).

Mar. sept.—Prope Tanger! (*Salzm. Exsicc.* sub *A. rosea*)! (*Webb Herb.*)! In monte Djebel Kebir (*spec. ex Herb. Schsb.*)!

Ar. Geog. Peninsula Iberica. Ditio Tingitana.

Ipse auctor speciei et el. Scringe in *DC. Pr. l. c.* affinitatem proximam cum *A. tenuifolia* viderunt adeo ut de diversitate specifica dubitarunt. Facie est revera similis, nisi petalis emarginatis calyce longioribus, sed structura omnino diversa. In hac capsula *Arenariæ* genuinæ valvis profunde bidentatis, dum *A. tenuifolia* in § *Alsine* (capsulæ valvis integris) retinenda. Ego semper vidi petala calyce sesquilingiora, nec breviora ut dantur in auct. citatis.

A. TRINERVIA (*L., DC. Pr.* i. 412), = *Mœhringia trinervia* (*Clairv. Man. Herb.* 150, et plurim. auct. recent.).

Mar. sept.—In monte Djebel Kebir pr. Tanger! In monte Beni Hosmar supra Tetuan!

Ar. Geog. Europa et Asia temperata. In umbrosis reg. Mediterraneæ Europæ et Asiæ et in Macaronesia infrequens. In Africa boreali rara.

A. PROCUMBENS (*Vahl, Symb.* i. 50, tab. 33; *DC. Pr.* i. 413), = *A. geniculata* (*Poir. Dict.* vi. 365), = *Rhodalsine procumbens* (*Gay, Ann. Sc. Nat.* ser. 3, x. 25).

Mar. sept.—Prope Tetuan ad ostium fluminis (1851)! Inter Tetuan et Ceuta!

Mar. oecid.—Mogador! (*Lowe Cat.*). Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Sat frequens in planitie et in regione inferiore Atlantis Majoris. Prov. Haha! Prov. Shedma! Prope Amsmiz! In convalle Aït Mesan usque 1400 m.! Prov. Ksima (*Coll. Indig. Mardochæus, Coss. Index.*)

Ar. Geog. Regio Mediterranea Europæ et Africæ borealis. Adest etiam in Macaronesia, sed in Asia Mediterranea desideratur.

Species satis ludibunda quoad formam foliorum et longitudinem sepalorum, sed varietates stabilire inutile duco.

A. TENUIFOLIA (*L., DC. Pr.* i. 405), = *Alsine tenuifolia* auct. plurim.

Mar. sept.—In monte Beni Hosmar supra Tetuan specimen unicum legi!

Ar. Geog. Europa media. Asia temperata. Regio Mediterranea omnis (exclusa Macaronesia).

— —, var. *GLANDULOSA*, = *A. viseidula* (*Thuill. Par.* 219).

Mar. merid.—In saxosis pr. Seksaoua! Milhain!

Ar. Geog. Fere per totam arcem speciei imprimis calidiorem diffusa.

ARENARIA FASCICULATA (*Gouan, Ill.* p. 30; *Jacq. Austr.* tab. 182;

DC. Pr. i. 407, non Sibth. et Sm. Fl. Gr. v. 30), = *Alsine* Jaquinii (Koch, Syn. Fl. Germ. 125, et auct. plurim.).

Mar. merid.—In regione subalpina Atlantis Majoris in convalle Ait Mesan a 2200 m. usque 2500 m. !

Ar. Geog. Europa media, in regione Mediterranea rarius obvia. Citatur in Albania a eel. Grisebach et in montibus Velebith a eel. Visiani. In latere meridionali Alpium Europæ haud infrequens, sed in regionem inferiorem vix descendit.

Nomen triviale *A. fasciculata* a Gouan prius publici juris factum. Quid sit *A. fasciculata*, L. Spec. Pl. ed. 12, iv. 733, eruere non potui. Planta Floræ Græcæ est *Alsine Smithii* (Fzl. Alsin. 57), = *Alsine lancifolia* (Vis. Ic. Fl. Dalm.), = *Alsine fasciculata* (Maly, Enum. Anstr. 295, et Vis. Fl. Dalm. iii. 177, tab. xxxiv.). A sp. nostra differt sepalis trinervibus, foliis 5-7-nervibus, cæterisque notis gravioribus.

Specimina nostra a typo Europæo parum diversa sunt sepalis minus inæqualibus. Radix annua vel biennis, vel interdum (?) perennans.

A. SETACEA (Thuill. Par. ed. 2, 220; DC. Pr. i. 407), = *Alsine* setacea (M. K. et auct. plurim.). Var. ATLANTICA, nob., cui simillima *A. erythrosepala* (Boiss. Diagn. Pl. Or. ser. 1, viii. 98, et Fl. Or. i. 679).

Mar. merid.—In regione subalpina et alpina Atlantis Majoris. In jugo Tagherot usque 3100 m. ! In monte Djebel Tezah, J. D. H. !

Ar. Geog. Species polymorpha per montes regionis Mediterraneæ ex Atlante usque Persiam, et inde in Sibiriam Uralensem diffusa. In Europa media obvia, fit rarior occidentem versus. Deest in Macaronesia, in montibus peninsulæ Ibericæ (?) et in Alpibus Europæ.

Specimina nostra pluribus notis inter *A. setaceam*, Thuill., et *A. fasciculatam*, Gouan, media sunt, et re vera illa, nisi radice perenni, ab hac ægre dignoscitur. Sepala in specim. nostris basin versus obscure trinervia sunt, præsertim in statu juniore; inflorescentia fere *A. setaceæ* typicæ, sed interdum cymoso-corymbosa; folia trinervia, breviora, adpressa, apice cartilagineo prædita.

Cel. Boissier (Fl. Or. i. 680) sex formas seu varietates *A. setaceæ* enumeravit, quibus, pæce tanti viri, addendæ sunt *A. erythrosepala*, Boiss., et *A. libanotica*, Boiss. His adjungenda erit *Arenaria filifolia* (Forsk. Fl. Æg.-Arab. p. 211) quæ monente beato J. Gay est *Alsine Schimperii*, Hochst., stirps in montibus Abyssiniæ et Arabiæ obvia. Nostræ plantæ valde proxima est No. 983, Balansa, Pl. d'Algérie (*Alsine setacea*, M. K., var., see. Cosson). Specimina a monte Djebel Tougourt, Bal. Pl. d'Alg. No. 856 (*Alsine setacea*, var. *pubescens*, Coss.), gaudent habitu diverso et facie potius *A. juniperinæ*. De his novissimis in notula MS. b. Gay scripsit "ab *A. setacea* sine dubio distinctissima." Frustra tamen notas diagnosticas quæsivi. Semina (in spec. nostris nimis immatura) nullomodo ab *A. setacea* diversa videntur.

ARENARIA VERNA (L., DC. *Pr.* i. 405), = *Alsine verna auct. plurim.*, var. *BRACHYPETALA*, petalis calyce paullo brevioribus oblongo-ellipticis.

Mar. merid.—In regione subalpina et alpina Atlantis Majoris. In jugo Tagherot 2500–3000 m. ! In monte Djebel Tezah circ. 2500 m. !

Ar. Geog. Europa media. Asia media et borealis. Montes regionis Mediterraneæ (exclusa Macaronesia).

BUFFONIA (L. *Gen.* 168).

B. TENUIFOLIA (L. *Sp.* 179, et *Herb. ex. b. J. Gay*), = *B. annua* (DC. *Fl. Fr.* iv. 768, et *Prod.* i. 388).

Mar. merid.—Ad radices Atlantis Majoris sat frequens. Pr. Ourika ! Reraya ! Gurguri ! Seksaoua ! et in convalle Aït Mesan !

Ar. Geog. Gallia australis. Hispania. Græcia et (*ex Led.*) prov. Transcaucasicæ.

Radix interdum fit biennis. Specimen legi in distr. Reraya ramis anni præcedentis jam emarcidis, glomerulis floralibus in axillis foliorum præditis, ramis hornotinis inflorescentia racemosa typica donatis.

SAGINA (L. *Gen.* 236).

S. PROCUMBENS (L., DC. *Pr.* i. 389), var. *PARVIFLORA*, nob.

Mar. merid.—In regione subalpina Atlantis Majoris. In jugo Tagherot versus 2600 m. ! In monte Djebel Tezah circa 2500 m. !

Ar. Geog. Europa fere omnis et Asia temperata. Montes regionis Mediterraneæ. America borealis et australis.

Præter flores minores specimina nostra typo Europæo videntur omnino conformia.

S. MARITIMA (*Don in Eng. Bot.* tab. 2195; DC. *Pr.* i. 389).

Mar. sept.—In rupibus juxta Cap. Spartel ! In litore freti Herculis juxta Ceuta !

Ar. Geog. Europa media et australis in maritimis. Specimina Africana (nostris exceptis) non vidi.

S. APETALA (L., DC. *Pr.* i. 389).

Mar. sept.—Tanger ! In monte Beni Hosmar supra Tetnan !

Ar. Geog. Europa media. Regio Mediterranea omnis (inclusa Macaronesia).

S. ciliata (*Fr. Nov. Fl. Succ.* 59; *Rchb. Ic. Fl. Germ.* fig. 4956), = *S. patula* (*Jord. Obs. Pl. Fr.* p. 23).

Mar. occid.—Casa Blanca !

Forsan prioris var., sed nota diagnostica ex sepalis supra capsulam maturam adpressis, nec patulis crueiformibus, videtur satis constans et cultura immutata. Characteres cæteri plane variabiles. Haud sine dubio pro subspecie enumeravi.

S. LINNÆI (*Presl, Rel. Hænk.* ii. 14; *Boiss. Fl. Or.* i. 663), = *S. saxa-*

tilis (*Wimm. Fl. Schles.* 76; *Koch, Syn. Fl. Germ.* 119), = *Spergula saginoides* (*L., DC. Pr.* i. 394).

Mar. merid.—In regione subalpina Atlantis Majoris. In convalle Aït Mesan usque 2500 m.!

Ar. Geog. Montes Europæ fere totius, Asiæ borealis et Americæ borealis. Hucusque in Africa non indicata.

SPERGULA (*L. Gen.* 586).

S. ARVENSIS (*L., DC. Pr.* i. 394).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Ar. Geog. In arvis Europæ fere totius. Hinc inde in regione Mediterranea (inclusa Macaronesia), in India orientali et in Abyssinia.

S. PENTANDRA (*L., DC. Pr.* i. 394).

Mar. merid.—Misra ben Kara haud procul urbe Mogador!

Ar. Geog. In cultis Europæ et regionis Mediterraneæ (inclusa Macaronesia).

Semina videntur omnino diversa ab illis *S. arvensis*; sed non desunt formæ intermediæ, et hæc forsau melius pro subspecie habenda.

SPERGULARIA (*Pers. Syn.* i. 504; *B. et H. f. Gen. Pl.* i. 125).

Arenarias stipuliferas a cæteris Linnæanis Persoon *loc. cit.* primus separavit, et ad genus novum (*Spergularia*) constituendas proposuit. Nomen igitur *Lepigonum* a plurimis receptum, consentientibus Benthams, Hooker, Boissier, rei botanices magistris, prætermisi.

S. MARINA (*All. Fl. Ped.* ii. 114, sub *Arenaria*; *Wahlb. Fl. Gothob.* 47; et *Kindb. Mon. Lepig.* 18, sub *Lepigono*), = *Arenaria rubra* β. *marina* (*L. Fl. Suec.* p. 152), = *Arenaria marginata* (*DC. Fl. Fr.* iv. 793), = *Spergularia marginata* (*Boiss. Fl. Or.* i. 733), = *A. media* (*DC. Pr.* i. 401, non *L.*).

Mar. oecid.—Casa Blanca! Mogador!

Ar. Geog. Species amphigea per orbem terrarum (exclusis reg. polaribus) in maritimis diffusa, rarius in terris continentalibus subsalsis obvia.

De *L. media* (*L. Sp. Pl.* 606) conf. *Kindberg, loc. cit.* Nomen hodie inextricabile delendum.

S. FIMBRIATA (*Boiss. et Reut. Diagn. Pl. Or.* ser. 2, i. 94; *Salzm. Exsicc. MSS.*; *Kindb. Mon. Lepig.* 21, sub *Lepigono*).

———, var. α, TYPICA. Facies *S. macrorhizæ* (*Gr. et God. Fl. Fr.* i. 276), ramis elongatis sæpissime prostratis, mucrone in foliorum apice ipsorum diametrum æquante vel superante. Semina majore pro parte aptera, cætera margine fimbriata.

Mar. sept.—Tanger in maritimis! (*Boiss. et Reut. loc. cit.*).

Mar. oecid.—Casa Blanca!

———, var. β, CONDENSATA. Foliis in ramis brevioribus approximatis, hirtulis, brevissime mucronulatis, floribus numerosis in panicula

brevi condensatis, sepalis densius glanduloso-hirsutis, seminibus sæpe (non semper) omnibus apteris, stipulis argenteis maximis.}

Mar. occid.—In insula Mogador hanc formam insignem legimus !

— —, var. γ , TENUE. Ramis brevioribus tenuibus, erectis vel adscendentibus, foliis sæpius brevioribus mucrone ipsorum diametrum bis terve longiore armatis, seminibus apteris et fimbriato-marginatis commixtis.

Mar. occid.—In arenosis juxta Aïn el Hadjar !

Mar. merid.—In regione inferiore, præsertim versus radices Atlantis Majoris sat frequens. Legimus in prov. Shedma ! prope Seksaoua ! Milhaïn ! et alibi vidimus.

Permuitis capsulis scrutatis semina fimbriata nonnulla semper detexi sed in ditione nostra semina aptera certe frequentiora, et in var. β nostra semina fimbriata sat rara. Stipulæ elongatæ nitentes in hac specie insignes. Notæ diagnosticæ in hoc genere parum firmæ, sed hæc inter formas probe distinctas enumeranda videtur.

Ar. Geog. In Hispania Gaditana rara, in territorio Maroccoano verisimiliter communis, et in Macaronesia nascitur, nec, quoad sciam, alibi detecta.

SPERGULARIA RUBRA (L., et DC. Pr. i. 401, sub *Arenaria*; Pers. Syn. i. 504; Boiss. Fl. Or. i. 732), = *Lepigonum rubrum* (Fries, Fl. Halland. p. 76; Kindb. Mon. Lepig. 40).

Mar. merid.—Prov. Shedma, J. D. H. !

Ar. Geog. Europa fere tota (exclusa arctica). Regio Mediterranea omnis (inclusa Macaronesia). Asia temperata.

— —, var. PINGUIS (Fenzl, in Led. Fl. Ross. ii. 167), = *Lepigonum salinum* (Fries, Nov. Fl. Suec. Mant. iii. 34; Kindb. Mon. Lep. 36).

Mar. sept.—In maritimis juxta Ceuta !

Mar. occid.—In maritimis prope Mogador !

Mar. merid.—In prov. Shedma !

— —, var. SPERGULOIDES, nob., = *Lepigonum sperguloides* (Fisch. et Meyer, Ind. Sem. Hort. Petrop. 1853, ex Kindb. Mon. Lep. 39).

Distinguitur panicula racemosa, pedunculis subaphyllis, seminibus in spec. nostris nigrescentibus nitidis auriculato-excavatis.

Mar. occid.—Legi cum var. priore in maritimis prope Mogador !

Notæ diagnosticæ ab auctoribus, et præsertim a cl. Kindberg in Monographia allatæ, ad species plurimas hujus gregis firmandas, fallaces videntur. Stirpes enim summopere variabiles sunt *Spergulariæ* omnes; et quot species definire credunt auctores, tot formas intermedias natura præbet.

S. DIANDRA (Guss. Fl. Sic. Pr. i. 515, sub *Arenaria*; Boiss. Fl. Or. i. 733), = *Arenaria salsuginea* (Bunge in Led. Fl. Alt. ii. 163, Ic. tab. 409, flores errore lutei), = *Spergularia salsuginea* (Fenzl, in Led. Fl. Ross. ii. 166; Gren. et God. Fl. Fr. i. 275), = *Lepigonum salsugineum* (Kindb. Mon. Lepig. 42), = *Spergularia patens* (Hochst. in Schimp. Pl. Arab. Exs. No. 193, ex Boiss. l. c.).

Mar. oecid.—Casa Blanca!

Mar. merid.—Frequens in saxosis aridis regionis inferioris. Prov. Shedma! Mesfioua! Misra ben Kara!

Ar. Geog. Species Mediterranea, verisimiliter per totam regionem ealidiorem diffusa. In Macaronesia nondum detecta. In desertis Asiæ mediæ etiam obvia.

Eodem pacto quo congeneres hæc species etiam ludibunda est. Folia sæpe “obtusiuscula” (*Kindberg*), interdum “mucronata” (*Boiss.*), “glabra” (*Fenzl.*), vel sæpius puberula (*spec. nostra*); sepala occurrunt “margine tenuissime scariosa” (*Kindberg*), vel margine scariosa dorso herbaceo æquilata; stipulæ sæpissime pro genere minimæ, interdum longiores et nitentes. Semina in spec. nostris semper ad calcem diagnosis Boissierianæ “triquetro-obovata, aptera, scabriuscula, margine incrassata,” nec margine elevata angustata cristata sicut ea descripsit et delineavit *Kindberg l. c.* Nomen Gussonianum antiquius a *Kindberg* perperam prætermisum. Flores sunt revera sæpissime, sed non semper, diandri, sed nomen triviale nisi plane erroneum non mutandum. Stirps non semper in locis salsis nascitur, et eodem pacto nomen *S. salsuginea* haud exacte convenit.

POLYCARPON (*L. Gen.* 105).

P. TETRAPHYLLUM (*L.*, sub *Mollugine*; *L. fil. Suppl.*; *DC. Pr.* iii. 376).

Mar. sept.—Pr. Cap. Spartel! Inter Tanger et Tetuan! Juxta Tetuan (1851)!

Mar. oecid.—Casa Blanca! Mazagan! Mogador (*Lowe*).

Mar. merid.—Sat frequens in regione inferiore præsertim ad radices Atlantis Majoris. Legimus in prov. Shedma! Prov. Mtonga ad Aïn Tarsil! In convalle Ourika! In convalle Amsmiz circ. 1400 m.!

Ar. Geog. Europa media. Regio Mediterranea omnis (inclusa Macaronesia). Per orbem terrarum hinc inde cum seminibus allatum.

P. alsinefolium (*Biv. St. Rar. Sic. Man.* iii. 7, sub *Hagea*; *DC. Pr.* iii. 376), var. *LEIOSPERMUM*, *nob.*

A *P. alsinefolio*, cui simillima, differt insigniter var. nostra seminibus lævibus dorso leviter canaliculatis. A *P. peploidi* differt imprimis radice annua, nec perenni, sepalis acutis, nec obtusis subcucullatis.

Mar. oecid.—In rupibus insulæ Mogador! et in maritimis haud procul Mogador! Aïn el Hadjar!

Mar. merid.—In distr. Mesfioua haud procul urbe Mogador! Juxta Amsmiz spec. unicum legit J. D. II.!

Ar. Geog. *P. alsinefolium* est totius reg. Mediterraneæ (inclusa Macaronesia) civis, procul a litore maris rarius obvium.

Var. nostræ proximum est *P. arabicum* (*Boiss. Diagn. Pl. Or.* ser. 1,

x. 13). Hoc est etiam *P. tetraphyllo* arcte affine et cum *P. alsinefolio* sicut subspecies enumerandum.

POLYCARPON BIVONÆ (*J. Gay in Duchartre, Rev. Bot.* ii. 572; *Walp. Ann. Bot. Syst.* i. 82), var. *HERNIARIOIDES*, nob., = *P. herniarioides* (*Ball in Journ. Bot.* 1873, 301).

Mar. merid.—In convalle Aït Mesan a 2300 m. usque 2500 m.! In monte Djebel Tezah circ. 2400 m. (*J. D. II.*)!

Cel. Cosson certum me fecit stirpem Atlanticam fore nūl nisi var. *P. Bivonæ*, Gay, sp. polymorphæ in Europa rarissimæ, sed in Algeria e regione calidiore usque subalpinam late diffusæ. Var. nostra differt ramis brevibus condensatis, cymis paucifloris, laciniiis calyce inis obtusis apice paululum inflexis. Stamina in spec. ex Aït Mesan 3, in illis ex Djebel Tezah cæterum simillimis 5, semina nimis immatura. Calycis lacinia prorsus cecarinatæ, exteriores linea tenui dorso notatæ.

P. Bivonæ, monente b. Gay, est ipsissima *Hagea polycarpoides* (Biv. St. Rar. Sic. Man. ii. No. 3) a Candolle et Bertoloni erronee ad *P. peploides*, DC., relata.

Ar. Geog. Africa boreali-occidentalis. Sicilia. Occurrit etiam ex cel. Gay, *l. c.*, in insulis Canariensibus.

LÆFLINGIA (*L. Gen.* 52).

L. HISPANICA (*L., DC. Pr.* iii. 380; *Schousb. Gew. Marok.* 12).

Mar. occid.—In arenosis ad ripam fluvii prope Mogador! (*Schsb. l. c.*). Prope Aïn el Hadjar ad radices montis Djebel Hadid!

Ar. Geog. Per regionem Mediterraneam imprimis occidentalem diffusa, sed infrequens; deest in Italia continentali et in Græcia, necnon in Macaronesia, in Oriente rarissime occurrit.

POLYCARPIA (*Lam. Journ. Hist. Nat.* ii. 8, ex *DC. Pr.* i. 371; *Webb, Phyt. Can.* i. 156, nomen emendatum).

P. NIVEA (*Ait. Hort. Kew.* ed. 1, i. 286, et *Willd. Sp. Pl.* i. 1119, sub *Achryranthe*; *Webb, Spic. Gorg.* p. 104), = *P. gnaphalodes* (*Schousb. Gew. Marok.* 104, sub *Illecebro*; *Poir. Dict. Suppl.* iv. 473; *DC. Pr.* iii. 373), = *P. microphylla* (*Car. Anal. Cienc. Nat.* iii. 25), = *P. glauca* (*Chr. Sm. in Tuck.* p. 250, ex *Webb l. c.*), = *P. candida* (*Webb, Phyt. Can.* i. 158, tab. 21).

Mar. occid.—In arenosis maritimis pr. Mogador! (ubi jam legerant Schousboe et Broussonnet et Balansa).

Nomen triviale *nivea* prioritatem gaudet et a b. Webb rite restitutum.

Ar. Geog. Species Canariensis et ius. Cap. Viridis, deest in Madeira, et ultra plagam Mogadorensis in terris continentalibus nondum visa.

PORTULACACEÆ (*Juss. Gen.* 313; *Endl. Gen.* 940, ordo reformata et nomen emendatum).

PORTULACA (*L. Gen.* 602).

P. OLERACEA (*L., DC. Pr.* iii. 353).

Mar. occid.—Saffi! Mogador (*Lowe Cat.*).

Mar. merid.—Seksaua (*J. D. H.*)! et alibi visa in ruderatis.

Ar. Geog. Planta cosmopolita per regiones temperatas calidiores totius fere orbis diffusa.

MONTIA (*L. Gen.* 101).

M. FONTANA (*L., DC. Pr.* iii. 362).

Mar. merid.—In regione subalpina Atlantis Majoris. Legimus in eonvalle Aït Mesan vix florentem cire. 2500 m.!

Ar. Geog. Herba per Europam omnem et Asiam temperatam diffusa. Nec in Macaronesia, nec in Algeria indicatur.

TAMARISCINÆ (*Desv. Diss. Inst. Gall. Leet.* ex *DC. Pr.* iii. 95).

TAMARIX (*L. Gen.* 375).

T. AFRICANA (*Poir. Voy.* ii. 189; *DC. Pr.* iii. 95).

Mar. sept.—Tetuan (*Webb Herb.*)! Inter Tetuan et Centa!

Ar. Geog. Regio Mediterranea occidentalis. Deest in Macaronesia.

T. GALLICA (*L., DC. Pr.* iii. 96; *Schousb. Gew. Marok.* 131).

Mar. sept.—Tanger! (*Schousb. spec.*)! Tetuan (*Webb Herb.*)!

Mar. occid.—Mogador! (*Lowe Cat.*).

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

Specimen a nobis prope Tanger lectum, *T. Canariensi* simile, non differt a specie. Proxime affinis *T. Anglica*, Webb, habitat secus litora Atlantica Galliae occidentalis. In Anglia vix indigena est.

T. speciosa (*Ball in Journ. Bot.* 1873, 301).

Frutex 8–10-pedalis; cortex fusco-purpureus, nitidus; racemi coëtanei in ramis lignosis et in hornotinis subsessiles; bractea ovata, minus quam in *T. gallica* aetate; sepala oblongo-rotundata, rarius ovata subaetata; petala cito caduca; flores pentandri; capsula (pro grege) longa, pulchre rosea.

Mar. merid.—Copiose secus torrentem in distr. Mesfiona! ad 600 m. Frutex elegans speciosa, herba viridi, fructu roseo, a *T. Gallica* et *T. Africana* facie valde diversa.

Sicut subspecies hic enumerata, forsitan melius cognita pro sp. omnino diversa retinenda.

T. ARTICULATA (*Vahl, Symb.* ii. 48, tab. 32; *DC. Pr.* iii. 96).

Mar. merid.—Vidimus sine fl. vel fr. in planitie haud procul urbe Marocco, ramus sterilis tantum lectus!

Ar. Geog. Regio desertorum ; verisimiliter ex Marocco meridionali usque fines Indiæ Orientalis diffusa.

HYPERICINÆ (DC. *Théor. Elém.* p. 214).

HYPERICUM (*L. Gen.* 902).

II. PERFORATUM (*L., DC. Pr.* i. 549).

Mar. sept.—Tanger (*ex Herb. Schsb. spec.*)! (*Salzm. Exsicc.*)! Tetuan (1851)! Specimen ehen vermibus erosum!

Mar. merid.—Ad radices Atlantis Majoris. Tasseremont! Aït Mesan versus 1250 m.! Amsmiz! Tamarut (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Europa omnis (exclusa arctica). Regio Mediterranea fere omnis (inclusa Macaronesia). In America boreali videtur spontaneum.

Specimina ex Mar. meridionali habent lacinias calycinas longiores quam in typo Europæo.

II. UNDULATUM (*Schousb. in Willd. Enum.* p. 811), = *H. quadrangulum*, var. ϵ (*Choisy in DC. Pr.* i. 548), = *H. bæticum* (*Boiss. Voy. Esp.* i. 114, tab. 34).

Mar. sept.—Vidi in Herb. Kewensi specimen Tingitanum a Salzmann receptum cui scheda "*H. tetragonum*, var. *acutangulum*," *H. bætico*, Boiss., satis conforme.

Ar. Geog. Peninsula Iberica. Africa borealis. Madeira. Insul. Azoriæ (*H. decipiens*, Wats.).

Suadente cel. Trevirano (in *Hyp. Geu. Animad.* p. 9) nomen antiquum Schousbœanum *H. bætico* restitui, et, donec experientia melius docebit, velut speciem distinctam enumeravi.

II. AUSTRALE (*Ten. Syll. Fl. Neap.* 385; *Gren. et God. Fl. Fr.* i. 315), = *H. repens* (*Poir. Voy.* ii. 224; *Choisy in DC. Pr.* i. 548, non *L.*, nec *Boiss. Fl. Or.* i. 801).

Mar. sept.—In monte Beni Hosmar supra Tetuan! In monte Djebel Dersa pr. Tetuan (*Webb Herb.*)!

Ar. Geog. Regio Mediterranea occidentalis (quoad peninsulam Ibericam incertum). Forma affinis in Gallia occidentali.

Quid sit *H. repens*, *L.*, dijudicare non audeo. Suadente viro gravissimo, cel. Boissier, erit idem ac *H. perforatum* δ . *punctatum* (*Choisy in DC. Prod.*), = *H. alpestre* (*Stev. Verz.* p. 95), = *H. linarioides* (*Bosse in Ott. et Dietr. Gartenzeit.* iii. 90). Specimina plurima Orientalia quæ vidi a stirpe Gallica (*H. repens*, *DC. Pr.*) certe diversa sunt, et propius accedunt *H. hyssopifolio*, *Vill.* Hæc novissima est species valde polymorpha quacum cel. Boissier (*Fl. Or.* i. 799) plurimas formas ab ipso olim pro speciebus novis propositas semel conjunxit. Dubium mihi adest an notæ differentiales quibus vir celeb. *H. repens* suum ab *H. hyssopifolio* distinxit stabiles sint : quoad rationem

calyceis et capsulæ maturæ hæc certe variabilis est. *H. repens* (Sibth. et Sm. Fl. Gr. tab. 775, non L.) est species Cretensis, *H. trichocaulon* (Boiss. & Held. Diagn. Pl. Or. ser. 1, viii. 110). Hoc mihi valde proximum *H. australi*, Ten., videtur, sed sec. auctores in sectione *Millepora*, Spach, collocatur, dum species nostra inter *Taniocarpia*, Spach, enumeranda est.

Hic notandum est specimina *H. australis* ex Italia meridionali et Sicilia quæ vidi (e. g. spec. ab ipso Tenore in Herb. Gay, hodie Kewensi), et Algeriensiis plurima a me scrutata cum spec. Gallicis et Maroccanis hand adamussim convenire. Pluribus notis priora inter *H. australe* et *H. linearifolium* (Vahl, Symb. i. 65), = *H. linearifolium* (DC. Prod. et auct. plurim.) media sunt. Re vera characteres quibus distinguitur species Tenoreana, scil. folia latiora, sepala margine glandulis paucis sessilibus, nec numerosis pedicellatis, prædita, non gravis momenti sunt, et hæc melius ad subspeciem reducenda erit.

HYPERICUM TOMENTOSUM (L., DC. Pr. i. 551).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. occid.—Aïn el Hadjar! Prov. Haha prope Mogador! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Prov. Shedma! Mskala! Mtouga!

Ar. Geog. Regio Mediterranea occidentalis ex insulis Sicilia et Melita ad litora Atlantica. Deest in Macaronesia.

H. pubescens (Boiss. *Elench.* 42; *Walp. Rep.* i. 383), = *H. suberosum* (*Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Schousb. spec. ex Herb.*)! (*Webb Herb.*)! Tetuan prope urbem ed ad ostium fluminis, 1851!

Mar. occid.—Casa Blanca! Mazagan!

Mar. merid.—Sat frequens in regione inferiore. Aïn Beida! Mesfioua! Mzouda! Sektana (*R. et Fr. spec. in Hb. Kew.*)!

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

Tam *H. tomentosum* quam *H. pubescens*, quod pro subspecie teneo, in reg. meridionalibus fiunt glabriora et minora.

Specimina typica *H. pubescentis* omnino diversa ab *H. tomentoso* videntur, sed non desunt formæ intermediæ (e. g. spec. prope Alemque in Estremadura Lusitanica a Bar. de Paiva lecta, in Herb. Kew. asservata). Spec. nostra ex Mesfioua etiam subintermedia sunt.

H. COADUNATUM (*Chr. Smith in Buch, Besch. Canar.* 153; *Webb, Phyt. Canar.* i. 45, tab. 4), var. *ATLANTICUM*, nob.?

Mar. merid.—Nondum florens legimus in convalle Aït Mesan circ. 1700 m. s. m.!

Folia et caulis foliaceus cum spec. Canariensibus adamussim conveniunt, sed spec. nostra nimis incompleta. Cel. Cosson benevole communicavit spec. in monte Djebel Ouensa ab indigeno Ibrahim lecta, cui scheda "*H. atlanticum*, Coss. MSS.," verisimiliter huc etiam

referenda. Donec eel. auctor meliora docebit, hoc mihi videtur nimis proximum *H. coadunato*. A sp. Canariensi parum distant *H. Naudinianum* (Coss. et Dur. in Bull. Soc. Bot. Fr. ii. 308) stirps Algeriensis, et *H. caprifolium* (Boiss. El. 41; Voy. Esp. tab. 35) Hispaniæ australis incola; et hæc forsân sicut subspecies vel varietates ad *H. coadunatum* ducenda.

Ar. Geog. Species in insulis Canariensibus; varietas in Atlante Majori; formæ proximæ affines in Algeria et Hispania.

HYPERICUM PERFOLIATUM (*L., Sibth. et Sm. Prod. Fl. Gr.* ii. 116; *Gris. Spicil.* i. 224), = *H. ciliatum* (*Lam. Dict.* iv. 170; *DC. Pr.* i. 552; *Boiss. Fl. Or.* i. 812), = *H. dentatum* (*Lois. Fl. Gall.* p. 499; *DC. Pr.* i. 552), = *H. montanum* (*Desf. Fl. Atl.* ii. 216, non *L.*, nec aliorum).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)! Tetuan (*Webb Herb.*)! Inter Tetuan et Ceuta!

Ar. Geog. Regio Mediterranea fere omnis.

Cel. Grisebach nomen Linnæanum ab auctoribus recentioribus prætermisum vindicavit. Specimen in Herb. Linn. est omnino *H. ciliatum*, Lam.

MALVACEÆ (*R. Br. Cong.* p. 8).

MALOPE (*L. Gen.* 841).

M. MALACOIDES (*L., DC. Pr.* i. 429), var. **STIPULACEA**, = *M. stipulacea* (*Cav. Anal. Cienc. Nat.* iii. 74; *DC. Pr.* i. 429).

Mar. sept.—Frequens circa Tanger! Tetuan (1851)!

Mar. occid.—? Mogador sec. Broussonnet in Cav. loc. cit. B. Broussonnet interdum plantas originis diversæ semel miscuisse certe scimus. Hanc speciem legit in ditione Tingitana, ubi frequenter occurrit. Nemo recentiorum eam vidit in vicinitate urbis Mogador.

Ar. Geog. Species per regionem Mediterraneam (exclusa Maearonesia) late diffusa; varietas in Africa boreali-occidentali et in Hispania australi obvia est.

M. TRIFIDA (*Cav. Diss.* ii. 85, tab. 27; *DC. Pr.* i. 429).

Mar. sept.—Tanger (*Schousb. spec.*)! (*Salzm. Exsicc.*)! Tetuan (*Webb Herb.*)!

Ar. Geog. Hispania Bætica et ditio Tingitana.

ALTHÆA (*L. Gen.* 839).

A. HIRSUTA (*L., DC. Pr.* i. 437), var. **GRANDIFLORA** (*Ball in Journ. Bot.* 1873, 302).

Differt a typo corolla subduplo majore, purpureo-rosea, stipulis longioribus, angustioribus, subemneatis, nec ut in typo ovatis, foliis magis divisis, laevius angustioribus, planta tota minus hirsuta.

Mar. merid.—Specimina manea legimus in prov. Mtouga! et Shedma!

Melius explorata forsitan sicut species nova distinguenda. In herb. beati J. Gay (nunc Kewensi) adsunt specimina duo prope Monspelium lecta, quibus adjecit in scheda "*Althæa mutica*, Gay MSS. (1824)." In his corolla magna ut in nostra, sed stipulæ et folia omnino *A. hirsuta*.
Ar. Geog. *A. hirsuta* per Europam mediam et regionem fere omnem Mediterraneam (exclusa Macaronesia) diffusa est.

LAVATERA (*L. Gen.* 842).

L. TRIMESTRIS (*L., DC. Pr.* i. 438).

Mar. sept.—Frequens pr. Tanger! (*Salzm. Exsicc.*)! Tetuan (1851)!
 Mar. merid.—Sat frequens ad radices Atlantis Majoris. Mesfiona!
 Tasseremont! Ourika! Sektana (*R. & Fr. spec. in Herb. Kew.*)!
Ar. Geog. Per regionem Mediterraneam (exclusa Macaronesia) late diffusa; in Algeria rara.

— —, var. *MALVÆFORMIS*, *nob.*

Floribus minoribus pallide purpureis, foliis subrotundatis, superioribus vix angulatis. Herba tota longius pilosa.
 Mar. merid.—Distr. Reraya versus 1100 m.!

L. OLBIA (*L., DC. Pr.* i. 438).

Mar. sept.—In monte Beni Hosmar pr. Tetuan nondum florentem legimus!
 Ex indumento et calyceibus fructibusque emareidis anni præcedentis potius ad typum quam ad var. sequentem referenda.

— —, var. *HISPIDA*, = *L. hispida* (*Desf. Fl. Atl.* ii. 118, tab. 171; *DC. Pr.* i. 438).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! In rupibus pr. Cap. Spartel!
 In montibus Djebel Dersa et Beni Hosmar (*Webb Herb.*)!
Ar. Geog. *L. Olbia*, enim var. *L. hispida*, in umbrosis et rupestribus regionis Mediterraneæ occidentalis habitat, nunquam a mari remota. Deest in Macaronesia. Species proxime affinis (*L. unguiculata*, *Desf.*) Græciam, Siciliam, Cyprum et Cretam ineolit.

L. MARITIMA (*Gouan, Ill.* p. 46, tab. 11; *DC. Pr.* i. 439).

Mar. merid.—Prope Aïn el Hadjar!
Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

L. CRETICA (*L., DC. Pr.* i. 439), = *L. sylvestris* (*Brot. Fl. Lus.* ii. 227, non *Cyr.*), = *Malva pseudolavatera* (*Webb, Phyt. Can.* i. 29), = *M. hederæfolia* (*Vis. Fl. Dalm.* iii. 205).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Tetuan (1851)! Inter Tetuan et Ceuta!

Mar. oecid.—Casa Blanca! Rabat (*R. et Fr. spec. in Herb. Kew.*)! Mogador! (*Lowe Cat.*; *Balansa*; *Coss. Index*).

Mar. merid.—Pluries visa; legimus prope Amsmiz! et in ruderalis urbis Maroeo!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). In Anglia australi advena. In America australi ex Europa allata.

— —, var. *ACUTILOBA*, *nob.* Differt foliis acute lobatis, lobis sinu acuto vel rectangulo divisim margine serratis, fol. superioribus angustioribus interdum subhastatis, tota planta glabriuscula, et statura paululum minore.

Mar. occid.—Casa Blanca! Mazagan! Prope Mogador!

Nisi vidissem formas intermedias inter spec. ex insula Teneriffa et nostra Maroccana pro subspecie nova hanc varietatem habucri.

Tam in typo quam in var. nostra segmenta involucri interdum fere ad basin divisa. Species est ambigua, inter *Malvas* et *Lavateras* fere media. Est planta polymorpha aspectu variabilis, sed in sententiam cl. Lowe (*Man. Fl. Mad.* i. 64) *L. sylvestrem*, Brot., a sp. Linneana distinctam esse convenire nequeo.

MALVA (*L. Gen.* 841).

M. HISPANICA (*L., DC. Pr.* i. 431).

Mar. sept.—Tanger (*Schsb. spec.*)! (*Salzm. Exsicc.*)! (*Webb Herb.*)! Tetuan (1851)!

Mar. occid.—Casa Blanca! Djebel Hadid (*R. et Fr. spec. in H. Kew.*)! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Frequens in regione inferiore et ad radices Atlantis Majoris. Mesfioua! Tasseremout! Ourika! Mtouga! Seksaoua! Mskala!

Ar. Geog. Hispania. Africa boreali-occidentalis.

M. SYLVESTRIS (*L., DC. Pr.* i. 432).

Mar. sept.—Tanger (*Webb Herb.*)!

Mar. merid.—Haud raro in irrigatis reg. inferioris et ad radices Atlantis Majoris. Sheshaoua! Ourika! Amsmiz! Milhaïn! et in convalle Aït Mesan circ. 1300 m.!

Ar. Geog. Europa omnis, exclusa arctica. Sibiria. Regio Mediterranea fere omnis; desideratur in Macaronesia, sed occurrit var. sequens in Madeira.

— —, var. *MAURITIANA* (*Boiss. Fl. Or.* i. 819), = *M. mauritiana* (*L., DC. Pr.* i. 432; *Lowe, Man. Fl. Mad.* i. 66).

Mar. sept.—Specimen nondum florens legit in monte Beni Hosmar pr. Tetuan J. D. H.! fere certe huc referendum.

M. ROTUNDIFOLIA (*L., DC. Pr.* i. 432), = *M. vulgaris* (*Fries, Nov. Fl. Suec.* p. 219).

Mar. merid.—In regione media Atlantis Majoris. Legimus in convalle Aït Mesan circ. 1600 m.! et supra Arround circ. 2400 m.!

Ar. Geog. Europa omnis, exclusa arctica. Regio Mediterranea, præsertim orientalis, in montanis.

— —, var. ? *INTERMEDIA*, *nob.*

Mar. occid.—Prope Mogador frustula incompleta a locustis erosa legi-

mus ! Horum specimen unum est fere exacte *M. rotundifolia* ; gaudet petalorum unguibus barbatis, et carpellis lævibus brevissime puberulis. Spec. alterum defloratum habet carpella rugosa, vel superne lævia et infra (sub calyce) rugosa. An foret vera varietas, seu potius lusus *M. parvifloræ* in regione litorali frequentis ?

Malva parviflora (L., DC. Pr. i. 433).

Mar. occident.—Casa Blanca ! Mazagan ! Mogador ! (Lowe Cat.).

Mar. merid.—Frequens in calidioribus subhumidis. Sheshaoua ! Mskala ! Sektana ! In ruderalis urbis Marocco !

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia).

Notæ diagnosticæ quibus *M. rotundifolia*, L., *M. borealis*, Wallm., *M. parviflora*, L., et *M. microcarpa*, Desf., discernuntur certe variabiles sunt ; *M. parviflora* tamen (eius var. levis est *M. microcarpa*) pluribus notis facile dignoscitur, et vera subspecies est. De hac specie et sequente confer Lowe (Man. Fl. Mad. i. 65). Vir accuratissimus hanc cum *M. boreali* conjungit, sed verisimiliter ob diversitatem carpellorum, a *M. rotundifolia* omnino diversam judicat. Sed carpella etiam interdum ludunt in hac et in *M. nicænsi*, et nota ex eorum superficie diagnosin securam non præbet.

M. NICÆNSIS (All. Fl. Ped. ii. 40 ; DC. Pr. i. 433).

Mar. sept.—Tanger (mense Junio florentem) ! Tetuan (1851) !

Mar. merid.—In ruderalis urbis Marocco !

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

Specimina Tingitana habent carpella glabrescentia parum rugosa.

LINEÆ (DC. Théor. Elém. i. 217).

RADIOLA (Gm. Syst. i. 298).

R. LINOIDES (Gmel. l. c. ; DC. Pr. i. 428), = *Linum radiola* (L., Schousb. Gew. Marok. 139).

Mar. sept.—In monte Djebel Kebir prope Tanger (Schousb. l. c.).

Ar. Geog. Europa fere tota, imprimis occidentalis. In reg. Mediterranea line inde. Deest in Macaronesia.

LINUM (L. Gen. 389).

L. GALLICUM (L., DC. Pr. i. 423 ; Schousb. Gew. Marok. 136).

Mar. sept.—Tanger (Schsb. l. c.).

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). Abyssinia.

L. STRICTUM (L., DC. Pr. i. 424 ; Schousb. Gew. Marok. 137).

Mar. sept.—Tanger (Schsb. spec.) ! Tetnan 1851 !

Mar. occident.—Casa Blanca ! Mazagan ! Mogador ! Djebel Hadid !

Mar. merid.—Frequens in regione inferiore et ad radices Atlantis Majoris. Mesfoua ! Tasseremont ! Seksaona ! Onrika (R. et Fr. in Herb. Kew.) !

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia). Abyssinia. Forma corymbulis densis axillaribus (*L. spicatum*, Pers.) est potius lusus quam varietas; fere ubique cum specie simul crescit.

LINUM SETACEUM (Brot. *Phyt. Lus.* fasc. i. No. 21; *DC. Pr.* i. 423), = *L. tenuifolium* (Schousb. *Gew. Mar.* 135, non *L. nec Lam.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*, sub “*L. bicolor*, Schsb.”) ! (*Schsb. l. c.*). Tetuan (1851) !

Mar. occid.—Mazagan !

Mar. merid.—Prov. Shedma ! Prov. Mtouga !

Ar. Geog. Peninsula Iberica. Marocco.

Valde proximum sequenti, ejus mihi videtur forma minor parviflora, inflorescentia sæpius paniculato-racemosa.

L. TENUE (*Desf. Fl. Atl.* i. 280, tab. 81; *DC. Pr.* i. 423), = *L. virgatum* (*Schousb. Gew. Mar.* 136), = *L. melianthum* (Brot. *Fl. Lus.* i. 484) ? = *L. scabrum* (*Kunze in Flora*, 1846, 654), et verisimiliter = *L. ramosissimum* (*Willk. in Bot. Zeit.* 1847, 240, non *Cl. Gay*).

Mar. occid.—Casa Blanca ! Mogador ! (*Lowe Cat.*). Aïn el Hadjar ! In monte Djebel Hadid ! Prov. Iaha pr. Mogador ! (*Schsb. l. c.*).

Mar. merid.—Frequens in prov. Shedma prope Aïn Oumast ! Mskala ! et alibi.

Ar. Geog. Africa boreali-occidentalis. Peninsula Iberica.

Species variabilis, quoad staturam et inflorescentiam valde ludibunda.

Inflorescentia sæpius paniculato-corymbosa, sed inter specimina eodem loco nata occurrit lusus paniculato-racemosus, floribus secundis, a cl. Kunze pro specie nova (*L. scabrum*) descriptus.

L. CORYMBIFERUM (*Desf. Fl. Atl.* i. 279, tab. 80; *DC. Pr.* i. 424), var. *MUNBYANUM* (*Boiss. et Reut. Pag.* 24).

Mar. sept.—Tanger ! (*Schousb. Herb.* sub “*L. corymbiferum*”), *Coss. spec.* ! (*Webb Herb.*) ! (*Salzm. Exsicc.* sub “*L. virgatum*”) !

Mar. merid.—Sat frequens ad radices Atlantis Majoris, rarius in convallibus interioribus. Legimus in distr. Mesfioua ! prope Mzonda ! et in convalle Aït Mesan !

Ar. Geog. Africa boreali-occidentalis; forsitan etiam in Lusitania—specimen ex collibus herbosis Estremadura (*Welwitsch*, No. 80, in *Herb. Kew.*, sub *L. meliantho*, Brot.) propius huic quam *L. tenui* accedit.

Affinitas inter stirpem nostram et *L. corymbiferum* nimis areta, et nisi auctoritate celeb. Boissier et Cosson devinctus, vix pro varietate habuerim, dum formæ intermediæ plurimæ adsunt. Quoad notas diagnosticas quibus *L. Munbyanum* a *L. corymbifero* distinguitur, monendum est iconemet descriptionem in ‘*Fl. Atlantica*’ parum accuratas esse. In omnibus hujus gregis speciebus margo foliorum cartilaginea sæpius læviuscula vel punctis prominentibus asperula, sed interdum ex denticulis minutis serrulata, denticulis tamen mere cartilagineis et semper antrorsum, nec retrorsum versis. Verba igitur ecl. Desfon-

taines, "folia retrorsum aspera," certe erronea, et folia superiora in icone citata vere dentata in vivo nunquam visa. Quoad inflorescentiam hæc in *Linis* omnibus hujus gregis sat variabilis, et differentię parvi momenti. Ratio longitudinis stigmati et staminum potius ex *dimorphismo* quam ex caractere specifico hæret? Inter specimina prope Tanger mense Junio lecta video folia, fere exacte ut in *L. corymbifero* Algeriensi, lanceolata margine serrulata, et calycees nullomodo diversos. *L. corymbiferum*, ejus specimen authenticum in herb. beati Desfontaines accuratè scrutavi, differt a *L. Munbyano* foliis latioribus cartilagineo-serrulatis magis confertis, corymbo minus laxo. In *L. Munbyano* a loco classico prope Oran folia multo angustiora margine læviusecula, radix nunquam fit lignosus ut in *L. corymbiferum*. Nostra planta in scipibus et dumetis sæpius obvia, gaudet ramis usque 4-5-pedalibus, et minus dense foliosis. Videtur sæpius bienne, interdum perennans.

Hic notandum est *L. corymbiferum* et *L. tenue* proxime affinia esse, adeo ut illud potius pro subspecie quam pro typo distincto habendum esse videtur. Specimina facile invenies aspectu toto cœlo diversa: sed adsunt permultæ formæ intermediæ; et qui has accuratè observaverit facile credet *L. corymbiferum* esse progeniem *L. tenuis* factam biennem et vegetiorem. Hæ species ad sect. *Syllinum*, Gris., petalorum unguibus cohærentibus, referendæ. Vidimus sæpe petala caduca simul cohærentia adeo ut corollam monopetalam simulabant.

LINUM BICOLOR (*Schousb. Mar.* 135; *DC. Pr.* i. 428). Deest in versione operis Schousbœi lingua Germanica edita.

Mar. sept.—Tanger (*Schsb. ex DC. loc. cit.*).

Quid sit nescio. Ex brevi diagnosi in *DC. Prod. l. c.* videtur hybridum *L. setaceo-angustifolium*?

L. ANGUSTIFOLIUM (*Huds. Fl. Angl.* 134; *DC. Pr.* i. 426).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Mar. occid.—Casa Blanea!

Mar. merid.—Mzouda!

Ar. Geog. Europa occidentalis. Regio Mediterranea omnis (inclusa Macaronesia).

Hæc species, ut jam monuit cel. Boissier, non est perennis, sed sæpe perennans. In Madeira, teste Lowe, semper annuum est.

L. PERENNE (*L. Sp. Pl.* 397), = *L. anglicum* (*DC. Pr.* i. 427).

Mar. sept.—? In monte Beni Hosmar supra Tetuan legi specimen unicum nondum florens et ideo omnino incertum!

Ar. Geog. Europa media. In regione Mediterranea rarius occurrit.

L. SUFFRUTICOSUM (*L., DC. Pr.* i. 427).

Mar. sept.—Tanger (*Schousb. Herb. ex Coss. spec.*)!

Ar. Geog. Gallia australis. Peninsula Iberica. Africa borealis.

ZYGOPHYLLÆ (*R. Br. Gen. Rem.* p. 13).TRIBULUS (*L. Gen.* 532).

T. TERRESTRIS (*L., DC. Pr.* i. 703).

Mar. occid.—Mazagan! Casa Blanca (*R. et Fr. sp. in Herb. Kew.*)!

Mar. merid.—Prov. Shedma!

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia). Occurrit ad Prom. B. Spei, an introductus?

ZYGOPHYLLUM (*L. Gen.* 530).

Z. FONTANESII (*Webb, Phyt. Can.* i. 17, tab. 1; *Walp. Rep.* i. 496), =Z. Webbianum (*Coss. Bull. Soc. Bot. Fr.* ii. 364, in obs. ad *Z. cornutum*).

Mar. occid.—In rupestribus maritimis prope Saffi!

Ar. Geog. Hucusque nonnisi in Ins. Canariensibus et Cap. Viridis visum.

Beatus Webb hoc a *Zygophyllo albo* Ægyptiaco diversum esse non dubitavit, sed, forsan ex synonymo in *Buch. Phys. Besch.* Canar. Ins. citato in errorem ductus, *Z. album*, Desf. Fl. Atl. esse idem ac stirpem Canariensem, et a *Z. albo*, Linn., alienum existimavit. Priori igitur nomen *Fontanesii* imposuit. Cel. Cosson (in *Bull. Soc. Bot. Fr. loc. cit.* et in vol. iii. 705) differentias *Zygophyllorum* hujus gregis accurate definivit. Monente viro perspicaci *Z. album*, Desf., est ipsissimum *Z. album*, L., cujus synonymon est *Z. proliferum*, Forsk. Hoc per regionem Mediterraneam orientalem ex Cilicia et Arabia Petraea usque in regnum Tuncetanum diffusum est. Regionem Saharensis incolit *Z. cornutum*, Coss., et in eadem regione, sed verisimiliter rarius, occurrit *Z. Geslini*, Coss. Hoc novissimum *Z. albo* proxime affine est, forsan sicut subspecies enumerandum. Nostram stirpem occidentalem, nonnisi in litore Atlantico obviam, Desfontaines non cognovit; unde nomen triviale *Fontanesii* in *Webbium* mutari voluit cel. Cosson. Error tamen auctoris ad scientiam falsificandam non aptus, et per multæ species nomine botanicorum ipsis alienorum decorantur, quamobrem nomen *Fontanesii* non mutavi.

FAGONIA (*L. Gen.* 531).

F. CRETICA (*L. DC. Pr.* i. 704).

Mar. sept.—Tetuan (*Webb Herb.*)!

Mar. occid.—Saffi! In monte Djebel Hadid haud procul Mogador!

Mar. merid.—Sat frequens in planitie, et rarior in regione inferiore Atlantis Majoris. Prov. Iaha! Prov. Shedma! Prov. Mzouda! In convalle Aït Mesan usque 1300 m.! Seksaoua!

Ar. Geog. Regio Mediterranea imprimis occidentalis (inclusa Macaronesia). In Oriente nonnisi in insulis Creta et Cypro detecta.

In spec. pr. Seksaoua lectis capsulæ sunt tota superficie glanduloso-

hirtulæ et folia aliquid minora, quibus notis ad *F. sinaicam*, Boiss., accedit; petala tamen, capsulæ et semina sunt omnino *F. creticæ*.

GERANIACEÆ (*DC. Fl. Fr.* iv. 83).

GERANIUM (*L. Gen.* 832).

G. MALVÆFLORUM (*Boiss. et Reut. Pug.* 27; *Walp. Ann.* iv. 395), = *G. tuberosum*, var. *debile* (*Ball, in Journ. Bot.* 1873, 302).

Rhizoma horizontale crassum (ex tuberibus concatenatis) caules erectos edens; folia radicalia petiolo longo retrorsum vel patenter piloso suffulta, cum caulinis inferioribus ambitu pentagona, 5-7-partita, segmentis pinnatipartitis, laciniis oblongo-linearibus integris vel parce et obtuse dentatis, superiora breviter petiolata, segmentis angustioribus, omnia utrinque adpresse pilosa, subtus glaucescentia, nervis prominentibus; petioli cum caule ex pilis longioribus brevioribusque intermixtis hirti; sepala cis *G. tuberosi* subduplo majora, oblongo-ovalia, adpresse hirsuta, margine villosa, apice abrupte mucronata, mucrone cito marcescente; petala magna sepalis triplo longiora, profunde emarginata, sinu obtuso, purpureo-lilacina, venis saturatioribus picta; rostrum hirtum in stylo attenuatum. Fructum maturum nondum vidi.

Mar. merid.—In regione subalpina Atlantis Majoris. In convalle Aït Mesan supra Arround circ. 2300 m., et in monte Djebel Tezah usque 2600 m. nondum florens legimus.

Ar. Geog. *G. tuberosum* per regionem Mediterraneam omnem (exclusa Macaronesia) diffusum est. *G. malvæflorum* nascitur in Africa boreali-occidentali et in Hispania Granatensi.

Ex radice a clar. G. Maw in hortum suum allata specimina pulchra florentia mihi benevole misit, et ipse viva vidi. In his rhizoma fere cylindricum incrassatum, caules pedales et ultra, flores eos *Malvæ sylvestris* æmulantes. Ex descriptione (loco citato) planta a Boissier et Reuter prope Tlemsen lecta nostræ fere exacte conformis. Specimina quæ habui a Jamin (No. 139), prope Alger lecta, videntur multo propiora *G. tuberoso*, et ejus meram varietatem sistunt. Plantam nostram esse progenies *G. tuberosi* non dubito, sed notæ diagnosticæ satis graves ut pro subspecie habeatur.

G. FYRENAICUM (*L., DC. Pr.* i. 643).

Mar. merid.—In regione media et inferiore Atlantis Majoris. In convalle Aït Mesan 1500-1800 m. ! In convalle Amsmiz 1400-1800 m. ! Prope Sektana (*R. et Fr. in Herb. Kew.*) !

Ar. Geog. Europa omnis, exclusa arctica, rarius in regione Mediterranea. Oriens. Deest in Macaronesia.

G. MOLLE (*L., DC. Pr.* i. 643).

Mar. sept.—Tanger ! Cap. Spartel !

Mar. merid.—Mskala! In convalle Aït Mesan supra Arround circ. 2200 m.!

Ar. Geog. Europa omnis (exclusa arctica). Regio Mediterranea tota (inclusa Macaronesia).

GERANIUM ROTUNDIFOLIUM (*L., DC. Pr. i. 643*).

Mar. sept.—Cap. Spartel! Tetuan (1851)! In monte Beni Hosmar supra Tetuan!

Mar. occid.—Casa Blanca! Aïn el Hadjar haud procul Mogador!

Mar. merid.—Frequens in regione inferiore et in reg. media Atlantis Majoris. Mesfioua! Tasseremout, *J. D. H.*! Amsmiz! In convalle Aït Mesan 1300–1500 m.! In monte Djebel Tezah usque 1900 m., *J. D. H.*!

Ar. Geog. Europa fere omnis (exclusa arctica). Regio Mediterranea omnis (inclusa Macaronesia). Asia temperata.

— — —, var. FOLIIS PALMATIPARTITIS, *nob.*

Mar. occid.—Aïn el Hadjar haud procul Mogador!

Folia profundius divisa, segmentis oblongis obtusis.

G. DISSECTUM (*L., DC. Pr. i. 643*).

Mar. merid.—Sheshaona! Sektana! Amsmiz!

Ar. Geog. Europa omnis (exclusa arctica). Regio Mediterranea (inclusa Macaronesia). Asia temperata.

G. LUCIDUM (*L., DC. Pr. i. 644*).

Mar. sept.—In monte Beni Hosmar supra Tetuan!

Mar. merid.—In regione inferiore et subalpina Atlantis Majoris. In convalle Ourika circ. 1400 m.! In convalle Aït Mesan circ. 2200 m.!
In monte Djebel Tezah ad 2000 m.!

Ar. Geog. Europa omnis (exclusa arctica). Regio Mediterranea (inclusa Macaronesia). Siberia Uralensis.

G. ROBERTIANUM (*L., DC. Pr. i. 644*).

Mar. sept.—Tanger! In rupibus calcareis pr. Tetuan!

Ar. Geog. Europa omnis (exclusa arctica). Regio Mediterranea fere omnis (inclusa Macaronesia). Siberia Altaica. In America boreali advena?

— — — β , PURPUREUM (*DC. Prod. l. c.*), = *G. purpureum* (*Vill. Dauph. iii. 174*).

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Ourika!
In convalle Aït Mesan 1300–1400 m.!

Ar. Geog. Regio Mediterranea (inclusa Macaronesia). Sæpe occurrit in convallibus calidioribus Alpium Europæ. Est forma australis *G. Robertian* per reg. Mediterraneam cum specie obvia.

ERODIUM (*L'Hérit. in Ait. Hort. Kew. ed. 1, ii. 414*).

E. PETRÆUM (*Willd. Sp. Pl. iii. 625; DC. Pr. i. 645*), var. = *E. cheilanthifolium*, *Boiss. El. ?*

Mar. sept.—? In monte Beni Hosmar supra Tetuan, nondum florens et ideo ineertum!

Ar. Geog. Hispania. Gallia australis. Ditio Tetuanensis?

Monente eel. Boissier in 'Flora Orientali' (i. 887) hoc ab *E. trichomauefolio*, L'Hér., differt pube cinerea non glandulosa, foliis ambitu ovato-oblongis nec oblongo-lanceolatis, et petalis superioribus maculatis. Specimina nostra rhizomate et foliis exemplaribus Hispanicis *E. cheilanthifolii* sat congrua, sed segmenta foliorum magis acuta. Characteres *Erodiorum* hujus gregis non bene definiti sunt, et forsan plurimæ species auctorum in unam redueendæ.

E. ATLANTICUM (*Coss. in Bull. Soc. Bot. Fr. xx. 244*).

Descriptionem el. auct. proferre juvat:—

“Planta *perennis*, caudice crasso sæpius plurieipite, cæspitosa, e speciminibus pluribus suppetentibus *acaulis*, pedunculos radicales plures emittens. *Folia* omnia radicalia rosulata, *mollia*, *deuse pubescenti-subtomentosa*, ambitu ovata vel ovato-oblonga, basi cordata, longe petiolata, *lobata vel lobulata lobis obtusis* inæqualiter crenulato-dentatis. Stipulæ latæ ovatæ. *Pedunculi radicales* foliis vix vel subdimidio longiores, pubescentes *pilis longis patentibus* permixtis pilis brevissimis glandulosis, 4–7-flori, pedicellis elongatis. Involneri bracteolæ membranaceæ, ovato-suborbiculatæ, pubescenti-villosæ, pedicellis multoties breviores. Flores majusculi. *Sepala* longe pilosa, *oblonga abrupte mucronata mucrone subulato* (sepalorum mucronibus alabastra coronantibus), exteriora æqualiter 5-nervia. *Petala pallide purpurascencia* venis pieta, calyce subduplo longiora, ungue intus barbato, *subinæqualia*, superiora late obovata latiora, inferiora paulo longiora obovato-subeuneiformia. *Staminium fertilium filamenta* in parte inferiore membranaceo-dilatata, *edentula*, superne subulata, sterilium subdimidio breviora membranacea, lanceolata, inferne dorso pubescentia. Fructus rostrum 25–30 millim. longum, *carpellis oblongis inferne attenuatis*, pilis patulis hispidis, processu rostrali interne setis rigidulis sensim decreescentibus et superne evanidis donato, *foveolis apicalibus latis suborbiculatis, sub foveola plica destitutis*.”

Mar. merid.—In monte Djebel Orguis supra Imintenont legit Balansa anno 1867! In monte supra Seksaona circ. 1300 m.! In rupibus prope Milhau!

Monente cel. Cosson, “*E. Atlanticum* juxta *E. Boissieri*, Coss. in herb. (*E. asplenoides*, Boiss. Voy. Esp. 123. non Willd.; Boiss. Exsicc. Hisp.; Bourgeau, Pl. Esp. n. 1100; Pedro del Campo ed. Bourgeau, 1852,

n. 20, excl. syn. *Geranium asplenioides*, Desf. Atl. ii. 109, t. 168), quod sepalis mucronatis quoque gaudet, collocandum, sed distinctum foliis mollibus lobatis lobulatisve, non trisectis, pubescentia pedunculorum longiore, fructus rostro subdimidio brevior, carpellis minoribus. Habitu *E. montanum*, Coss. et DR. (in Bull. Soc. Bot. iii. 737), refert, sed differt foliis omnibus radicalibus minus profunde lobatis, sepalis abrupte mucronatis, non muticis, petalis subinæqualibus.”

ERODIUM CICUTARIUM (L., sub *Geranio*; *L'Hérit. in Ait. Hort. Kew.* ed. 1, ii. 414; *DC. Pr.* i. 646).

Mar. sept.—Tanger versus Cap. Spartel! Tetuan (1851)! In monte Beni Hosmar!

Ar. Geog. Europa omnis (exclusa arctica). Regio Mediterranea (inclusa Macaronesia). Asia media. In temperatis totius orbis introductum.

Species polymorpha mire modo ab auctoribus, et imprimis a b. Delile (Ind. Sem. Hort. Monspel. 1838), vexata; lusus enim fere innumeri, et limites vix aut ne vix definiendi. Viri graviores qui de lana caprina certare nolunt, et pro speciebus distinctis notas diagnosticas stabiliores quæsierunt, parum feliciter desudarunt. Natura invita a vinculis definitionum effugit. Ut sileam de forma et indumento foliorum, de colore et magnitudine petalorum, radix variat tenuis aut crassa, sæpissime annua, sed interdum perennans; carpella gaudent rostro brevior vel longior, glabro vel scabro; immo cotyledones ludunt cordatæ integræ vel trilobæ vel quinquelobæ. Ut morem gererem rei Botanices magistris formas eximias Maroccanas (attamen inter se ægre definiendas) hic enumeravi.

— —, var. γ , *DC. Pr. l. c.*, = *Geranium chærophyllum* (*Cav. Diss.* iv. 226, tab. 95, fig. 1), = *Erodium cicutarium* (*Willd. Sp. Pl.* iii. 629, non aliorum), = *E. Salzmanni*, *Del. loc. cit.*, = *E. viscosum* (*Salzm. Exsicc.* 1823). Et vix distinguendum *E. scandicinum*, *Del. loc. cit.*

Mar. sept.—Prope Tanger sat frequens! (*Salzm. Exsicc.*)!

Mar. occid.—Casa Blanca!

— —, var. *MACULATUM*, = *E. maculatum* (*Salzm. Exsicc.*), et verisimiliter = *E. atomarium* (*Del. loc. cit.*).

Mar. sept.—Prope Tanger sat frequens! (*Salzm. Exsicc.*)! Cap. Spartel!

— —, var. *BIPINNATUM*, = *Geranium bipinnatum* (*Cav. Diss.* iv. 273, tab. 126), = *G. numidicum* (*Poir. Voy.* ii. 201), = *G. æthiopieum* (*Lam. Dict.* ii. 655, errore quoad nomen specificum; conf. *Poir. l. c.*).

Mar. occid.—In arenosis pr. Mogador! Aïn el Hadjar! Prov. Haha pr. Mogador! Mazagan!

Mar. merid.—Frequens in saxosis regionis inferioris. Mskala! Seksaoua! Mtouga!

Hæc est forma solennis regionis ealidioris. Distinguitur statura minore, foliis sæpino bipinnatifidis laciniis linearibus obtusiusculis glabris vel puberulis, pedunculis sæpissime, præsertim in maritimis, bifloris, interdum 3-6-floris, rostro carpellorum tenui elongato glabriusculo vel scabro, petalis minoribus subæqualibus colore saturatiore. Specimen unicum prope Mazagan lectum, a cæteris valde diversum, videtur esse lusum ex solo uberiore natus.

Erodium Jacquinianum (*Fisch. et Mey. Ind. Sem. H. Petrop. ix. Suppl. p. 11; Boiss. Fl. Or. i. 890*), = *E. staphylinum* (*Bert. Fl. It. vii. 185*), = *E. tenuisetum* (*Gren. et God. Fl. Fr. i. 311*).

Facies valde proximum *E. cicutario*, var. *bipinnato*, quocum Bertoloni, loc. cit., conjunxit, sed dignoscitur carpellorum foveolis plicæ destitutis et cotyledonibus pinnatifidis. Specimina nostra tamen accurate scrutata incertum me fecerunt. In *E. cicutario* typico foveola carpellorum subtus plicæ concentrica prædita est. In varietatibus meridionalibus, et præsertim in var. *bipinnato*, plicæ externa minus profunda et interdum subobsoleta. In *E. Jacquiniano* plicæ omnino obsoleta. Non desunt tamen exemplaria incertæ sedis, et inter specimina ex regione montana (Aït Mesan) adest exemplar carpellis fere exacte *E. cicutarii*, scil. plicæ manifesta sub foveola notatis.

Mar. occident.—Prope Mogador! et præsertim in Insula!

Mar. merid.—In convalle Aït Mesan a 1400 m. usque 2400 m.!

Ar. Geog. Occurrit sparsim in regione Mediterranea calidiore (exclusa Macaronesia).

Specimina ex regione montana ostendunt foliorum lacinias latiores, et aspectu *E. cicutario* typico propius accedunt.

E. ROMANUM (*L.*, sub *Geranio*; *Willd. Sp. Pl. iii. 630; DC. Pr. i. 647*)?

Mar. sept.—In monte Djebel Dersa prope Tetuan (*Webb Herb.*)!

Specimina nondum fructifera mihi incerta. Pedunculi 6-10-flori. An foret *E. cicutarii*, var. *maculati* lusum in statu juniore?

Ar. Geog. *E. romanum* in Gallia australi et Italia frequens, fit rarius in Græcia et in Oriente. Decet in peninsula Iberica, et in Macaronesia.

E. MOSCHATUM (*L.*, sub *Geranio*; *Willd. Sp. Pl. iii. 361; DC. Pr. i. 647*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Cap. Spartel! visum prope Tetuan!

Mar. occident.—Aïn el Hadjar haud procul Mogador!

Mar. merid.—In rudertis urbis Marocco!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia). Europa occidentalis ex Hibernia usque fretum Herculis. Abyssinia. In America australi advena.

——, var. *DISSECTUM*, foliorum segmentis pinnatifidis lobis dissectis.

Mar. merid.—In prov. Shedma !

ERODIUM CICONIUM (*L.*, sub *Geranio* ; *Willd. Sp. Pl.* iii. 629 ; *DC. Pr.* i. 646).

Mar. occid.—Casa Blanca !

Mar. merid.—Prov. Shedma prope Aïn Oumast ! Mtouga ! Seksaoua !

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

E. BOTRYS (*Cavan. Diss.* iv. tab. 90. fig. 2 ; *Bert. Amœn. Ital.* 35 ; *DC. Pr.* i. 647).

Mar. sept.—Tanger !

Ar. Geog. Regio Mediterranea occidentalis ex Italia usque Macaronesian. Ager Byzantinus, nec alibi in Oriente.

E. LACINIATUM (*Cavan. Diss.* iv. 228, tab. 113. fig. 3, sub *Geranio* ; *Willd. Sp. Pl.* iii. 633 ; *DC. Pr.* i. 646).

Mar. occid.—Mogador ! In regione litorea prov. Ilaia (mense Aprili) ! Aïn el Hadjar (mense Junio) !

Ar. Geog. Regio Mediterranea calidior (inclusa Macaronesia).

E. pulverulentum (*Cavan. l. c.* tab. 125. f. 1) est lusus hujus speciei quoad formam foliorum et indumentum valde ludibundæ. Specimina mense Aprili lecta omnia pulverulenta. Quæ legimus ineunte Junio post imbres mensis præcedentis ostendunt folia ex ramis novellis enata viridia, vix puberula. Radix interdum perennans. Folia interdum eis *E. botryis* simillima. Carpellorum foveola semper plica destituta.

E. MALACOIDES (*L.*, sub *Geranio* ; *Willd. Sp. Pl.* iii. 639 ; *DC. Pr.* i. 648).

Mar. sept.—Tanger (*Webb Herb.*) ! Inter Tanger et Tetuan ! Prope Tetuan visum !

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—Occurrit passim in regione inferiore et ad radices Atlantis Majoris. Prov. Shedma pluribus locis ! In ruderatis urbis Marocco ! Mesfioua ! Reraya ! In convalle Aït Mesan circ. 1300 m. ! Amsmiz ! Seksaoua ! Mtouga !

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia). Gallia occidentalis.

E. CHIUM (*L.*, sub *Geranio* ; *Willd. Sp. Pl.* iii. 634 ; *DC. Pr.* i. 647).

Mar. sept.—Tanger ! Cap. Spartel ! Tetuan !

Ar. Geog. Regio Mediterranea, imprimis occidentalis (inclusa Macaronesia). Deest in Asia Mediterranea. Formæ arete affines in Ægypto et in Palæstina.

E. litoreum (*Léman, in DC. Fl. Fr.* iv. 843, et *Prod.* i. 648).

Mar. sept.—Inter Tanger et Tetuan !

Mar. occid.—Aïn el Hadjar haud procul Mogador !

Mar. merid.—In prov. Shedma prope Castellum Gubernatoris !

Ar. Geog. Gallia australis. Hispania. Africa boreali-occidentalis.

Hoc pro subspecie præcedentis enumeravi. Distinguitur foliis profundius divisis lobis angustioribus. Nota diagnostica ex filamentis staminum sterilium *ciliatis* ut fertur in *E. chio*, *glubris* in *E. malacoidi* et *E. litoreo*, mihi plane instabilis videtur. In fructu nullum discrimen detegere potui.

ERODIUM GLAUCOPHYLLUM (L., *Sp. Pl.* 952, et *Cavan. Diss.* iv. 221, tab. 92. fig. 2, sub *Geranio* ; *Ait Hort. Kew.* ed. 1, ii. 416 ; *DC. Pr.* i. 648).

Mar. merid.—In planitie saxosa aridissima inter Sheshaoua et Aïn Oumast legit cl. G. Maw !

Ar. Geog. Regio desertorum ex Arabia et Mesopotamia per Ægyptum et Algeriam usque planitiem Maroccanam.

In iconc Cavanillesii citata et in plurimis speciminibus a nobis visis folia subintegra crenata. Specimina Maroccana referunt plantam vegetiorem subbipedalem foliis majoribus longioribus, sæpius subtrilobis donatam. Carpella apice plica duplici cincta.

E. GUTTATUM (*Desf. Fl. Atl.* ii. 113, tab. 169, sub *Geranio* ; *Willd. Sp. Pl.* iii. 636 ; *DC. Pr.* i. 648).

Mar. merid.—Prov. Shedma, in saxosis aridis pr. Aïn Oumast ! Seksaoua ! In convalle Aït Mesan in reg. inferiore Atlantis Majoris circ. 1200 m., *G. Maw spec.* !

Ar. Geog. Africa borealis. Hispania Granatensis.

Facie priori subsimile, sed plurimis notis probe distinctum. Carpella non vere foveolata, in spec. Maroccanis undique favosa et setis rigidis fulvis obsita. In spec. Algeriensibus carpella video plus minusve rugosa et setas pauciores.

OXALIS (*L. Gen.* 582).

O. CORNICULATA (L., *DC. Pr.* i. 692).

Mar. sept.—Inter Tanger et Tetuan ! Tetuan (1851) !

Mar. merid.—In rudcratis urbis Marocco ! In convalle Aït Mesan circ. 1800 m., *J. D. H.* !

Ar. Geog. Herba amphigea per Europam mediam, regionem omnem Mediterraneam, necnon per Indiam orientalem et Americam borealem calidiorem late diffusa.

In territorio Maroccano admodum rara.

O. SERICEA (*L. fl, Suppl.* ; *DC. Pr.* i. 695), var. *PILOSA*, nob. = *O. cernua* (*Thunb. Diss. Ox.* tab. 2. fig. 2), = *O. libyca* (*Viv. Fl. Lib. spec.*).

Mar. sept.—In vicinitate urbis Tanger, certe advcna, hodie facta spontanea ! Occurrit sæpissime flore pleno.

Ar. Geog. In Africa australi orta per regionem Mediterraneam plurimis

locis diffusa, seil. in insul. Canariensibus, Madeira, Corsica, Zacynthio, pr. Smyrnam, in agro Tingitano, et alibi in Africa boreali.

Oxalis sericea, L. fil., *O. cernua*, Thunb., et *O. compressa*, Jacquin, sunt stirpes Capenses arcte affines. Prima dignoscitur indumento brevissimo, herba subglabra, stylis abbreviatis; secunda indumento sericeo-piloso, stylis longitudine intermediis; tertia indumento sericeo-piloso stylis elongatis. Qui seriem formarum scrutabit facile has in unam speciem conjungendas fatebitur. Characteres quibus *O. libyca* distinguitur omnino fallaces. Bulbilli e collo radicis orti non desunt (etsi rari) in speciminibus Capensibus. Calli versus apicem sepalorum non semper adsunt in exemplaribus Mediterraneis, nec semper desunt in spec. Capensibus.

RUTACEÆ (*Juss. Gen.* 296).

RUTA (*L. Gen.* 523).

R. MONTANA (*Clus. Hist.* ii. 136; *Vill. Dauph.* iii. 582; *DC. Pr.* i. 710), = *R. tenuifolia* (*Desf. Fl. Atl.* i. 836).

Mar. occid.—Prope Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Ad radices Atlantis Majoris, Ourika! Seksaoua! Tassgittelt (*R. & Fr. spec. in Herb. Kew.*)! et in planitie; prov. Shedma!

Ar. Geog. Regio Mediterranea fere omnis, exclusa Macaronesia.

R. CHALEPENSIS (*L. Mant.* i. 69), var. α , = *R. angustifolia* (*Pers. Syn.* i. 404; *DC. Pr.* i. 710).

Mar. sept.—Prope Tetuan in rupibus! et in monte Beni Hosmar! ubi eam olim legit P. B. Webb.

Mar. merid.—In regione media Atlantis Majoris a 1600 m. usque 2400 m.! In rupibus Seksaoua! Aït Mesan!

Ar. Geog. Regio Mediterranea, imprimis occidentalis. Deest in Oriente (*Boiss. Fl. Or.* i. 922) et in Macaronesia.

— — —, var. β , BRACTEOSA (*Boiss. loc. cit.*), = *R. bracteosa* (*DC. Pr.* i. 710).

Mar. sept.—Prope Tanger in monte Djebel Kebir, *J. D. H.*! Inter Tetuan et Ceuta!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia). Non desunt formæ inter has varietates mediæ. Spec. nostrum ex Seksaoua sub var. α enumeratum forsitan melius sisteret var. *intermediam*.

HAPLOPHYLLUM (*A. Juss. Mem. Mus.* xvi. 464).

H. BROUSSONNETIANUM (*Coss. in Bull. Soc. Bot. Fr.* xx. 244).

Descript. in verbis cel. auctoris hue refero.

“Radix simplex vel parce ramosa, descendens, indurato-perennans. Caules subsolitarii vel plures, erecti, simplices vel superne parce ramosi, inferne glabrescentes, superne vix pubescentes et sparsim ver-

ruculoso-glandulosi. *Folia* subcoriacea, conferta, glabriuscula, *omnia indivisa*, oblongo-obovata, inferne attenuata, obtusa, viridia, subtus pallidiora, punctato-glandulosa. *Inflorescentia* pluriflora cymosocorymbiformis, *laxiuscula* ramis patulis. Calyx deciduus, minutus. 5 partitus, lobis ovatis obtusiusculis, dorso et præsertim ad marginem pubescentibus. Petala (ex alabastro deprompta) flava, ovato-suborbiculata, glandulis minimis sparsis punctulata, obsolete trinervia. Filamenta libera, in alabastro ultra medium complanato-liguliformia. *Ovarium* in disco brevissimo annulari supra partem connatam carpidorum expanso insidens, e carpellis tot quot sepala efformatum, glabrum, haud verrucosum, vix glanduloso-punctatum, *carpellis 4-ovulatis*. Stylus filiformis, carpellorum partem liberam longe excedens. *Capsula* campanulato-subglobosa, lævis, *carpellis* superne liberis, *in parte libera divergentibus*, *acuminato-rostratis* rostro brevi *arcuato-patente*, introrsum dehiscen- tibus. Semina (immatura tantum nota) compressa, reniformia, reticulato-rugosa.”

Mar. occid.? —“ Mogador ” (*Broussonnet in Herb. Bouchet, ex Cosson, l. c.*).

Mar. merid.—Ad radices Atlantis Majoris ad Keira legit fructiferum mense Majo 1867 Balansa.

Monente cel. auctore, “ *H. Broussonnetianum* carpellis 4-ovulatis, dehiscentibus, rostro eglanduloso terminatis, foliis indivisis, filamentis liberis juxta *H. pumilum*, Boiss. (*Diagn. Or. ser. 1, i. 61, et Fl. Or. i. 928*), collocandum, sed ab illo et cæteris speciebus affinibus distinctissimum carpellis divergentibus acuminato-rostratis rostro patente.”

Certum me feci b. Broussonnet plantas nonnullas e terris interioribus habuisse, et cum exemplaribus in agro Mogadorensi lectis semel commiscuisse. Locus natalis *H. Broussonnetiani* Mogadorensis a botanicis posterioribus non verificatus mihi valde incertus videtur.

PEGANUM (*L. Gen. 601*).

P. HARMALA (*L., DC. Pr. i. 712*).

Mar. occid.—Prope Mogador ! (*Lowe Cat.*).

Mar. merid.—Frequens in regione inferiore præsertim in planitie arida. Prov. Shedma ! visum in vicinitate urbis Marocco ! in prov. Mtouga ! et alibi.

Ar. Geog. Regio Mediterranea fere omnis, deest tamen in Italia continentali, in Gallia australi et in Macaronesia ; deinde ex Hungaria per Rossiam meridionalem in Asiam mediam diffusum.

Herba, ut videtur, locustis invisâ. Post transitum insectorum voracium inter cæteras stirpes a foliis prorsus denudatas, hanc solam illæsam pluries vidimus.

CELASTRINEÆ (*R. Br. Gen. Rem. 22*).CELASTRUS (*L. Gen. 270*).

C. SENEGALENSIS (*Lam. Dict. i. 661* ; *DC. Pr. ii. 8*), var. *EUROPÆUS*, = *C. europæus* (*Boiss. El. 46* ; *Walp. Rep. i. 533*), = *Catha europæa* (*Boiss. Voy. 127, tab. 33*).

Mar. occid.—In monte Djebel Hadid haud procul Mogador !

Mar. merid.—Ad radices Atlantis Majoris prope Tasseremout ! Ourika !

Ar. Geog. Species per regiones tropicales orbis veteris, ex India orientali usque oras Atlanticas Africæ æquatorialis late diffusa. Inde, ut videtur, per territorium Maroccanum Hispaniam australem attingit.

Cl. Oliver, in *Flor. Trop. Africa*, identitatem hujus et plurimarum specierum ab auctoribus descriptis cum *C. senegalensi*, *Lam.*, primus indicavit ; species valde ludibunda est, sed habitus semper fere idem.

RHAMNEÆ (*R. Br. Gen. Rem. 22*).ZIZYPHUS (*Juss. Gen. 380*).

Z. LOTUS (*L.*, sub *Rhamno* ; *Lam. Dict. iii. 316* ; *DC. Pr. ii. 19* ; *Schousb. Gew. Marok. 87*).

Mar. occid.—In prov. Haha haud procul Mogador ! sed in reg. litorali non obviat.

Mar. merid.—In saxosis incultis reg. inferioris fere ubique. Legimus pluries in prov. Shedma ! prope urbem Marocco ! et alibi !

Ar. Geog. Africa borealis. In Europa australi calidiore (Græcia, Sicilia, peninsula Iberica) rarius occurrit. In Arabia australi ad Mare Rubrum (*Schweinfurth*).

In incultis aridissimis, ubi vix ulla alia planta arborea, nutrimentum sibi consequitur, et dum incolarum incuria vitam concedit, fit arbor 20-pedalis, trunco crasso, diametro bipedali et ultro.

RHAMNUS (*L. Gen. 265*).

R. ALATERNUS (*L.*, *DC. Pr. ii. 23* ; *Schousb. Gew. Marok. 87*).

Mar. sept.—In monte Djebel Kebir pr. Tanger ! (*Schsb. l. c.*).

Mar. occid.—Prope Mogador (*Lowe Cat.*).

Mar. merid.—Frequens in regione inferiore Atlantis Majoris. Legimus in convalle Ourika ! in convalle Amsmiz ! Aït Mesan circa 1400 m. !

In monte Djebel Tezali 1600 m. !

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

R. OLEOIDES (*L.*, *DC. Pr. ii. 24* ; *Schousb. Gew. Marok. 87*).

Mar. sept.—Tanger (*Webb Herb.*) ! Prope Cap. Spartel ! In rupibus calcareis juxta Tetuan ! In monte Beni Hosmar ! (*Webb Herb.*) !

Ar. Geog. Regio Mediterranea a Græcia usque peninsulam Ibericam et Imperium Maroccanum. Deest in Oriente et in Macaronesia.

— —, var. *ANGUSTIFOLIA* (*Lange, MSS. in Herb. Kew.*). Differt a typo habitu toto, ramis longioribus minus divaricatis, foliis longioribus. Cum spec. Hispanico a cl. Lange misso omnino convenit.

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Ourika !
Distr. Reraya circa 1200 m. !

Ar. Geog. Forma Hispanica et Maroccana.

— —, var. *AMYGDALINUS*, = *B. amygdalinus* (*Desf. Fl. Atl. i. 198 ; DC. Pr. i. 25*).

Differt foliis crassioribus, coriaceis, subtus non reticulatis, floribus binis in quovis fasciculo.

Mar. occid.—In monte Djebel Hadid ! et in collibus prov. Haha haud procul Mogador !

Facie sp. Canariensi (*R. integrifolio, DC.*) haud absimilis.

Hanc formam nonnisi ex Africa boreali vidi.

RHAMNUS LYCIOIDES (*L., DC. Pr. ii. 25*).

Mar. merid.—Ad radices Atlantis Majoris. In convalle Amsmiz circ. 1300 m. ! Seksaoua !

Ar. Geog. Hispania. Africa boreali-occidentalis.

AMPELIDEÆ (*Humb., Bonpl. et Kth. Nov. Gen. v. 223*).

VITIS (*L. Gen. 284*).

V. VINIFERA (*L., DC. Pr. i. 633*).

Hodie ab incolis parce et rarius culta, verisimiliter olim a Romanis ubique in Mauritania introducta. Ad radices Atlantis Majoris pluribus locis, e. g. prope Amsmiz ! et Gurguri ! videtur spontanea in dumetis et saxosis.

Ar. Geog. Ex Asia occidentali orta, per regionem Mediterraneam diu culta, et sæpissime subsponsanea facta est.

SAPINDACEÆ (*Juss., B. et H. Gen. Pl. i. 389*).

Subordo *ACERINEÆ* (*B. et H. l. c. 391*).

ACER (*L. Gen. 1155*).

A. MONSPESSULANUM (*L., DC. Pr. i. 594*).

Mar. merid.—In regione media Atlantis Majoris, in convalle Aït Mesan a 1600 m. usque 1800 m. !

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia). Ramis occurrit in calidioribus Europæ mediæ.

Spec. nostra flore et fructu carent, sed folia et rami omnino conformes.

ANACARDIACEÆ (*Lindl. Introd. ed. ii. 166*).

PISTACIA (*L. Gen. 1108*).

P. ATLANTICA (*Desf. Fl. Atl. ii. 364 ; DC. Pr. ii. 64*).

Mar. merid.—(*Schousb. in Herb.*) *Coss. Index*. Vidimus sat frequentem

in regione inferiore; sæpius fit arbuseula 12-18-pedalis, sed in locis humidis juxta fontes, ubi verisimiliter culta, fit arbor grandis; e. g. ad Aïn Beida, trunci diametro 2-3-pedali!

Pistaciæ e grege *P. Terebinthi*, L., extricatu difficiles, et forsā melius cognitæ plurimæ ad varietates reducendæ. *P. atlantica* genuina est Africæ borealis indigena civis. Planta Græca (monente cel. Boissier) ad *P. muticam*, F. M., referenda est.

PISTACIA LENTISCUS (L., DC. Pr. ii. 65).

Mar. sept.—Frequens in maritimis. In monte Djebel Kebir prope Tanger! Prope Tetuan! et versus Ceuta! pluribus locis visa.

Mar. occid.—Mogador (*Lowe Cat.*). In monte Djebel Hadid!

Mar. merid.—Ad radices Atlantis Majoris et in convallibus calidioribus. Prope Ourika! Aït Mesan usque 1600 m.!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

RHUS (*L. Gen.* 369).

R. OXYACANTHA (*Cavan. Anal. Cienc. Nat.* iii. 36), = R. oxyacanthoides (*Dum. Cours Bot. Cult.* vi. 662), = R. dioica (*Brouss. in Willd. Enum.* 325; DC. Pr. ii. 71).

Mar. occid.—In agro Mogadorensi (*Broussonet, ex DC. et Willd. l. c.*). In monte Djebel Hadid fructiferam legimus incunte Junio!

Specimina nostra ad formam typicam foliis viridibus trifidis, foliolorum dentibus obtusis, cortice fusco, referenda.

—— —, var. foliolis omnibus profunde et acute dentatis.

Mar. merid.—Specimen unicum prope Seksaoua legit J. D. II.!

—— —, var. ZIZYPHINA, = R. zizyphina (*Tin. Pl. Rar. Gen. Sic. Pug.* p. 8).

Mar. occid.—Prope Agadir (*Coll. Indig.* 1871)!

—— —, var. ALBIDA, = R. albida (*Schousb. Gew. Mar.* 128; DC. Pr. ii. 70), = R. oxyacantha (*Schousb. in Act. Soc. Hafn.*; DC. Pr. ii. 71, non Cav.), = R. cratægiformis (*Pers. Syn.* i. 362).

Mar. occid.—Prope Mogador in arenosis! (*Lowe*, sub *R. oxyacantha*, spec. in herb. Kew. nostro conforme)!

Ar. Geog. Africa boreali-occidentalis. Varietates in Sicilia et in Oriente occurrunt.

R. oxyacantha est species summopere variabilis, per regionem Mediterraneam calidiorem late dispersa, nullibi frequens. In Europa nonnisi in Sicilia; in Macaronesia in sola insula Teneriffa detecta. Monente ipso Schousboe *loc. cit.* var. *albida* nostra, quæ primo aspectu valde diversa foliolis leviter crenatis, subintegerrimis, canis, cortice albido, ramis inermibus, culta in Horto Hafniensi “exuit canitiem et tomentum, foliaque evaserunt glabra et magis crenata quam in planta spontanea.”

RHUS PENTAPHYLLA (*Desf. Fl. Atl.* i. 267, tab. 67; *DC. Pr.* ii. 72).
 Mar. occid.—Prope Mogador (*R. et Fr. in herb. Kew.*)!
 Mar. merid.—In regione inferiore haud infrequens. Prov. Shedma!
 Mzouda! Prope Kcira, *J. D. H.*! Seksaoua!
Ar. Geog. Africa borealis. Sicilia.

CORIARIÆ (*DC. Prod.* i. 739).

CORIARIA (*L. Gen.* 458).

C. MYRTIFOLIA (*L., DC. Pr.* 739).
 Mar. sept.—Tanger! (*Schsb.*) *Coss. Index.*
Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia),
 Græciæ dubia civis, forsân errore indicata?

LEGUMINOSÆ (*Juss. Gen.* 345).

ANAGYRIS (*L. Gen.* 509).

A. FÆTIDA (*L., DC. Pr.* ii. 99).
 Mar. occid.—Prope Agadir (*Coll. Indig.* 1871)!
 Mar. merid.—Sat frequens in regione inferiore. Prov. Shedma! Aïñ
 Tarsil! Prov. Haha! Ad ruinas urbis Tasagkimat (*Schousb. Herb.*)
Coss. spec.!
 — —, var. seu lusus foliolis latioribus.
 Mar. merid.—Specimen unicum sine flore vel fructu legit G. Maw pr.
 Tasseremout, fere media inter typum et *A. latifoliam* (Brouss. in Willd.
 Enum. 489) stirpem Canariensem quæ mihi ad varietatem reducenda
 videtur.
Ar. Geog. Regio Mediterranea calidior. Forma arete affinis in Maca-
 ronesia.

LOTNONIS (*Eckl. et Zeyh. Enum.* p. 179; *DC.* sub *Ononide*).

L. MAROCCANA (*Ball, in Journ. Bot.* 1873, 302). (Plate XV.)

Planta perennis, ramosa, prostrata; herba tota sericea; folia petiolata,
 trifoliolata; foliola obovata, apiculata, basi cuneata, brevissime stipi-
 tata; stipulæ 1–2 foliolatæ; flores sæpissime solitarii, brevissime pe-
 dicellati; calyx demum campanulatus, purpurascens, basi striis validis
 notatus; vexillum alas longiuscule superans, extus cum carina insig-
 niter sericeum; legumina latiuscula, glabra, 8–10-ovulata.
 Mar. merid.—In regione inferiore Atlantis Majoris prope Tasseremout!
 Ourika! Aït Mesan! Amsmiz! a 1000 m. ad 1400 m.
 Species probe distincta, a *L. lupinifolia*, Boiss. (sub *Leobordea*), dif-
 fert imprimis foliis tri- nec quinquefoliolatis; a *L. cytisoide* toto
 habitu et florum structura recedit; in hac nempe vexillum carinæ
 subæquale, in nostra carinam et alas longe superat.

LUPINUS (*L. Gen.* 865).

L. HIRSUTUS (*L., DC. Pr.* ii. 407).

Mar. sept.—Tanger! (*Webb Herb.*)! Inter Tanger et Tetuan!

Ar. Geog. Regio Mediterranea (exclusa Macaronesia).

L. PILOSUS (*L., DC. Pr.* ii. 407).

Mar. merid.—In glareosis torrentis Onrika! Amsmiz!

Ar. Geog. Oriens. Africa borealis. Deest in Europa austro-occidentali. (conf. *Boissier, Fl. Or.* ii. 27). Species arcte affinis (*L. varius*, *L.*) in peninsula Iberica.

L. pilosus ad subspeciem *L. varii* reducendus videtur, dum notæ diagnosticæ parum stabiles. Legumina in spec. nostris 3–4-sperma, semina minute asperula non scabrida.

L. ANGUSTIFOLIUS (*L., DC. Pr.* ii. 407).

Mar. sept.—Tanger! (*Webb Herb.*)!

Ar. Geog. Regio Mediterranea omnis. In Madeira obvius, sed in Ins. Canariensibus non indicatus.

Pedicelli non vere ebracteolati ut fertur, bracteolæ enim semper adsunt ad squamas membranaceas cito marcescentes reductæ. *L. linifolius*, Roth, dignoscitur floribus minoribus bracteolatis, calycis labio inferiore subtrifido, nec ut in *L. angustifolio* integro, et insuper seminibus multo minoribus globulosis maturitate badiis umbrino et flavo variegatis, nec maculis albidis vel flavis variegatis. Cel. Boissier tamen (in *Fl. Or.* ii. 29) *L. angustifolio* calycis labium inferius sub-2–3-denticulatum, semina grisea maculis albis lincisque irregularibus fulvis marmorata, tribuit; dum *L. linifolium*, Roth, pro synonymo *L. reticulati*, Desv., habet. In hac specie (sec. Boissier *l. c.*) labium calycis inferius subintegrum, semina ovato-globosa albida nigro-reticulata, maculis saturationibus nigro-punctulatis obsita. Grenier et Godron (*Fl. Fr.* i. 366) tres species Europæas hujus gregis adesse suspicantur, et *L. linifolium* (Roth et Guss. *Fl. Sic.* ii. 269) a *L. reticulato*, Desv., differre credunt seminibus badiis vix nigro-maculatis. In spec. nostris vix florentibus characteres a seminibus latent. Labium calycis inferius superiore multo longius basi utrinque denticulo parvo auctum est. *Lupini* plerique cum seminibus cerealibus per regionem Mediterraneam dispersi, soli et cœli vicissitudini obnoxii, valde variabiles sunt, et characteres stabiles difficillime detegendi.

L. LUTEUS (*L., DC. Pr.* ii. 407).

Mar. sept.—Prope Tanger et in monte Djebel Kebir! (*Webb Herb.*)! Inter Tetuan et Ceuta!

Ar. Geog. Peninsula Iberica. Italia. Africa borealis. In Madeira advena.

ARGYROLOBIUM (*Eckl. et Zeyh. Enum.* 184).

A. LINNÆANUM (*Walp. in Linnæa*, xiii. 506), = *Cytisus argenteus* (L., *DC. Prod.* ii. 156).

Mar. sept.—In rupibus calcareis pr. Tetuan! In monte Djebel Dersa pr. Tetuan (*Webb Herb.*)!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

—— —, var. FALLAX, = *A. fallax* (*Ball, in Journ. Bot.* 1873, 302).

Stirpem hanc pro subspecie *A. Linnæani* loc. cit. enumeravi; hodie mihi videtur melius ad varietatem reducenda. Differt a typo floribus minoribus, foliorum foliolis latioribus obtusioribus, et præsertim calycis labio inferiore inæqualiter trifido, lacinia media lateralibus multo angustiore, nec subæquali, fere ut in *A. calycino*, M. B. (sub *Cytiso*). Semina in quovis legumine 6–10, sæpissime 8.

Mar. merid.—Legimus in convalle Aït Mesan sat frequens a 1400 m. ad 1600 m. Specimen unicum legi supra Arround circ. 2100 m.

A. stipulaceum (*Ball, in Journ. Bot.* 1873, 302).

Fruticulus nanus, suberectus, ramosus; folia primaria longe petiolata; flores brevissime pedicellati, solitarii seu geminati in vertice ramulorum; stipulæ et bracteolæ ad basin calycis majores et latiores quam in *A. Linnæano*, cito marcescentes; calyx basi laxus, labii inferioris laciniiis anguste lanceolatis subæqualibus, labio superiore paullo longioribus; legumen subtorulosum, 5–6-spermum; semina flava.

Mar. merid.—In regione inferiore Atlantis Majoris. Prope Seksaoua! et verisimiliter huc referenda spec. nimis incompleta prope Tasseremout et in convalle Amsmiz lecta!

Forsan pro specie omnino distincta vindicanda erit, sed characteres citati in hoc genere instabiles videntur.

A. MICROPHYLLUM (*Ball, in Journ. Bot.* 1873, 303).

Perenne, 3–6-pollicare; herba tota pube brevi rigida incana; pedunculi uniflori, axillares et terminales, folia parva breviter petiolata superantes; stipulæ minimæ; calyx coloratus, ad basin usque bisectus, labio superiore profunde bifido, inferiore ad quartam partem tridentato, dente medio subulato lateralibus lanceolatis quidquam brevioribus; flores parvi, calycem parum excedentes, bicolores (vexillo fulvo, carina citrina); legumen 4–5-spermum.

Mar. merid.—In regione inferiore Atlantis Majoris. Supra Seksaoua circa 1100 m.!

Argyrolobio nullo, nisi forte *A. unifloro*, Denc. (sub *Cytiso*), comparandum; sed ab hoc eximie differt calycis structura et floribus duplo majoribus.

ADENOCARPUS (*DC. Fl. Fr. Suppl.* 549).

A. TELONENSIS (*Lois. Fl. Gall.* ed. i. 446, sub *Cytiso*; *Rob. Cat. Toulon*, 24; *Boiss. Voy. Esp.* 418, tab. 43; *DC. Fl. Suppl.* 550, ex parte),

= *A. grandiflorus* (*Boiss. Bibl. Gen.* 1836; *Gren. et God. Fl. Fr.* i. 363).

Mar. sept.—In saxosis ealearcis supra Tetuan!

Ar. Geog. Gallia Mediterranea. Hispania. Ditio Tetuanensis.

III. A. P. De Candolle hanc speciem cum *A. commutato*, Guss., commisit; sed nomen triviale *telonensis*, a Loiseleur jam primum propositum, non ideo mutandum. Nomen *grandiflorus* infaustum, dum flores non raro minores quam in *A. divaricato*, L'Hér. Cel. Boissier (*Fl. Or.* ii. 34) optimo consilio sub *A. divaricato*, L'Hér., formas plurimas nulla nota stabili distinguendas conjunxit. Nomen Lamarekianum *parvifolius* recentius est et omnino ineptum, dum folia magnitudine variabilia.

ADENOCARPUS ANAGYRIFOLIUS (*Coss. in Balansa, Pl. Maroc.* 1867, et in *Bull. Soc. Bot. Fr.* xx. 240).

Sequitur descriptio Cossoniana:—

“*Frutex* elatus, ramosus, dumosus, *glaber*, ramis divergentibus vix striato-angulatis cortice pallide virente. *Folia* Anagyrin foetidam referentia, in genere *ampla*, longe petiolata, trifoliolata, *glabra*, *foliolis nitidulis* oblongis *vix complicatis*, mucronulatis, nervo medio subtus prominente. *Stipulae* citissime deciduae, in speciminibus suppetentibus evanidae. *Racemi* eanlem et ramos terminantes, *laxiusculi*, demum elongati, rachii puberulo-sericea demum glabra, *pedicellis* erecto-patulis calyce paulo longioribus *bibracteolatis bracteolis citissime deciduis*. *Calyx* puberulo-sericeus, *eglandulosus*, campanulato-obconicus bilabiatus, *labio* superiore bipartito lobis ovato-triangularibus, *inferiore apice tridentato dentibus brevibus* lanceolato-linearibus subaequalibus vel medio paulo longiore. Corolla lutea, vexillo ascendente obovato-suborbiculato extus pubescenti-sericeo, alis carinaque subaequilongis glabris. Ovarium lineare, glandulis subpedicellatis obsitum. Stylus filiformi-subulatus basi crassior, supra basin abrupte areolato-refractus. Stigma minutum, terminale, capitellato-depressum. Legumen 3–4 centim. longum, pedicello multoties longius, lineari-oblongum, plano-compressum, subtorulosum, glandulis fuscescentibus subpedicellatis obsitum.”

Mar. merid.—Satis frequens in convallibus Atlantis Majoris, unde in planitiem in glareosis torrentium descendit. Balansa hanc stirpem insignem primum detexit in glariosis Oued Ghagaia. Legimus in convalle Aït Mesan a 1400 m. usque 1800 m.! In convalle Amsniz! prope Ourika! et prope pagum Amsniz! Ex montibus Djebel Ouensa et Djebel Afougneur attulit indigenus Ibrahim et specimina mihi benev. comm. eel. Cosson.

Ab *A. divaricato*, L'Hér., insigniter differt foliis glabris nitidulis, pro genere amplis, *Anagyrin foetidam* referentibus, dentibus labii calycini inferioris brevioribus, et praesertim racemis elongatis (4–8-pollicariibus) ramis strictis erectis, vix striato-angulatis, glabris.

CALYCOTOME (*Link in Schrad. Journ.* ii. pars 2, p. 50).

C. VILLOSA (*Poir. Voy.* ii. 207, sub *Spartio*; *Link. loc. cit.*), = *Spartium lanigerum* (*Desf. Fl. Atl.* ii. 135), = *Cytisus lanigerus* (*DC. Pr.* ii. 154).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Prope Tetuan!

Ar. Geog. Regio Mediterranea ealidior (exclusa Macaronesia).

—— —, var. INTERMEDIA, *nob.*, = *C. intermedia* (*Salzm. Exsicc.* sub *Cytiso*; *Presl, Bot. Bemerk.* 51; *Walp. Ann.* i. 223).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Versus Cap. Spartel! Inter Tanger et Tetuan! In monte Beni Hosmar pr. Tetuan (*Webb Herb.*)!

Nimis vicina præcedenti, vix sicut subspecies enumeranda.

GENISTA (*L. Gen.* 859; *Benth. et H. fil. Gen. Pl.* i. 482).

G. MONOSPERMA (*Lam. Dict.* ii. 616; *DC. Pr.* ii. 150; *Boiss. Voy. Esp.* p. 144, sub *Retama*), var. CLUSII, *nob.*, = *Spartium Clusii* (*Spach, Ann. Sc. Nat.* ser. 2, xix. 290, tab. 16. fig. 3), = *S. Webbii* (*Spach, loc. cit.* fig. 4).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Inter Tanger et Tetuan!

Mar. occid.—Casa Blanca! Mogador! (*Lowe Cat.*).

Mar. merid.—Frequens in regione inferiore, sed fugit loca arida saxosa. Legimus in prov. Shedma! Prope Keira! Reraya (*R. et Fr. spec. in Herb. Kew.*)!

—— —, var. BOVEI, = *Spartium Bovei* (*Spach, loc. cit.* 297). Forma grandiflora.

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)!

Ar. Geog. *G. monosperma* Africam borealem et peninsulam Ibericam incolit, et proxime affinis est *G. Rætam* (*Forsk. Fl. Æg.-Arab.* 214) species orientalis in Ægypto-Syria et Arabia obvia.

Notæ quibus plurimæ s. d. species hujus sectionis distinguuntur mihi videntur nimis incertæ et variabiles.

G. DASYCARPA (*Coss. in Ind. Pl. Maroc.* sub *Retama*, sin. descript.), = *Genista*, sp. nov.! *nobis, MSS. in herb.*

Mar. merid.—Copiose crescit in convalle Aït Mesan usque 1700 m. s. m.!

Nuper cel. Cosson spec. florifera et fructifera in Atlante Majore ab indigeno Ibrahim leeta nobis benevole communicavit. Descriptionem viri optimi mox prodituram prævenire nolui

Legimus etiam in convalle Amsmiz spec. nondum florentia, speciem novam ab affinibus valde diversam sistentia!

G. TRICUSPIDATA (*Desf. Fl. Atl.* ii. 138, tab. 183; *DC. Pr.* ii. 148).

Mar. merid.—Seksaua! Ourika (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Africa boreali-occidentalis.

GENISTA TRIDENS (*Cavan. Anal. Cienc. Nat.* iv. 59; *DC. Pr.* ii. 148),
= *G. gibraltarica* (*Lowe Cat. non DC.*).

Mar. sept.—? Tanger (*Broussonnet, ex Cavan. l. c.*).

Mar. occid.—Legimus in monte Djebel Hadid hand procul Mogador!
Vidi specimen prope Mogador a b. Lowe lectum sub *G. gibraltarica* in
herb. Kew.!

Mar. merid.—Satis frequens ad radices Atlantis Majoris. Mesfiona!
Tasseremout! Ourika!

Cel. Cosson certum me fecit plantam a nobis in provinciis meridionali-
bus lectam ad *G. tridentem*, Cav., referre deberi. Floribus remotis in
racemis laxis dispositis ab affinibus statim dignoscitur. De loco
natali prope Tanger indicato valde dubito. Broussonnet nimis facile
plantas originis diversæ semel commiscuit, et specimen ex provinciis
meridionalibus forsitan Cavanillesio misit.

Stirps Imperii Maroccani propria.

G. GIBALTARICA (*DC. Pr.* ii. 148).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Schsb. spec. comm. Cosson*)!

Ar. Geog. Hucusque nonnisi in utroque litore Freti Herculis detecta.

G. juniperina (*Spach, Ann. Sc. Nat.* ser. 3, ii. 275; *Walp. Rep.*
v. 478).

Mar. sept.—In rupibus prope Tetuan (1851)! In monte Djebel Dersa
supra Tetuan (*Webb Herb.*)!

G. juniperina stirps tantum in locis indicatis cognita, *G. gibraltarica*
proxima, gaudet habitu proprio, sed notæ diagnosticæ non gravis mo-
menti, et hæc potius ad subspeciem reducenda mihi videtur.

G. TRIACANTHOS (*Brot. Fl. Lus.* ii. 89, *Phyt. Lus.* tab. 54; *DC. Pr.*
ii. 147), var. *GALIOIDES* (*Spach loc. cit.*)

Mar. sept.—In monte Djebel Kebir prope Tanger admodum frequens et
ludibunda! Tanger! (*Webb Herb.* in scheda, *G. triacanthos*, var. *ga-*
lioides manu cel. Spach addita)! (*Salzm. Exsicc.*)!

Ar. Geog. Peninsula Iberica. Mauritania Tingitana.

In collibus supra Algeciras legi specimina a Tingitanis non distin-
guenda.

G. FEROX (*Desf. Fl. Atl.* ii. 136, tab. 182, sub *Spartio*; *Poir. Dict.*
Suppl. ii. 708; *DC. Pr.* ii. 147), var. *MICROPHYLLA*, nob.

Speciminibus Algeriensibus conformis nisi foliis multo minoribus pro
longitudine latioribus.

Mar. occid.—In monte Djebel Hadid hand procul Mogador!

Mar. merid.—In prov. Shedma!

Ar. Geog. Africa borealis.

Specimina fructifera tantum vidimus. Planta ulterius inquirenda.

G. MYRIANTHA (*Ball, in Journ. Bot.* 1875, 303).

Frutex 4-6-pedalis, ramosissimus; herba tota subglaberrima; ramulo-

rum hornotinorum debilium folia sessilia, anguste lanceolata, annotinorum obovato-oblonga, omnia parvula, 1-foliolata, carnosula, obtusa vel interdum mucronulata; stipulæ minutissimæ bidenticulatæ; flores numerosissimi versus apicem ramulorum conferti; bracteolæ 2 marcescentes in quovis pedunculo; calycis dentes superiores triangulares tubo lævi subæquilongi, tres inferiores æquales, omnes apice ciliolati; carina vexillo æqualis, concolor, pallide flava; alæ saturatæ luteæ. Legumen non vidimus.

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Amsmiz circa 1300 m.!

A *G. scorpio*, DC., differt habitu, glabritie, floribus bicoloribus, calycis tubo lævi nec striato. Facie prior *G. lucidæ*, Camb., sed in hac carina vexillum valde superat, calyx profundius fissus.

GENISTA FLORIDA (*L.*, *DC. Pr.* ii. 151), var. MAROCCANA, *nob.*

Differt a typo pubescentia sericea copiosiore, foliis minoribus angustioribusque, floribus minoribus et calyce pro magnit. corollæ minore. Frutex erectus, 4–8-pedalis.

Mar. merid.—In regione inferiore Atlantis Majoris. Legimus in glareosis convallis Ourika, et iterum in convalle Amsmiz a 1000 m. ad 1500 m.!

Hanc stirpem pro subspecie *G. floridæ* olim habui, sed characteres citati tantum varietatem sistunt.

Notæ diagnosticæ quibus distinguuntur *G. polygalæfolia*, DC. Prod., et *G. leptoclada*, J. Gay, mihi videntur nimis variabiles. His etiam addenda est velut subspecies *G. floridæ*, *G. oretana*, Webb. *G. florida* est imprimis peninsulæ Ibericæ civis.

G. PSEUDOPILOSA (*Coss. Not. Pl. Mid. Esp.* p. 102; *Walp. Ann.* iii. 847)?

Mar. merid.—Ex distr. Reraya ad radices Atlantis Majoris specimen unicum florens attulerunt cl. Rein et Fritsch! Vidi in herb. Kew.; sed ob defectum fructus species incerta manet.

Ar. Geog. Hispania. Algeria.

G. CLAVATA (*Poir. Dict. Suppl.* ii. 717; *Spach, Ann. Sc. Nat.* ser. 3, iii. 144; *DC. Pr.* ii. 146).

Species valde ludibunda cujus varietates quatuor enumeravit cl. Spach, *loc. cit.* Hæ omnes in ditione Tingitana obviæ, sed limitibus certis vix definiendæ.

—— —, var. α , = *Spartium capitatum* (*Cav. Anal. Cienc. Nat.* 1801, iv. p. 63; *Schousb. in Herb.*), = *Genista umbellata*, var. β . capitata (*DC. Pr.* ii. 147).

Mar. sept.—Tanger (*Schsb. spec.*)! (*Salzm. Exsicc.* sub *Spartio capitato*)!

Mar. occid.—In agro Mogadorensi (ex *DC. Pr. l. c.*). Adest in herb.

Kew. spec. ex herb. Gouan cum scheda MS. Broussonnet "ex Mauritania."

— —, var. β . WEBBIANA (*Spach, l. c.*; *Walp. Rep.* v. 499).

Mar. sept.—Tanger (*Webb Herb.*)! Inter Tanger et Tetuan!

— —, var. γ . GOUDOTIANA (*Spach, l. c.*; *Walp. l. c.*).

Mar. sept.—"Crescit prope Tingidem," ex *Spach, l. c.*

— —, var. δ . CASUARINOIDES (*Spach, l. c.*; *Walp. l. c.*).

Mar. sept.—Prope Tanger (*Webb Herb.*)! El Araiseli (*R. & Fr. spec. in herb. Kew.*)!

Ar. Geog. Species in Imperio Maroccano et peninsula Iberica australiore obvia.

Spartium molle (Cav. Anal. Cicne. Nat. 1801, iv. 57), = *G. mollis* (DC. Pr. ii. 146), ex agro Mogadorensi a Broussonnet allatum, est verisimiliter forma quædam *G. clavata*. Specimina Mogadorensia non vidi, nec vidit ecl. Cosson.

Genistæ specimen nondum florentem et omnino incertum legit in convalle Aït Mesan celeb. J. D. Hooker. Planta tota subglaberrima; rami erecti stricti parce foliati; folia simplicia, exstipulata, linearilanceolata, juniora retrorsa, ex pulvinula inconspicua orta. Planta ulterius inquirenda.

ULEX (*L. Gen.* 881).

U. BÆTICUS (*Boiss. El.* p. 30; *Webb, Ot. Hisp.* p. 39; *Walp. Rep.* i. 626), var. *CONGESTUS*, nob., = *U. scaber* (*Kunze in Flora*, 1846, 696), var. δ . *congesta* (*Webb, Ot. Hisp.* p. 40, tab. 32 B).

Mar. sept.—In monte Beni Hosmar pr. Tetuan! eodem fere loco ubi olim legerat *P. B. Webb*.

Ar. Geog. Hispania australis. Africa boreali-occidentalis.

Characteres quibus *U. Bourgaeanus*, *Webb*, et *U. scaber*, *Kunze*, ab *U. bætico* separantur mihi videntur omnes variabiles, et eodem pacto *Ulices* fere omnes hujus gregis locis diversis exortos pro sp. diversis haberes. Nomen triviale *scaber* haud aptum. *Ulices* omnes affines plus minusve pubescentes vidi, non vero scabros.

U. africanus (*Webb, Ot. Hisp.* p. 38, tab. 29; *Walp. Ann.* iv. 464), var. *DELESTREI*, *Webb, l. c.*

Mar. sept.—Prope Djemma el Ghazaouat ad fines Mauritaniae Tingitanæ (*Delestre, Webb, loc. cit.*).

Ar. Geog. Algeria boreali-occidentalis.

Stirps mihi tantum ex descript. et icone Webbia cognita. *U. bætico* valde proxima, et non sine dubio pro subspecie enumerata.

U. BOIVINI (*Webb, It. Hisp.* p. 51; *Walp. Rep.* i. 626), var. *SALZMANNI*, nob., = *Nepa Salzmanni* (*Webb, Ot. Hisp.* p. 31, tab. 26; *Walp. Ann.*

iv. 468), = *Ulex genistoides* (*Lindl. Bot. Reg. tab. 1452; Salzm. Exsicc. pro parte, non Brot. nec Clem.*).

Mar. sept.—In monte Djebel Kebir pr. Tanger mense Junio florentem legi! Tanger (*Salzm. Exsicc. sub U. genistoidi*)!

— —, var. *MEGALORITES, nob.*, = *Nepa megalorites* (*Webb, Ot. Hisp. p. 30, tab. 25; Walp. Ann. iv. 467*), = *U. genistoides* (*Salzm. Exsicc. pro parte*).

Mar. sept.—Tanger (*Schousb. spec.*)! eandem formam nondum florentem legi mense April in monte Djebel Kebir! Specimina nullomodo a typo Webbiano diversa legi mense Junio florentia in collibus prope S. Roque haud procul Gibraltar!

Ar. Geog. Africa Tingitana. Hispania.

Qui descriptiones et icones Webbianas sub oculis tenet species tres hic semel conjunctas omnino diversas esse judicavit. Notæ diagnosticæ evidentes non desunt; sed qui plantas in vivo accurate scrutabit sententiam facile mutabit. Dentes calycini, bracteolæ et petala omnia variabilia sunt, et vix invenies plantas locis diversis enatas omnino conformes. *N. Salzmanni* Webb est inter *U. Boivini* et *M. megaloritem* pluribus notis media. Salzmann certe varietates nostras semel confudit, et ambo in herbariis sub *U. genistoide* suo obviæ sunt.

CYTISUS (*L. Gen. 877; Benth. et H. fil. Gen. Pl. i. 484*).

C. BÆTICUS (*Webb, It. Hisp. 52; Ot. Hisp. p. 45, tab. 40; Walp. Rep. i. 627, sub Sarothamno; Steud. Nomencl.*), = *Sarothamnus Gaditanus* (*Boiss. et Reut. Diagn. Pl. His. No. 15; Walp. Rep. i. 628*), = *S. arboreus* (*Boiss. Voy. Esp. 400, non Desf. sub Spartio*).

Mar. sept.—Tanger (*Webb Herb.*)! (*Salzm. Exsicc.*)! In collibus inter Tanger et Tetuan!

Mar. occid.—Mogador (*Lowe Cat., sub C. arboreo, Desf.*)?

Ar. Geog. Hispania australis. Africa boreali-occidentalis.

— —, var. *MACRANTHUS, nob.*

Mar. sept.—In clivo septentrionali montis Djebel Kebir legi florentem mense Aprili!

Arbuseula speciosa 10-pedalis et ultra, floribus duplo majoribus longius pedicellatis, foliorum petiolis longioribus insignis, sed aliis notis non a typo diversa.

C. ARBOREUS (*Desf. Fl. Atl. ii. 131, tab. 177, sub Spartio; DC. Pr. ii. 154*), = *Sarothamnus arboreus, Webb, non Boiss. Voy.*

Mar. merid.—Prope Tasseremout fructiferum legimus incunte Majo!

Specimina nostra foliis et ramis apice denudatis typo Algeriensi similima sed legumina aliquantulum diversa, illis *C. arborei* et *C. scoparii*, Lk., fere intermedia—juniora pilosa pilis subpatentibus, demum glabrescentia, breviora quam in *C. scopario*. In nostris legumina utrinque

æqualiter dehiscencia; in *C. scopario* vidi semper suturam superiorem serius et difficile dehiscentem.

Ar. Geog. Africa borealis.

C. scoparius, var. *glabratus* (Link in Buch, Beschr. Ins. Can. p. 156) in insula Teneriffa a Chr. Smith lectus et a recentioribus non detectus, est forsitan forma *C. arborei*, ut suspicatus est el. Lowe (Man. Fl. Mad. 123).

CYTISUS BALANSÆ (Boiss., *Diagn. Pl. Or.* ser. 2, ii. 7, sub *Sarothamno*), var. ATLANTICUS (Ball, in *Journ. Bot.* 1873, 303).

Flores solitarii, axillares in ramis hornotinisi nermibus, vel rarius in ramis annotinis glabris striatis pungentibus paucifoliatis; pedunculi bracteolati ad basin foliolis 2-3 obverse lanceolatis, obtusis, sericeis, ex pulvinula inerassata ortis, fulerati; calyx parce pilosus campanulatus vix bilabiatus, labio superiore vix emarginato, inferiore brevissimo dentibus 3 subobsoletis; corolla glaberrima; vexillum carinam incurvo-falcatam et alas paulo superans; stylus apice non incrassatus, magis quam in affinibus (*C. purgante* et *C. Hænseleri*) exsertus.

Mar. merid.—In regione superiore Atlantis Majoris. Aït Mesan! usque in juga Tagherot! Djebel Tezah! a 2200 m. ad 3000 m. et ultra.

Ar. Geog. *C. Balansæ* typicus montes Algerienses (Djebel Aures) incolit.; sp. affines in Hispania et Gallia obviæ.

Species hujus gregis—scilicet *C. purgans*, L. (sub *Spartio*), *C. Hænseleri*, Boiss., et *C. Balansæ*, inter sectiones *Sarothamnum* et *Spartocytisum* transitum præbentes, proxime affines sunt, et ulterius inquirendum an notæ diagnosticæ satis constantes. In var. nostra flores magnitudinis *C. purgantis*, sed forma carinæ valde diversa.

C. ALBIDUS (DC. *Cat. H. Monsp.* 101, et *Prod.* ii. 155).

Mar. oecid.—Mogador inter monticulos arenæ mobilis! In monte Djebel Hadid!

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Amsmiz! Prope Mzouda! Prope Seksaona!

Ar. Geog. Stirps Maroccana verisimiliter etiam in Teneriffa indigena. Olim in Horto Orotavensi introductus, locus natalis hodie incognitus. Confer Webb, *Phyt. Can.* ii. p. 47.

C. TRIFLORUS (L'Hér. *Stirp.* 184; DC. *Pr.* ii. 154).

Mar. sept.—Tanger (Webb *Herb.*)! In monte Djebel Kebir!

Ar. Geog. Regio Mediterranea imprimis occidentalis. Deest in Oriente et in Macaronesia.

C. CANDICANS (L., et DC. *Pr.* ii. 145, sub *Genista*; DC. *Fl. Fr.* iv. 504), = *Teline candicans* (Webb, *Phyt. Can.* ii. 36).

Mar. sept.—Tanger! (Schousb. *herb.*) Coss. sp.! In monte Djebel Kebir! Cap. Spartel!

Ar. Geog. Regio Mediterranea occidentalis (inclusa Macaronesia). Deest in Oriente et verisimiliter in Græcia.

CYTISUS CININNATUS, *nobis*. Frutex ramis lignosis; flores ad apicem ramorum eongesti subcapitati; folia trifoliolata, breviter petiolata, foliis lanceolato-ellipticis, margine inflexis, utrinque tomentosis, præsertim in pagina superiore pilis cincinnatis (spiraliter arete convolutis) dense obteectis, adeo ut oculo non fortiter armato videntur foveolata; vexillum vix emarginatum, extus cum calyce sericeo-villosum; earina minus quam in sp. affinibus falcata; stylus lituiformis, apice rectus, demum exsertus. Flores saturate flavi vel aurantiaci? Bracteolæ infra basin calycis ad fasciculam pilorum sericeorum reductæ.

Mar. merid.—Specimen unicum in prov. Mtouga a ell. Rein et Fritsch lectum in herb. Kew. vidi et descripsi!

Species pluribus notis ad *Cytisos* Canarienses (*Teline*, Webb.) aecedit, et præsertim *C. congesto* (Willd., sub *Spartio*), sed ab his omnibus differt inflorescentia, foliis majoribus, indumento singularissimo, bracteolis ad pilos reductis nec foliaceis. Fructus nondum visus.

C. HOSMARIENSIS (*Coss. in Bull. Soc. Bot. Fr. xx. 245*, sub *Genista*). Planta structura et habitu in grege *C. candicantis* (*Teline*, Webb) reponenda. Ccl. auctor hanc in genere *Genista* collocare mavult; nobis (ducibus Benth. et H. fil. in *Gen. Pl.*) fas sit eum *Cytisis* enumerare in societate specierum præcedentium. Descriptionem Cossonianam proferre juvat.

“*Frutex* ut videtur elatus, ramosus, *ramis* striatis *inermibus*, vetulis cortice rimoso cinerascete, novellis patentibus vel subdeflexis cortice virescente sericeo-pubescente, foliorum pulvinis demum modice incrassatis. *Folia* in genere ampla, *petiolata*, *3-foliolata*, alterna foliis præsertim pagina inferiore sericeo-pubeseentibus, obovatis, breviter petiolatis, obtusis, muticis vel brevissime mucronulatis, terminali majore. *Stipulæ minutæ, lineares*, marcescenti-persistentes. *Flores magni, apice ramulorum 6–8, in racemos subcorymbosos patentes dispositi, bracteis minutissimis ovatis obtusis, pedicellati pedicellis sub calyce bibracteolatis bracteolis minutissimis ovatis obtusis. Calyx* cum pedicellis bracteolisque dense sericeo-pubescent, amplus, campanulatus, *bilabiatus, labiis tubo subtriplo longioribus*, superiore inferius subæquante bipartito lobis ovato-oblongis acuminatis, *inferiore apice tridentato* dentibus ovato-lanceolatis acuminatis. Corolla flava, calyce subdimidio vel subtriente longior, vexillo porrecto, amplo, ovato-suborbiculato, apice emarginato, dorso pubescenti-sericeo, alas subæquante vel paulo superante, alis carina paulo longioribus, isti subæquilatis, cultriformi-oblongis, obtusis, glabris, carina cultriformi-oblonga, obtusa, dense sericeo-villosa. *Ovarium* calyce brevius, sericeo-villosum, lineari-lanceolatum, *6-ovulatum*, in stylum elongatum apice arcuato-ascendens sensim attenuatum. Stigma minutum, capitellatum. Legumen ignotum.”

Mar. sept.—In monte Beni Hosmar prope Tetuan a P. B. Webb anon 1827 detectus!

Pace optimi amici nomen triviale scribendum *Hosmariensis* nec *Osmar-*

riensis dum litera vocalis ab incolis fortiter aspiratur. Monente cel. auctore species hæc in § *Teline* collocanda. "Habitum et calycis fabricam affinis *C. heterochroa* (Webb in Bourg. Pl. Esp. 1852, n. 1719 c) in Hispaniæ regno Valentino prope Moxente (Bourgeau) et in Arragonia australi (Wilkomm) lecto, sed eximie distincta foliis majoribus obovatis, floribus in racemos pedunculatos dispositis, bracteolis minutissimis calyci subcontiguis, non linearibus a calyce remotis calycibus amplioribus densius pubescentibus, dentibus labii calycini inferioris latoribus brevioribusque, carina dense sericeo-villosa, non glabra, ovario sericeo-villoso, non glabro, etc."

Legimus in eodem monte ramos *Cytisi* cujusvis foliis novellis sericeo-villosis, vix tamen huc referendi!

CYTISUS LINIFOLIUS (*L. et DC. Pr.* ii. 146, sub *Genista*; *Lam. Dict.* ii. 249).

Mar. sept.—Tanger (*Webb Herb.*)! (*Salzm. Exsicc.*)! In monte Djebel Kebir!

Ar. Geog. Regio Mediterranea occidentalis nunquam à mari remotus; ex Gallia meridionali usque Macaronesiam.

C. TRIDENTATUS (*L. et DC. Pr.* ii. 151, sub *Genista*; *Benth. et II. fil. Gen. Pl.* i. 484), var. LASIANTHA, = *Genista lasiantha* (*Spach, Ann. Sc. Nat.* ser. 3, iii. 147).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! In monte Djebel Kebir!

Ar. Geog. Species est peninsulæ Ibericæ civis; var. nostra in utroque litore Freti Herculis obvia est. Legi spec. Tingitanis omnino conformia in collibus supra Algeciras.

Qui stirpem nostram cum *G. stenoptera*, Spach, comparabit facile hanc pro specie rite distincta habebit, sed in magna copia speciminum ex Hispania et Lusitania provenientium tot et tantas formas intermedias inveniet ut species quinque Spachianas in unam solam Linnæanam conjungendas esse fatebitur. Varietas nostra a cæteris vexillo sericeo, nec ulla alia nota, distinguitur.

C. FONTANESII (*Spach, Ann. Sc. Nat.* ser. 3, iii.), = *Spartium biflorum* (*Desf. Fl. Atl.* ii. 133, tab. 179), = *Genista biflora* (*DC. Pr.* ii. 146).

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Aït Mesan circ. 1250 m.! Supra Seksaoua 1400 m.!

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

Specimina nostra Algeriensibus conformia, sed flores sæpe 3, interdum 4, in apice ramorum.

ONONIS (*L. Gen.* 883).

O. NATRIX (*L., DC. Pr.* ii. 159).

Mar. merid.—In planitie haud procul urbe Marocco (*G. Maw spec.*)!

Pr. Sektana (*R. et Fr. spec. in herb. Kew.*)!

— — —, var. TOMENTOSA (*Boiss. Fl. Or.* ii. 59).

Mar. merid.—Prope Milhain!

— — —, var. MICROPHYLLA (*Boiss. l. c.*), = *O. ramosissima*, *Desf.*

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Tetuan (*Webb Herb.*)!

Mar. oeeid.—In arenosis prope Mogador! verisimiliter huc *O. natrix* β (Lowe Cat.) prope Mogador indicatus.

Mar. merid.—In provincia Shedma!

Ar. Geog.—Regio Mediterranea omnis (inclusa Macaronesia). Europa media.

O. Natrix est stirps quam maxime variabilis in plurimas falsas species ab auctoribus, etiam gravissimis, dilaniata. Varietates a celeberrimis Boissier et Webb propositæ ægre definiendæ. Exemplaria quæ vidi ex Africa boreali *O. pictam*, Dsf., referunt. Varietas nostra *tomentosa* verisimiliter eadem ac var. Boissieriana. Parum differt ab *O. cuspidata*, Dsf., ejus specimina authentica non vidi.

Ononis angustissima (Lam. Dict. i. 502; Webb. Phyt. Can. ii. 23), = *O. longifolia* (Willd. Enum. 750; DC. Pr. ii. 159).

Mar. oeeid.—Mogador in arenosis! (Lowe Cat.). In provincia Haha haud procul Mogador! Agadir (Coll. Indig. 1871)!

Ar. Geog. Stirps Canariensis et Mogadorensis.

Hæc mihi videtur proles *O. natrix* foliis angustis apice argute serratis, et stipulis elongatis apice setaceis. Notas diagnosticas a beato Webb loc. cit. "cultura immutatis" laudatas vir optimus nusquam indicavit. Auctoritati magistrorum obsecutus hanc sicut subspeciem enumeravi.

O. viscosa (L., DC. Pr. ii. 160), var. ? FRUTICESCENS (Ball, in Journ. Bot. 1873, 304).

Differt insigniter radice ramisque inferioribus lignosis, legumine et seminibus fere intermediis inter *O. viscosam* et *O. siculam*. Corolla sat magna calycem subduplo excedens. Vexillum luteum rubro stratum.

Mar. merid.—Specimen unicum legi in distr. Reraya, in reg. inferiore Atlantis Majoris!

O. sicula, Guss., ab affini *O. viscosa* differt præsertim legumine angustiore polyspermo, seminibus maturis 12–20, ellipsoideo-reniformibus, pallidis; dum in *O. viscosa* semina multo majora sunt, 3–6, compressa, atro-fusca. Nostra planta melius eognita speciem novam forsan sistet, forsan nexum inter species supra memoratas.

Ar. Geog. *O. viscosa* typica peninsulam Ibericam, Galliam australem et Italiam incolit; hujus var. vel subspecies *O. breviflora*, DC., etiam in Græcia et Oriente obvia est.

O. porrigens (Salzm. Exsicc.), = *O. fœtida* (Schsb. herb.; DC., Pr. ii. 161), = *O. viscosa*, Coss. MSS.

Mar. sept.—Tanger! (Salzm. Exsicc.)! (Schsb. herb. spec.)!

Hanc ego pro subspecie habui dum notis plurimis ab *O. viscosa* insigniter differt. Foliola nisi in foliis infimis ad basin caulis semper acuta, minute et æqualiter serrulata; flores quam in *O. viscosa* multo majores roseo-purpurei, nec flavi vexillo apice purpurascente; legumen latius, brevius, vix glanduloso-pubescent; semina paululum majora, oculo armato minute granulata. Canles bipedales et ultro, erectius-

culi, parum ramosi. Ccl. Cosson mihi benevole concessit specimina ex herb. Schousboe quæ formam minorem hujus stirpis referunt. Præter flores minores hæc a spec. nostris non diversa sunt.

ONONIS SICULA (*Guss. Prod. Fl. Sic.* ii. 387; *DC. Pr.* ii. 160).

Mar. merid.—Distr. Mesfiona! Prope Ourika (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Per regionem Mediterraneam calidiorē (exclusa Macaronesia) late dispersa, sed non ubique obvia.

Præter notam ex seminibus supra relatam, differt ab *O. viscosa* pedicellis supra aristam brevissimis.

O. POLYPHYLLA (*Ball in Journ. Bot.* 1873, 304). (Plate XVI.)

Annua, pumila, e basi ramosa; herba tota virens, glanduloso hirta; folia numerosa, approximata, trifoliolata, foliolo medio stipitato, lateralibus basi disjunctis, omnibus angustis apice dentatis; stipulæ elongatæ, demum subscariosæ vaginantes; pedunculi uniflori, aristati, folia et stipulas vix æquantes; vexillum segmentis calycinis æquale, apice rubescens; legumen oblongum, latitudine sua fere duplo longius, lacinias calycinas lanceolato-cuneatas vix superans.

Mar. merid.—In planitie ad radices Atlantis Majoris distr. Mesfiona! et versus Tasseremout.

Habitu ad *O. pubescentem*, L., accedit, etsi statura multo minor; præter folia et stipulas angustiores differt pedunculis aristatis et legumine oblongo, nec breviter ovali-rhomboideo.

O. PUBESCENS (L., *DC. Pr.* ii. 160).

Mar. occid.—Casa Blanca! In monte Djebel Hadid (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio Mediterranea (exclusa Macaronesia).

O. CINTRANA (*Brot. Phyt. Lus.* 138, tab. 57; *DC. Pr.* ii. 160).

Mar. sept.—Tanger (*Schsb. Herb. ex Coss. spec.*)! (*Webb Herb.*)! (*Brouss. spec. in Herb. Kew.*)! (*Salzm. Exsicc.*)!

Ar. Geog. Nonnisi in Lusitania et in ditione Tingitana detecta.

O. MAWEANA (*Ball. in Journ. Bot.* 1873, 304).

Annua, e basi ramosa, parce glanduloso-hirta; folia infima et suprema unifoliolata, cætera trifoliolata, foliolo medio majore stipitato, omnia obovato- vel oblongo-lanceolata, apice rotundata (rarius rectusa), argute denticulata; stipulæ inferiores clongatæ, superiores dilatatæ ovatæ acutæ argute denticulatæ; pedunculi biflori, exaristati, foliis paulo longiores, infra medium furcati, sub anthesi erecti; calycis segmenta tubum bis excedentia, angusta, quinquenervia, nervo medio subobsoleto, nervis lateralibus duobus utrinque approximatis, prominulis; corolla lacinias calycinas parum excedens, insigniter bicolor, vexillo pallido striato, carina (cum alis) apice aurantiaca. Legumen nondum visum.

Mar. sept.—In ditione Tiugitana, juxta stagna subsalsa, detexit oculatissimus G. Maw, botanices et præsertim floriculturæ fautor egregius. Species ut videtur omnino distincta, nec ulli proxima nisi forte *O. hebecarpæ*, Webb. Ab hac differt imprimis pedunculis infra medium, nec apice tantum, furcatis, stipulis ter quaterve majoribus, foliis angustioribus, et cet. *O. biflora*, Dsf., longius distat pedunculo aristato, apice tantum furcato, floribus pendulis, statura multo majore. Facie similiores sunt *O. gibraltarica*, Boiss., *O. brachycarpa*, DC. Prod., *O. nutricoides*, Duf., quæ omnes præter alias notas pedunculis 1-floris aristatis distinguuntur.

ONONIS ATLANTICA (*Ball, in Journ. Bot.* 1873, 304), = *O. spinulosa* (*Coss. in herb.*). (Plate XVII.)

Pereunis, basi suffruticosa, ramis erectis; herba tota pube brevi glandulosa vestita; foliorum subsessilium foliola omnia sessilia crassiuscula, obverse cuneata, apice tautum denticulata; stipulæ brevissimæ; pedunculi rigidi, uniflori, fructiferi supra aristam brevem spinescentem reflexi; calycis laciniæ subæquales, tubo brevissimo ter longiores; petala flava, vexillo carina sesquilongiore; legumen breviter stipitatum, e basi obliqua oblongum, calyce ter longiore; semina pauca minute granulata.

Mar. merid.—In regione inferiori Atlantis Majoris. In convalle Amsmiz detexit cel. J. D. Hooker, circa 1250 m.! In monte Djebel Ouensa legit indigenus Ibrahim et spce. mihi benevole comm. cel. Cosson. Ab *O. cenisia*, L., recedit herba glandulosa, foliis subsessilibus, pedunculorum arista rigida spinescente, calyce profundius fisso, petalis flavis nec purpurascentibus, et præsertim legumine stipitato, angustiore, et seminibus minoribus minute et undique granulatis, nec inæqualiter tuberculatis.

Propior est forsau *O. rigida*, Kze., sed in hæc foliola obovata circuitu toto, nec apice tantum, dentata, stipulæ longiores insigniter dentatæ; hujus legumen a me non visum, nec descripsit cl. auctor.

Hanc esse *O. cenisia*, Coss. MS., a Balansa in Atlante lecta, in *Journ. Bot.* l. c. mira hallucinatione deceptus dubitavi. Vir oculatissimus species probe diversas confundere nequit. Specimina *O. cenisiæ* genuinæ in Atlante Majore ab indigeno Ibrahim pariter lecta debeo humanitati cel. Cosson.

O. PENDULA (*Desf. Fl. Atl.* ii. 147, tab. 191; *DC. Pr.* ii. 162).

Mar. sept.—Tanger (*Webb Herb.*)! Cap. Spartel (*Schousb. herb.* sub "*O. reclinata*," L.). *Coss. sp.*!

Mar. occid.—Casa Blanca! Mazagan! Mogador (*Lowe, spec. in Herb. Kew.* sub "*O. laxiflora*")!

Specimina nostra prope Casa Blanea lecta ab *O. Schounii* (*DC. Pr.* ii. 162) non diversa sunt. Hæc tamen vix pro varietate habenda, dum nulla adest nota diagnostica.

— —, var. *BROUSSONNETII*, = *O. Broussonnetii* (*DC. Pr.* ii. 161).
 Mar. occid.—In arenosis prope Mogador! ubi verisimiliter Broussonnet olim legebat spec. ab ill. Candolleo descripta.

Hæc est potius lusus æstivalis quam vera varietas. Spec. nostra ineunte Junio lecta gaudent foliis angustis glabrescentibus, stipulis subcoalitis persistentibus, floribus in racemum subaphyllum dispositis, interdum breviter pedunculatis, interdum pedunculis inferioribus elongatis subpollicaribus. Confer de evolutione sp. affinis *O. laxifloræ* observationes beati P. B. Webb (*Phyt. Can.* ii. 27).

Ononis laxiflora (*Desf. Fl. Atl.* ii. 146, tab. 190; *DC. Pr.* ii. 161).

Mar. merid.—In saxosis aridis prope Aïn Oumast, pr. Shedma! In prov. Mtouga!

Hæc est *O. pendulæ* arcte affinis; differt pedunculis laxè corymbosis quam in typo longioribus, nec ad apicem ramorum congestis, laciniis calycinis angustioribus, stipulis profundius denticulatis, seminibus paulo majoribus sub lente tuberculatis nec minute granulatis, et meo sensu pro subspecie enumeranda est.

Ar. Geog. *O. pendula* et *O. laxiflora* cives indigenæ Africæ boreali-occidentalis, paulo extra fines Africanas dispersæ sunt. *O. pendula* in Hispania Granatensi et in Sicilia, *O. laxiflora* in Macaronesia obvia est.

O. pendula et *O. reclinata* sunt re vera proximæ species, sed in sententiam cl. Lowe, qui has in unam speciem conjunxit (*Man. Fl. Mad.* i. 130), convenire nequeo. *O. pendula* non solum differt magnitudine partium omnium dupla, sed habitu toto et seminibus.

O. RECLINATA (*L.*, *DC. Pr.* ii. 162).

Mar. sept.—Tanger (*Webb Herb.*)!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). Europa occidentalis ex Scotia usque fretum Herculeum.

O. mollis, Savi, et *O. Cherleri*, *DC.*, non *L.*, sunt meræ formæ hujus speciei.

O. FILICAULIS (*Salzm. Exsicc.*; *Boiss. Voy. Esp.* 433, tab. 46; *Walp. Rep.* i. 625).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Schsb. Herb.*)! (*Webb Herb.*)!

Ar. Geog. Hispania Granatensis. Africa boreali-occidentalis.

Legi prope San Roque specimina Tingitanis omnino conformia!

O. ANTIQUORUM (*L.*, *Rchb. Fl. Exc.* p. 517, et *Ic. Pl. Crit.* i. tab. 14), = *O. spinosa*, var. γ . glabra (*DC. Pr.* ii. 163).

Mar. merid.—Sat frequens in planitie et ad radices Atlantis Majoris. Mesfioua! Mtouga! In convalle Amsniz circ. 1200 m.!

Ar. Geog. Regio Mediterranea (exclusa Macaronesia).

In planta Maroecana rami semper tenues glabri vel superne tenuiter
 LINN. JOURN.—BOTANY, VOL. XVI. 2 G

glanduloso-pubescences. Spinæ elongatæ sæpius geminæ nitidæ albæ. Folia superiora unifoliolata.

ONONIS MITISSIMA (L., DC. Pr. ii. 163).

Mar. sept.—Tanger! (Salzm. Exsicc.)! (Webb Herb.)!

Mar. merid.—Aïn Tarsil, prov. Mtouga (J. D. H.)!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia).

O. SERRATA (Forsk. Fl. Æg.-arab. p. 130; DC. Pr. ii. 163), = O. Dehnhartii (Ten. Fl. Nap.), = O. diffusa (Ten. Prod. Fl. Nap. p. 41; DC. Pr. ii. 163).

Mar. sept.—Tanger in arenosis maritimis (Schsb. Herb.).

Mar. oecid.—Agadir (Coll. Indig. 1871)!

— — —, var. MINOR. Flores et fructus sublimidio minores; foliola lateralia ad laciniâs lineares brevissimas reducta, interdum earentia. Semina granulata paulo minora.

Mar. oecid.—Ad radices montis Djebel Hadid haud procul Mogador!

Mar. merid.—In distr. Mesfoua!

Specimina ex Mesfoua formam fere intermediam referunt, et his similia sunt specimina Webbiana ex insula Lancerotta.

De O. diffusa, Ten., sententiæ virorum gravissimorum discordes. Præeunte eel. Boissier (Pug. Pl. Nov. Afr. Bor. p. 33, et Fl. Or. ii. 63), et observatis plurimis exemplaribus ex regno Neapolitano a Gussone, Parlatores aliisque receptis, habui O. diffusam, Ten., pro forma O. serrata, vix aut ne vix pro varietate habendam. Vidi ex Ægypto superiore specimina var. minori nostræ similia.

Ar. Geog. O. serrata, Forsk., per regionem Mediterraneam calidiorem a Persia australi usque Macaronesiam diffusa est.

O. COSSONIANA (Boiss. et Reut. Pug. p. 33; Walp. Ann. iv. 463), = O. diffusa (Coss. Pl. Crit. 35, non Ten.).

Mar. sept.—Tanger! (Schsb. Herb.). Coss. spec. (Salzm. Exsicc., sub "O. tingitana" et iterum sub "O. gibraltarica," spec. in herb. Kew.)! (Webb Herb.)!

Ar. Geog. Hispania Bætica. Imper. Maroccanum.

Huic proxima est O. hirta, Desf.; sed hæc satis differt seminibus dimidio minoribus, flore fructuque minoribus, stipulis minimis, racemo foliato, cæterisque notis.

O. TOURNEFORTII (Coss. Pl. Crit. 34; Walp. Ann. iii. 845).

Mar. oecid.—Mogador in arenosis!

Specimina quæ vidit in herb. Desfontaines eel. auctor cum scheda "O. variegata, Tanger," verisimiliter a Broussonnet ad Mogador lecta sunt. Nemo recentiorum hanc speciem prope Tingidem vidit, etsi in altero litore freti Hereulei obvia sit.

Ar. Geog. Species Hispaniæ Bætiæ et agri Mogadorensis propria.

O. ALOPECUROIDES (L., DC. Pr. ii. 163), var. TRIFOLIOLATA (Coss.

Pl. Crit. p. 33), = *O. Salzmanniana* (*Boiss. et Reut. Pug.* 34; *Walp. Ann.* iv. 463), = *O. monophylla* (*Boiss. Voy. Esp. et DC. Prod.* ii. 163, *non Desf. Fl. Atl.*).

Mar. sept.—Tanger (*Schousb. Herb.*, sub *O. variegata* et *O. Mauritanica*)! (*Webb Herb.*)! (*Salzm. Exsicc.* sub *O. alopecuroidi*)!

Mar. occid.—Mogador (*DC. loc. cit.*).

Vidi specimen Lusitanicum in herb. Hooker Tingitanis conforme, cui scheda "*O. paradoxa*, Welw. MSS."

Ar. Geog. *O. alopecuroides* per regionem Mediterraneam (exclusa Macaronesia) late dispersa est. Varietas nostra in peninsula Iberica et Imp. Maroccano obvia, alibi non indicatur.

Ex spec. in herbario Candolleano asservato et ex descriptione citato plane apparet *O. monophyllam*, DC., a stirpe Fontanesiana diversam esse, et ipsissimam var. nostram referre. Specimina ex agro Mogadorensi a DC. citata verisimiliter a Broussonnet lecta.

Folia sunt magna pro parte unifoliolata, sed superiora in racemo foliaceo plerumque trifoliolata.

ONONIS MINUTISSIMA (*L., DC. Pr.* ii. 164).

Mar. sept.—Prope Tetuan (1851)! In monte Djebel Dersa prope Tetuan (*Webb Herb.*)!

Ar. Geog. Regio Mediterranea occidentalis a Dalmatia usque fretum Herculeum.

TRIGONELLA (*L. Gen.* 898).

T. MONSPELIACA (*L., DC. Pr.* ii. 183).

Mar. merid.—In regione inferiore et in convallibus calidioribus Atlantis Majoris. Mesfona! Prov. Shedma pr. Mskala! Prov. Mtonga!
In convalle Aït Mesan circ. 1300 m.!

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia).

T. POLYGERATA (*L., DC. Pr.* ii. 184).

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Amsmiz circ. 1400 m.!

Ar. Geog. Gallia australis. Hispania. Marocco.

— —, var. ATLANTICA (*Ball, in Journ. Bot.* 1873, 305). Differt a typo legumine dimidio brevior, rugis transversis arenatis nec valde elongatis, stipulis latioribus, seminibus (nondum maturis) nigropunctatis.

Mar. merid.—In regione superiore Atlantis Majoris. In convalle Aït Mesan a 2100 m. ad 2600 m.

MEDICAGO (*L. Gen.* 899).

M. LUPULINA (*L., DC. Pr.* ii. 172).

Mar. merid.—Mesfona! In convalle Aït Mesan 1200–1400 m.!

Ar. Geog. Europa omnis (exclusa arctica). Asia temperata. Regio omnis Mediterranea (inclusa Macaronesia). Abyssinia.

Specimina nostra formam inter typum et *M. Cupanianam* (Guss.¹ Fl. Sic. ii. 362) mediam referunt. Hæc novissima in regione Mediterranea sæpe occurrit, certe ad meram varietatem reducenda.

MEDICAGO SUFFRUTICOSA (*Ram. in DC. Fl. Fr.* iv. 541; *DC. Pr.* ii. 173).

Mar. merid.—In regione subalpina et alpina Atlantis Majoris. In convalle Aït Mesan supra Arround, et in jugo Tagherot a 2200 m. usque 3000 m.! In monte Djebel Tezah legit J. D. H.!

Ar. Geog. Hucusque in Pyrenæis tantum nota. Sp. affinis *M. leiocarpæ*, Benth., occurrit etiam in montibus Hispaniæ australis.

Nisi pedunculis paucifloris (plerumque bifloris) specimina nostra Pyrenæis omnino conformia.

M. HELIX (*Willd. Sp. Pl.* iii. 1409).

— —, var. α , *LÆVIS*, = *M. lævis* (*Desf. Fl. Atl.* ii. 213; *DC. Pr.* ii. 174).

Mar. sept.—Prope Tanger! (*Webb Herb.* sub *M. tornata*)! Inter Tetuan et Ceuta forma subinermis!

Mar. occid.—Ex insula Mogador (*Balansa spec.*)!

— —, var. β , *SPINULOSA* (*Moris, Fl. Sard.* i. 438, tab. 39. fig. B).

Mar. occid.—In maritimis prope Mogador et præsertim in insula!

Ar. Geog. Regio Mediterranea occidentalis calidior (inclusa Macaronesia). Deest in Gallia et in Italia superiore et media.

Pariter ac plurimæ hujus generis difficillimi hæc est species polymorpha et formæ extricatu parum faciles. Ccl. Moris in Flora Sard. l. c. species Sardoas optime illustravit, et mihi certiore fecit numerum spirarum ad species delimitandas parum valere. In hac nempe variant 2, 3 vel 4. Mirum videtur in insula exigua juxta urbem Mogador cl. Balansa formam lævem, spinis omnino destitutam, die 25 Aprilis 1867 legisse, dum nos post quatuor annos die 27 Aprili 1871 nonnisi legumina spinulosa vidimus in plantis quæ locustis superstiterunt. Ardua est sententia et naturæ scrutatoribus remittenda utrum inter plantas ejusvis varietatis in eodem loco creescentes causas externas—cœli temperiem variabilem, frequentiam vel varietatem animalium nocivorum, etc.—agere ut una alterave varietas magis floreat, an progeniem ejusdem parentis ab iisdem causis modificatam, alterna vice legumina lævia vel spinulosa proferre credendum sit. In *M. helice* spiræ sæpissime, sed non semper, dextrorsum versæ, dum in *M. litorali* fere semper sinistrorsæ sunt. Hæc nota *M. littoralis* β , *inermis*, quæ interdum *M. helici* similis est, facile dignoscitur.

M. ORBICULARIS (*All. Fl. Ped.* No. 1150; *DC. Pr.* ii. 174).

Mar. occid.—Casa Blanca!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia). Abyssinia.

MEDICAGO TORNATA (*Willd. Sp. Pl.* iii. 1409; *DC. Pr.* ii. 175)?

Specimina florentia sed nondum fructifera in ditione Tingitana lecta huc retuli, sed hodie mihi omnino incerta videntur.

Species tamen in Marocco septentrionali inquirenda.

M. LITTORALIS (*Rohde in Lois. Not.* 118; *DC. Pr.* ii. 177).

— —, var. *a* (*Moris, Fl. Sard.* i. 439, tab. 40. fig. A), = *M. cylindracea* (*DC. Cat. Monsp. et Prod.* ii. 178).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Prope Cap. Spartel! Inter Tanger et Tetuan!

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

— —, var. *γ*, LONGEACULEATA (*Moris, l. c.* p. 440), = *M. littoralis*, var. *α* (*Ser. in DC. Pr. l. c.*).

Mar. occid.—Frequens in ditione Mogadorensi. Hue verisimiliter *M. littoralis* (Lowe Cat.) referenda est. Legi prope Mogador! juxta Aïn el Hadjar! et in prov. Haha! haud procul Mogador!

Mar. merid.—Prope Ourika! specimen leguminibus intermediis!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia).

Var. *β*, *inermis*, *Moris, loc. cit.* in Marocco nondum detecta est. Hæc sec. cel. *Moris* est eadem ac *M. striata*, Bast. et DC. Prod. Sed in *M. striata* monentibus Gren. et God. (*Fl. Fr.* i. 387) spiræ dextrorsæ, dum in *M. littorali* fere semper sinistrorsæ. Inter specimina plurima agri Mogadorensis tantum duo spiris dextrorsum versis detexi. Beatus Lowe (*Man. Bot. Mad.* 163) hanc cum sequente conjunxit. Re vera proximæ sunt species, sed mihi videntur satis diversæ, nec transitum unquam observavi.

M. TRIBULOIDES (*Desr. in. Lam. Dict.* iii. 635; *DC. Pr.* ii. 178; *Moris, Fl. Sard.* tab. 41).

Mar. sept.—In maritimis prope Tanger! Inter Tanger et Tetuan!

Mar. occid.—Casa Blanca! Agadir (*Coll. Indig.* 1871)!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia).

— —, var. LONGEACULEATA, *nob.* Leguminis aculeæ ipsius diametrum subæquantes attenuatæ.

Mar. sept.—Prope Tanger!

Facie valde diversa, sed differentias graves non video. In spec. Maroccanis spiræ *M. tribuloides* fere semper dextrorsum versæ, dum monentibus Gren. et God. in Gallia sinistrorsum vertuntur, et hujus sententiæ mihi fidem faciunt exemplaria plurima ex Gallia, Hispania, Italia, et Oriente. Inter specimina Algeriensia video nonnulla spiris sinistrorsis, altera dextrorsis. Præter alias notas *M. tribuloides* differt a grege *M. turbinatæ* pedunculis aristatis.

M. MARINA (*L., DC. Pr.* ii. 176).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! In litore Atlantico a Cap. Spartel meridiem versus!

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia). Gallia occidentalis.

MEDICAGO GERARDI (*Willd. Sp. Pl.* iii. 1415; *DC. Pr.* ii. 179).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia). Europa media.

Hæc est fere certe *M. polymorpha*, var. *rigidula*, L., = *M. rigidula* (Desr. in Lam. Diet. iii. 634, vix Willd.). De *M. rigidula* auctores gravissimi dissenserunt. Nomen *M. Gerardi* ab omnibus receptum, etsi recentius, nullomodo mutandum. Affinitatem proximam *M. Gerardi* et *M. turbinatæ* optime indicavit ccl. Boissier in 'Flora Orientali,' ii. pp. 100, 101.

M. TURBINATA (*Willd. Sp. Pl.* iii. 1409; *DC. Pr.* ii. 175; *Boiss. Fl. Or.* ii. 100), var. *ACULEATA* (*Moriš, Fl. Sard.* i. 445, tab. 45 B).

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Amsmiz circ. 1200 m.! Shedma!

In hac legumen apice minus ovoideum potius doliiforme, pubescens, nervo dorsali prominente, aculei breves.

— —, var. *MURICATA*, *Boiss.*, = *M. muricata* (*Benth. Cat. Pyr.* 102; *Guss. Fl. Sic. Prod.* ii. 574, vix Willd.), = *M. neglecta* (*Guss. loc. cit.* 575).

Mar. sept.—Tanger! Tctuan (*Webb Herb.*)! In monte Beni Hosmar, *J. D. H.*!

Mar. occid.—Frustulum huc referendum inter plantas prope Mogador lectas inveni! Casa Blanca!

Mar. merid.—Videtur sat frequens in regione inferiore et in convallibus calidioribus Atlantis Majoris. Mesfioua! Tasscremout! Reraya! In convalle Aït Mesan!

Ar. Geog. *M. turbinata* regionem Mediterraneam imprimis occidentalem incolit. Fertur in Oriente, sed spec. non vidit ccl. Boissier, et deest in Macaronesia. Var. *muricata* in Gallia australi, Hispania et in Imp. Maroccano haud infrequens.

Duce ccl. Boissier *M. muricatam*, Benth., ad var. *M. turbinatæ* reduxi, sed a typo magis differt quam var. *aculeata*, Moriš, præsertim legumine subgloboso, spiris centralibus majoribus, et aculeis longioribus. Spiræ fere semper (vel semper) sinistrorsum versæ. Inter specimina juniora prope Tanger lecta pedunculis bifloris donata, pedunculum unicum triflorum observavi.

M. SPHÆROCARPA (*Bert. Amæn. Ital.* 91; *DC. Pr.* ii. 180).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia), sed infrequens.

MEDICAGO DENTICULATA (*Willd. Sp. Pl.* iii. 1414; *Benth. Cat. Pyr.* 103; *DC. Pr.* ii. 176).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. occid.—Mogador (*Lowe Cat.*)?

Mar. merid.—In convalle Amsmiz!

Ar. Geog. Europa media. Asia temperata et Japonia. Regio Mediterranea omnis (inclusa Macaronesia). India boreali-occidentalis. Abyssinia.

M. lappacea (*Desr. in Lam. Dict.* iii. 637; *Gren. et God. Fl. Fr.* i. 390), var. **PENTACYCLA**, = *M. pentacycla* (*DC. Cat. Monsp.* 124, et *Prod.* ii. 177).

Mar. sept.—Tanger pluribus locis! (*Salzm. Exsicc.*)! (*Webb Herb.*)! Tetuan!

Mar. occid.—Casa Blanca!

Mar. merid.—Prov. Shedma! In ruderatis urbis Marocco!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia).

Auctores gravissimi—*Webb in Phyt. Can.* ii. 64, *Boissier in Fl. Or.* ii. 103—*M. pentacyclam* inter varietates *M. denticulatæ* enumerarunt; et eadem sententia placuit cel. Morisio (*Fl. Sard.* i. 446). Nonnullis *M. lappacea*, *Desr.*, est eadem ac *M. denticulata*, *W.*, aliis est fere ipsissima *M. pentacycla*. Permultis exemplaribus accurate scrutatis, et observationibus cel. *Gren. et Godr. l. c.* rite perpensis, *M. denticulatam* cum var. *M. apiculata* pro typo speciei habui dum *M. lappaceam*, cum var. *pentacycla*, pro subspecie enumeravi. Sunt re vera plantæ proxime affines et notæ diagnosticæ a *Gren. et God.* laudatæ non semper certæ. *M. lappacea* distinguitur pedunculis 5-8-floris nec 2-3-floris, leguminis spiris mediis æqualibus, aculeis divaricatis plerumque multo brevioribus, nec patentibus, seminibus reniformibus, nec oblongis umbilico leviter depressis. Mihi inter specimina Maroccana, Italica et Hispanica *M. pentacyclæ* occurrunt, sed rarius, exemplaria pedunculis quinquefloris. In *M. denticulata*, fructu minore semper donata, aculei fere semper divaricati, dum in *M. lappacea* sunt sæpissime patentés. In iconc Morisiana (*Fl. Sard.* tab. 48), *M. lappacea* nostra (in Italia media sæpia obvia) sat bene delineatur, sed fig. 4 ejusdem tabulæ (legumine extenso aculeis brevibus divaricatis) ad plantam alienam (*M. denticulatam*) certe spectat. Quoad notam diagnosticam e seminibus deductam ego sæpius vidi semina *M. pentacyclæ* oblonga vix emarginata, sed vercor ne hæc nota incerta sit. Vidi specimina seminibus intermediis. Differentiæ inter *M. lappaceam* et *M. pentacyclam* parvi momenti sunt. Prior gaudet leguminibus aliquid minoribus sæpissime trieyclis. In *M. pentacyclæ* exemplaribus minoribus depauperatis legumina trieycla haud raro obvia sunt, sed normalia semper spiris quinque donata.

In spec. nostris Maroccanis spiræ semper sinistrorsum versæ. Vidi spec.

duo Tingitana pedunculis breviter aristatis. Nota igitur ex pedunculis non aristatis haud omnino certa.

MEDICAGO MACULATA (*Willd. Sp. Pl.* 1414; *DC. Pr.* ii. 179).

Mar. sept.—Tanger (*Webb Herb.*)! Tctuan (1851)!

Ar. Geog. Europa media. Regio Mediterranea, sed non ubique. Deest in Macaronesia.

M. LACINIATA (*All. Fl. Ped.* No. 1159; *DC. Pr.* ii. 180).

Mar. occid.—Mazagan!

Mar. merid.—Satis frequens in regione inferiore. Legimus pluries in prov. Shedma! in planitie prope urbem Marocco! in distr. Mesfioua!

Ar. Geog. Regio Mediterranea imprimis calidior (inclusa Macaronesia). Deest in Græcia, Asia Minore, et in Italia australi. Frequens in Africa boreali ultramontana.

Specimina Maroccana quæ ccl. Cosson ex Oasi Akka communicavit, Algeriensibus et Tunetanis omnino conformia sunt. Nostra invicem ludunt aculeis fere duplo longioribus. Pedunculi aristati fere semper uniflori. Legumen maturum in pedunculo semper solitarium. Semina pro genere numerosa, 7-9, oblongo-reniformia.

M. MINIMA (*Lam. Dict.* iii. 636; *DC. Pr.* ii. 178).

Mar. merid.—Ad radices Atlantis Majoris et in convallibus calidioribus. Mesfioua! Aït Mesan! et alibi visa.

Ar. Geog. Europa media. Regio Mediterranea omnis (inclusa Macaronesia). Abyssinia.

Specimina Maroccana ad var. *γ. longisetam* (*DC. Pr. l. c.*) referenda sunt. Sed inutile mihi videtur varietates ex sola longitudine aculeorum (in omnibus Medicaginibus admodum variabili) desumptas enumerare.

M. INTERTEXTA (*All. Fl. Ped.* No. 1152; *Webb, Phyt. Can.* ii. 62), = *M. echinus* (*DC. Fl. Fr.* iv. 546, et *Prod.* ii. 181; *Moris, Fl. Sard.* tab. 52), = *M. muricata* (*Salzm. Exsicc. sec. Webb*).

Mar. sept.—Tanger (*Webb Herb.*)! (*Salzm. Exsicc.*)!

Ar. Geog. Italia. Sicilia. Africa borealis. Macaronesia.

MELILOTUS (*Juss. Gen.* 356).

M. INDICA (*All. Fl. Ped.* No. 1121; *Moris, Fl. Sard.* tab. 56), = *M. parviflora* (*Desf. Fl. Atl.* ii. 192; *DC. Pr.* ii. 187).

Mar. occid.—Mazagan! Mogador frequentissime! (*Lowe Cat.*). Prov. Haha haud procul Mogador!

Mar. merid.—Sat frequens in regione inferiore Atlantis Majoris. In convalle Ourika! Aït Mesan 1200-1400 m.!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). In India

orientali et China (an spontanea?). In America boreali et Cap. Bon. Spei introducta.

MELILOTUS SULCATA (*Desf. Fl. Atl.* ii. 193; *DC. Pr.* ii. 189).

Mar. merid.—Prope Misra Ben Kara!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

— —, var. MAJOR, *Camb.*, = *M. compacta* (*Salzm. in Guss. Prod.*

Fl. Sic. ii. 485), = *M. intermedia* (*Boiss. Voy. Esp.* 475).

Mar. sept.—Tanger! (*Webb Herb.*)! (*Salzm. Exsicc.*)!

Mar. occid.—Casa Blanca! Mazagan!

Ar. Geog. Per totam regionem Mediterraneam hinc inde occurrit.

Specimen ad Mazagan lectum differt stipulis omnibus subintegerrimis.

TRIFOLIUM (*L. Gen.* 896).

T. ANGUSTIFOLIUM (*L.*, *DC. Pr.* ii. 189).

Mar. sept.—Tanger! (*Webb Herb.*)! Tetuan (1851)!

Mar. occid.—Casa Blanca! Djebel Hadid haud procul Mogador (*R. et Fr. spec. in herb. Kew.*)!

Mar. merid.—Prope Amsmiz!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

T. ARVENSE (*L.*, *DC. Pr.* ii. 190).

Mar. sept.—Tanger (*Webb Herb.*)!

Mar. occid.—Casa Blanca!

Ar. Geog. Europa omnis (exclusa arctica). Regio omnis Mediterranea (inclusa Macaronesia). Sibiria Uralensis. Abyssinia.

— —, var. PRESLIANUM, *nob.*, = *T. preslianum* (*Boiss. Diagn. Pl. Or. ser. 1*, ii. 25).

Mar. merid.—Prope Ourika (*R. et Fr. spec. in herb. Kew.*)!

Ar. Geog. Asia Minor. Africa borealis. Forsan alibi in reg. Mediterranea.

Nisi fallor vidi in Oriente formas intermedias.

T. LIGUSTICUM (*Balb. in Act. Acad. Ital.* i. 192; *DC. Pr.* ii. 191).

Mar. sept.—Tanger (*Schsb. Herb.*)!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

T. LAPPACEUM (*L.*, *DC. Pr.* ii. 191).

Mar. sept.—Tanger (*Schsb. Herb.*)! (*Salzm. Exsicc.*)! (*Webb Herb.*)!

Mar. occid.—Casa Blanca!

Mar. merid.—Ad radices Atlantis Majoris. Mesfioua! Amsmiz!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia).

T. BOCCONI (*Savi, Obs. Trif.* 37; *DC. Pr.* ii. 192).

Mar. sept.—Djebel Kebir prope Tanger (*Schsb. Herb.* sub "*T. gemellum*" et "*T. biceps*")!

Mar. occid.—Casa Blanca!

Ar. Geog. Regio Mediterranea imprimis occidentalis (inclusa Macaronesia). Europa occidentalis. Deest in Asia Mediterranea.

In spec. nonnullis Gallieis dentes calycini minus inæquales; in nostris et in plurimis a me scrutatis dentes cal. inferiores superioribus duplo longiores et corollam excedentes.

TRIFOLIUM ATLANTICUM (*Ball in Journ. Bot.* 1873, 305). (Plate XVIII.)

Annum, pumilum, e basi ramosum; herba tota molliter villosa; folia in ramis prostratis subæqualia, longiuscule petiolata, foliolis obovato-euneatis vix denticulatis; stipulæ superiores dilatatæ, breviter acuminatæ, insigniter nervosæ; capitula gemella, cylindracea, sessilia, in ramis terminalia; calycis dentes villosi subæquales, corollam pallide rubentem vix attingentes; semina minima, orbiculata, subcompressa, radicle non prominula.

Mar. merid.—In regione subalpina Atlantis Majoris. Legi in convalle Aït Mesan! circ. 2100 m.

Ab affini *T. Bocconi*, Savi, distinguitur imprimis ramis debilibus, nec strietis, indumento villosa, foliis omnibus longiuscule petiolatis, nec inferioribus breviter petiolatis superioribus subsessilibus, calycis dentibus mollibus subæqualibus, nec rigidis inæqualibus, seminibus orbicularibus, nec reniformi-ovoideis, radicle prominula donatis.

T. SCABRUM (*L.*, *DC. Pr.* ii. 192).

Mar. sept.—Tanger (*Schsb. Herb.*), *Coss. spec.*! Inter Tanger et Tetuan! Tetuan (1851)!

Mar. occid.—Mazagan!

Mar. merid.—In umbrosis regionis inferioris. Sheshaoua! Amsmiz!

Ar. Geog. Europa media. Regio omnis Mediterranea (inclusa Macaronesia).

T. MARITIMUM (*Huds. Fl. Angl.* 284; *DC. Pr.* ii. 192).

Mar. sept.—Tanger (*Schousb. Herb.* sub "*T. squarrosus*"), *Coss. spec.*! (*Webb Herb.*)! Prope Cap. Spartel! In regione collina inter Tanger et Tetuan!

Ar. Geog. Europa occidentalis. In regione Mediterranea late dispersa sed infrequens. In Macaronesia nonnisi in ins. Madeira.

T. BRACTEATUM (*Schousb. in Willd. Enum.* 792; *DC. Pr.* ii. 195), = *T. leucophæum* (*Schousb. MSS. icon. ined.*) forma floribus albis.

Mar. sept.—Tanger (*Schsb. Herb.*, "In collibus aprieis inter *Chæmaropem humilem*"), *Coss. spec.*! et iterum (sub "*T. leucophæum*, Schsb.")! (*Webb Herb.*, sub "*T. pallidum*, W. K. ? In rupestribus Tingitanis")! Eandem plantam legit Schousboe in monte Djebel Kebir.

Cel. *Coss.* hoc pro lusu *T. pratensis*, L., habet; mihi tamen, licet dubitative, pro specie vel saltem pro subspecie enumerandum videtur. A *T. pratensi* differt stipulis chartaceis, a nervis parallelis prominentibus eximie striatis, sub capitulo valde expansis, fere rotundatis,

apice acuminato-setaceis, et calycis laciniis magis inæqualibus. Plantæ nostræ var. levis est *T. baticum* (Boiss. Voy. Esp. Append. 725; Walp. Ann. ii. 349). Hoc parum differt caulibus elatioribus, habitu laxo, calycis dente inferiore superioribus subduplo-, nec sesquiflongiore.

Ar. Geog. Species in utroque litore freti Hereulei obvia; nec alibi?

TRIFOLIUM CHERLERI (L., DC. Pr. ii. 196).

Mar. sept.—Tanger (*Schsb. Herb.*), *Coss. spec.*!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

T. STELLATUM (L., DC. Pr. ii. 197).

Mar. sept.—Tanger! Tetuan! (*Webb Herb.*)!

Ar. Geog. In tota regione Mediterranea frequentissimum, hucusque nonnisi in Marocco septentrionali detectum.

T. GLOMERATUM (L., DC. Pr. ii. 198).

Mar. sept.—Tanger! Djebel Kebir (*Schsb. Herb.*), *Coss. spec.*!

Mar. merid.—Ad radices Atlantis Majoris. Mesfioua! Ourika!

Ar. Geog. Europa occidentalis. Regio omnis Mediterranea (inclusa Macaronesia). Deest in Græcia et Macedonia.

— —, var. *CONDENSATUM*, nob. Forma pumila, condensata, pauciflora, calycis dentibus et stipulis magis herbaceis, pilosiusculis.

Mar. merid.—In regione subalpina et alpina Atlantis Majoris. In convalle Aït Mesan et in jugo Tagherot a 2400 m. usque 3000 m.!

T. STRICTUM (L., DC. Pr. ii. 198), = *T. lævigatum* (*Poir. Voy.* ii. 219; *Desf. Fl. Atl.* ii. 195, tab. 208).

Mar. sept.—In monte Djebel Kebir prope Tanger (*Schousb. Herb.* sub "*T. lucidum*"), *Coss. spec.*!

Ar. Geog. Europa media, in Gallia et Hungaria. Regio Mediterranea sed non ubique. Deest in Asia Mediterranea, in Græcia et in Macaronesia.

T. REPENS (L., DC. Pr. ii. 198).

Mar. sept.—Inter Tanger et Tetuan (*J. D. H.*)! Tetuan (1851)!

Mar. merid.—In regione inferiore Atlantis Majoris. Prope Ourika (*R. et Fr. spec. in herb. Kew.*)! Aït Mesan circ. 1500 m.! Taradut (*R. et Fr. spec.*)!

Ar. Geog. Europa omnis. Sibiria. America borealis. In omni regione Mediterranea late diffusa, sed in calidioribus infrequens.

T. HUMILE (*Ball, in Journ. Bot.* 1873, 305).

Perenne, pumilum, cæspitosum; rhizoma lignosum, ramosum, pedunculos subpollicares petiolis subæquales edens; foliorum foliola obovata, minute denticulata; stipulæ basi seariosæ, elongatæ, in apicem viridem sensim attenuatæ; capitulum ex pedicellis eiliatis braeteolatis demum reflexis laxum; calycis paræ eiliati dentes inæquales tubo 10-nervi breviores, nervis intermediis sub sinu bifurcatis, unde dentes

trinerves; corolla (pro grege) parva; legumen sessile, biovulatum; semina compressa, suborbicularia, hilo breviter emarginata, lævissima.

Mar. merid.—In regione alpina Atlantis Majoris. Legi in jugo Tagherot ad 3000 m.!

Proximum *T. cæspitosum*, Reyn., differt stipulis cuspidato-acuminatis, nec apicem versus attenuatis, pedicellis erectis, nec reflexis, bractcolis brevioribus, calycis dentibus brevioribus. Nostra species fere media est inter *T. cæspitosum*, Reyn., et *T. Parnassi*, Boiss. et Sprun., a quo differt, præter alias notas, seminibus lævissimis nec minute granulatis.

Melius observatæ hæ formæ ad subspecies forsan reducendæ, quibus ad sociandum erit *T. pallescens*, Schreb. In Alpibus Helveticis, Sabaudis, et Pedemontanis characteres differentiales inter *T. cæspitosum* et *T. pallescens* satis stabiles ego semper observavi, sed in Alpibus Carinthiæ et Carniolæ formæ intermediæ mihi obviæ sunt. Simili modo specimina *T. Parnassi* accuratius explorata inter se discrepant. Specimina ex monte Parnasso ostendunt dentes calycinos valde inæquales, quorum superiores tubo longiores sunt; spec. ex monte Taygeto gaudent contra dentibus brevissimis; spec. denique ex monte Oleno prioribus intermedia sunt.

Si pro subspeciebus unius ejusdemque speciei has formas habemus, nomen *T. cæspitosum*, antiquius et sat congruum, pro typo retinendum.

TRIFOLIUM ISTHMOCARPUM (*Brot. Phyt. Lus.* 148, tab. 61; *DC. Pr.* ii. 201), = *T. mauritanicum* (*Salzm. Exsicc.*).

Mar. sept.—Tanger! (*Schousb. Herb.* sub "*T. rubicundum*" et "*T. erythraeum*"), *Coss. spec.*! (*Webb Herb.*)! (*Salzm. Exsicc.*)!

Mar. occid.—Casa Blanca!

Ar. Geog. Africa boreali-occidentalis. Peninsula Iberica. Sicilia.

Foliola argute serrulata, apice rotundata vel acuta; stipulae vaginantes parte libera lanecolato-subulatæ; flores in capitulo sessiles, basi minute bracteolati; dentes calycis ex nervo prominulo subulati. Flores in spec. Algarbiensibus et Algeriensibus videntur albidī, in Maroccanis semper rubri vel carnei.

T. SUBTERRANEUM (*L., DC. Pr.* ii. 202).

Mar. sept.—Tanger! In monte Beni Hosmar prope Tctuan!

Ar. Geog. Europa occidentalis. Regio Mediterranea omnis (inclusa Macaronesia).

T. SPUMOSUM (*L., DC. Pr.* ii. 202).

Mar. occid.—Casa Blanca!

Ar. Geog. Regio Mediterranea (exclusa Macaronesia).

Legumina in spec. nostris semper tetrasperma, in rostrum viride attenuata, rostro longe exserto.

TRIFOLIUM RESUPINATUM (L., DC. Pr. ii. 202).

Mar. sept.—Tanger! (*Webb Herb.*)! Tetuan ad ostium fluminis (1851)!

Mar. merid.—Ad radices Atlantis Majoris. Mesfioua! Amsmiz! Reraya!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia). Gallia occidentalis.

T. FRAGIFERUM (L., DC. Pr. ii. 202).

Mar. sept.—Tanger (*Salzm. Exsicc.* sub "*T. macrocephalum*")! In monte Djebel Kebir! Tetuan (1851)!

Ar. Geog. Europa omnis (exclusa arctica). Regio Mediterranea (inclusa Macaronesia).

T. TOMENTOSUM (L., DC. Pr. ii. 203).

Mar. sept.—Tanger! Inter Tanger et Tetuan!

Mar. merid.—In regione inferiore et media Atlantis Majoris. Mesfioua! In convalle Aït Mesan circ. 2000 m. (*J. D. H.*)!

Ar. Geog. Regio omnia Mediterranea (inclusa Macaronesia).

T. PROCUMBENS (L., DC. Pr. ii. 205), var. CAMPESTRE (*Seringe in DC. Pr. l. c.*), = T. campestre (*Schreb. in Sturm D. Fl. fasc. 16*), = T. agrarium (*Poll. Fl. Pal. ii. 342*; *Savi, Bot. Etr. iv. 48*; *Gren. et God. Fl. Fr. i. 423*; non *Schreb.*). Confer Benth. in *Brit. Fl.*

Mar. sept.—Tetuan! (*Webb Herb.*)!

Mar. occid.—Mogador (*Lowe Cat.*). Casa Blanca!

Mar. merid.—Ad radices Atlantis Majoris. Mesfioua! Onrika!

Ar. Geog. Europa omnis (exclusa arctica). Regio Mediterranea (inclusa Macaronesia). Abyssinia.

ANTHYLLIS (L. Gen. 864).

A. GERARDI (L., DC. Pr. ii. 169).

Mar. sept.—Tanger (*Webb Herb.*)! (*Salzm. Exsicc.*)! In monte Djebel Kebir!

Ar. Geog. Regio Mediterranea occidentalis a Corsica et Sardinia usque Lusitaniam. Deest in Macaronesia.

A. VULNERARIA (L., DC. Pr. ii. 170).

Mar. sept.—Prope Tetuan, et in monte Beni Hosmar (*Webb Herb.*)!

Mar. merid.—In regione inferiore et subalpina Atlantis Majoris. In convalle Aït Mesan usque 2400 m.!

In spec. nostris flores pallide citrini. Folia primaria simplicia, caulina 4-6-juga, foliolo terminali maximo vel cæteris conformi.

Ar. Geog. Europa omnis (exclusa arctica). Regio Mediterranea fere omnis (exclusa Macaronesia). Abyssinia.

— —, var. DILLENII, = A. Dillenii, *Schult.*

Mar. occid.—Djebel Hadid hand procul Mogador (*R. et Fr. spec. in herb. Kew.*)!

Mar. merid.—In regione inferiore et ad radices Atlantis Majoris. Tassremout! Seksaona! Aït Mesan 1200-1500 m.!

ANTHYLLIS TEJEDENSIS (*Boiss. Bibl. Gen.* 1838, *Elench.* 58; *Walp. Rep.* i. 634).

Mar. sept.—In summis rupibus montis Beni Hosmar prope Tetuan !

Ar. Geog. Hispania Granatensis. Montes Tetuanenses.

Specimina nostra nondum florentia sed Hispanicis adamussim conformia.

A. HAMOSA (*Desf. Fl. Atl.* ii. 151; *DC. Pr.* ii. 171).

Mar. occid.—Haud procul Mogador, juxta Aïn el Hadjar !

Ar. Geog.—Africa borealis. Hispania Granatensis.

Unicum specimen inveni, verisimiliter depauperatum, foliis bijugis foliolis angustis; tubo calycino gracili, multo minus hirsuto quam in spec. Algeriensibus, et dentibus magis inæqualibus. Planta ulterius inquirenda.

A. TETRAPHYLLA (*L., DC. Pr.* ii. 171), = *Physanthyllis tetraphylla* (*Boiss. Voy. Esp.* 162).

Mar. sept.—Tanger ! (*Webb Herb.*) ! Tetuan !

Mar. merid.—Ad radices Atlantis Majoris et in convallibus inferioribus.

Ourika ! Seksaoua ! Sektana ! In convalle Aït Mesan !

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

DORYCNIUM (*Vill. Fl. Dauph.* iii. 416).

D. RECTUM (*L., sub Loto; Ser. in DC. Pr.* ii. 208).

Mar. sept.—Tanger !

Mar. merid.—Juxta Gurguri prope Amsniz !

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia).

—— —, var. PAUCIFLORUM, *nob.* Capitulis multo minoribus, paucifloris.

Mar. merid.—Inter Ourika et Reraya ! Prope Ourika (*R. et Fr. spec. in herb. Kew.*) !

LOTUS (*L. Gen.* 879).

L. EDULIS (*L., DC. Pr.* ii. 209).

Mar. sept.—Tanger (*Schsb.*) *Coss. spec.* ! Tetuan (1851 et 1871) ! In monte Djebel Dersa prope Tetuan (*Webb Herb.*) !

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia).

L. ORNITHOPODIOIDES (*L., DC. Pr.* ii. 209).

Mar. sept.—Tanger ! (*Salzm. Exsicc.*) ! (*Webb Herb.*) ! Tetuan (1851) !

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

L. ARENARIUS (*Brot. Fl. Lus.* ii. 120; *DC. Pr.* ii. 210), = *L. anran-tiacus* (*Boiss. Voy. Esp.* 174, tab. 53), = *L. canescens* (*Kunze, Chlor. Hisp.* No. 534).

Mar. sept.—Tanger ! (*Salzm. Exsicc.*) ! (*Webb Herb.*) !

Mar. occid.—Casa Blanca ! Mazagan ! Mogador (*Lowe Cat.*) In prov.

Haha haud procul Mogador !

Mar. merid.—In provinciis meridionalibus cisatlanticis, valde frequens in planitie et ad radices Atlantis Majoris. Legimus pluries in prov. Shedma! Mesfioua! Amsmiz! Ourika!

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

— —, var. WEBBII, nob., = *L. dumetorum* (*Webb, MSS. ined., et in Bourgeau Pl. Can.*).

Mar. occid.—In arenosis prope Mogador! Aïn el Hadjar! Agadir (*Coll. Indig. 1871*)!

Mar. merid.—In saxosis aridissimis circa Aïn Oumast, et alibi in prov. Shedma!

Ar. Geog. Macaronesia. Marocco meridionali-occidentalis.

Differt a specie habitu procumbente, herba hirsutissima subincana, foliis minoribus; sed characteres graviores satis constantes non video. *L. arenarius* est species valde indibunda, et formæ intermediæ non desunt. Hinc etiam referenda sunt (nisi fallor) specimina nana exilia pauciflora a nobis in distr. Mesfioua lecta. Nomen *dumetorum*, saltem quoad spec. Maroccana, haud aptum, dum hæc varietas semper in arenosis vel saxosis solo arso obvia est.

Legi in umbrosis prope Sheshaoua specimen incompletum forsitan ad *L. arenarium* referendum, leguminibus elongatis polyspermis, pedicellis fructiferis omnibus insigniter recurvis, segmentis calycinis paulo longioribus et acutioribus.

LOTUS CRETICUS (*L., DC. Pr. ii. 211*).

Mar. sept.—In arenosis inter Tetuan et Ceuta!

Mar. occid.—Mazagan in arenosis maritimis! Mogador in arenosis!

Ar. Geog. Regio Mediterranea (exclusa Macaronesia).

L. commutatus (*Guss. Fl. Sic. Prod. ii. 545*), = *L. Salzmanni* (*Boiss. et Reut. Pug. 37; Walp. Ann. iv. 477*).

Mar. sept.—Tanger in rupibus maritimis! Tanger in arenosis (*Salzm. Exsicc.*)! (*Boiss. et Reut. l. c.*). Tanger (*Schsb. Herb. sub "L. creticus"*) *Coss. spec.*! In rupibus maritimis ad Cap. Spartel!

Mar. occid.—Casa Blanca! Agadir (*Coll. Indig. 1871*)!

Ar. Geog. Africa boreali-occidentalis. Hispauia Gaditana. Algarve. Sicilia.

Specimina Sicula non vidi, sed descriptio Gussoniana omnino conformis. Planta Salzmanniana a *L. cretico* differt habitu, caulibus suberectis rigidis basi sublignosis, pedunculis longioribus crassioribus, floribus majoribus pallidioribus. Sistit forsitan subspeciem, sed vereor ne in typum per formas intermedias transeat.

L. CYTISOIDES (*DC. Fl. Fr. et Prod. ii. 211*, et verisimiliter *L.*), = *L. Allionii* (*Desr. Journ. Bot. iii. 77; Gren. et God. Fl. Fr. i. 433*), = *L. creticus*, var. *cytisoides* (*Boiss. Fl. Or. ii. 165*).

Mar. sept.—Tetuan (1851 et 1871)! In monte Beni Hosmar prope Tetuan!

Mar. oeeid.—? In monte Djebel Hadid haud procul Mogador (*J. D. H.*)!?

Mar. merid.—In convalle Ait Mesan eire. 1300 m.! Prope Ourika (*R. et Fr. spec. in herb. Kew.*)!

Ar. Geog. Species variabilis per totam fere regionem Mediterraneam diffusa. Deest in Macaronesia.

LOTUS MAROCCANUS (*Ball, in Journ. Bot.* 1873, 306).

Radix lignosa, e collo ramosissima, caules adscendentes vel ereeti, subæquales; folia breviter petiolata, petiolo compresso insigniter 3-nervi; foliola sessilia, fol. inferiorum obovata, superiorum rhombeo-lanceolata, omnia acuminata; stipulae conformes; pedunculi terminales, rarius laterales, 1–4-flori, folio florali cæteris conformi suffulti; pedicelli reflexi, articulati; calycis dentes subæquales, subulati, tubo nervis 5 atrovirentibus notato subduplo longiores, sinu rotundato divisi; stylus dentatus; legumen (pro genere) longum, nitidum, torulosum, ineurvum; semina viridia, ovoidea.

Mar. oeeid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In glareosis ad radices Atlantis Majoris frequens. Tasseremout! Ourika! Reraya! Amsmiz! Seksaoua!

Species insignis, nulli proxima. Tota planta, præter legumen, molliter villosa.

L. HISPIDUS (*Desf. Cat. H. Par.* 190; *DC. Pr.* ii. 212), = *L. odoratus* (*Sims, Bot. Mag.* tab. 1233; *DC. Pr.* ii. 214).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)! In monte Djebel Kebir (*Schsb. Herb.* sub “*L. odoratus*”), *Coss. spec.*!

Ar. Geog. Europa occidentalis. Regio Mediterranea occidentalis (inclusa Maearonesia). Fertur in Græcia, sed deest in Boiss. ‘*Fl. Orientali.*’

Valde affinis *L. angustissimo*, L., et forsā melius sicut hujus subspecies enumerandus.

L. PARVIFLORUS (*Desf. Fl. Atl.* ii. 206; *Seringe, in DC. Pr.* ii. 208, sub *Dorycnio*), = *L. microearpus* (*Brot. Fl. Lus.* ii. 219; *Ser. in DC. Pr.* ii. 209, sub *Dorycnio*).

Mar. sept.—Tanger (*Schsb. Herb.* sub “*L. humilis*” et “*L. pilosus*”), *Coss. spec.*! (*Webb Herb.*)!

Mar. oeeid.—Rabat (*R. et Fr. spec. in herb. Kew.*)! Casa Blanca!

Ar. Geog. Regio mediterranea imprimis occidentalis (inclusa Maearonesia). Deest in Asia Mediterranea.

L. LAMPROCARPUS (*Boiss. Diagn. Pl. Or.* ser. 1, ix. 33; *Walp. Ann.* ii. 356)?

Mar. merid.—Mesfiona in fruticetis juxta torrentem!

Ar. Geog. In Græcia et Asia Mediterranea late diffusa.

Specimina nostra leguminibus omnino immaturis non satis certa. Spec. Heldreichianis ad Phalerum Atticæ lectis bene conveniunt nisi corollis in sicco aurantiacis, nec ut in his virescentibus, et legumine (immaturo) fusco, nec cinereo-nitente.

LOTUS TINGITANUS (*Boiss. Diagn. Pl. Or. ser. 1, ii. 36; Walp. Rep. ii. 854*).

Mar. sept.—“In arenis Tingidis in Regno Maroccano” (specimen in herb. Pavon, hodie Boissier, a Schousboeo missum), *Boiss. l. c.*

Species ex descript. insignis, floribus solitariis, legumine calycem parum superante, mihi omnino incerta et ditionis nostræ dubia civis.

In herbario b. Schousboe non adest, nec viderunt botanici plurimi qui agrum Tingitanum postea perlustrarunt.

L. TETRAGONOLOBUS (*L., DC. Fl. Fr. iv. 553*), = *Tetragonolobus purpureus* (*Mænoch, Meth. 164; DC. Pr. ii. 215*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Tetuan (*Webb Herb.*)!

Ar. Geog. Regio Mediterranea (exclusa Macaronesia).

L. SILIQUOSUS (*L., DC. Pr. ii. 215*, sub *Tetragonolobo*).

Mar. merid.—In umbrosis irrigatis juxta Sheshoua aliis stirpibus Europæ mediæ adsociata!

Ar. Geog. Europa media. Per regionem Mediterraneam (exclusa Macaronesia) dispersa, sed infrequens.

L. CONJUGATUS (*L., DC. Pr. ii. 215*, sub *Tetragonolobo*).

Mar. sept.—Tanger (*Schsb. Herb.*), *Coss. spec.*!

Ar. Geog. Oriens. Hispania. Ager Tingitanus. Varietas in Sicilia legumine subalato.

SECURIGERA (*DC. Fl. Fr. iv. 609*).

S. CORONILLA (*DC. l. c. et Prod. ii. 313*).

Mar. sept.—Tanger!

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

Genus *Securigera* inter *Loteas* et *Coronilleas* fere medium tenet. Duce cel. Boissier (*Fl. Or. ii. 176*) has post *Loteas* enumeravi, et ab *Hedysareis* in serie lineari separavi.

SCORPIURUS (*L. Gen. 886*).

S. SULCATA (*L., DC. Pr. ii. 308*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Tetuan (*Webb Herb.*)!

Mar. occid.—Casa Blanca!

Mar. merid.—Mesfioua!

Ar. Geog. Regio Mediterranea calidior (inclusa Macaronesia). Abyssinia. Decet in Gallia, Italia, Græcia, et Asia Minore.

Scorpiurus subvillosa (L., DC. Pr. ii. 308).

Mar. sept.—Inter Tanger et Tetuan ! Tetuan (1851)!

Ar. Geog. Regio Mediterranea omnis (inclusa Maearonesia).

Nimis vicina *S. sulcatæ* et sicut subspecies enumeranda.

S. VERMICULATA (L., DC. Pr. ii. 308).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Regio Mediterranea occidentalis inclusa Maearonesia, sed potius rara. Fertur in Attica (*Sibth.*), et in Tauria (*M. B.*), sed specimen Orientalia Boissier non vidit.

ORNITHOPUS (L. Gen. 884).

O. COMPRESSUS (L., DC. Pr. ii. 311).

Mar. sept.—Tanger ! (*Webb Herb.*)!

Mar. occid.—Rabat (*R. et Fr. spec. in herb. Kew.*)! Casa Blanea !

Ar. Geog. Regio Mediterranea omnis (inclusa Maearonesia). Gallia occidentalis.

O. ISTHMOCARPUS (*Coss. Not. Crit.* 16 ; *Walp. Ann.* iii. 849).

Mar. sept.—Tanger ! (*Webb Herb.*)! In monte Djebel Kebir (*Salzm. Exsicc.*, sub "*O. intermedius* ")!

Ar. Geog. Peninsula Iberica. Ager Tingitanus.

Confer Boiss. et Reut. Pug. in Addend. p. 130. Nil dubitandum b. Brotero species duas probe diversas sub *O. sativo* (Fl. Lus. ii. 160) confudisse—scilicet *O. isthmocarpum*, Coss., et *O. roseum*, L. Dufour. Prior distinguitur legumine valde arcuato, inter semina in isthmos elongatos contracto, et in rostrum elongatum uncinatum desinente. *O. sativus* (Gren. et God. Fl. Fr. i. 499, et plurimorum auctorum) est certe idem ac *O. roseus* (L. Duf. Ann. Sc. Nat. ser. 1, v. 84). Hunc, præeunte Seringe in DC. Pr. ii. 312, habeo pro forma majore *O. perpusilli*, L.

O. EBRACTEATUS (*Brot. Fl. Lus.* ii. 159 ; *DC. Pr.* ii. 311, sub *Astrolobio*).

Mar. sept.—Tanger ! (*Webb Herb.*)! (*Salzm. Exsicc.*)! In monte Djebel Kebir (*Schsb. Herb.*), *Coss. spec.*! Cap. Spartel!

Ar. Geog. Regio Mediterranea omnis, imprimis occidentalis. Europa occidentalis. In Oriente rarus.

CORONILLA (L. Gen. 883).

C. GLAUCA (L., DC. Pr. ii. 309).

Mar. sept.—In rupibus montis Beni Hosmar pr. Tetuan florentem legimus!

Ar. Geog. Regio Mediterranea. Deest in Oriente et in Maearonesia.

Specimina nostra stipulis angustis lanceolato-acuminatis certe ad *C. glaucam* referenda. Folia fere omnia apice lata, emarginata, in sinu plerumque mucronata.

CORONILLA PENTAPHYLLA (*Desf. Fl. Atl.* ii. 171; *DC. Pr.* ii. 309).
Mar. merid.—In regione inferiore Atlantis Majoris. Aït Mesan circ.
1300 m.!

Ar. Geog. Africa boreali-occidentalis.

C. pentaphylla *C. glauca*, L., simillima, et nisi stipulurum forma vix
dignoscenda, forsan melius sicut subspecies enumeranda.

C. JUNCEA (L., *Lam. Dict.* ii. 121; *DC. Pr.* ii. 309).

Mar. merid.—Ex prov. Ksima attulit Indig. Mardocheus. *Coss. Index.*

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

Forma leguminis in hac specie post Lamarek ab auctoribus vel male descripta vel prætermissa. In multis exemplaribus diligenter scrutatis ego vidi legumina semper verbis Lamarekianis conformia, “Gousses menues, articulées, légèrement comprimées, et munies dans la longueur de deux petits rebords opposés,” nec tetragona ut habent Gren. et God. *Fl. Fr.* i. 496.

C. RAMOSISSIMA (*Ball in Journ. Bot.* 1873, 306).

Suffrutex ramosissimus, 1-2-pedalis; rami substriati; folia brevina, imparipinnata, 2-3-juga, foliolis breviter petiolulatis, crassiusculis, cuneatis, apice truncatis emarginatis, mucronulatis, margine non cartilagineis, infimis cauli approximatis; stipulae membranaceae, e basi lata ovato-acuminatae; pedunculi 7-10-flori, folia longe superantes; flores pro genere parvi; calycis dentes breves, apice interdum ciliolati; vexilli lutei unguis calyce paulo longior; legumen tenue, leviter arcuatum, vix compressum, ex suturis incrassatis insigniter biangulatum.

Mar. merid.—Frequens in regione inferiore Atlantis Majoris. Distr. Reraya 1000-1200 m.! Aït Mesan circ. 1400 m.!

Hanc in *Journ. Bot.* l. c. habui pro subspecie *C. juncea*, L. Melius observata mihi videtur ab illa probe distincta. Forma foliolorum omnino diversa, et foliola infima ramo approximata (saepè sessilia), notas graves differentiales in hoc genere suppetunt. His addenda sunt legumina tenuiora longiora minus compressa, et stipulae majores basi latiores.

C. MINIMA (L., *Vill. Fl. Dauph.* iii. 396, non *DC. Pr.* ii. 309), = *C. coronata* (*DC. Pr.* ii. 310).

Mar. merid.—In convalle Aït Mesan circ. 1600 m.!

Ar. Geog. Europa media praesertim occidentalis. Hispania. Atlas Major.

C. VIMINALIS (*Salisb. Lond.* 13; *DC. Pr.* ii. 309), = *C. pulchra* (*Ball in Journ. Bot.* 1873, 307).

E basi suffruticosa rami tenues, flexuosi, herbacci, elongati (inter frutices 9-10-pedales); folia 4-10-juga; foliola petiolulata, ovato-cuneata vel oblongo-elliptica, apice saepius emarginata, vix mucronulata, infe-

riora eauli approximata, non amplexentia; stipulæ parvæ, albæ, ovales, obtusæ; flores in pedunculis axillaribus 4-8, magni, colore variabiles, interdum flavescentes, vexillo linea rubra notato, interdum albi vel dilute purpurei, alis saturatioribus; calyx quam in sp. affinis multo major, campanulatus, dentibus latis brevissimis vel subobsoletis, latere superiore basin versus subinflatus; petala subæqualia, omnia latiora quam in affinis; legumen longissimum (nondum maturum decimetrum superans), 10-15-articulatum, ex stylo elongato torto persistente eximie rostratum; semina subcylindrica, compressiusecula.

Mar. occid.—Mogador (*Salisb. l. c.*).

Mar. merid.—In regione inferiore et ad radices Atlantis Majoris. Infra Tasseremout inter frutices! Supra Seksaoua in rupibus! Supra Milhain! Aouina prov. Shedma (*Coss. spec.*)!

Ex descriptione nimis incompleta, et pluribus notis incongrua, specimina nostra ad *C. viminalem* non retuli, et pro typo sp. novæ perperam descripsi. Nomen ipsum *viminalis* infaustum, dum rami tenues flexuosi hyeme pereunt, et nisi fruticibus aliove sustentaculo fulcrati breves evadunt. Flores in spec. nostris, ex albo purpurei; in spec. ad Mogador lectis dicuntur flavescentes; confer iconem Salisburianam. Foliola in spec. nostris ad 17, in spec. Mogadorensibus usque 21.

Sec. Salisbury *l. c.* planta ex seminibus facile propagata, in hortis plurimis Anglicis suo tempore culta, serius ut videtur amissa, et hodie paucis cognita.

Pulchra species, decus floræ Maroccanæ, nullæ aliæ valde proxima. *C. grandiflora*, Boiss., quæ floribus æmula, est species herbacea, calyce diverso, et stipulis majoribus ovatis mucronatis, nec parvis ovalibus obtusis, a nostra bene distincta; *C. varia*, L., floribus multo minoribus 12-18 nec 4-8 in pedunculo gaudens, differt insigniter calycis dentibus triangularibus, acuminatis, legumine multo brevior, 4-6-articulato.

In herb. Gouan (hodie Kewensi) exstat specimen unicum *Coronillæ viminalis* cui adieitur in scheda manu ignota "*Coronilla gibraltarica* de Broussonnet. Ex Horto Barcinouensi, 1797. Affinis *C. variæ* an non eadem?" Semina proculdubio ex agro Mogadorensi a cl. Broussonnet allata, et cum Gibraltariis commixta.

CORONILLA REPANDA (*Lam. Ill. tab. 631, et Dict. iv. 580; Poir. Voy. ii. 215, sub Ornithopo; DC. Pr. ii. 311, sub Astrolobio; Boiss. Diagn. Pl. Or. ser. 2, ii. 35*), = *Ornithopus lotoides* (*Viv. ex DC. Pr. loc. cit.*).

Mar. sept.—Prope Cap. Spartel (*Schsb. Herb.*), *Coss. spec.*!

Ar. Geog. Occurrit rarius in regione Mediterranea occidentali.

C. SCORPIOIDES (*L., sub Ornithopo; M. et K., Deutschl. Fl. v. 201*), = *Astrolobium scorpioides* (*DC. Pr. ii. 311*).

Mar. occid.—Casa Blanca! Agadir (*Coll. Indig. 1871*)!

Mar. merid.—Sat frequens in regione inferiore. Mesfoua! Tasseremout (*G. Maw*)! Milhain! Aït Mesan circ. 1300 m.!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

HIPPOCREPIS (*L. Gen.* 885).

H. SCABRA (*DC. Pr.* ii. 312).

Mar. merid.—Prope Seksaoua!

Ar. Geog. Hispania. Africa borealis.

Specimina nostra ex legumine nimis immaturo haud certa.

H. scabram genuinam vidi ex Cerro Negro pr. Matritum (*Bourg. Pl. Esp.* 1854) et ex Djebel Tougourt pr. Batna (*Balansa*, 1853, No. 924). Cætera specimina quæ vidi a Bourgeau et Reuter sub hoc nomine distributa omnes ad speciem omnino diversam (*H. Salzmanni*) referenda. *H. scabra* est species perennis (erronee in *DC.* (I) notata), ex pilis brevibus adpressis scabriuseula, et calyce brevi donata.

H. ATLANTICA (*Ball in Journ. Bot.* 1873, 307).

Fruticulus, radice et caulibus lignosis, ramos prostratos foliaceos edens; herba tota pilis albis brevibus vestita; folia 6–10-juga; foliola approximata, petiolulata, obtusissima, fol. infer. apice retusa; stipulæ minutæ, membranaceæ, ovatæ, obtusæ; pedunculi terminales, breves, 3–6-flori; calyceis labio superiore apice bidenticulato, inferiore 3-dentato, dentibus subæqualibus; legumen (immatuum) areuatum, scabrum, sinibus profundis orbicularibus excavatum.

Mar. merid.—In regione alpina Atlantis Majoris. In jugo Tagherot circ. 2700 m.! Fere eodem loco specimen legerunt el. Rein et Fritsch!

Ab affini *H. scabra*, *DC.*, differt foliis multijugis, foliolis petiolulatis fere oboordatis, pedunculis brevioribus et stipulis obtusis.

H. MULTISILIKUOSA (*L., DC. Pr.* ii. 312).

Mar. merid.—Saksaoua! Tasseremout (*G. Maw*)!?

Ar. Geog. Per regionem Mediterraneam omnem (inclusa Macaronesia) dispersa.

Specimen unicum ex Tasseremout nimis immaturum et ideo incertum.

— —, var. MAJOR, = *H. minor* (*Munby, Fl. de l'Alg.* 80; *Walp. Ann.* i. 245).

Mar. merid.—Ad radices Atlantis Majoris et in planitie. Amsmiz! Mskala! Aït Mesan circ. 1250 m.!

Nomen *H. minor* infaustum; hæc est enim *H. multisiliquosæ* forma major, pedunculis quam folium duplo longioribus, nec folio subæqualibus, legumine sublato recto vel parum areuato, nec insigniter enervato. Inter *H. multisiliquosam* et *H. Salzmanni*, quæcum b. Munby (*l. c.* in *Add p.* 116) hanc infeliciter conjunxit, locum medium tenet, sed notis

gravioribus ad priorem accedit. In spec. Algeriensibus calyces hirsuti, in nostris tantum margine ciliati.

HIPPOCREPIS SALZMANNI (*Boiss. et Reut. Pug.* 42). In Walpers Ann. prætermisssa.

Differt a proxima *H. multisiliquosa* statura multo majore, floribus duplo majoribus, legumine sesquilatiore, et præsertim calycis tubo fere duplo longiore dentibus lanceolatis sinu angustiore separatis, nec cuspidato-acuminatis sinu rotundato divisis. Herba subglaberrima pilis raris præsertim in pedunculo adspersa. Stipulæ fere conformes, sed parce nec eximie ciliatæ.

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Boiss. et Reut.*) ex *loc. cit.*
Prope Cap. Spartel!

Mar. occid.—? Aïn el Hadjar haud procul Mogador! Specimen unicum exile, calycis dentibus quam in typo brevioribus.

Ar. Geog. Hispania Gaditana. Imper. Maroccanum.

PSORALEA (*L. Gen.* 894).

P. BITUMINOSA (*L., DC. Pr.* ii. 219), quacum monente Boiss. conjungendæ sunt *P. palæstina* (*Gouan. Ill.* 51) et *P. plumosa* (*Rehb. Fl. Exens.* 869).

Mar. sept.—Tetuan (*Webb Herb.*, sub *P. palæstina*)!

Mar. merid.—Frequens ad radices Atlantis Majoris. Mesfioua! Tasseremout! Ourika! Mzouda! Seksaoua! In convalle Aït Mesan!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia).

Specimina nostra inter formam Europæam vulgarem et *P. palæstinam* ambigua.

P. DENTATA (*DC. Pr.* ii. 221), = *P. americana* (*L. Sp. Pl.* 1075).

Mar. sept.—Vidi specimina prope Tanger lecta, sed notulam perdididi.

— —, var. CALYCIUS VILLOSIS, = *P. polystachya* (*Poir. Dict. Suppl.* iv. 587) ex *DC. l. c.*

Mar. occid.—Prope Alcazar (*DC. loc. cit.*).

Ar. Geog. Species originis dubiæ, verisimiliter non Americana (nomen igitur Linnæanum delendum); in cultis Hispaniæ australis, Mauritaniæ Tingitanæ et Madeiræ rarius obvia. Olim legi in cultis prope Coronil Hispaniæ Gaditanæ.

Psoralea bracteata (*L., DC. Pr.* ii. 218), ex agro Mogadorensi a Broussonnet olim allata, est species Capensis, proculdubio cultura introducta, hodie non invenienda.

COLUTEA (*L. Gen.* 880).

C. ARBORESCENS (*L., DC. Pr.* ii. 270).

Mar. sept.—Tetuan (*Webb Herb.*)!

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—In regione inferiore Atlantis Majoris. Aït Mesan circ. 1250 m.!

Ar. Geog. Europa media calidior. Regio Mediterranea omnis (exclusa Macaronesia).

ASTRAGALUS (*L. Gen.* 892).

Ordinem specierum a eel. Bunge in opere classico "*Astragali Species Gerontogæ*" stabilitam secutus sum.

A. EPIGLOTTIS (*L., DC. Pr.* ii. 290).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Schsb. Herb.*), *Coss. spec.*!
Tetuan (*Webb Herb.*)!

Mar. merid.—Ad radices Atlantis Majoris. Tasseremout! Amsmiz!
Sektana!

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia).

A. SESAMEUS (*L., DC. Pr.* ii. 288), = *A. malacensis* (*Salzm. Exsicc.*)

Mar. merid.—In convalle Aït Mesan circ. 1200 m.!

Ar. Geog. Regio Mediterranea (exclusa Macaronesia).

A. PENTAGLOTTIS (*L., DC. Pr.* ii. 287).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Schsb. Herb.*), *Coss. spec.*!
Tetuan!

Mar. merid.—Tasseremout (*G. Maw*), spec. benev. communicatum!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

A. HAMOSUS (*L., DC. Pr.* ii. 290).

Mar. sept.—Tanger! (*Webb Herb.*)!

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Ad fines prov. Ilaha et Shedma (*G. Maw spec.*)!

Ar. Geog. Regio Mediterranea omnia (inclusa Macaronesia).

A. SOLANDRI (*Lowe, in Hook. Journ. Bot.* viii. 294, et *Man. Fl. Mad.* i. 188; *Bge. Astrag.* ii. 16), = *A. canescens* (*Solander, in Herb. Banks.*; *Buch. Cat.*; non *DC.*).

Mar. occid.—Sat frequens in arenosis prope Mogador!

Ar. Geog. Madeira. Plaga Mogadorensis.

Speciminibus ex insula Madeira a Lowe et Masson lectis bene convenit.

Specimina ab Ehrenberg prope Rosetta Ægypti lecta (a eel. Bunge "*A. trimestris*, L." notata) variant a nostris stipulis angustioribus, calycis dentibus brevioribus, legumine minus piloso et seminibus numerosioribus. Nostra planta, *A. hamoso* valde proxima, gaudet dentibus calycinis tubum subæquantibus, stipulis majoribus quam in stirpe Maderensi, superioribus sæpius lanceolatis, inferioribus late ovatis, acuminatis, interdum basi folio opposita coalitis. Legumen tamen ab illo *A. hamosi* semper diversum compressum et dorso late canaliculatum. Ipse eel. Bunge hanc speciem inter sectiones suas *Buceras* et *Harpi-lobus* mediam esse fatetur, et, pace tanti viri, plane nos docet *Har-*

pilobum sistere sectionem nimis artificialem, melius cum *Bucerate* conjugendam. An *A. Solandri* melius sicut subspecies *A. trimestris* collocandus sit experientia docebit. Mirum est ell. Broussonnet et Schousboe hanc plantam in arena Mogadorensi non invenisse. An serius, casu aliquo—opere hominum vel avium—ex insula Madeira hue allata?

ASTRAGALUS REINII, *nobis*, nov. sp.

Planta caulescens, ramosa, habitu ad § *Onobrychium* accedens; pili simplices basi fixi; folia imparipinnata, 12–15-juga, pilis albis obsita, juniora serieco-villosa; foliola oblongo-elliptica obtusissima; stipulæ liberæ, cuneato-acuminatæ; pedunculi pauciflori striati rigidi foliis subduplo longiores; calyx subcylindricus, demum ruptus, dentibus lineari-acuminatis tubo dimidium brevioribus instructus; flores; legumen breviter pilosum, compressum, arcuatum, dorso anguste canaliculatum, biloculare, sutura dorsali carinatum, ex stylo persistente recurvo apice hamatum.

Mar. merid.—In regione alpina Atlantis Majoris. Ex jugo Tagherot specimen unicum attulerunt ell. Rein et Fritsch hodie in herb. Kew. conservatum!

In sectione *Harpilobo*, Bge., collocandus, ab affinibus differt habitu et legumine valde compresso, dorso anguste canaliculato, parum arcuato, seminibus pro grege paucis, 8–10.

A. BÆTICUS (*L., DC. Pr.* ii. 191).

Mar. sept.—Tanger (*Webb Herb.*)! Djebel Dersa pr. Tetuan (*Webb Herb.*)!

Mar. occid.—Casa Blanca!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia).

A. LUSITANICUS (*Lam. Dict.* i. 312; *Boiss. Fl. Or.* ii. 267), = *Phaca Bætica* (*L., DC. Pr.* ii. 273), = *Erophaca Bætica* (*Boiss. Voy. Esp.*).

Mar. sept.—Inter Tetuan et Ceuta! Tetuan (*Webb Herb.*)!

Mar. occid.—Casa Blanca! In monte Djebel Hadid! In eodem monte legerunt R. et Fr. (*spec. in Herb. Kew.*);

Ar. Geog. Regio Mediterranea præsertim calidior, sed non ubique; deest in Italia, Gallia, Hispania boreali et in Macaronesia.

Species ab affinibus valde diversa, a cel. Bunge in opere classico prætermissa, a cel. Boissier in *Fl. Orientali*, ii. 210, in sectione propria (*Erophaca*) collocata. Specimina omnia quæ ipse legi, et quæ vidi ex monte Djebel Hadid flore et fructu destituta sunt.

A. LANIGERUS (*Desf. Fl. Atl.* ii. 181, tab. 202; *DC. Pr.* ii. 302), var. GLABRESCENS, *nob.*

Mar. occid.—Prope Agadir (*Coll. Indig.* 1871)! Spec. fructifera tantum visa.

Mar. merid.—In saxosis aridissimis prope Aïn Oumast, prov. Shedma'

Ar. Geog. *A. lanigerus* est species Africæ borealis propria. Fertur in Oriente sed verisimiliter errore antiquorum. Deest in Boissier, Fl. Orientali.

Planta non est acaulis. In var. nostra ramos vel caudiculos plurimos breves e collo radiceis emittit; pedunculi dum pluriflori foliorum 6–8-pollicarium longitudinem dimidium superant; stipulæ villosissimæ; foliola 25–35, breviter petiolulata, viridia, subglaberrima; racemus fructifer clongatus; flores brevissime pedicellati; calyx membranaceus demum ruptus, dentibus tubo brevioribus; legumen in statu juniore glaberrimum, maturum parce et breviter pilosum, insigniter rostratum, magis quam in spec. Algeriensibus inflatum. Tam in nostris quam in pluribus spec. Algeriensibus a me scrutatis calycis dentes lanceolati tubi $\frac{2}{3}$ longitudinem attingentes, nec breves cuneati tubi $\frac{1}{4}$ partem vix æquantes, ut in icone Fontanesiana supra citata.

Var. nostra cum var. *subglabrato* (DC. Pr. l. c.), ab icone in Buxbaum, Cent. iii. 38, fig. 2, desumpta, non confundenda. Hæc est *A. tragacanthoides* (Willd. Sp. Pl. iii. 1323), planta a cel. Willdenow tantum ex icone cognita, et a stirpe Fontanesiana certe diversa. De hac siluerunt celeb. Bunge et Boissier, et verisimiliter lapsu calami adjecit ill. DC. literas “(v. s.)”

ASTRAGALUS GLAUX (L., DC. Pr. ii. 288).

Mar. sept.—Tanger (*Schsb. Herb.*), *Coss. spec.!* (*Salzm. Exsicc.*)!

Mar. merid.—Prov. Shedma pluribus locis!

Ar. Geog. Hispania. Africa boreali-occidentalis. In Gallia australi olim inventa, a recentioribus non visa.

— —, var. *ROSTRATA*, *nob.* Legumine in rostrum conicum sensim attenuato, rostro apice recurvo.

Mar. merid.—In convalle Aït Mesan circ. 1300 m., et iterum supra Arround alt. 2000 m.!

Specimen florigerum nanum prope Mogador lectum forsitan ad hanc speciem referendum.

A. ATLANTICUS (*Ball in Journ. Bot.* 1873, 306).

A proximo *A. narbonensi* (Gouan, Ill. 49; DC. Pr. ii. 294), et ab *A. africano* (Bgc. Astr. No. 415), differt præsertim structura florum. In sp. nostra carina angustior, sensim, nec abrupte, curvata, alas superans et vexillo subæquilonga, alæ basi profundius auriculatæ. Calycis tubus sesquilingior, molliter villosus, dentibus minus inæqualibus, tubo ipso paulo brevioribus instructus. Bracteolæ duplo longiores quam in *A. narbonensis* spec. Hispanicis. Foliorum rachis et caulis plus minusve patenti-villosa, foliola viridia subglaberrima. Stipulæ lineari-subulatæ villosæ, longiores et angustiores quam in *A. narbonensi*.

Mar. merid.—Legimus florentem in convalle Aït Mesan alt. circ. 1200 m.!

Legumen nondum visum melius docebit affinitatem hujus stirpis speciosæ. Ut mihi videtur simul ac affini *A. africano*, Bge., pro subspecie sub typo *A. narbonensis* habenda est.

Ar. Geog. *A. narbonensis* in Hispania et Gallia australi obvia; *A. africanus* in Algeria; subspecies nostra in Atlante Majore.

ASTRAGALUS OCHROLEUCUS (*Coss. in Ind. Pl. Maroc.*, sine descriptione).

De hac planta a cel. Cosson mox describenda amplius hic referre nolo.

Mar. merid.—Frustulum fructiferum ex regione superiore Atlantis Majoris attulerunt ell. Rein et Fritsch!

A. CHLOROCYANEUS (*Boiss. et Reut. Pug.* 39; *Walp. Ann.* ii. 395).

Mar. sept.—Tanger (*Schsb. Herb.*), *Coss. spec.*! Tetuan!

Ar. Geog. Hispania. Africa boreali-occidentalis.

Ab hoc distinguere nequeo *A. Cossoni* (Bunge, *Astrag.* ii. 192). Ambo sunt *A. monspessulano*, L., valde proximi, et sicut subspecies (forsan sicut varietates?) enumerandi. Alia forma proxime affinis est *A. monspessulanus*, Desf. non L., cui in scheda (in herb. Kew. conservata) cel. Bunge adjecit “nova sp. *A. monspessulanus* (sensu latiore) per regionem Mediterraneam occidentalem (exclusa Macaronesia) diffusa est, nec deest in calidioribus Europæ mediæ.

A. INCURVUS (*Desf. Fl. Atl.* ii. 182, tab. 203; *DC. Pr.* ii. 204).

Mar. merid.—Legi fructiferum in convalle Aït Mesan circ. 1200 m.!

Ar. Geog. Hispania. Africa borealis.

BISERRULA (*L. Gen.* 893).

B. PELECINUS (*L.*, *DC. Pr.* ii. 307).

Mar. occid.—Casa Blanca!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia).

GLYCYRHIZA (*L. Gen.* 882).

G. FÆTIDA (*Desf. Fl. Atl.* ii. 170, tab. 199; *DC. Pr.* ii. 248).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! El Araïsch (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Africa borealis.

HEDYSARUM (*L. Gen.* 887).

H. CAPITATUM (*Desf. Fl. Atl.* ii. 177; *DC. Pr.* ii. 341, non *Burm.*).

Mar. sept.—Tanger!

Ar. Geog. Regio Mediterranea calidior. Deest in Oriente et in Macaronesia.

H. FLEXUOSUM (*Desf. Fl. Atl.* ii. 176, an *L. Sp. Pl.* 1058? an *DC. Pr.* ii. 341?).

Mar. sept.—Tanger! (*Webb Herb.*)! In collibus inter Tanger et Fondak!

Ar. Geog. Africa boreali-occidentalis. Hispania.

HEDYSARUM MEMBRANACEUM (*Coss. in Bull. Soc. Bot. Fr.* xx. 246). Sequitur descriptio Cossoniana:—

“Planta plus minus pube adpressa sericea, *perennis* inferne *fruticosa*, l vel $1\frac{1}{2}$ m. alta, ramosa, laxe dumosa, ramis erectis sæpius confertis indurato-rigidis striato-angulatis. *Folia* petiolata, imparipinnata, 4-7-juga, foliolis minutis, supra glabrescentibus, infra pube adpressa subsericeis, oblongis, petiolulatis, margine arefactione sæpius subinvolutis, obtusis, mucronulatis vel subretusis. *Stipulæ* membranaceo-scariosæ, fusæ, cito evanidæ, *inter se basi et cum petiolo adnatæ*. *Racemi* axillares 3-4-flori, *laxi*, folio sæpius breviores, floribus, circiter 15 millim. longis, primum erectis dein patulis, pedicellis calycis tubo brevioribus sub calyce minute bibracteolatis bracteolis linearibus fuseis. *Calyx* pube adpressa subsericeus, tubo campanulato, *dentibus* e basi triangulari linearibus *tubo subtriplo brevioribus*, superioribus distantibus. *Corolla* glabra, pallide lilacino-purpurascens, marcescens sed fructu maturo decidua, *vexillo* oblongo-obovato *carina* *via brevior*, alis oblongis *carina* triente brevioribus, *carina latere inferiore sensim arcuata*. Stamen vexillare liberum, cætera in vaginam membranaceam superne fissam coalita in quarta parte superiore libera filiformia. *Legumen* 2-4 centim. longum, adpresse et minutissime puberulum, *stipitatum* stipite calyce longiore, bi-articulatum sæpe abortu ad articulum unicum reductum, inter articulos angustius, *articulis* 9-12 millim. latis *plano-compressis*, *disco* oblongo reticulatim nervoso *aculeis destituto*, ad semen tantum convexiusculo, *margine membranaceo latitudinem dimidiam disci excedente*. Semina subreniformia, estrophiolata.”

Mar. merid.—Juxta pagum Keira (750 m.) in alluviis et inter segetes legit floriferum et fructiferum Balansa mense Majo 1867. Eodem mense, anno 1871, nos vidimus hanc speciem insignem satis frequentem in glareosis torrentium et alibi ad radices Atlantis Majoris, sed semper alabastris florigeris nondum explicatis.

Monente cel. auctore:—“*II. membranaceum* inter omnes species generis insigne stipularum fabrica, racemis brevibus laxifloris, dentibus calycis brevibus, *carina sensim arcuata*, leguminibus complanatis latissime membranaceo-marginatis.”

ONOBRYCHIS (*Gærtn. Fr.* 318).

O. SATIVA (*Lam. Fl. Fr.* ii. 652; *DC. Pr.* ii. 344). = Hedysarum Onobrychis, L.

Mar. sept.—Tanger (*Webb Herb.*)! Specimen ineertum fructu carens.

— —, var. PSEUDOSUPINA. Caulibus basi proeumbentibus adscen-

dentibus, dentibus calycinis tubo ter longioribus, vexillo carina vix longiore.

Mar. sept.—Prope Cap. Spartel!

Ar. Geog. *Onobrychis sativa* per Europam median et Asiam temperatam late diffusa, in reg. Mediterranea minus frequenter occurrit. Adsunt tamen formæ affines plurimæ sicut subspecies enumerandæ. In ditione Maroccana videtur admodum rara nec colitur in provinciis a nobis exploratis. Genus omne in Macaronesia deest.

ONOBRYCHIS ERIOPHORA (*Desv. Journ. Bot.* 1814; *DC. Pr.* ii. 345).

Mar. sept.—Tetuan!

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

O. CRISTA GALLI (*Lam. Fl. Fr.* ii. 652; *Boiss. Fl. Or.* ii. 528; non *DC. Pr.* ii. 346), = *Hedysarum Crista Galli* (*L. Syst.* 563; *Sibth. et Sm. Fl. Gr.* tab. 724), = *O. trilophocarpa* (*Coss. et Dur. in Balansa, Pl. d'Algér.* 1852, no. 351).

Mar. occid.—In prov. Haha haud procul Mogador!

Mar. merid.—Juxta Aïn Oumast, prov. Shedma!

Ar. Geog. Species imprimis Orientalis, per Africam borealem usque littora Oceani diffusa, sed verisimiliter infrequens. Fertur in Italia et in Gallia australi, sed vix spontanea, aut non genuina.

De synonymia hujus speciei confer Boissier (*Diagn. Pl. Or.* ser. 1, ix. 108), et Cosson (*Bull. Soc. Bot. Fr.* iv. 17).

EBENUS (*L. Gen.* 895).

E. PINNATA (*Desf. Act. Soc. Hist. Nat. Par.* i. 21, tab. 3; *DC. Pr.* ii. 350).

Mar. occid.—Aïn el Hadjar! Djebel Hadid! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Sat frequens in regione inferiore. Aïn Oumast! Tasseremout! Ourika! Seksaua! Keira (*Balansa*), *Coss. Index*.

Ar. Geog. Africa boreali-occidentalis.

Ludit plus minusve villosa, pedunculis brevioribus vel longissimis.

VICIA (*L. Gen.* 873).

V. ONOBRYCHOIDES (*L.*, *DC. Pr.* ii. 358).

Mar. merid.—In convalle Aït Mesan circ. 2200 m.!

Ar. Geog. Per Europam australem late diffusa, in convallibus meridionalibus Alpium ultra 1500 m. ascendens, et in calidioribus interdum obvia. Africa boreali-occidentalis.

—— —, var.? Stipulis linearibus minimis vix 2 millimetra longa, doribus minoribus a *V. onobrychoide* diversa videtur. Desunt legumina.

Mar. merid.—Legimus in ditione Reraya circ. 1100 m.

Planta ulterius inquirenda.

Vicia elegans (Guss. *Prod. Fl. Sic.* ii. 438)?

Mar. merid.—Legi spec. unicum in ditone Mesfoua!

V. elegans, Guss., species mihi non satis nota, ex descriptione et spec. (an authentica?) quæ vidi in grege *V. onobrychoidis* collocanda videtur, et spec. nostro sat bene convenit. In pl. nostra foliola mucronata angustiora quam in *V. onobrychoide*, stipulæ angustiores, cirri magis ramosi, flores æquilongæ sed tenuiores, calycis dentes longiores. Styli barbati nee pubescentes stirpem nostram e grege *V. Craccæ* sejungunt. Cel. Boissier in *Fl. Orientali V. elegantem* ad var. *V. variæ* reduxit, et sp. Gussonianam in grege *V. Craccæ* colloavit.

V. ALTISSIMA (Desf. *Fl. Atl.* ii. 163).

Mar. sept.—Tanger (*Webb Herb.*)!

Ar. Geog. Italia australis eum insulis. Dalmatia. Africa borealis.

V. BIFLORA (Desf. *Fl. Atl.* ii. 166, tab. 197; *DC. Pr.* ii. 360).

Mar. sept.—Inter Tetuan et Centa!

Ar. Geog. Africa borealis.

Speciminibus paucifloris *V. pseudocraccæ*, Bert., similis; sed præter alias notas *V. biflora* gaudet stylo apice circumcirca hirsuto, dum in *V. pseudocracca* stylus ad latus inferius apice barbatus.

Planta nostra a typo Fontanesiano paululum variat. Folia glabrescentia, parce pilosa, apice longiuscule cirrata, 3–6-juga; foliola sæpius 10, sæpe alterna, oblongo-elliptica, mucronulata; stipulæ semisagittatæ lobo unilaterali limbum subæquante; pedunculi sæpissime biflori, aristati, foliis fere dimidio breviores; calycis breviter pilosi dentes inæquales, inferiores subulati tubo æquilongi, superiores breves triangulares; vexillum earina paullo longius. Flores in pedunculo rarius 3 vel 1.

V. biflora (Webb, *Phyt. Can.* tab. 65, B) est species aliena = *V. albicans*, Lowe.

V. GLAUCA (Presl, *Del. Prag.* i. 37; *Moris, Fl. Sard.* i. 565, tab. 60; *Walp. Rep.* i. 713), var. *RERAYENSIS* (Ball in *Journ. Bot.* 1873, 339).

Primo intuitu a typo parum differt floribus minoribus pallide lilaeinis, earina apice atropurpurea. Notas differentiales graviores video in calyce valde inæquali, dentibus inferioribus subulatis tubo longioribus, superioribus vix tubum dimidium æquantibus.

Mar. merid.—In regione superiore Atlantis Majoris. In convalle Aït Mesau a 2400 m. ad 2600 m.! In jugo Tagherot specimen legerunt ell. R. et Fr. hodie in herb. Kew conservatum!

Legumen immaturum pilosiusculum. In spec. ex Sicilia et Algeria video suturam leguminis pilosam.

Legumina in spec. a Tagherot quam in typo paululum latiora. Semina 4–6 ferunt.

Ar. Geog. *V. glauca* est species Africæ borealis, in Europa nonnisi in insulis Siciliæ et Sardinia detecta.

VICIA ATROPURPUREA (*Desf. Fl. Atl.* ii. 164; *DC. Pr.* ii. 359).

Mar. sept.—Tanger! (*Webb Herb.*)! (*Salzm. Exsicc.*)! Cap. Spartel!

Ar. Geog. Regio Mediterranea occidentalis (inclusa Macaronesia); deest in Græcia.

V. DURANDI (*Boiss. Diagn. Pl. Or.* ser. 1, ix. 116; *Walp. Ann.* ii. 398).

Mar. sept.?—"Specimina in Mauritania ad Tingidem legit Durand in herb. Pavon. hodie Boissier."

Dubia eivis. Cel. Cosson, qui herbarium proprium Durand sedule compulsi, nec in illo nec in aliis herbariis antiquioribus specimina Maroccana vidit.

? V. LENS (*Coss. et Germ. Fl. Par.*), = *Ervum* Lens (*L., DC. Pr.* ii. 366).

Colitur prope Tanger et in provinciis meridionalibus, sed nec indigena nec (quod sciam) facta spontanea.

V. ERVIFORMIS (*Boiss. Voy. Esp.* 550), = *Ervum* vicoides (*Desf. Fl. Atl.* ii. 168, tab. 198; *DC. Pr.* ii. 366).

Mar. sept.—Tetuan ad ostium fluminis (*Webb Herb.*)! Ibidem in calcareis juxta urbem!

Ar. Geog. Africa boreali-occidentalis. Hispania Bætica.

V. PUBESCENS (*DC. Cat. Hort. Monsp.* 109, et *Prod.* ii. 367, sub *Ervo*; *Boiss. Fl. Or.* ii. 596), = *Ervum* Loiseleurii (*M. B. Fl. Taur. Cauc. Suppl.* 475).

Mar. sept.—Tetuan (*Webb Herb.*)!

Præter indumentum differt a *V. tetrasperma*, cui valde proxima, calycis laciniis duplo longioribus lanceolato-subulatis. Pro subspecie habenda?

Ar. Geog. Hæc est forma meridionalis per reg. Mediterraneam omnem (inclusa Macaronesia) late diffusa.

V. SATIVA (*L., DC. Pr.* ii. 360).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! In lapidosis laud procul Cap. Spartel! Tetuan! In monte Beni Hosmar!

— — —, var. ANGUSTIFOLIA (*Seringe, in DC. Pr. l. c.*), = *V. angustifolia* (*Roth, Tent. Fl. Germ.* i. 310).

Mar. merid.—In regione inferiore et subalpina Atlantis Majoris. Aït Mesan alt. 1400 m., et iterum supra Arround eire. 2300 m.!

— — —, var. ABYSSINICA (*Bak. in Fl. Trop. Afr.* ii. 172), = *V. abyssinica* (*Alefeld in Bonplandia*).

Mar. merid.—Aït Mesan, alt. 1300 m.!

Speciminibus Schimperianis in herb. Kew. omnino convenit nisi legumine (unico) in nostra planta subtoruloso. Folia, fere ut in *V. lathyroidi*, L., 2-4-juga; stipulæ superiores semisagittatæ dente subulato

auetæ, inferiores profunde laciniato-dentatæ, dentibus acutis subulatis; semina fusca, lævia.

Legumina in spec. Schimperianis supra citatis, a planta disereta et chartæ agglutinata, mihi videntur incerta et forsan ex sp. aliena orta.

— — —, var. *CORDATA* (*Boiss. Fl. Or.* ii. 575), = *V. cordata* (*auctorum, an Wulf.?*)

Mar. occid.—Agadir (*Coll. Indig.* 1871)! Casa Blanca (*J. D. H.*)!

Ar. Geog. *V. sativa*, verisimiliter in regione Mediterranea indigena, est species polymorpha per orbem temperatum veteribus notum late diffusa, et in Americam cum seminibus introducta. Formæ plurimæ ab auctoribus pro speciebus propositæ parum stabiles et varietatibus intermediis connexæ sunt.

VICIA LUTEA (*L., DC. Pr.* ii. 363).

Mar. sept.—Tanger! (*Webb Herb.*)! (*Salzm. Exsicc.*)!

Mar. occid.—Casa Blanca (*J. D. H.*)!

— — —, var. *HIRTA* (*Boiss. Fl. Or.* ii. 570), = *V. hirta* (*Balb. Miscell. Alt. Pers. Syn.* ii. 308).

Mar. sept.—Inter Tanger et Tetuan! Tetuan (*Webb Herb.*)!

— — —, var. γ , *MURICATA* (*Seringe, in DC. Pr. l. c.*), = *Vicia vestita* (*Boiss. Elench.* 67, et *Voy. Esp.* tab. 57; *Walp. Rep.* i. 717).

Mar. sept.—Tanger (*Webb Herb.*)!

Specimen a me prope Algeiras lectum Reuterianis ex eodem fere loco provenientibus omnino conforme est. Floribus violaceis et legumine dense vestito var. hæc a typo recedit, sed in stirpe tam ludibunda hæ notæ non gravis momenti. Nomen *muricata* a Seringe allatum haud omnino aptum, sed prioritatis jure gaudet.

— — —, var. *NITIDA*, *nobis*.

Flores magni pallide purpurei, legumen, in statu juniore pilis nonnullis basi non tubereulatis adpersum, maturum omnino glabrum. In frustulo a eel. H. fil. prope Sektana lecto, verisimiliter ad plantam nostram referendo, flores ochroleuci fere ut in typo. Stipulæ in spec. ex Agadir minimæ, semina fere nigra, ellipsoidea, compressa.

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Sektana!

Var. nostra pluribus notis ad *V. lævigatam* (*Sm. E. Bot.* tab. 483) accedit. Icon, verisimiliter quoad stipulas falsa, cum nulla planta recentioribus cognita convenit; ex lusu quodam *V. luteæ* in plaga marina prope urbem Weymouth ex regione Mediterranea olim allato, verisimiliter desumpta est.

Ar. Geog. *V. lutea* est species polymorpha per Europam mediam et regionem Mediterraneam omnem (inclusa Macaronesia) late diffusa.

LATHYRUS (*L. Gen.* 872).

L. LATIFOLIUS (*L., DC. Pr.* ii. 370),

Mar. sept.—Tetuan (*Webb Herb.*) !

Ar. Geog. Europa media et austro-occidentalis.

L. APHACA (*L., DC. Pr.* ii. 372).

Mar. sept.—Tanger !

Mar. merid.—In regione inferiore Atlantis Majoris. Aït Mesan 1300 m. !

Ar. Geog. Europa media. Regio Mediterranea omnis (inclusa Macaronesia).

L. SPHÆRICUS (*Retz. Obs.* iii. 39 ; *DC. Pr.* ii. 372), = *L. angulatus* (*L. Herb. et Mant. Sec.* 443, *non Spec. Pl.* 1031).

Mar. merid.—In convalle Aït Mesan 1200–1400 m. !

In speciminibus minoribus cirri ad aristam brevem reducti.

Ar. Geog. Regio Mediterranea omnis, in cultis Europæ mediæ hinc inde obvia.

L. CÆRULESCENS (*Boiss. et Reut. Pug.* 43 ; *Walp. Ann.* iv. 530).

Mar. sept.—Tanger versus montem Djebel Kebir (*Boiss. et Reut. l. c.*).

Specimina non vidi ; notæ diagnosticæ quibus a *L. setifolio* distinguitur mihi videntur nimis leves. Pro subspecie enumerandus ?

L. SATIVUS (*L., DC. Pr.* ii. 373), var. *AMPHICARPOS*, *Coss.*

Mar. sept.—Tanger ! (*Schb. Herb.*), *Coss. spec.* !

Hic est forsitan *L. amphicarpos* (*L. Sp. Pl.* 1029 ; *DC. Pr.* ii. 373). *L. amphicarpos* (*Sibth. et Sm. Fl. Gr. tab.* 693) est tamen species diversa (*L. blepharicarpos*, *Boiss.*), herba hirta, leguminibus ciliatis, et seminibus tuberculatis, facile dignoscenda. Species Linnæana inter hanc et nostram ambigua est et nomen hodie prætermittendum.

Ar. Geog. *L. sativus* per regionem omnem Mediterraneam diffusus est, et in Abyssinia nascitur—an indigenus ?

L. FISSUS (*Ball in Journ. Bot.* 1873, 332).

Annuus, nanus, caule debili ramoso, 4-angulato ; folia 2–3-juga, cirro simplici munita ; stipulæ foliolis dimidio breviores, semisagittatæ ; pedunculi uniflori, aristati, foliis æquilongi ; calycis dentes subulati, subæquales, tubo 4–5 longiores ; flores parvi (jam emarceidi videntur rubelli) ; legumen dispermum, breviter stipitatum, ovatum acutum, apice obliquo subrostratum.

Mar. merid.—Ad radices Atlantis Majoris—in rupibus apricis supra Seksaoua detexit J. D. H. !

Huic similis sed non affinis est *L. saxatilis* (*Vent. et DC. Pr.* ii. 380, sub *Orobo* ; *Vis. Fl. Dalm.* iii. 330), = *L. ciliatis*, *Guss.* In hoc calyx et legumen omnino diversa. Affinitas vera, nisi fallor, cum *L. sativo*.

L. CICERA (*L., DC. Pr.* ii. 373).

Mar. occid.—Agadir (*Coll. Indig.* 1871) !

Mar. merid.—Misra Ben Kara!

Specimina nostra sistunt formam majorem leguminibus et foliolis longioribus.

Ar. Geog. Europa media in cultis, vix indigenus. Regio omnis Mediterranea (inclusa Macaronesia).

LATHYRUS ANNUUS (L., *DC. Pr.* ii. 373).

Mar. sept.—Tanger! Tetuan (*Webb Herb.*)!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

L. TINGITANUS (L., *DC. Pr.* ii. 374).

Mar. sept.—Tanger! Tanger et Tetuan (*Webb Herb.*)!

Ar. Geog. Regio Mediterranea occidentalis. Sardinia. Hispania Bætica. Africa boreali-occidentalis. Macaronesia.

L. CLYMENUM (L., *DC. Pr.* ii. 375), = *L. purpureus* (*Desf. Ann. Mus.* xii. 56), forma fol. latioribus, = *L. tenuifolius* (*Desf. Fl. Atl.* ii. 160).

Mar. sept.—Tanger (*Webb Herb.*)! (*Salzm. Exsicc.*)! In monte Djebel Kebir! Prope Cap. Spartel (formæ latifolia et angustifolia)! Tetuan! (*Webb Herb.*)! Inter Tetuan et Ceuta!

Ar. Geog. Per regionem Mediterraneam fere omnem (inclusa Madera), imprimis occidentalem, late diffusus.

L. articulatus (L., *DC. Pr.* ii. 375).

Mar. sept.—Tanger! (*Webb Herb.*)!

Mar. merid.—Distr. Mesfioua haud procul urbe Marocco!

Ar. Geog. Regio Mediterranea occidentalis fere ubiquæ.

Notæ diagnosticæ quibus hic a *L. Clymeno* distinguitur mihi videntur nimis instabiles. Confer Gren. et Godr. *Fl. Fr.* i. 479, et Lowe, *Man. Fl. Mad.* i. 216. In *L. articulato* foliola sæpius latiora ex mucrone apiculari prominente videntur cuspidata, sed in forma angustifolia (rarius obvia) foliola omnino ut in *L. Clymeno*. Differentiam inter stylos a Gr. et Godr. indicatam ego frustra quæsivi. Semina *L. Clymeni* a variis auctoribus “ovata truncata,” “ovoidea truncata,” “compressa,” “subgloboso-quadrangula parum compressa” dicuntur, et ea *L. articulati* “multo majora,” “magis compressa,” “velutina vel subvelutina.” Re vera forma et magnitudo seminum variant, et semina *L. articulati* oculo fortiter armato non vere velutina. In speciminibus multis a me scrutatis video semina semper magis compressa. Quoad hili longitudinem hanc video variabilem, a quinta ad octavam partem peripheriæ seminis. Legumina *L. articulati* majora subtorulosa, nervis suturæ dorsali parallelis plus minusve prominentibus, inde interdum canaliculata, interdum plano-carinata. Lieet dubitative, pro subspecie proposui.

L. OCHRUS (L., sub *Piso*; *DC. Fl. Fr.* iv. 578, et *Prod.* ii. 375).

Mar. occid.—Casa Blanca!

Ar. Geog. Regio Mediterranea fere omnis (inclusa Macaronesia).

PISUM (*L. Gen.* 870).

P. ELATIUS (*M. Bieb. Fl. Taur.-Cauc.* ii. 151).

Mar. sept.—Tanger (*Webb Herb.*)! An introductum?

Ar. Geog. Per regionem Mediterraneam imprimis orientalem late diffusum. In Gallia occidentali prope Nantes obvium, sed verisimiliter introductum. Deest in Macaronesia.

CERATONIA (*L. Gen.* 1167).

C. SILIQUA (*L., DC. Pr.* ii. 486).

Mar. sept.—Tanger!

Mar. oecid.—Mogador (*Lowe Cat.*), verisimiliter eulta.

Mar. merid.—Sat frequens in regione inferiore et in convallibus interioribus Atlantis Majoris. In convalle Ourika! Aït Mesan usque 1500 m.!

Ar. Geog. In regione Mediterranea ealidiore. In Macaronesia non spontanea.

Alibi semper in maritimis indicatur; in Imp. Maroccano meridionali loca a mare remota non fugit, et fit arbor grandis diametro 3-6-pedali.

Circa Tanger colitur in hortis et fit arbor elata, 40-pedalis et ultra, *Algarobia glandulosa* (Torr. in *Ann. Lye. N. York*, ii. tab. 2; Walp. *Rep.* i. 861, sub *Prosopide*). A cel. auctore dicitur frutex vel arbuseula.

ACACIA (*Willd. Spec. Pl.* 1049).

A. GUMMIFERA (*Willd. Sp. Pl.* iv. 1056; *DC. Pr.* ii. 455), = A. coronillæfolia (*Desf. Cat. Hort. Par.* 207; *DC. loc. cit.*).

Mar. oecid.—Prope Mogador (*Brouss.*), *Willd.*

Mar. merid.—Vidimus plerisque locis in regione inferiore, non tamen frequentissime. In prov. Haha et Shedma sat frequens! Legumina matura cel. Cosson habuit et mihi benevole dedit.

Floret verisimiliter mense Augusto, et fructus hieme matureseunt. Mensibus Aprili et Majo vidimus plantam semper flore et fructu earentem. Hæc species, vel alia proxime affinis, colitur in hortis Tingitanis et fit arbor humilis, vix 20-pedalis, sed pl. spontaneam semper vidimus arbuseulam, 6-8-pedalem, truneo diametro vix 4-pollicari. *Acaciæ gummiferæ* Africæ borealis inter se valde affines studio accurato earent. Confer Schweinf. in *Linn.* 1867.

A. FARNESIANA (*L.*, sub *Mimosa*; *Willd. Sp. Pl.* iv. 1083; *DC. Pr.* ii. 461), = *Vachellia Farnesiana* (*W. et Arn. Fl. Pen. Ind.*).

Mar. merid.—In hortos urbis Marocco introducta et hodie facta subspontanea!

Ar. Geog. Patria incerta: per regiones tropicales et subtropicales late dispersa, sed ubique, ut videtur, cultura introducta.

Albizzia Julibrissin (Willd. et DC. Pr. ii. 469, sub *Acacia*) circa Tanger colitur sed nullibi spontanea. Species est orientalis, in Persia, nec ultra versus occidentem, indigena.

ROSACEÆ (*Juss. Gen.* 334).

PRUNUS (*L. Gen.* 620).

P. DOMESTICA (*L., DC. Pr.* ii. 533).

Mar. merid.—Fructu nondum maturo legimus ineunte Majo in irrigatis juxta Sheshaoua, verisimiliter ab indigenis cultam! Ejusdem speciei formam sylvestrem (?) legimus sine fructu in convalle Ourika circ. 1100 m.!

Ar. Geog. Species orientalis ab antiquo culta et in calidioribus reg. Mediterraneæ interdum facta spontanea.

P. PROSTRATA (*Labill. Dec. Pl. Syr.* iv. tab. 6; *DC. Pr.* ii. 538, sub *Ceraso*), = *P. humilis* (*Colla, Herb. Ped.* ii. 293).

Mar. merid.—In regione alpina et subalpina Atlantis Majoris. In monte Djebel Tezah a 2200 m. usque 3000 m. In convalle Aït Mesan et in jugo Tagherot a 2100 m. usque 2900 m.!

Ar. Geog. In montibus excelsis regionis Mediterraneæ calidioris. Decet in Italia continentali, in Pyrenæis et in Macaronesia.

Cel. Boissier distinxit sub “var. *concolor*” formam hujus sp. foliis subtus glaberrimis, utrinque fere concoloribus. In spec. nostris folia ludunt concoloria, vel subtus pallida et in statu juniore tomentosula, vel subtus albo-tomentosa fere ut in *Cotoneastro*.

Folia variant interdum toto circuitu serrulata, interdum basin versus integerrima apice inciso-serrata. Inde *Cerasus incisa* (Boiss. Fl. Or. ii. 648) est species valde dubia, vix notis stabilibus dignoscenda. Valde affines sunt etiam *C. diffusa* (Boiss. et Hausskn. in Fl. Or.) et *C. tortuosa* (Boiss. et Hausskn. eodem loco); sed *Microcerasi* orientales omnes studio ulteriore et observatione in vivo egent.

RUBUS (*L. Gen.* 652).

R. FRUTICOSUS (*L., DC. Pr.* ii. 560), var. *DISCOLOR*, = *R. discolor* (*Weihe et Nees, Rub. Germ.* 46, tab. 20).

Mar. sept.—Tanger! (*Webb Herb.*)!

Mar. merid.—Sat frequens ad radices Atlantis Majoris. In convalle Ourika! (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Hæc est forma *R. fruticosi* regionis Mediterraneæ totius solennis.

R. DEBILIS (*Ball in Journ. Bot.* 1873, 332). In § *Dalibarda* collocandus? (Plate XIX.)

Prostratus, ramosissimus, ramis debilibus; folia ovalia crenata, utrinque

viridia, integra vel basi lobata, vel ternata foliolis lateralibus petiolulatis lobo terminali multo minoribus; stipulæ filiformes; flores in spec. nostris emarceidi; achenia (infertilia) omnino sicca, apice acuta, stylo caduco.

Mar. merid.—Ad radices Atlantis Majoris. Legimus in convalle Ourika! circ. 1100 m.

Genus Linnæanum *Dalibarda*, a celeb. Benth. et Hook. ad sectionem *Rubi* reductum, in America boreali et meridionali (ad fretum Magellanicum) et Nepalia detectum, etiam Africæ borealis, ut videtur, indigenum est; sed specimina nostra manca observatione ulteriore in loco natali egent.

POTENTILLA (*L. Gen.* 634).

P. reptans (*L., DC. Pr.* ii. 574).

Mar. merid.—In umbrosis irrigatis regionis inferioris et ad radices Atlantis Majoris. Sheshaoua! Amismiz! Reraya circ. 1200 m.! Keira!

Ar. Geog. Europa omnis (excl. arctica). Asia temperata. Regio omnis Mediterranea (inclusa Macaronesia). Abyssinia.

In spec. nostris folia brevius petiolata quam in Europæis.

AGRIMONIA (*L. Gen.* 607).

A. odorata (*Mill. Dict.* No. 3; *DC. Pr.* ii. 587).

Mar. sept.—Tanger! Tetuan (*J. D. H.*)!

Ar. Geog. Per Europam mediam et australem hinc inde obvia. Specimina Africana (exceptis Maroccanis) non vidi.

Characteres videntur satis stabiles; sed hæc certe proles *A. Eupatoriæ* et melius sicut subspecies enumeranda.

POTERIUM (*L. Gen.* 1069).

P. sanguisorba (*L., DC. Pr.* ii. 594; *Boiss. Fl. Or.* ii. 783), = *P. dictyocarpum* (*Spach, Ann. Sc. Nat.* ser. 3, v. 34; *Walp. Ann.* i. 283).

Mar. merid.—In convalle Aït Mesan alt. 1600 m.! (*R. et Fr. spec. in Herb. Kew.*)! Achenia margine valde prominente subalata.

Ar. Geog. Europa omnis (exclusa arctica). Asia temperata. Montes regionis Mediterraneæ (exclusa Macaronesia).

P. verrucosum (*Ehrenb. Cat. Pl. Hort. Berol.* 1829; *Spach, loc. cit.* 39; *Walp. Ann.* i. 284), = *P. mauritanicum* *α* (*Boiss. in Kel. Fl. Calp.* 219).

Mar. sept.—Tanger!

Mar. merid.—Tasseremont! Seksaoua!

— — — *β*, *MAGNOLII* = *P. Magnolii* (*Spach, l. c.* 38; *Walp. Ann.* i. 284), = *P. mauritanicum β* (*Boiss. in Kel. Fl. Calp. l. c.*).

Mar. sept.—Tanger!

Mar. occid.—Casa Blanca! forma intermedia, fere *P. alveolosum* (Spach, l. c. 38; Walp. l. c. 284). Aïn el Hadjar haud procul Mogador! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In convalle Aït Mesan!

Ar. Geog. Species verisimiliter per totam regionem Mediterraneam diffusa.

Hæc species, miro modo ludibunda, in multas s. d. species ab auctoribus recentioribus dilaniata, variat insigniter indumento, habitu, statura, quin etiam fructu tetragono majori vel minori, sphærico, vel ovali, vel obovato, angulis cristatis, vel tuberculatis, vel subobsoletis, faciebus verrucosis, tuberculis obtusis vel acutiusculis. Speciminibus permultis scrutatis formas intermedias tantas vidi ut vix varietates, ne dicam species diversas, sistere videntur. *Sanguisorba mauritanica* (Desf. Fl. Atl. ii. 142) est species incerta, teste Spach, ex *P. Fontanesii*, Spach, et *P. verrucoso*, Ehrh., composita; nomen igitur non conservandum.

POTERIUM MULTICAULE (*Boiss. et Reut. Pug.* 44; *Walp. Ann.* iv. 665).

Mar. sept.—In monte Djebel Kebir prope Tanger! (*Boiss. et Reut. l. c.*). Juxta Cap. Spartel! In monte Beni Hosmar prope Tetuan!

Habitu ad *P. ancistroides*, fructu ad *P. verrucosum* accedit, sed hinc usque formas intermedias non vidi, et, licet dubitative, pro specie probe distincta enumeravi.

Ar. Geog. Hispania Bætica. Ditio Tingitana.

P. ANCEPS (*Ball in Journ. Bot.* 1872, 332).

Perenne, multicaule, ramulis foliaceis; folia radicalia in pagina inferiore caulina sæpius utrinque sericeo-villosa; capitula parva globosa; achenia pulverulento-tomentosa, ancipitia, ad angulos laterales alata, ad anteriorem et posteriorem costis subobsoletis vix prominentibus prædita.

Mar. merid.—In regione subalpina Atlantis Majoris. In convalle Aït Mesan supra Arround alt. 2500 m.! In monte Djebel Tezah 2500–2600 m.!

Invito animo novam proposui speciem in hoc genere speciebus incertis nimis onusto; sed achenia in speciminibus nostris ab omnibus mihi cognitis prorsus diversa. Habitu ad *P. ancistroides* paululum accedit, sed, præter alias notas, ramulis foliaceis et foliis sericeo-villosis insigniter differt.

P. ANCISTROIDES (*Desf. Fl. Atl.* ii. 346, tab. 251; *DC. Pr.* ii.).

Mar. merid.—In convalle Amsmiz in rupibus apricis, alt. 1400 m.!

Ar. Geog. Africa boreali-occidentalis. Hispania

Spec. nostra Algeriensibus (prope Oran lectis) multo minora; sed achenia conformia, ex angulis subobsoletis fere fusiformia, faciebus insigniter reticulato-rugosa.

POTERIUM, sp., an P. DURIÆI (*Spach, loc. cit.* 33; *Walp. Ann.* i. 282)?

Mar. merid.—In rupibus aridis supra Milhain!

Specimen unicum legi Majo decedente fructu maturo jam amisso.

Folia elongata, foliolis paucis remotis petiolulatis profunde inciso-dentatis; caulis bipedalis, tetragonus, ramosus, ramis tenuibus parce foliatis. Capitula terminalia.

ROSA (*L. Gen.* 631).

R. SEMPERVIRENS (*L., DC. Pr.* ii. 597).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Tetuan!

Mar. merid.—Prope Tasseremout! Ourika (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio Mediterranea, imprimis occidentalis. Deest in Asia Mediterranea et in Macaronesia.

Sec. cel. Crepin, qui *Rosas* nostras adnotationibus pretiosis ornavit, spec. a Tasseremout sistunt formam cui botanici recentiores nomen *R. scandens* imposuerunt.

R. CANINA (*L., DC. Pr.* ii. 613).

Mar. merid.—Tasseremout! Verisimiliter huc referendum spec. prope Sektana a ell. Rein et Fritsch lectum in herb. Kew. conservatum!

Ar. Geog. Europa omnis (exclusa arctica). Asia temperata. Regio omnis Mediterranea (inclusa Macaronesia).

—— —, var. SUBINERMIS, *nob.*

Mar. merid.—In convalle Aït Mesan 1200–1500 m.! et forsan huc spec. incompleta in convalle Amsmiz lecta!

Rami floriferi sæpissime (non semper) subinermes. Legimus florentem et in eadem planta fructus maturos segmentis calycinis jam amissis.

De hac stirpe scripsit cel. Crepin:—"Probablement une forme du *canina*, mais si les sépales se relèvent sur le fruit, je la classerais parmi les *Montanæ* comme *R. Reuteri*, var., ce qui après tout n'est qu'une forme du *R. canina*. Celle-ci ressemble extrêmement à une espèce suisse, *R. exannata*."

R. MICRANTHA (*Sm. Eng. Bot.* tab. 2490; *Boiss. Fl. Or.* ii. 687), = *R. rubiginosa*, var. π , *micrantha* (*DC. Pr.* ii. 617).

Mar. sept.—Tanger (*Webb Herb.*, sub "*R. rubiginosa*")!

—— —, var. ATLANTICA, *nob.* Rami antiquiores steriles, aculeis adnatis basi latis conformibus armati, juniores subinermes, vel omnino inermes, glaberrimi; foliola minute duplicato-serrata, subtus parce pilosa glandulosa; rachis foliorum et pedunculi glandulis stipitatis crebris, et aciculis paucis obsiti.

Mar. merid.—Mesfioua! Amsmiz!

Ar. Geog. *R. micrantha* per Europam mediam occidentalem diffusa, in regione Mediterranea videtur perrara. Specimina nulla vidi.

ROSA SERAPHINI (*Viv. Fl. Cors. Sp. Nov.* 8; *DC. Pr.* ii. 625), = *R. sicula* (*Tratt. Ros.* ii. 68; *DC. Pr.* ii. 624), = *R. sieula*, var. *Serafini* (*Wallr.*).

Mar. merid.—In regione subalpina et alpina Atlantis Majoris haud infrequens. In convalle Aït Mesan, supra Arround, circ. 2500 m.! In monte Djebel Tezali 3000 m. et ultra!

Hæc species cum *R. micrantha* a cel. Grenier et Godron sicut varietates *R. graveolentis*, Gr. et Godr., enumerantur. *R. Seraphini* in Apennino Italiæ mediæ mihi sæpe obvia, statura nana, aculeis creberrimis et pedunculis glabris a *R. micrantha* videtur satis diversa, nec transitum observavi.

PYRUS (*L. Gen.* 626).

P. CYDONIA (*L., Benth. et II. f. Gen. Pl.* i. 626), = *Cydonia vulgaris* (*Pers. Syn.* ii. 40; *DC. Pr.* ii. 638).

Mar. sept.—Prope Tanger in fruticetis et dumetis!

Ar. Geog. Regio Mediterranea; in Oriente certe indigena, etiam in Europæ australis locis montosis, a cultura remotis, sæpe obvia, et verisimiliter indigena.

CRATÆGUS (*L. Gen.* 622).

C. OXYACANTHA (*L., DC. Pr.* ii. 628).

Mar. sept.—Prope Tanger in monte Djebel Kebir! El Araïsch (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Europa fere omnis (exclusa arctica). Asia temperata. Regio Mediterranea omnis (exclusa Macaronesia).

— —, var. *FOLIIS OBOVATO-CUNEATIS*.

Mar. sept.—Inter Tanger et Tetuan!

Stirps aspectu singulari. Folia circuitu obovata, basi cuneata, apice incisodentata, inferiora subtriloba; pedunculi glaberrimi; fructus biovulatus. In eodem corymbo fructus maturi, flores petalis expansis, et nondum expansi semel crescunt.

SAXIFRAGÆ (*Juss. Gen.* 235).

SAXIFRAGA (*L. Gen.* 559).

S. GLOBULIFERA (*Desf. Fl. Atl.* i. 342, tab. 96; *DC. Pr.* iv. 31), var. *GIBALTARICA* (*Seringe, in DC. Pr. l. c.*), = *S. gibraltarica* (*Boiss. et Reut. Pug.* 47).

Mar. merid.—In regione media et superiore Atlantis Majoris. In convalle Aït Mesan circ. 1500 m., et iterum supra Arround alt. 2400 m.! In convalle Amsmiz circa 1700 m.! In monte Djebel Tezali usque 3000 m.!

Ar. Geog. Africa boreali-occidentalis. Hispania australis.

Notæ differentiales quibus *S. gibraltarica*, B. et R., et *S. granatensis*, B. et R., dignoscuntur videntur omnino fallaces. Specimina Hispanica

ex locis diversis provenientia inter se plus minusve differre fatendum est; hæc tamen in varietates, ne dicam species diversas, separare non potui. Nota diagnostica ex gemmis elongatis acutis axillaribus substipitatis, vel ovatis obtusis terminalibus sessilibus, omnino incerta. Vidi gemmas utrasque in sureulis sterilibus unius ejusdemque plantæ. Planta Algeriensis sæpissime gaudet foliis majoribus glabrioribus longius petiolatis, et calycis segmentis angustioribus, sed non desunt specimina *S. granatensi*, B. et R., omnino conformia, gemmis sessilibus terminalibus. Botanicis quibus placet *S. oranensem*, Munby, *S. granatensem* et *S. gibraltariam* pro speciebus rite distinctas vindicare, *S. globulifera*, Dsf., plane evadit, et nec in herbariis nec in rerum natura ultro invenienda est.

Saxifraga Maweana (*Baker in Gard. Chron.* 1871, p. 1355).

Cæspitosa, multicaulis, basin versus vestigiis foliorum emareidorum vestita; sureuli florigeri erecti, superne subaphylli, basin versus gemmas axillares stipitatas gerentes; folia inferiora longiuscule petiolata, basi subcordata, triloba, lobo medio integro vel tridentato, lateralibus inciso-dentatis, caulina bracteiformia lineari-oblonga; flores breviter pedicellati, laxè paniculati, panicula sæpius pauciflora; calycis segmenta oblongo-ovata, obtusiuscula, tubo longiora; petala magna 8-9-linearia. Pedicelli et calyx basin versus parce glanduloso-pilosi, cæterum herba tota glaberrima. Styli erecti, non divergentes, longiusculi, petalis paulo breviores.

Mar. sept.—In monte Beni Hosmar legerunt P. B. Webb anno 1827, et G. Maw anno 1869; iterum a nobis lecta anno 1871! Vidi specimina in herb. Webb sub *S. globulifera*, Dsf.

Ex descriptione valde proxima *S. globulifera*, sed facie valde diversa. Floribus magnis speciosis *S. granulata* æmula, *S. maderensi*, Don, subsimilis. Hæc autem differt herba viscidula, calycis segmentis acutis, foliis magis divisis, caule foliaceo, et stylis divergentibus.

S. TRIDACTYLITES (*L., DC. Pr.* iv. 35).

Mar. sept.—Tetuan! In monte Beni Hosmar!

Mar. merid.—In convalle Aït Mesan supra Arround alt. 2400 m.!

Ar. Geog. Europa fere omnis (inclusa arctica). Regio omnis Mediterranea (exclusa Macaronesia).

S. GRANULATA (*L., DC. Pr.* iv. 35).

Mar. sept.—In vertice montis Beni Hosmar prope Tetuan!

Mar. merid.—In regione media Atlantis Majoris. In convalle Aït Mesan circ. 2300 m. In monte Djebel Tezah 2200–2400 m.!

Ar. Geog. Europa fere omnis. Africa borealis. Montes Asiæ mediæ.

S. granulata est species polymorpha in varietates bene definitas laudè faile distinguenda; sed a botanicis in species plurimas instabiles dilaniata. Specimina nostra Atlantica spec. in mont. Himalaya a celeb. J. D. Hooker lectis fere exacte conformia sunt.

- S. pauciculata*, Cav. (see. specimina Willkommiana), *S. bulbosa*, Höchst., *S. glaucescens*, Reut., quin etiam *S. græca*, Boiss. et Held., et *S. elegantula*, Boiss. MSS., omnes ad typum reducendæ videntur. *S. corsica*, Gren. et Godr., paululum magis differt, sed inter hanc et typum transitum præbent specimina Hispanica ex Sierra de Alcoy (*S. granulata*, var. *ramosissima*, Coss., = *S. Cossoniana*, Boiss. et Reut. Diagn. Pl. Or.).
- S. Arundana*, Boiss., cum var. *carpetana*, = *S. carpetana*, Boiss. et Reut., sistit speciem inter *S. granulata* et *S. bulbiferam* collocandam.

RIBES (*L. Gen.* 281).

- R. GROSSULARIA* (*L., Boiss. Fl. Or.* ii. 845), = *R. Uva crispa* (*L., DC. Pr.* iii. 478), var. *ATLANTICA*, nob. Ovario glaberrimo, calycis segmentis pilosis erectis, non reflexis.
- Mar. merid.—In regione subalpina et alpina Atlantis Majoris. In convalle Aït Mesan circ. 2000 m. et ultra! In vertice montis Djebel Tezali ultra 3200 m.!
- Planta Atlantica ad formam *spinosissimam* auct. accedit, sed ovario glabro et calyce erecto insigniter diversa est. Folia parva in pagina inferiore eximie nervosa, tomentosula, superne glabra. Specimina paulo infra summum verticem montis Djebel Tezali lecta omnino prostrata, ramis subpedalibus, foliis profundius divisis insignia, sed notis gravioribus non diversa. Fructus in convalle Aït Mesan glaberrimus, vel demum setis paucis glandulosis adpersus.
- Ar. Geog. R. grossularia*, per Europam mediam (exclusa aretica) diffusa, habitat etiam in montibus regionis Mediterraneæ. Deest in Macaronesia et in montibus Hispaniæ meridionalis. Forma arete affinis in Sibiria Altaica.

CRASSULACEÆ (*DC. Bull. Philom.* 1801, p. 1).

TILLEA (*L. Gen.* 177).

- T. MUSCOSA* (*L., DC. Pr.* iii. 381).
- Mar. sept.—Tanger (*Webb Herb.*)! (*Salzm. Exsicc.*)!
- Ar. Geog. Europa media.* Regio Mediterranea fere omnis (inclusa Macaronesia). In Asia Mediterranea nondum detecta, sed forma affinis (subspecies?) *T. trichopoda*, Fenzl, locum tenet.

COTYLEDON (*L. Gen.* 578).

- C. UMBILICUS* (*Sm. Eng. Bot.* tab. 325), = *Umbilicus pendulinus* (*DC. Pl. Grass. et Prod.* iii. 400).
- Mar. sept.—Tanger (*Webb Herb.*)! In monte Djebel Kebir! In clivo maritimo haud procul Cap. Sparte!
- Ar. Geog. Europa occidentalis.* Regio Mediterranea imprimis occidentalis (inclusa Macaronesia).

Planta occidentalis refert *C. umbilicum*, var. β (L. Sp. Pl. 615); varietas α , L. l. c., ad speciem orientalem (*C. erectum*, DC.) spectat.

— —, var. HORIZONTALIS (Lowe, *Man. Fl. Mad.* i. 323), = *C. horizontalis* (Guss. *Ind. Sem. Hort. Pan.* 1826, 4; DC. *Pr.* iii. 400, sub *Umbilico*).

Mar. sept.—Tanger! (*Webb. Herb.*)! (*Schsb. Herb.* inter radices plantæ alienæ a cel. Cosson comm.)! (*Boiss. et Reut.* sub *Umbilico gaditano*).

Mar. occid.—Casa Blanca! Mogador (Lowe). In monte Djebel Hadid! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In aridissimis regionis inferioris et in regione inferiore Atlantis Majoris usque 1800 m. Aïn Oumast, prov. Shedma! Mesfoua haud procul urbe Marocco! Aït Mesan! In monte Djebel Tezali circ. 1800 m.!

Ar. Geog. Prædiligit regionem Mediterraneam occidentalem a Sicilia usque in insulam Maderam, nec deest in Oriente.

Formæ extremæ *C. umbilici* et *C. horizontalis* satis diversæ videntur; vidi tamen seriem formarum intermediarum, et stirpem Gussonianam velut subspeciem enumerare non potui. Specimina *C. umbilici* prope Cap. Spartel lecta et *C. horizontalis* ab Agadir allata vix distinguenda. *Umbilicus gaditanus* (Boiss. et Reut. *Pug.* 45) est species pessima, notis instabilibus et levis momenti fulcrata.

COTYLEDON MUCIZONIA (*Ortega, De Nov. Stirp. Cotyled.* 1772), = *C. hispida* (*Lam. Dict.* ii. 141; *Desf. Fl. Atl.* i. 359), = *Umbilicus hispidus* (DC. *Pr.* iii. 399).

Mar. occid.—Mogador (Lowe).

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis. Teneriffa (ex spec. in herb. Candolleano)? sed dubia civis, dum cl. Broussonnet plantas patriæ diversæ haud raro semel miscuit.

Nomen antiquius, lege prioritatis coactus, invitus restitui.

C. HISPANICA (L., *Benth. et H. f. Gen. Pl.* i. 659), = *Pistorinia hispanica* (DC. *Pr.* iii. 399).

Mar. merid.—In collibus aridissimis inter Aïn Oumast et Sheshaoua (lusus corollis læte flavis, tubo demum vix rubescente)! Distr. Mesfoua (forma corollis brevioribus ochroleucis, tubo demum pallide rubescente)!

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

C. hispanica est species summopere ludibunda. Qui specimina typica cum *Pistorinia Salzmanni*, Boiss., comparabit hanc pro specie omnino diversa certe habebit; sed inter eas tot et tantæ adsunt formæ intermediæ ut limites certos frustra quæret. Specimina Hispanica typica gaudent corolla rubella vel purpurea, tubo cylindrico, longitudine non valde variabili, limbo patente; dum specimina nostra Maroccana exhibent corollam flavam vel pallide ochroleucam, tubo superne paululum et vix ampliato, inter plantas simul crescentes longitudine ratione

2 ad 3 diverso, limbo subpatulo vel erectiusculo, folia ut in typo brevissima tenuia. Specimina *P. Salzmanni* prope Algeciras lecta habent corollam flavovirentem tubo fauce valde ampliato, limbo subpatulo vel erecto, folia multo majora et crassiora. In speciminibus Tingitanis *P. Salzmanni* corolla initio pallide rubella fit rubro-purpurea et tubus minus dilatatus. Denique subspecies nostra *C. Cossoniana* ex notis infra indicatis videtur omnino diversa; vereor tamen ne formis intermediis hæc quoque in typum redeat.

— — —, var. *SALZMANNI*, nob., = *Pistorinia Salzmanni* (*Boiss. Voy. Esp.* tab. 63, B), = *P. breviflora* (*Boiss. Elench.* 74, non *Coss.*), = *P. intermedia* (*Boiss. et Reut. Diagn. Pl. Or.* ser. 2, ii. 60), forma corolla longiore.

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)!

Ar. Geog. Hispania. Africa boreali-occidentalis.

In spec. Tingitanis corollæ tubus longitudine variabilis, dum longiores sistunt *P. intermedium*, B. et R.

Cotyledon Cossoniana (*Ball in Journ. Bot.* 1873, 332), = *Pistorinia brachyantha* (*Coss. Index, in Bull. Soc. Bot. Fr.* 1875, 18), = *P. breviflora* (*Coss. in Bull. Soc. Bot. Fr.* xx. 247, non *Boiss. Elench.* 74).

Descriptio *Cossoniana l. c.* exactissima quoad plantam Mogadorensensem; sed *Pistorinias* vix duas invenies locis diversis enatas inter se conformes. Specimina nostra, sicut fit in *C. hispanica*, valde ludibunda, sed sequentibus notis dignoscenda.

Planta pumila, glanduloso-viscida; inflorescentia cymoso-corymbosa; pedunculi tenues, sæpius longiusculi; calycis segmenta quam in *C. hispanica* duplo longiora, lanceolato-attenuata, apice interdum subulata, tubi corollini dimidiam vel tertiam partem attingentia; corolla parva, pallide vel rarius saturate purpurea, subcampanulata, ad tertiam vel dimidiam partem usque fissa, lobis erectiusculis ovatis vel oblongis apice cuspidatis, tubo abbreviato; stamina corollæ subæqualia. Hue, nisi fallor, referenda sunt spec. Algeriensia ad Oued Sidi el Kebir prope Blidah lecta, a cel. Cosson *P. Salzmanni* nominata.

Mar. occid.—Casa Blanca! Mazagan! Aïn el Hadjar! Mogador!

Mar. merid.—In prov. Shedma!

Pro specie omnino distincta habuerim nisi vidissem specimina ex Casa Blanca pluribus notis ad *C. hispanicam* accedentia.

Cel. Cosson nos monet hanc esse inter *Pistoriniam* et *Umbilicum* auct. forma corollæ intermedium, et confirmavit sententiam cell. Benthani et Hooker, qui hæc ad genus Linæanum *Cotyledon* revocarunt.

SEDUM (*L. Gen.* 579).

S. RUBENS (*L.*, sub *Crassula*; *DC. Pr.* iii. 405).

Mar. sept.—In monte Beni Hosmar prope Tetuan exemplar unicum legi!

Ar. Geog. Europa media. Regio Mediterranea imprimis occidentalis (inclusa Macaronesia). Deest in Asia Mediterranea.

SEDUM CÆSPITOSUM (*Cavan. Icon.* tab. 69, sub *Crassula*; *DC. Pr.* iii. 405).

Mar. occid.—Casa Blanca!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

S. MODESTUM (*Ball in Journ. Bot.* 1873, 333). (Plate XX.)

Annuum, nanum; caulis inferne nudus, glaberrimus; folia pauca, plana, late spathulata, approximata, glaberrima; rami floriferi axillares et terminales, hirtuli; cyma depauperata, subaphylla; flores hexameri vel heptameri; sepala cum pedicellis ex pilis simplicibus hirtula, ad tertiam partem usque coalita; petala oblongo-lanceolata, acuminata, pallide albo-rubentia, nervo prominulo dorso pilifero sæpius apiculata semina minutissima.

Mar. merid.—In fissuris saxorum et ad radices arborum in regione inferiore et media Atlantis Majoris. Legimus in convallibus Ourika! et Aït Mesan! et in monte Djebel Tezah! a 1100 m. ad 1900 m.!

Mtougā (*R. et Fr. spec. in Herb. Kew.*)!

Species distinctissima cui proximum, sed longo intervallo, foret *S. alsinifolium*, All.

S. DASYPHYLLUM (*L., DC. Pr.* iii. 406).

Mar. merid.—(*R. et Fr. spec. in Hort. Kew.*)!

Ar. Geog. Europa media. Regio Mediterranea imprimis occidentalis. Deest in Asia Mediterranea et in Macaronesia.

— — —, var. *OBLONGIFOLIUM*, *nob.* Foliis oblongis nec breviter obovatis insignē. Planta spontanea nitida glaberrima in Horto Kewensi culta glandulas paucas in pedicellis ostendit.

Mar. sept.—In monte Beni Hosmar prope Tetuan!

— — —, var. *GLANDULIFERUM*, *Moris*, = *S. glanduliferum* (*Guss. Prod. Fl. Sic.* i. 519).

Mar. sept.—In monte Beni Hosmar prope Tetuan (nondum florens)!

Mar. merid.—Frequens in regione inferiore ad radices Atlantis Majoris, et rarius occurrit in convallibus interioribus. Seksaoua! Milhaïn! Mzouda (*J. D. H.*)! In convalle Amsmiz alt. 1400 m. In convalle Aït Mesan circ. 2000 m.!

Ab hoc vix differt *S. corsicum*, Duby.

S. BREVIFOLIUM (*DC. Rapp. Voy.* 1808, et *Prod.* iii. 406), var. *INDURATUM*, *Coss.*, = *S. nudum* (*Schousb. Herb.*; non *Ait.*), = *S. turgidum*, (*Salzm. Exsicc. Cors.*; non *Urv. Enum.*).

Mar. sept.—Tanger in rupibus (*Webb Herb.*)! Djebel Kebir (*Schsb. Herb.*, ex *Coss. spec.*)! Prope Cap. Spartel!

Ar. Geog. Gallia. Hispania. Corsica. Ager Tingitanus.

SEDUM ALBUM (L., DC. Pr. iii. 406).

Mar. merid.—Prov. Haha alt. 750 m. (*Balansa*), Coss. *Index*.

Ar. Geog. Europa fere omnis (exclusa arctica). Asia temperata. Regio Mediterranea omnis (exclusa Macaronesia).

— — — β , MICRANTHUM (DC. Pr. iii. 406), = *S. micranthum* (Bast. in DC. Fl. Fr. v. 523; Gren. et Godr. Fl. Fr. i. 623).

Mar. merid.—Saksaona! Milhaïn! et alibi in reg. inferiore visum!

Ar. Geog. Habitat cum specie, præsertim in calidioribus: deest in Europa media.

Cel. Boissier in Fl. Orient. ii. 781, *S. micranthum* ad synonymon *S. albi* reduxit. Formæ intermediæ sæpe obviæ sunt, sed hoc sæpe in locis calidioribus solum prævalet, et facie satis diversum est ut pro varietate habeatur.

— — —, var. GLANDULIFERUM. Foliis pubescenti-glandulosis differt fere eodem modo quo *S. glanduliferum*, Guss., a *S. dasphyлло*, L., sed in nostro pedicelli et calyces glabri.

Mar. merid.—Gurguri prope Amsmiz circ. 900 m.! In convalle Amsmiz circ. 1400 m.!

S. ACRE (L., DC. Pr. iii. 407).

Mar. sept.—In montibus prope Tetuan anno 1869 legit cl. G. Maw, et specimen in horto cultum communicavit!

Mar. merid.—In regione media Atlantis Majoris, a 1600 m. usque 2500 m. In convalle Amsmiz 1600-1700 m.! In monte Djebel Tezah circ. 2300! In ascensu versus jugum Tagherot, 2500 m.!

Ar. Geog. Europa omnis (exclusa arctica). Sibiria. Montes regionis Mediterraneæ (exclusa Macaronesia).

S. PRUINATUM (Brot. Fl. Lus. ii. 209), = *S. elegans* (Lej. Fl. Spa. i. 205), = *S. reflexum*, var. γ (DC. Pr. iii. 408).

Mar. sept.—In monte Beni Hosmar anno 1869 legit cl. G. Maw, et in horto coluit; nos iterum legimus nondum florens anno 1871! Spec. cultum habui a cl. Maw.

Ar. Geog. Europa media. Peninsula Iberica. Ditus Tetuanensis. In regione Mediterranea media et orientali verisimiliter deest.

Spec. nostra exsiccata cum authenticis comparata satis conformia. Spontanea primo aspectu *S. reflexo* similia, sed flores omnino *S. elegantis* auct. Oculo fortiter armato folia exsiccata ex papillis minutis hyalinis asperula. *S. pruinatum*, ex spec. authenticis Lusitanicis, est omnino idem ac *S. elegans*. Brotero plantam vivam non vidit, sed spec. descripsit a Link communicata. Descriptio Broteriana idonea, et nomen ab ipso impositum jure prioritatis gaudet.

S. ALTISSIMUM (Poir. Dict. iv. 634; DC. Pr. iii. 408).

Mar. sept.—In ditioe Tingitana legerunt cl. Rein et Fritsch (spec. in *Herb. Kew.*)!

Mar. merid.—Ourika (*J. D. H.*)!

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia).

SEDUM HISPIDUM (*Desf. Fl. Atl.* i. 361; *DC. Pr.* iii. 409, *non Poir.*).

Mar. sept.—In monte Beni Hosmar prope Tetuan, specimen nanum nondum florens caule exsiccato anni præcedentis recognoscendum!

Ar. Geog. Africa boreali-occidentalis.

SEMPERVIVUM (*L. Gen.* 612).

S. ATLANTICUM (*Ball in Journ. Bot.* 1873, 333; *Bot. Mag.* tab. 6055).

E grege *S. tectorum* et sicut subspecies enumerandum; differt a typo notis sequentibus:—folia rosularum obovato-oblonga, apice breviter mucronata, utrinque (in planta spontanea) breviter glanduloso-pubescentia, margine ciliata; folia caulina lanceolato-oblonga, superiora glanduloso-pubescentia; calyx fere ad basin fissus; stamina omnia perfecta, virginea obscure viridia (nec rubella); squamæ hypogynæ minutissimæ glandulosæ; carpella divergentia, basi toro elevato suffulta, et a verticillis exterioribus (squamis exceptis) omnino discreta, staminibus æqualia, petalis $\frac{2}{3}$ breviora.

Mar. merid.—Rosulas foliorum legimus in convalle Aït Mesan, anno 1871; ex his, post annos duos, caules floriferi in Horto Kewensi enati sunt.

In tanta copia formarum pro speciebus novis a botanicis hodiernis, et imprimis a cl. Jordan, descriptorum notæ differentiales plane evadunt. Inter tabulas Jordanianas nullam video plantæ nostræ omnino conformem, et invito animo nomen novum proposui.

MONANTHES (*Haw. Rev. Pl. Succ.* 68).

M. ATLANTICA (*Ball in Journ. Bot.* 1873, 333), = *M. muralis* (*H. f. in Bot. Mag.* ser. 3, No. 333, tab. 5988); *non* *Petrophytes muralis* (*Webb in Bourgeau, Plant. Canar.* No. 269 et 1284, *nec* *C. Bolle in Bonplandia*, vii. 244), = *Sedum surculosum* (*Coss. Index in Bull. Soc. Bot. Fr.*).

Mar. merid.—In regione superiore Atlantis Majoris. Legi in monte Djebel Tezah alt. circ. 2400 loco uliginoso juxta nives deliquescentes! Djebel Ouensa (*Coll. Indig. Ibrahim*), *Coss.*, sub *Sedo surculoso*!

Pace amici celeberrimi stirpem nostram cum planta Canariensi supra memorata in unam speciem confundere nequeo. Ratio partium floralium est omnino diversa; calyx brevior dentibus triangularibus; petala late ovato-acuminata, nec linearilanceolata ("linearia," Bolle, *l. c.*), calycem duplo superantia, nec paulo longiora; staminodia (squamæ perigynæ) parvula, ima basi carpellorum adhærentia et ter breviora, nec magna, margine imbricata, carpellis paulo breviora ("coronam conspicuam laxam formantia," Bolle, *l. c.*); stamina longiora petalis

opposita, calyce duplo longiora sunt, et petalis subæqualia, series altera petalis alterna fit demum subæquilonga (ut videtur in tabula Webbiana *Petrophyis agriostaphis*, Phyt. Canar. tab. 36, C); carpella denique omnino *nitida*, nec *undique papillosa*, latiora sunt et apice stylifero demum divaricata. His addas characteres levioris momenti: scil., flores pentameri vel rarius hexameri, folia in ramis florigeris sparsa, et in planta spontanea tantum basi rosulata, calyx semper *glaberrimus* nec glandulis rubris hirtus, pedicelli subglaberrimi sed interdum pilis paucis glandulosis adpersi.

Propior erit *P. agriostaphis*, Webb, et forsan *P. tilophila*, Bolle (nomen barbarum!), species montanæ, sed ambæ a nostra satis superque diversæ. De speciebus Canariensibus quarum 5 jam descripsit beatus P. B. Webb, et 4 addidit clar. C. Bolle, judicent qui eas in loco natali vivas videbunt. In sicco extricatu nimis difficiles sunt. In planta nostra ovula pauca adscendentia versus basin follieni ad suturam ventralem affixa sunt, ut jam innotuit celeb. J. D. Hooker loco citato. Si eadem se habet structura in sp. affinibus, foret character genericus non parvi momenti.

Tabula Fitchiana in Bot. Mag. l. c., mendis nonnullis non caret; stamina longiora petalis alternantia, nec opposita, et nimis brevia ostendit; folia caulina (in planta spontanea) longiora sunt et basin versus attenuata, carpella matura in vivo sunt magis inflata, nec superne sub stylo sensim attenuata.

Cel. Cosson, verisimiliter ob squamas perigynas parvulas et flores sæpe pentameros, plantam nostram inter *Seda* collocavit; sed, ut mihi videtur, invita natura, nisi genus *Monanthes* pro subgenere *Sedi* habendum sit. In *Monanthe* numerus partium floralium quam maxime variabilis; nil mirum igitur si dantur etiam species pentameræ. Genus videtur satis diversum, calycis basi disciformi, stigmatibus capitellatis, ovulis adscendentibus, et habitu singulari a *Sedo* dignoscendum.

DROSERACEÆ (DC. *Théor. Elém.* 214).

DROSOPHYLLUM (*Link, in Schrad. N. Journ.* 1806, p. 2).

D. LUSITANICUM (*Link, l. c.; DC. Pr.* i. 320), = *Drosera lusitanica*, L. Mar. sept.—Prope Tanger in monte Djebel Kebir (*Salzm. Exsicc.*)! (*Schsb. Herb.*), *Coss. spec.*! (*Webb Herb.*)! Prope Cap. Spartel! In saxosis calcareis prope Tetuan!

Ar. Geog. Species propria peninsulæ Ibericæ et Mauritaniæ Tingitanæ.

HALORAGEÆ (*R. Br. Gen. Rem.* 17).

CALLITRICHE (*L. Gen.* 13).

C. VERNALIS (*Kütz. in Linnæa*, vii. 175), = *C. verna* (L., *DC. Pr.* iii. 70), pro parte.

Mar. sept.—Tanger!

Ar. Geog. In temperatis et reg. frigidioribus hemisphæræ borealis fere ubique. In regione Mediterranea minus frequens.

MYRIOPHYLLUM (*L. Gen.* 1066).

M. SPICATUM (*L., DC. Pr.* iii. 68).

Mar. sept.—? Ex Mauritania, verisimiliter Tingitana (*Salzmann, sec. Webb, Phyt. Can.* ii. 4).

Ar. Geog. In aquis stagnantibus et lente fluentibus hemisphæræ borealis (exclusa aretica et tropicali), et ad prom. Bon. Spei.

MYRTACEÆ (*R. Br. Gen. Rem.* 14).

MYRTUS (*L. Gen.* 617).

M. COMMUNIS (*L., DC. Pr.* iii. 239).

Mar. sept.—In collibus Tingitanis! El Araiseh (*R. et Fr. spec. Hort. Kew.*)!

Ar. Geog. Regio Mediterranea omnis. In Macaronesia nonnisi in insula Madeira.

In horto Præfecti juxta pagum Amsmiz ad radices Atlantis vidimus varietatem singularem *M. communis* ramis erassioribus velutinis, foliis magnis $1\frac{1}{2}$ –2-pollicaribus, præsertim secus nervum medium pilosiusculis, demum subglaberrimis, non lucentibus, dense confertis, florum pedicellis quam folia dimidio brevioribus. De origine hujus varietatis nil certi comperire potui, sed verisimiliter ex hortis urbis Marocco introducta est. In itinere per provincias meridionales Imperii *M. communis* spontanea non nobis obvia fuit.

LYTHRARIÆ (*Juss. Dict. Sc. Nat.* xxvii. 453).

PEPLIS (*L. Gen.* 446).

P. NUMMULARIÆFOLIA (*Lois. Not.* 74, 1810, et *DC. Pr.* iii. 81, sub *Lythro*; *Jord. Obs. Fragm.* iii. 85. tab. 5. fig. D), = *P. erecta* (*Req. in Benth. Cat. Pl. Pyr.* sine diagnosi, 1826; *Gr. et Godr. Fl. Fr.* i. 598; *Boiss. Fl. Or.* ii. 742), = *P. biflora* (*Salzm. Exsicc. Tingit.* 1825; *DC. Pr.* iii. 77), = *P. tithymaloides* (*Bertol. Fl. It.* iv. 233).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Nec a nobis nec ab aliis botanicis recentioribus in agro Tingitano visa.

Ar. Geog. Regio Mediterranea imprimis occidentalis (exclusa Macaronesia). In Oriente nonnisi prope Byzantium indicata. In Italia et Græcia desideratur.

Nomen antiquius restitui, dum hæc est certissime *Lythrum nummulariæfolium*, *Lois. l. c.*, et fere certe etiam *Pers. Syn.* ii. 8, quamvis ob descriptionem maneam nomen Persoonio nontribuendum. Nomen *erecta* (ineptum, dum sp. affines pariter erectæ), siue nota diagno-

stica editum, recentius, et nulla auctoritate gaudens, omnino rejiciendum. Nomen Salzmannianum infelix, dum flores sæpiissime solitarii.

LYTHRUM (*L. Gen.* 604).

L. HYSSOPIFOLIUM (*L., DC. Pr.* iii. 81).

Mar. sept.—In monte Djebel Kebir!

Mar. occid.—Casa Blanca! Mazagan!

Ar. Geog. Herba amphigea in temperatis fere totius orbis obvia, per omnem regionem Mediterraneam diffusa.

Specimina pr. Mazagan lecta sunt diffusa basi prostrata; cætera omnia erecta.

—, var. *THYMIFOLIUM*, = *L. thymifolium* (*DC. Pr.* iii. 81, fors. *L. et auct. plurim.*; non *Moris, Fl. Sard.*).

Mar. sept.—Prope Cap. Spartel (*Schousb. Herb.*), *Coss. spec.*!

Ar. Geog. Regio Mediterranea fere omnis. In Madeira? Rossia australis.

Specimina a cel. Cosson benevole communicata referunt formam intermedium inter typum et *L. thymifolium* auct. In collibus humidiusculis prope San Roque haud procul Gibraltar legi formas plurimas, quibus transitus a typo in *L. thymifolium* plane demonstratur. Specimina nonnulla habent folia anguste linearia et calyces 8-dentatos, altera a *L. hyssopifolio* nullomodo distinguenda sunt, altera intermedia.

L. flexuosum (*Lag. Gen. et Sp. Nov.* 16; *DC. Pr.* iii. 82), = *L. acutangulum* (*Lag. l. c.*), = *L. Græfferi* (*Ten. Fl. Nap.* iv. 256; *DC. Pr. l. c.*), = *L. alatum* (*Presl, non Pursh*), = *L. Preslii* (*Guss. Pl. Rar.* 188), = *L. hyssopifolium* (*L., Herb., non Sp. Pl.* 642; *D'Urv. Enum.* 52).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Inter Tanger et Tetuan! Prope Tetuan (1851)!

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—Sat frequens in irrigatis et locis humidiusculis regionis inferioris. Legimus ad Aïn Beida inter Sheshaoua et urbem Marocco! in distr. Mesfioua! et prope Ourika!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia).

A *L. hyssopifolio* facile distinguitur floribus duplo triplove majoribus læte roseis, sed hæ notæ diagnosticæ parvi momenti et parum stabiles. Hoc sæpius basi prostratum radicans, dum *L. hyssopifolium* sæpius sed non semper erectum. Folia latiora basi subcordata; sed similia vidi etiam in *L. hyssopifolio*. Pro subspecie enumeravi.

L. SALZMANNI (*Jord. Obs. Fragm.* v. 42, tab. 2. fig. B), = *L. bibracteatum* (*Salzm. in DC. Pr.* ii. 81), = *L. tribracteatum* (*Salzm. Exsicc. Tingit.*; *Spreng. Syst. Cur. Post.* 190), = *L. thymifolium* (*Sibth. et Sm.*

Pr. Fl. Gr. i. 321 ; *Moris, Fl. Sard.* ii. p. 71 ; *L. Sp. Pl.* 642 ; quoad synon. *Barrelieri*), = *L. thymifolium*, var. β (*DC. Pr.* ii. 81).

Mar. sept.—Prope Tanger a b. Salzmann detectum (*Webb Herb.*)!

Ar. Geog. Species per regionem Mediterraneam (exclusa Macaronesia) late diffusa, nullibi frequens, sed verisimiliter a botanicis plurimis prætervisa.

Videtur probe distincta species, nec eum *L. thymifolio* confundenda.

Nomen Salzmannianum *bibracteatum* ineptum, dum sp. omnes affines floribus bibracteatis gaudent ; *tribracteatum* erroneum, dum bractææ semper binæ.

LYTHRUM SALICARIA (*L., DC. Pr.* iii. 82), var. TOMENTOSUM (*DC. Pr. l. c.*), = *L. tomentosum* (*Mill. Dict. sec. Boiss.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Species per hemisphæram fere omnem borealem temperatam, et usque in Australiam diffusa, fit rarior in reg. Mediterranea calidiore, ubi var. nostra sat frequens, sed in ditone Maroccana hæc etiam rara, a nobis nullibi visa. Tam species quam varietas deest in Macaronesia.

In Marocco colitur hinc inde *Lawsonia alba* (*Lam. Dict.* i. 106), = *L. inermis*, *L.*, et *L. spinosa*, *L.* ; sed nec indigena nec spontanea facta.

PUNICA (*L. Gen.* 618).

P. GRANATUM (*L., DC. Pr.* iii. 3).

Mar. sept.—Prope Tetuan videtur spontanea !

Mar. occid.—Casa Blanea ! Mogador (*Lowe Cat.*).

Mar. merid.—Colitur frequenter in prov. meridionalibus et verisimiliter per totum Imperium et sæpe videtur spontanea, sed pro specie vere indigena vix habenda. Mtouga (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Ex India boreali-occidentali olim allata ; hodie in regione Mediterranea sponte provenit.

ONAGRARIÆ (*Juss. Ann. Mus.* iii. 315).

EPILOBIUM (*L. Gen.* 471).

E. HIRSUTUM (*L. Sp. Pl.* 494, pro parte ; *DC. Pr.* ii. 42).

Mar. occid.—Mogador (*Lowe Cat.*).

Ar. Geog. Europa omnis (exclusa aretica). Asia temperata. Per regionem Mediterraneam (exclusa Macaronesia) dispersa, sed infrequens. Indieatur etiam in Abyssinia et ad prom. Bon. Spei.

E. PARVIFLORUM (*Schreb. Spic. Fl. Lips.* ; *DC. Pr.* iii. 43), = *E. molle* (*Lam. Dict.* iii. 475), = *E. pubescens* (*Schousb. Gew. Marok.* 171).

Mar. sept.—Tanger (*Schsb. l. c.*).

Ar. Geog. Europa omnis (exclusa aretica). Regio omnis Mediterranea (inclusa Macaronesia).

E. TETRAGONUM (*L., DC. Pr.* iii. 43 ; *Schousb. Gew. Marok.* 171).

Mar. sept.—Tanger ! (*Schsb. l. c.*).

Ar. Geog. Per hemisphæram borealem (exclus. reg. arcticis et zona tropica) late diffusum; in regione Mediterranea omni (inclusa Macaronesia) satis frequens. Ad prom. Bon. Spei, et in insula Tasmania obvium.

Stirps valde ludibunda in species plurimas ab auctoribus dilaniata. Harum nonnullæ videntur satis constantes et sient subspecies enumerandæ.

— —, var. ? TINGITANUM, = *E. tingitanum*, *Salzm. Exsicc.*

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Forma insignis, floribus duplo triplove majoribus *E. hirsuti* æmula, caule $1\frac{1}{2}$ –3-pedali, inferne pallide fuseo nitido, superne pubescente, lineis obscuris, vix tetragono, ramoso, foliis oblongis basin versus attenuatis, etiam inferioribus sessilibus, mediis latiusculis apice ovato-obtusis, supremis angustis obtusiusculis. Semina matura non vidi. Pluribus notis ad *E. Tournefortii* (Michx. in Bull. Bot. Soc. Fr. 1855, 731; Boiss. Fl. Or. ii. 748) accedit, sed in hoc caules ex lineis elevatis subalati.

CUCURBITACEÆ (*Juss. Gen.* 393).

CITRULLUS (*Schrad. in Eckl. et Zeyh. Enum.* 279).

C. COLOCYNTHIS (*L., DC. Pr.* iii. 302, sub *Cucumi*; *Schrad. Linnæa*, xii. 414).

Mar. oecid.—Prope Saffi (*J. D. H.*)!

Mar. merid.—In saxosis aridissimis regionis inferioris satis frequens et certe indigenus. Legimus pr. Sheshaoua! Mzouda! Mtouga! Prope Frouga (*G. Maw spec.*)!

Ar. Geog. Regio Mediterranea calidior. Reg. subtropica Africæ borealis et Indiæ orientalis. Ins. Canarienses et Cap. Viridis. In Europa nonnisi in Hispania Bætica et in insula Pantellaria detecta.

ECBALIUM (*Rich. in Dict. Class. Hist. Nat.* vi. 19).

E. ELATERIUM (*L., DC. Pr.* iii. 311, sub *Momordica*; *Rich. loc. cit.*).

Mar. sept.—Visum inter Tetuan et Ceuta!

Ar. Geog. Regio Mediterranea fere omnis, imprimis orientalis. Deest in Macaronesia.

BRYONIA (*L. Gen.* 1093).

B. DIOICA (*Jacq. Fl. Austr.* tab. 199; *DC. Pr.* iii. 307).

Mar. oecid.—Prope Mogador!

Mar. merid.—Sat frequens in dumetis et sepibus ad radiées Atlantis Majoris. Amsmiz! Aït Mesan 1300–1400 m.!

Ar. Geog. Europa media. Regio fere omnis Mediterranea (exclusa Macaronesia).

B. ACUTA (*Desf. Fl. Atl.* ii. 360; *DC. Pr.* iii. 308).

Mar. sept.—Tanger!

Mar. occid.—Agadir (*Coll. Indig.* 1871)! Prov. Haha haud procul Mogador!

Mar. merid.—Satis frequens in regione inferiore et ad radices Atlantis Majoris. Legimus in prov. Shedma! pr. Mzouda! et Seksaoua! Mtouga (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Africa boreali-occidentalis. Sicilia.

[CACTEÆ nullæ in ditione Maroccana indigenæ sunt; sed *Opuntia vulgaris*, Mill., olim in Africam borealem introducta, hodie spontanea facta, per territorium omne nobis cognitum frequens occurrit.]

FICOIDEÆ (*Juss. Gen.* 315).

MESEMBRYANTHEMUM (*L. Gen.* 628).

M. NODIFLORUM (*L., DC. Pr.* iii. 447).

Mar. occid.—Saffi! Mogador!

Mar. merid.—Satis frequens, præsertim in saxosis aridissimis regionis inferioris. Legimus prope Sheshaoua! Misra Ben Kara! Ourika! Prov. Shedma!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia).

M. CRYSTALLINUM (*L., DC. Pr.* iii. 448).

Mar. occid.—Mogador, præsertim in insula! Saffi!

Ar. Geog. In arenosis et rupestribus maritimis regionis Mediterraneæ calidioris (inclusa Macaronesia) late diffusa. Occurrit etiam in Africa australi extratropica.

AIZOON (*L. Gen.* 629).

A. CANARIENSE (*L., DC. Pr.* iii. 453).

Mar. merid.—Distr. Mesfioua haud procul urbe Marocco! Frouga (*G. Maw spec.*)!

Ar. Geog. Per regionem Mediterraneam calidiorem subtropicam dispersa, nec in Europa, nec versus litora Maris Interni se extendit. Indicatur in India boreali-occidentali, et ad prom. Bon. Spei.

A. HISPANICUM (*L., DC. Pr.* iii. 454).

Specimina ex Imperio Maroccano, verisimiliter a b. Schousboe communicata, exstant in herb. Webb sine loco natali!

Ar. Geog. Oriens. Hispania australis. Africa borealis.

UMBELLIFERÆ (*Juss. Gen.* 218).

HYDROCOTYLE (*L. Gen.* 325).

H. VULGARIS (*L., DC. Pr.* iv. 59; *Schousb. Gew. Marok.* 112).

Mar. sept.—Tanger (*Schsb. l. c.*).

Ar. Geog. Europa omnis (exclusa arctica). In regione Mediterranea rarior, in Oriente nonnisi in insulis Zacyntho et Creta detecta; desideratur in peninsula Iberica australiore et in Macaronesia.

ERYNGIUM (*L. Gen.* 324).

E. CAMPESTRE (*L., DC. Pr.* iv. 88; *Schousb. Gew. Marok.*).

Mar. sept.—Tanger (*Schsb. l. c.*). (*Salzm. spec. in herb. nost.*)!

Folia tantum visa haud procul Tanger!

Ar. Geog. Europa media. Regio Mediterranea fere omnis. Desideratur, sicut et omnes sp. hujus generis, in Macaronesia.

E. BOURGATI (*Gouan, Ill.* 7, tab. 3; *DC. Pr.* iv. 88), var. *ATLANTICUM* (*Ball in Journ. Bot.* 1873, 334). Differt a typo habitu graciliore, capitulis minoribus, involneri phyllis longioribus parce spinosis.

Mar. merid.—In regione superiore Atlantis Majoris. In monte Djebel Tezah circ. 2400 m.!

Ar. Geog. Species habitat in Pyrenæis et in montibus Hispaniæ; forma proxime affinis in Oriente.

Specimina nostra non florentia, caulibus fructiferis anni præcedentis prædita, et ideo haud omnino certa. Var. nostræ proximum est *E. Heldreichii* (*Boiss. Diagn. Pl. Or. ser. i. x.* 20), phyllis involueri earinatis, margine integris. Vidi tamen inter spec. *E. Heldreichii* phylla nonnulla margine parce spinulosa, et hoc pro varietate vel saltem pro subspecie habendum videtur.

E. DILATATUM (*Lam. Dict.* iv. 755; *DC. Pr.* iv. 88), = *E. amethystinum* (*Schousb. Gew. Marok.* 112).

Mar. sept.—Tanger (*Salzm. Exsicc.* sub "*E. amethystinum*")! (*Schsb. l. c.*).

Ar. Geog. Regio Mediterranea occidentalis. Deest in Oriente.

E. MARITIMUM (*L., DC. Pr.* iv. 89).

Mar. sept.—Tanger! (*Webb Herb.*)! Ex Maroeco septentrionali (*Brouss.*) see. Cavanilles.

Ar. Geog. In litoribus arenosis Europæ omnis temperatæ, Africæ borealis et Asiæ Mediterraneæ.

E. TRIQUETRUM (*Vahl, Symb.* ii. 46; *DC. Pr.* iv. 90).

Mar. occid.—Casa Blanca!

Mar. merid.—Satis frequens in regione inferiore. Sheshaoua! prope urbem Maroeco! Distr. Mesfioua! Amsmiz! Aïn Tarsil! Reraya! Ourika (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Africa boreali-occidentalis. Sicilia.

E. DICHOTOMUM (*Desf. Fl. Atl.* 226, tab. 55; *DC. Pr.* iv. 90).

Mar. sept.—Legi specimen unicum, caule fructifero anni prioris, in monte Beni Hosmar prope Tetuan!

Ar. Geog. Africa borealis ex Ægypto usque montes Tetuanenses. Hispania. Sicilia.

E. VARIIFOLIUM (*Coss. in Ind. Pl. Maroc.* 19), = *E. dichotomum*, var. *feariæfolium* (*Ball in Journ. Bot.* 1873, 334).

Mar. merid.—Legimus nondum florens, caulibus exsiccatis anni prioris,

in uliginosis Atlantis Majoris circ. 1700 m. in convalle Aït Mesau !
Ex monte Djebel Ouensa attulit indigenus Ibrahim et spec. comm.
Cosson !

Planta Atlantica mihi non satis nota, a ccl. Cosson innox describenda.

ERYNGIUM AQUIFOLIUM (*Cav. Anal. Cienc. Nat.* iii. 32 ; *DC. Pr.* iv. 91).

Mar. sept.—? Tanger (*Brouss., sec. Cav. l. c.*).

Ar. Geog. Hispania Gaditana. Mauritaniæ dubia civis.

A botanicis recentioribus nusquam in agro Tingitano visum, nec a sedulo
scrutatori b. Schousboe. Broussonnet forsân specimen Hispanicum
inter plantas Tingitanas misit ?

E. TRICUSPIDATUM (*L. Sp. Pl.* 337 ? ; *Desf. Fl. Atl.* i. 224 ; *DC. Pr.*
iv. 92 ; *Schousb. Gew. Marok.* 110).

Mar. sept.—In monte Djebel Kcbir et alibi pr. Tanger ! (*Salzm. Exsicc.*) !
(*Schsb. l. c.*) (*Webb Herb.*) ! Tetuan (1851) !

Mar. occid.—Casa Blanca ! Mazagan ! Agadir (*Coll. Indig.* 1871) !

Mar. merid.—Plurics visum, specimen legit G. Maw prope Tasseremout !

Ar. Geog. Regio Mediterranea occidentalis. Africa borealis. Hispania
australis. Sardinia. Sicilia. Græciæ dubia civis.

E. ILICIFOLIUM (*Lam. Dict.* iv. 757 ; *DC. Pr.* iv. 92).

Mar. occid.—Sallé (*Brouss. ex Cavanilles*). Saffi ! Aïn el Hadjar !
Djebel Hadid (*R. et Fr. spec. in Herb. Kew.*) ! Prov. Haha laud
procul Mogador ! Agadir (*Coll. Indig.* 1871) !

Mar. merid.—Occurrit frequentissime in regione inferiore, sæpius nanum,
interdum ramis plurimis 6-8-pollicaribus. Prov. Shedma pluribus
locis ! Aïn Tarsil !

Ar. Geog. Africa boreali-occidentalis. Peninsula Iberica.

E. TENUE (*Lam. Dict.* iv. 755 ; *DC. Pr.* iv. 92).

Mar. occid.—Mogador (*Brouss. ex Cavanilles*) (*R. et Fr. spec. in Herb.*
Kew.) ! Aïn el Hadjar !

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

HIPPOMARATHRUM (*Link, Enum. H. Berol.* i. 271,
non Rivin.).

H. PTEROCHLÆNUM (*DC. Pr.* iv. 237, sub *Cachryde* ; *Boiss. Ann. Sc.*
Nat. ser. 3, ii. 74), = *Cachrys sicula* (*L., Herb., ex Gay ; an Sp. Pl.*
355 ?).

Mar. sept.—Tanger ! (*Salzm. Exsicc.*, sub “ *Cachrys sicula* ”) ! (*Webb*
Herb.) !

Mar. merid.—? Prov. Mtonga (*R. et Fr. spec. in Herb. Kew.*) !

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis. Sardinia,
Deest, sicut etiam sp. affines, in Macaronesia.

H. cristatum (*DC. Pr.* iv. 238, sub *Cachryde* ; *Boiss. Fl. Or.* iv. 932),
= *Cachrys sicula* (*Sibth. et Sm. Fl. Gr. tab.* 278 ; *Urv. Enum.* 31 ; *non*
L.), var. BOCCONI = H. Bocconi (*Boiss. Ann. Sc. Nat.* ser. 3, ii. 74),

=*C. Libanotis* (*Salzm. Exsicc. et Schousb. Herb., non L.*), = *C. lævigata* (*Brouss., non Lam.,* sec. specimen ab ipso Broussonnet in *Herb. Kew.*), = *C. pungens* (*Jan.,* sec. spec. Genuense ab ipso Jan in *Herb. Kew.*, et verisimiliter *DC. Pr. iv. 238*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Schsb. Herb.*) *Coss. spec.!* (*Brouss. in Herb. Gouan, hodie Kew.*)!

Ar. Geog. *H. cristatum* in regione Mediterranea orientali, *H. Bocconi* in occidentali obvium, sed ob synonyma dubia loci natales plurimi incerti.

Hippomarathrum Libanotis (*L., et DC. Pr. iv. 237, sub Cachryde; Koch in litt., sec DC. loc. cit.*).

Mar. merid.—? Aïn Tarsil spec. nimis immaturum! et verisimiliter eadem planta, foliis tantum expansis, prope Sektana!

Ar. Geog. “Mauritania; Sicilia” (fide *DC. l. c.*).

Ab affinibus differt herba glauca glabra, foliis minus divisus segmentis breviter trifidis, involuero ex foliolis latoribus, inferne ex nervo prominulo bisulcatis, margine scabris, radiis dimidio brevioribus, integris linearibus crassiusculis, involucellis brevissimis. Spec. nostra, ob fructus defectum incerta, forsân ad sp. præcedentem referenda.

Hippomarathra omnia mihi nota valde ludibunda sunt et notæ differentiales parum stabiles. Optimo consilio, ut mihi videtur, cel. Moris plurimas formas sardoas in unam speciem (*Ægomarathrum siculum*, *Fl. Sard. ii. 176*) reduxit. Specimina fructifera in herbariis sæpe desunt, et diagnoses nimis arduæ et ineertæ fiunt. Nisi obstitisset auctoritas summorum virorum, qui species affines in sectiones generis proprias, immo in genera diversa collocarunt, *Hippomarathra* omnia Europæa et Africana boreali-occidentalia in unam speciem nomine triviali Linnaeo (*H. siculum*) reducere maluissem.

[Quid sit *Cachrys humilis* (*Schousb. Gew. Marok. 118*), prope Cap. Spartel cum *Crithmo maritimo* creescentem detecta, eruere non potui. Planta ulterius inquirenda, a nemini post Schousboe visa.]

CONIUM (*L. Gen. 336*).

C. MACULATUM (*L., DC. Pr. iv. 242*).

Mar. sept.—Tetuan (*Webb Herb.*)!

Mar. merid.—In hortis et ruderatis urbis Marocco!

Ar. Geog. Europa omnis (exclusa arctica). Regio Mediterranea omnis (inclusa Macaronesia). Asia temperata. Abyssinia.

SMYRNIUM (*L. Gen. 363*).

S. OLUSATRUM (*L., DC. Pr. iv. 247*).

Mar. sept.—Prope Tetuan!

Mar. merid.—Secus rivulum prope Amsmiz!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia). In Anglia et Gallia occidentali, olim introductum, nunc spontaneum.

BUPLEURUM (*L. Gen.* 328).

B. TENUISSIMUM (*L., DC. Pr.* iv. 127), var. **COLUMNÆ**, *Coss.*, = *B. Columnæ* (*Guss. Pr. Fl. Sic. Suppl.* 70).

Mar. sept.—Tanger (*Schsb. Herb.*), *Coss. spec.*!

Ar. Geog. Regio Mediterranea. Deest in Oriente (exceptis insulis Maris Interni), et in Macaronesia. In cultis Europæ mediæ usque in Sueciam australem diffusum, sed verisimiliter eum seminibus introductum.

B. procumbens (*Desf. Fl. Atl.* i. 230, tab. 56; *DC. Pr.* iv. 127; *Schousb. Gew. Marok.* 113).

Mar. merid.—In provincia Haha (*Schsb. l. c.*).

Ar. Geog. Africa borealis. Sardinia.

Nimis proximum *B. tenuissimo*, sicut subspecies, nisi forte varietas, enumerandum.

B. GLAUCUM (*Rob. et Cast. in DC. Suppl. Fl. Fr.* 515; *DC. Pr.* iv. 127), = *B. semicompositum* (*Sibth. et Sm. Fl. Gr.* tab. 261; *DC. Pr.* iv. 128, *vix L.*).

Mar. occid.—Mogador! Aïn el Hadjar! Prov. Haha, haud procul Mogador.

Mar. merid.—Ex prov. Shedma haud procul castello Gubernatoris!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

B. semicompositum (*L. Sp. Pl.* 342) est species ineerta, hodie vix extricanda. Planta est Hispanica, notis diagnosticis non defuita, *B. odontiti* affinis, et ideo a sp. nostra diversa.

B. ODONTITES (*L., DC. Pr.* iv. 129)?

“Mauritania” (*DC. Pr. l. c.*). Ditionis nostræ dubia eivis, sed verisimiliter non deest.

Ar. Geog. Regio Mediterranea imprimis orientalis. Spec. ex Africa boreali nondum vidi. Deest in Macaronesia.

B. PROTRACTUM (*Link et Hoffm. Fl. Port.* ii. 387; *DC. Pr.* iv. 129), = *B. rotundifolium* (*Schousb. Gew. Marok.* 112, *non L.*).

Mar. sept.—Tanger! (*Webb Herb.*)! (*Schsb. l. c.*).

Ar. Geog. Regio Mediterranea fere omnis. In Macaronesia nonnisi in ins. Madeira detectum. Gallia media et occidentalis.

B. PANICULATUM (*Brot. Fl. Lus.* i. 455; *DC. Pr.* iv. 131).

Mar. sept.—Tanger (*Schsb. Herb.*, sub “*B. nervosum*”), *Coss. spec.*!

Ar. Geog. Peninsula Iberica. Ditio Tingitana.

B. SPINOSUM (*L. fl., DC. Pr.* iv. 133).

Mar. merid.—In Atlante Majore valde frequens, a regione inferiore (e. g. Tasseremout, 1100 m.!) usque in reg. alpinam (in jugo Tagherot, 3500 m.!).

Ar. Geog. Africa boreali-occidentalis. Hispania. Corsica.

Cum *Alyso spinoso* adsociatum dumeta densa in clivo Atlantis efficit.

BUPLEURUM FRUTESCENS (L., DC. Pr. iv. 133).

Mar. sept.—Tetnan (1851)! (*Webb Herb.*)!

Ar. Geog. Hispania, Gallia australis. Africa boreali-occidentalis.

Specimen nostrum Hispanicis sat conforme, sed omnia quæ vidi Africana elatiora et minus ramoso-intricata sunt. Foliola involueri et involucelli ab auctoribus dicuntur *subulata*; sed in multis spec. a me scrutatis hæc semper lanceolata sunt, interdum subacuta, sæpius subacuminata, in spec. ex Djebel Zaghouan a cl. Kralik lecta plane acuminata sunt. Hæc nota et radiis filiformibus planta Kralikiana ad sequentem proxime accedit.

B. oligactis (*Boiss. Diagn. Pl. Or. ser. 2, ii. 84*), = *B. exaltatum*, *M. B.*, var. *oligactis* (*Coss. Ind. Pl. Maroc.*).

Mar. merid.—In monte Djebel Sidi Fars, alt. 1200 m. (*Balansa*) *Coss.*

Index. Specimen nimis immaturum et incertum legi prope Seksaoua!

B. oligactis, Boiss., planta Africæ borealis mihi non satis cognita, monente cel. auctore est valde proximum *B. frutescenti*, et ex descriptione subspeciem florum pedicellis paulo longioribus distinguendam sistit. Nota diagnostica ex forma foliolorum involueri et involucelli certe fallax. In *B. frutescente* pedicelli semper plus minusve inæquales. Video flores nonnullos subsessiles, alios pedicellis quam involucellum dimidio brevioribus fulcratos. Si bene novi *B. exaltatum*, *M. B.*, hoc a *B. oligactine* differt, præter alias notas, fructus jugis prominentibus, subalatis, uce obtusissimis subevanidis. Sententia cel. Boissier, qui *B. exaltatum* cum *B. Sibthorpiano* ad *B. gramineam* (Vill. Fl. Dauph. ii. 575) reduxit, mihi melius placuit; sed nomen Villarsianum Hosteano recentiori (*B. baldense*) anteponendum. Confer Fl. Or. ii. 847.

B. DUMOSUM (*Coss. et Bal. in Balans. Pl. Maroc. 1867*; *Coss. in Bull. Soc. Bot. Fr. xx. 249*).

Sequitur descriptio Cossoniana:—"Frutex circiter 1 m. altus, erectus, divaricatim ramosus, dumosus, caule ramisque vetustis lignosis cortice cinereo-fuscescente vel nigrescente, ramis florigeris annotinis frutescentibus striatis glabris. Folia parvula, alterna, pleraque in ramis axillaribus abbreviatis fasciculata, glaucescentia, coriacea, pereunantia, anguste oblonga, 3-nervia, nervis parallelis prominulis, venis transversalibus obsoletis, margine lævigata vel scabriuscula, etiam inferne laud albo-marginata, sessilia, apice abrupte mucronata mucrone recurvo. Umbellæ laxè paniculatæ, 3-4-radiatæ, radiis subæqualibus gracilibus suberectis. Involueri 3-4-phylli foliolis subæqualibus lanceolato-linearibus, radiis umbellæ multoties brevioribus. Involucello 3-5-phylla pedicellos longe excedentia, foliolis linearibus acuminatis. Pedicelli brevissimi fructu multoties breviores. Fructus oblongus, nitidulus, jugis vix prominentibus, valleculis levibus."

Mar. merid.—In montibus ad austro-occidentem urbis Maroc. in parte superiore montis Aït Ongourt prope Keïra, ad 1100 metr.; fructiferum legit mense Majo 1867 cl. Balansa. Spec. in herb. nost.!

Juxta *B. fruticescens*, L., collocandum, sed, monente cel. auctore, distinctissimum caule fere toto lignoso, foliis oblongis, non linearibus, plerisque in axillis fasciculatis, umbellulis 2-5-floris, non plurifloris.

Hæc est species præcox: eodem mense, anno 1871, *B. acutifolium* vix florens vidimus, et in *B. laterifloro* et *B. oblongifolio* alabastra umbellifera nondum expansa.

BUPLEURUM ACUTIFOLIUM (Boiss. *Elench.* 83; *Voy. Esp.* 246, tab. 71; *Walp.* ii. 397).

Mar. merid.—In convalle Aït Mesan 1400–1800 m. ! In convalle Am-smiz circ. 1500 m. !

Ar. Geog. Hispania australis. Atlas Major.

Specimina nostra immatura Hispanicis conformia sunt. Florum pedicellis ter quaterve longioribus (præter alias notas) a *B. fruticescente* egregie differt.

B. FOLIOSUM (Salzm. in DC. *Pr.* iv. 133).

Mar. sept.—Tanger (Salzm. *Exsicc.*) ! In monte Djebel Kebir (Schsb. *Herb.*), Coss. *spec.* !

Ar. Geog. Species propria agri Tingitani.

Hoc est *B. tortuosum* et *B. obliquatum*, Schsb. *Herb.* Variat foliis minoribus crebris vel majoribus magis distantibus; caulina superiora semper basi dilatata subamplexicaulia. Ab omnibus affinibus differt foliorum nervo dorsali prominente, parallelis nullis, lateralibus obsoletis.

B. CANESCENS (Schousb. *Gew. Marok.* 113; DC. *Pr.* iv. 133).

Mar. sept.—El Araïsch (*R. et Fr. spec. in Herb. Kew.*) !

Mar. occid.—Mogador (Schsb.) l. c. (*Brouss., sec. Cavan. Anal. Cienc. Nat.*) (*Lowe Cat.*). Magis frequens prope Aïn el Hadjar in dumetis et sepibus usque 9-10-pedale !

Ar. Geog. Species nonnisi in litore Atlantico Mauritaniae lucusque visa. Diagnosis Candolleana haud accurata. Folia nullomodo membranacea, sed coriacea, superiora sæpius breviter mucronata, nec obtusissima, involucri et involucelli foliola ovato-lanceolata acuta, nec oblonga obtusa. Confer Schousboe l. c., qui speciem optime descripsit. Valde proximum est *B. plantagineo* (Desf. *Fl. Atl.* i. 233, tab. 57). Ille differt herba non glaucescente, foliis inferioribus multo majoribus, ovato-lanceolatis, omnibus mucronatis, involucri foliolis anguste lanceolatis duplo longioribus. Planta nostra forsitan sicut subspecies enumeranda.

B. OBLONGIFOLIUM, nob.

Fruticosum ramosum, e trunco lignoso ramos longos flexiles cylindricos teretes lineolatos emittens; folia omnia oblonga apice ovalia, mucronata, basi vix attenuata, semiamplexicaulia, margine leviter, nervis 7-9 parallelis in pagina inferiore prominentibus notata; umbellæ laxè paniculatæ 3-5-radiatæ, radii tenues inæquales; involucri foliola pauca inæqualia linearilanceolata; umbellulæ 4-6; involucelli foliola lanceolata

acuta pedicellis brevibus (inæqualibus?) duplo longiora. Fructum non vidimus.

Mar. merid.—In convalle Aït Mesan circ. 1400 m.! Prope Seksaoua, spec. unicum legit J. D. H.!

Ex ramis anni præcedentis umbellis exsiccatis adhuc onustis, prodeunt rami novelli florigeri, superne umbellas terminales et laterales paucas ramulis brevibus suffultas gerentes. Caules sub lente lineolati, vix aut ne vix striati vel sulcati.

Planta adhuc male cognita cum specie nulla adhuc descripta confundenda. A *B. dumoso*, Coss., recedit ramis tenuibus flexilibus, foliis ter longioribus insigniter 7-9-nerviis. *B. montanum*, Coss., longius distat ramis hieme marcescentibus, nec perennantibus, foliis acuminatis 5-nerviis. Hujus, sicut et nostræ, speciei fructus mihi adhuc incognitus.

BUPLEURUM LATERIFLORUM (*Coss. Ind. Pl. Maroc.* 19).

Mar. merid.—Leguminus nondum florens in convalle Aït Mesan a 1300 m. usque 1600 m.! Ex monte Djebel Afougueur attulit indigenus Ibrahim (*Coss. l. c.*).

Species Atlantica, ut videtur, ab affinibus omnino distincta, ab auctore cel. mox describenda.

B. GIBALTARICUM (*Lam. Dict.* i. 520; *DC. Pr.* iv. 133), = *B. eoriaecum* (*L'Hérit., St. Nov.* fasc. 6; *Schousb. Gew. Marok.* 113), = *B. obliquum* (*Vahl, Symb.* i. 24).

“Hinc inde in fruticetis Imperii Maroceani” (*Schsb. l. c.*).

Ar. Geog. Africa boreali-occidentalis. Peninsula Iberica.

B. FRUTICOSUM (*L., DC. Pr.* iv. 133).

Mar. sept.—(*Schsb. Herb. ex Coss. spec.*)!

Ar. Geog. Per regionem Mediterraneam omnem (exclusa Macaronesia) dispersum, nullibi frequens.

APIUM (*L. Gen.* 367).

A. NODIFLORUM (*L. et Schousb. Gew. Marok.* 120, sub *Sio*; *Benth. et H. f. Gen. Pl.* i. 888), = *Helosciadium nodiflorum* (*Koch, Umb.* 126; *DC. Pr.* iv. 104).

Mar. sept.—Tanger! Tetuan (1851)!

Mar. occid.—Casa Blanca! Mogador! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In irrigatis juxta urbem Marocco!

Ludit sæpe petiolis basi dilatatis margine membranaceis. Confer var. β , *ochreatum* (*DC. Prod. l. c.*).

Ar. Geog. Europa media. Regio Mediterranea omnis (inclusa Macaronesia). Abyssinia.

A. GRAVEOLENS (*L., DC. Pr.* iv. 101; *Schsb. Gew. Marok.* 128).

Mar. sept.—See. *Schsb.* introductum. Legi juxta ostium fluminis Tetuanensis (1851)!

Mar. occid.—In fossis inundatis juxta Ain el Hadjar haud procul Mogador, spontaneum et forte indigenum!

Mar. merid.—Prov. Shedma (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. In Europa media hodie spontaneum in salinis maritinis, sed forsani olim introductum. Regio Mediterranea omnis a Beluchistan usque insulas Canarienses. Ad prom. B. Spei et in America australi introductum.

AMMI (*L. Gen.* 334).

A. MAJUS (*L., DC. Pr.* iv. 112), inclusis *A. glaucifolio* (*L., DC. Pr. l. c.*)
et *A. intermedio* (*DC. Pr.* iv. 113).

Mar. sept.—Tetuan (1851)!

Mar. occid.—Casa Blanca! Mogador (*Lowe*).

Mar. merid.—Frequens in regione inferiore. Shedma! (*R. et Fr. spec.*)!
Aïn Beida! Prope urbem Marocco! Tassereimout ad radices Atlantis
Majoris (*G. Maw spec.*)!

———, var. TENUIS, *nob.*, verisimiliter *A. Broussonnetii* (*DC. Pr.*
iv. 13).

Foliorum segmentis tenuioribus, umbellis minoribus, radiis filiformibus, involucri foliolis lineari-subulatis brevibus, sæpius integris, interdum trifidis, fructu juniore subasperulo. Specimina pumila hujus varietatis post imbres vernaes mense Majo cnata, foliorum segmentis lineari-bus, involucrio depauperato, interdum ad foliolum unicum reducto, facie *Ptychotidi* similia sunt.

Mar. occid.—Aïn el Hadjar! Djebel Hadid! Agadir (*Coll. Indig.* 1851)!

Mar. merid.—Prov. Shedma prope Mskala!

Ex descript. Candolleana *A. Broussonnetii* a pl. nostra vix differt fructu juniore pube adpressa rara adperso.

Ar. Geog. *Ammi majus* est species valde ludibunda per regionem Mediterraneam omnem late diffusa, etiam in Abyssinia obvia. In cultis Europæ mediæ hinc inde cum seminibus introducta, et subspontanea facta.

A. VISNAGA (*L., Schousb. Gew. Marok.* 116, sub *Dauco*; *Lam. Dict.* i. i. 132; *DC. Pr.* iv. 113).

Mar. sept.—Frequens circa Tanger! (*Webb Herb.*)! (*Schsb. l. c.*).

Ar. Geog. Regio Mediterranea fere omnis (inclusa Macaronesia).

DEVERRA (*DC. Coll. Mém.* v. 45), = *Pituranthos*
(*Viv. Fl. Lib. Spec.* 15, tab. 7).

D. SCOPARIA (*Coss. et Dur. in Bull. Soc. Bot. Fr.* ii. 248).

Mar. merid.—In convalle Amismiz alt. circ. 1600 m.!

Specimina nostra nondum florentia, umbellis emarcidis anni præcedentis munita, Algeriensibus conformia.

Ar. Geog. Africa boreali-occidentalis interior.

D. juncea, *nob.*

A *D. scopario*, cui proxima, differt plurimis notis gravioribus. Caules e basi lignosa erecti simplices, nec ramosi; folia omnia ad laciniam unicam rigidam reducta, basin versus in vaginam brevem membrana-

ceam amplexicaulem expansa; umbellæ terminalis radii 5–8 breves etiam fructiferi vix 6 mm. longi; involuero oligophyllo subpersistente, foliolo externo majore; involucellæ foliola ovato-lanceolata, margine membranacea; flores præsertim centrales brevissime pedicellati, pedicellis sub flore bracteolam latam cyathiformam gerentibus; petala sub anthesi arete inflexa, nec expansa; styli longiusculi divergentes, demum adpresse recurvi.

Mar. merid.—In rupibus supra Seksaoua!

Planta singularis, pro subspecie *D. scopariæ* proposita, sed melius cognita verisimiliter pro specie omnino diversa habenda erit. Primo intuitu caulibus junciformibus, foliis ad laciniam unicam reductis, umbella contracta a cæteris *Deverris* distincta. Bracteolam sub flore cyathiformem vidi in *D. scoparia*, nec in aliis speciebus mihi cognitis.

CARUM (*L. Gen.* 365).

C. PETROSELINUM (*L.*, sub *Apio*; *Benth. et H. f. Gen. Pl.* i. 891), = *Petroselinum sativum* (*Hoffm. Umb.* i. 78; *DC. Pr.* iv. 102, et *auct. recent.*).

Mar. merid.—Subspontaneum in irrigatis ad urbem Marocco!

Ar. Geog. Herba diu culta, patriæ incertæ, videtur hodie subspontanea in regione Mediterranea.

C. RIDOLFIA (*Benth. et H. f. Gen. Pl.* i. 891), = *Ridolfia segetum* (*Moris, Fl. Sard.* ii. 212; *Boiss. Fl. Or.* ii. 858), = *Anethum segetum* (*Urv. Enum.* 33; *DC. Pr.* iv. 186, verisimiliter non *L.*).

Mar. sept.—Tanger! (*Webb Herb.*)! Tetuan (1851)!

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Satis frequens in arvis regionis inferioris. Prov. Shedna, prope Mskala! et alibi visum.

Ar. Geog. Per regionem Mediterraneam (inclusa Macaronesia) calidiorum late diffusum.

C. Ridolfia, habitu diverso, involuero et involucella deficientibus, videtur *C. Petroselino* nimis diversum ut in eandem sectionem generis enumerandum sit. Potius, ut nobis videtur, sectionem propriam sistit juxta *Trachysperma* collocandam.

C. DICHOTOMUM (*L. Mant.* 58, et *DC. Pr.* iv. 123, sub *Pimpinella*; *Benth. et H. f. Gen. Pl.* 891), = *Ptychotis dichotoma* (*Coss. MSS.*).

Mar. merid.—Ad radices Atlantis Majoris, supra Seksaoua! prope Tasseremout!

Ar. Geog. Hispania centralis. Algeria interior.

Herba sæpissime annua, pumila subspithamæa, interdum, ut videtur, fit biennis, e basi ramosissima. Involucrum et involucella desunt. Ex fructu ovato, breviter hispidulo, jugis obsoletis, forsitan propior *Pimpinellæ* quam *Caro*. Folia trita gaudent odore *Apii graveolentis* fortissimo.

C. MAURITANICUM (*Boiss. et Reut. Pug.* 49; *Walp. Ann.* v. 65).

Mar. sept.—Tetuan in monte Beni Hosmar (folia tantum)! In monte Djebel Dersa (*Webb Herb.*)!

Mar. occid.—In collibus arenosis prov. Haha, in sylvis *Arganiæ* pluribus locis!

Mar. merid.—In reg. inferiore et media Atlantis Majoris. In convalle Aït Mesan circ. 1400 m.! In rupibus supra Seksaoua versus 1200 m.! In monte Djebel Tezah usque 2000 m. (folia tantum)!

Ar. Geog. Africa boreali-occidentalis.

C. mauritanicum a *C. incrassato* (Boiss. Voy. Esp. 711) distinguitur radiis fructiferis involucri vix incrassatis, stylopodio minore, et fructu brevior. His notis specimina ex sylvis arenosis *Arganiæ* prope Mogador intermedia sunt, et sensu nostro planta Mauritanica pro subspecie habenda est. *C. ferulæfolium* (Desf. Coroll. p. 55, tab. 43, sub *Bunio*; Boiss. Diagn. Pl. Or. ser. 1, x. 22), quocum conjungitur *C. divaricatum*, Koch, Syn. p. 315, differt involucri oligophyllo, carpophoro non incrassato, mericarpiis margine contiguis, stylopodio depresso, dentibus calycinis obsoletis. An hæ notæ diagnosticæ omnino constantes sint ulterius inquirendum est. In speciminibus nostris Atlanticis involucri interdum 3-4-phyllum, et carpophorum vix incrassatum.

CARUM TENUIFOLIUM (*Benth. et H. f. Gen. Pl. i. 891*), = *Geocaryum capillifolium* (*Coss. Notes critiques sur quelques Plantes du Midi de l'Espagne*, p. 113, et *Walp. Ann. iii. 900*, quoad plantam Hispanicam et Tingitanam), = *Bunium tenuifolium* (*Salzm. Exsicc., DC. Pr. iv. 118*); non *Bunium capillifolium*, *Bert.*, nec *Myrrhis tenuifolia*, *Guss.*

Mar. sept.—In monte Djebel Kebir prope Tanger (*Salzm. Exsicc.*)! specimen ex herb. Schousboe comm. cel. Cosson! Folia verisimiliter a nobis visa, sed nimis incerta.

Ar. Geog. Peninsula Iberica. Africa Tingitana.

De genere *Geocaryo* confer *Benth. et H. f. l. c.*: ab affinibus stylis rigidis erecto-divergentibus statim dignoscitur, sed invita natura a proximo *C. ferulæfolio*, Desf., genericè separatur. Planta Italica a Bertoloni et Gussone memorata certe ad *Conopodium* referenda est. An species genuina Cossoniana intra fines Italiæ detecta sit non satis eminet. Involucri in spec. nostro sæpius nullum, rarius monophyllum.

C. AMMOIDES (*L. Sp. Pl.*; *Gouan, Ill. 16*; *Schousb. Gew. Marok. 124*, sub *Seseli*; *Koch, Umbell. p. 124*, sub *Ptychotide*; *Benth. et H. f. Gen. Pl. i. 892*), = *Seseli verticillatum* (*Desf. Fl. Atl. i. 260*), = *Ptychotis verticillata* (*Dub. Bot. Gall. 235*; *DC. Pr. iv. 108*).

Mar. merid.—In collibus aridis prope urbem Marocco! “Raro legi in locis aridis” (*Schsb. l. c.*)

Ar. Geog. Species per regionem Mediterraneam occidentalem, exclusa Macaronesia, late diffusa, in Marocco, ut videtur, infrequens. Præter Græciam, ubi rarius occurrit, Orientem fugit.

Descriptio Linnæana, ut mos erat antiquorum, incompleta, sed satis conveniens, et ex loco natali certe huc referenda.

PIMPINELLA (*L. Gen.* 366).

P. TRAGIUM (*Vill. Fl. Dauph.* ii. 606; *DC. Pr.* iv. 121).

Mar. merid.—In regone media Atlantis Majoris. Spec. nondum florentia legimus in convalle Aït Mesan circ. 2000 m.! Specimina Atlantica florentia et fructifera benevole comm. cel. Cosson!

Ar. Geog. Species valde ludibunda per regionem omnem Mediterraneam (exclusa Macaronesia) late diffusa.

P. BUBONOIDES (*Brot. Fl. Lus.* i. p. 403; *DC. Pr.* iv. 121), var. *VILLOSA*, = *P. villosa* (*Schousb. Gew. Marok.* 125; *DC. Pr.* iv. 121).

Mar. sept.—In arenosis Tingitanis (*Schsb. l. c.*) spec. comm. Cosson! Folia mense Aprili in agro Tingitano pluries visa!

Mar. occid.—Mazagan!

Mar. merid.—Ad radices Atlantis Majoris pluribus locis. Sektana! Seksaoua! Milhain!

Ar. Geog. *P. bubonoides* in peninsula Iberica, Italia inferiore et Sicilia obvia est. *P. villosa*, nonnisi petalis villosis distinguenda, est forma propria ditionis Maroccanæ.

TINGUARRA (*Parlat. in Webb, Phyt. Canar.* ii. 156).

T. SICULA (*L., et DC. Pr.* iv. 155, sub *Athamantha*; *Benth. et II. f. Gen. Pl.* i. 897).

Mar. sept.—In monte Beni Hosmar prope Tetuan! (*Webb Herb.*)!

Mar. merid.—In collibus prope urbem Marocco (*G. Maw*)! In convalle Amsniz circ. 1300 m.!

Ar. Geog. Africa borealis. Sicilia. Italia australis. Baleares.

SCANDIX (*L. Gen.* 357).

S. PECTEN VENERIS (*L., DC. Pr.* iv. 221).

Mar. sept.—Frequens in agro Tingitano! (*Webb Herb.*)! Inter Tanger et Tetuan!

Mar. occid.—Casa Blanca! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In regione inferiore et in convallibus Atlantis Majoris. Prov. Shedma! Tasseremout! In convalle Aït Mesan usque 1400 m. prope Amsniz!

S. AUSTRALIS (*L., DC. Pr.* iv. 221; *Schousb. Gew. Marok.* 124).

Mar. sept.—In collibus regionis Tingitanæ (*Schsb. l. c.*). A nobis non visa.

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia).

FÆNICULUM (*Adans. Fam. Pl.* ii. 101).

F. VULGARE (*Gærtn. Fruct.* i. 105; *DC. Pr.* iv. 142), = *Anethum fœniculum* (*L., Schousb. Gew. Marok.* 125).

Mar. sept.—Visum pr. Tanger, forsan ex hortis aufuga!

Mar. occid.—Casa Blanca!

Mar. merid.—Mtouga!

— —, var. PIPERITUM (*Bertol. Am. Ital.* 21, sub *Anetho*; *DC. Pr. l. c.*).

Mar. occid.—Mazagan!

Mar. merid.—Satis frequens in regione inferiore. Prov. Shedma!

Ar. Geog. Species diu culta in Europa media; in regione Mediterranea, inclusa Macaronesia, late diffusa, et verisimiliter indigena.

Notæ diagnosticæ quibus *F. piperitum* et *F. dulce* a *F. vulgari* distinguuntur certe variabiles sunt. *F. piperitum* est forma australis, sed sæpe occurrunt specimina intermedia. Confer Lowe, *Man. Fl. Mad.* i. 359.

KUNDMANNIA (*Scop. Introd.* 115).

K. SICULA (*L.*, sub *Sio*; *DC. Pr.* iv. 143).

Mar. sept.—Tanger! Tetuan!

Mar. occid.—Casa Blanca! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In convalle Aït Mesan!

Ar. Geog. Regio Mediterranea occidentalis, exclusa Macaronesia, fere semper in zona litorali. In Oriente nonnisi in insulis Creta, Zaeyntho, Samo et Meli obvium.

MAGYDARIS (*Koch in litt.*; *DC. Coll. Mem.* v. 68).

M. TOMENTOSA (*Desf. Fl. Atl.* i. 249, sub *Cachryde*; *Koch in DC. Pr.* iv. 241).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Africa borealis. Sicilia. Sardinia.

M. PANACINA (*DC. Pr.* iv. 241), = *Cachrys ambigua* (*Salzm. Exsicc.*).

Mar. sept.—Tanger, folia tantum visa! (*Salzm. Exsicc.*)! Specimen ex herb. Schonsboe cui scheda "*Cachrys tomentosa*," Schsb. non *Desf. benev. comm. cel. Cosson*!

M. ambigua (*DC. Pr. l. c.*), = *Cachrys ambigua* (*Salzm. Exsicc.*), videtur esse nil nisi lusus levis *M. panacinæ*.

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

CRITHMUM (*L. Gen.* 340).

C. MARITIMUM (*L.*, *DC. Pr.* iv. 164; *Schousb. Gew. Marok.* 118).

Mar. sept.—Juxta Cap. Spartel! (*Schousb. l. c.*).

Ar. Geog. Secus oras oceani Atlantici a Scotia usque Macaronesiam, et in litoribus Maris Interni totius.

CENANTHE (*L. Gen.* 352).

Æ. CALLOSA (*Salzm. Exsicc.*; *DC. Pr.* iv. 137).

Mar. sept.—Prope Tanger in dumetis et in monte Djebel Kebir! (*Salzm. Exsicc.*)! (*Webb Herb.*)!

Ar. Geog. Stirps, ut videtur, ditionis Tingitanæ propria.

Species *Æ. pimpinelloidi* affinis, videtur tamen rite distincta fructu maturo basi et apice (nec basi tantum) calloso, involuero subnullo (nec polyphyllo), foliorum radicalium circuitu oblongo-ovatorum segmentis latioribus, minus profunde incis, isi sp. sequentis subsimilibus.

Æ. APIIFOLIA (*Brot. Fl. Lus.* i. 420; *DC. Pr.* iv. 137).

Mar. sept.—Prope Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)! In monte Djebel Kebir! Inter Tanger et Tetuan! Prope Tetuan secus flumen (1851)! Inter Tetuan et Ceuta!

Ar. Geog. Peninsula Iberica. Ditio Tingitana. Forma proxime affinis (subspecies?) in insula Madeira.

Æ. GLOBULOSA (*L.*, *DC. Pr.* iv. 138; *Schousb. Gew. Marok.* 120).

Mar. sept.—Tanger (*Salzm. Exsicc.*, sub “*Æ. calycina*”)! (*Schsb. l. c.*). (*Webb Herb.*)! Tetuan ad ostium fluminis (1851)!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Maearonesia).

SCLEROSCIADIUM (*Koch in litt.*; *DC. Coll. Mem.* v. 43).

S. NODIFLORUM (*Schousb. Gew. Marok.* 120, sub *Cenante*), = *S. humile* (*Koch, DC. l. c.* tab. 2. fig. F; *DC. Pr.* iv. 140).

Mar. occid.—In uliginosis prope ostium rivuli Oued Kseb prope Mogador! (*Schsb. l. c.*). (*Balansa spec.* 1867)!

Ar. Geog. Species propria Imper. Marocani austro-occidentalis.

Confer Beuth. et H. f. Gen. Pl. i. 1009, in Addendis. Pace celeberrimorum, qui specimina non viderunt, planta nostra a *Capnophyllo* (*Kruber*) non tantum specificè, quin etiam genericè distinguenda est. Fructus ovato-globosus, in vivo lævis, exsiccatu tenuiter scaber, nullemodo compressus, jugis dorsalibus elevatis albis lateralibus excedentibus; valliculæ profundæ univittatæ; commissura bivittatæ; carpophorum indivisum. Calycis dentes conici persistentes discreti. Styli in stylopodio elevato conico persistentes, demum divergentes. In *Capnophyllo* fructus ellipticus dorso insigniter compressus, jugis dorsalibus lateralibus multo minora, calycis dentes subobsoleti. In *C. africano* vittæ tenuissimæ, in *C. dichotomo* omnino obsoletæ.

In planta nostra umbella primaria inter folia radicalia subexscapa, sequentes in ramis lateralibus brevibus axillares vel terminales; involucri foliola parva angusta, tot quot radii, cito marcescentia; radii umbellæ valde inæquales; involucellæ foliola margine membranacea; flores in umbellula laterales breviter pedicellati, centrales subsessiles. Petala ex albo rubella. Radix verticalis indurata parè fibrillosa.

CAPNOPHYLLUM (*Gærtn. Fruct.* 32).

C. DICHOTOMUM (*Desf. Fl. Atl.* i. 245, tab. 66, sub *Conio*; *Lag. Gen. et Spec.* 13), = *Tordylium peregrinum* (*L. Mant.* 55), = *Kruberia leptophylla* (*Hoffm. Umb.* 104; *DC. Pr.* 199; *Bertol. Guss. et fere omn. auct. recent.*), = *Kruberia peregrina* (*Boiss. Fl. Or.* ii. 1027), = *Ulospermum dichotomum* (*Link, Enum. Hort. Bert.* i. 267).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! In elivo montis Djebel Kebir!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia) in zona litorali.

Nomen triviale Linnæanum, *Tordyllo* perperam adsociatum, jure prioritatis gaudens, sed ab auctoribus, posterioribus neglectum, prætermisi.

Planta Mediterranea a *Capnophyllo africano* affini invita natura in genus alienum segregata a Lagasea optimojure suo loco restituta est.

MEUM (*Jacq. Fl. Aust.* iv. 2).

M. ATLANTICUM (*Coss. MSS.*).

Mar. merid.—In monte Djebel Tezah, alt. eire. 2700 m. legi spec. foliis tantum donatum! cui nomen *M. nevadense*, Boiss., non sine dubio affixi. Postea cel. Cosson benevole communicavit spec. pariter foliaceum cui scheda *Meum? atlanticum*, Coss. Fere certe in grege *M. athamantici* colloecanda, species adhuc dubia est.

FERULA (*L. Gen.* 343).

F. COMMUNIS (*L., DC. Pr.* iv. 172).

Mar. sept.—? Specimen cui scheda "*Thaspia?*" ex monte Beni Hosmar prope Tetuan exstat in herb. Webb, et hujus frustulum mihi benevole concessit cel. Parlatore. Ex caule tereti cæterisque notis planta Webbiana certe ad *F. communem* sive *F. glaucam* referenda est, priori magis conveniens. In recensione herbarii Mauritaniæ beati Webb, cel. Cosson *F. sulcatam*, Desf., ex monte Beni Hosmar enumeravit. Hujus specimina non vidi in herb. Webb.

Mar. occid.—Prope Casa Blanca vidimus canles exsiccatos *Ferulæ* ejusdem, verisimiliter *F. communis*, ad hortorum sepimenta ab ineolis adhibitos.

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

F. TINGITANA (*L., DC. Pr.* iv. 173).

Mar. sept.—Prope Tanger!

Specimina duo legi mense Junio fructu nondum maturo, umbellis centralibus pedunculis 5–6-pollicaribus suffultis. Indunt involueris et involuellis nullis, vel in altero specimine involuero 3–4-phyllis, foliolis longiusculis setaceis, et involuellis polyphyllis foliolis brevibus setaceis.

Ar. Geog. Africa borealis. Oriens.

PEUCEDANUM (*L. Gen.* 339).

P. MUNBYI (*Boiss. Diag. Pl. Or.* ser. 2, ii. 89).

Mar. sept.—Specimina valde incompleta fructu nondum maturo prope Tingidem legit beatus Schousboe die 13 Septembris 1802. Frustula plantæ parum cognitæ benevole comm. eel. Cosson!

Ar. Geog. Nonnisi in Algeriæ prov. Oran et prope Tingidem hucusque detectum. Monente eel. Boissier folia fert Martio, flores Septembri vel Octobri, fructus Decembri. In grege *P. Chabraei* collocandum.

HERACLEUM (*L. Gen.* 345).

H. SPHONDYLIIUM (*L., DC. Pr.* iv. 192).

Mar. merid.—In regione alpina Atlantis Majoris. Folia legimus in jugo Tagherot eire. 2700 m.! Specimina fructifera in monte Djebel Afougneur ab indigeno Ibrahim lecta comm. eel. Cosson!

Ar. Geog. Europa media et borealis. Species vel formæ proxime affines in montibus regionis Mediterraneæ (excluis Macaronesia et Asia Mediterranea).

CORIANDRUM (*L. Gen.* 356).

C. SATIVUM (*L., DC. Pr.* iv. 250).

Mar. sept.—Prope Tanger inter segetes!

Ar. Geog. Species regionis Mediterraneæ orientalis, ut videtur, indigena, alibi in temperatis calidioribus inquilina, deest in Macaronesia.

BIFORA (*Hoffm. Umb.* 191).

B. TESTICULATA (*L., sub Coriandro; Spreng. in Schult. Syst.* vi. 38).

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Aït Mesan eire. 1300 m.!

Ar. Geog. Regio Mediterranea (exclusa Macaronesia).

DAUCUS (*L. Gen.* 333).

D. MURICATUS (*L., DC. Pr.* iv. 210).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Tetuan! (*Webb Herb.*)!

Mar. merid.—Sektana (*Rein et Fritsch, spec. in Herb. Kew.*)!

Ar. Geog. Regio Mediterranea occidentalis. Græciæ dubia civis. Deest in Macaronesia.

D. SETULOSUS (*Guss. in DC. Pr.* iv. 211), var.?

Mar. merid.—Ad radices Atlantis Majoris juxta Tasseremont eire. 1200 m.!

Ar. Geog. Species per regionem Mediterraneam orientalem diffusa. In Africa Tunetana legit Kralik, ex Africa boreali-occidentali (nostris exceptis) specimina non vidi.

Specimina nostra pilis longis mollibus vestita ostendunt umbellam primariam centralem pedunculo brevi pollicari suffultam, ramis laterali-

hinc spithamæis superatam. Forsan melius cognita pro specie diversa habenda erit.

DAUCUS PARVIFLORUS (*Desf. Fl. Atl.* i. 241, tab 60; *DC. Pr.* iv. 211).

Mar. sept.—Tanger!

Ar. Geog. Africa boreali-occidentalis. Fertur in insulis Canariensibus, sed spec. Bourgaëana *D. parviflori*, “No. 401. Ex insula Canaria” inter formas *D. Carotæ* enumeranda sunt. Confer Lowe, *Man. Fl. Mad.* i. 368. Specimina genuina ex Europa non vidi.

Huc verisimiliter *D. mauritanicus* (Schousb. *Gew. Marok.* 116).

D. CAROTA (*L.*, *DC. Pr.* iv. 211).

Mar. occid.—In arenosis pr. Mogador!

Ar. Geog. Europa omnis (exclusa arctica). Regio omnis Mediterranea. Sibiria. Abyssinia.

Qui *D. maximum* pro specie rite diversa habebit specimina nostra forsan inter hujus formas minores enumerabit.

D. maximus (*Desf. Fl. Atl.* i. 241; *DC. Pr.* iv. 212).

Mar. sept.—Tanger (*Webb Herb.*)!

Mar. occid.—Casa Blanca! Mogador (*Lowe*). Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Frequens in regione inferiore. Legimus prope Mskala in prov. Shedma! et pluribus locis prov. Mtouga!

Ar. Geog. Regio Mediterranea fere omnis. Specimina genuina ex Macaronesia non vidi.

D. maximus nullis notis diagnosticis stabilibus a *D. Carota*, specie quam maxime polymorpha, separatur, et inter hujus subspecies sine dubio enumeravi. Primo intuitu tamen a *D. Carota* Europæ mediæ valde diversus videtur. Caulis elatus 4–5-pedalis magis ramosus est, foliorum segmenta latiora, involuerum præ magnitudine umbellæ minus. Ludit radiis umbellæ plus minusve elongatis, petalis albis vel læte rubellis, exterioribus quam cætera multo majoribus, vel ea paulo superantibus. Umbellula centralis fere semper gaudet flore uno alterove abortivo atropurpureo, vel fere nigro. Confer Boiss. *Fl. Or.* ii. 1076.

D. GINGIDIUM (*L.*, *DC. Pr.* iv. 211), = *D. Hispanicus* (*Gouan, Ill.* p. 9), = *D. gummifer* (*Lam. Dict.* i. 634), = *D. maritimus* (*With. Arr. Brit.* 290, non *Lam.*, non *Gaertn.*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Mar. occid.—Saffi!

Ar. Geog. In litore Atlantico Europæ et Africæ borealis, et eirenum mare Mediterraneum late diffusus, præsertim in rupestribus.

Cl. Grenier et Godron (*Fl. Fr.* i. 669) *D. Gingidium* et *D. gummiferum* accurate definierunt, sed, ut mihi videtur, notæ diagnosticæ fallaces sunt, dum specimina intermedia nimis frequentia transitum præbent. Immo mihi valde dubium est, an hic a *D. Carota* satis distinctus, vel

melius ad subspeciem reducendus sit. Specimina nostra Tingitana nimis immatura cum spec. genuinis Calpensibus bene congruunt.

— —, var. *HISPIDUS*, *nob.*, = *D. hispidus* (*Desf. Fl. Atl.* i. 243, tab. 63; *DC. Pr.* iv. 217; *Schousb. Gew. Marok.* 117).

Mar. occid.—In arenosis prope Mogador (*Schsb.*).

Mar. merid.?—Prov. Mtonga! Specimen dubium a *D. maximo* petalis parvulis subæqualibus, radiis umbellæ fructiferæ vix inflexis diversum, et spec. Algeriensibus *D. hispidi* congruens.

D. hispidus a *D. Gingidio* (sensu latiore) nullis notis stabilibus separatur, et mihi videtur esse hujus forma procerior, in locis a mare remotis enata.

Sicut occurrit etiam in *D. Gingidio*, flores centrales interdum abortivi sunt et fiunt atropurpurei.

DAUCUS AUREUS (*Desf. Fl. Atl.* i. 242, tab. 61).

Mar. occid.—In maritimis prope Mazagan!

Ar. Geog. Per regionem Mediterraneam fere omnem (inclusa Macaronesia) diffusus, sed potius rarus.

Specimen nostrum unicum incompletum et ideo haud omnino certum.

D. CRINITUS (*Desf. Fl. Atl.* i. 242, tab. 62).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! (Ex herb. Schousboe sub "*D. verticillatus*") *Coss. spec.*!

Mar. occid.—Casa Blanca! Agadir (*Coll. Indig.* 1871)! folia tantum sed fere certa.

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

D. PUMILUS (*Gouan, Fl. Monsp.* 285, et *Schousb. Gew. Marok.* 115, sub *Caucalide*), = *D. muricatus*, var. *maritimus* (*L. Sp. Pl.* 349), = *D. maritimus* (*Gaertn. Fruct., non Lam. nec With.*), = *Orlaya maritima* (*Koch, Umbell.* 78; *DC. Pr.* iv. 209, et plurim. auct. recentiorum).

Mar. sept.—Tanger! (*Schsb. l. c.*) Cap. Spartel! Inter Tetuan et Ceuta!

Mar. occid.—Casa Blanca! Mazagan! Mogador!

Ar. Geog. Regio omnis Mediterranea in maritimis.

Ducibus celeb. Benth. et H. f. hanc speciem cum affinibus (*Orlaya*, Koch) ad *Daucum* revocavi.

CAUCALIS (*L. Gen.* 331; *Benth. et H. f. Gen. Pl.* i. 928).

C. LATIFOLIA (*L.*, sub *Tordyllo*; *Sm. Eng. Fl.* ii. 41), = *Turgenia latifolia* (*Hoffm. Umbell.* 59; *DC. Pr.* iv. 218).

Mar. merid.—Videtur satis frequens in convallibus inferioribus et ad radices Atlantis Majoris. Mesfioua! Inter Tasseremout et Ourika! Gurguri! In convalle Aït Mesan circ. 1400 m.!

Ar. Geog. Europa media. Asia temperata. Regio omnis Mediterranea (exclusa Macaronesia).

C. DAUCOIDES (*L.*, *DC. Pr.* iv. 216).

Mar. merid.—In convalle Amsmiz circ. 1500 m.!

Ar. Geog. Europa media et australis. Oriens. Ex Africa non vidi.

CAUCALIS LEPTOPHYLLA (L., DC. Pr. iv. 216).

Mar. merid.—Frequens in planitie et in regione media et inferiore Atlantidis Majoris. Prov. Shedma! Prope Ourika! Seksaoua! Reraya! In convalle Aït Mesan a 1200 m. usque 2200 m.! et in convalle Am-smiz a 1000 m. usque 1800 m.!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). In Europa media hinc inde introducta.

— —, var. HETEROCARPA, nob. Meriearpio exteriori longe aculeato sterili, interiore breviter muricato fertili.

Mar. occid.—Casa Blanea! Agadir (Coll. Indig. 1871)!

Eodem modo variat etiam *C. nodosa*. Confer DC. Prod. iv. 219. An ab hac satis differt *C. bifrons*, Coss. et Dur.?

C. cærulescens (Boiss. Elench. 97; Walp. Repert. ii 420).

Mar. merid.—In convalle Aït Mesan circ. 1300 m.! Prope Tasseremout (specimina nimis immatura)! Distr. Reraya!

Ar. Geog. Hispania Granatensis. Marocco meridionalis.

Specimina ex Reraya intermedia sunt. Forsan melius pro varietate *C. leptophyllæ* habenda.

C. NODOSA (L., sub *Tordyllo*; Huds. Fl. Angl. 114; Schousb. Gew. Marok. 116), = *Torilis nodosa* (Gaertn. Fruct. i. 82; DC. Pr. iv. 219).

Mar. sept.—Tanger (Schsb. l. c.). Cap. Spartel! Ex collibus inter Tanger et Tetuan! Prope Tetuan (1851)!

Mar. occid.—Ex prov. Haha prope Mogador!

Lusum umbellis pedunculatis prope Tanger legit J. D. H.!

— —, var. HETEROCARPA, nob.

Mar. occid.—Casa Blanea!

Ar. Geog. Species per Europam mediam et regionem omnem Mediterraneam late diffusa et indigena, in Americam australem aliasque terras longinquas introducta est. Var. *heterocarpa* cum specie hinc inde obvia est.

C. INFESTA (L., sub *Scandice*; Curt. Fl. Lond. vi. tab. 23), = *Torilis infesta* (Hoffm. Umb. 89; Boiss. Fl. Or. ii. 1082), = *Caucalis helvetica* (Jacq. H. Vind. iii. tab. 16), = *Torilis helvetica* (Gmel. Fl. Bad. i. 617; DC. Pr. iv. 219).

Mar. merid.—In hortis æe ruderatis urbis Marocco! Prope Mzouda (J. D. H.)!

Ar. Geog. Europa media. Regio omnis Mediterranea. Specimina ex Macaronesia quæ vidi, ad var. sequentem referenda sunt.

— —, var. NEGLECTA, = *Torilis neglecta* (Schultes, Syst. ii. 484; DC. Pr. iv. 218).

Mar. sept.—Legi florentem et fructiferam prope Tanger die 13 Junio 1871!

Hanc pro specie distincta habuerunt summi botanici A. P. De Candolle,

G. D. Koch et Boissier, dum accuratus Bertoloni (Fl. It. iii. 188) pro synonymo *C. infestæ* enumeravit. Distinguitur etenim statura majore, foliorum segmentis latioribus, petalis radiantibus majoribus, fructu majore, et stylis stylopodio pluries longioribus, nec paulo excedentibus. Hæ notæ omnes tamen, præsertim in Italia superiore, summo-pere variabiles, et formæ intermediæ sæpe obviæ sunt. Ut morem gererem magistris, plantam Tingitanam foliis et habitu *C. infestæ* simillimam, sed petalis exterioribus majusculis, fructu majore et stylis longis insignem, pro varietate *C. infestæ* enumeravi. Specimen unum habitu graciliore umbellis longe pedunculatis pauciradiatis ad sequentem accedit.

Caucalis heterophylla (Guss. Prod. Fl. Sic. i. 326 ; DC. Pr. iv. 219).

Mar. sept.—? Tanger (Webb Herb.)!

Ar. Geog. Hinc inde in regione Mediterranea.

Specimen Webbianum subdubium et, ut mihi videtur, inter varietates *C. infestæ* enumerandum. Cæterum sp. Gussoniana *C. infestæ* nimis affinis et pro subspecie enumeranda est.

THAPSIA (L. Gen. 361).

T. GARGANICA (L., DC. Pr. iv. 202 ; Schousb. Gew. Marok. 124).

Mar. sept.—Tanger! (Salzm. Exsicc.)! (Webb Herb.)! (Schsb. l. c.).

Inter Tanger et Tetuan!

Mar. occid.—Casa Blanca! Mogador (Schsb. l. c.).

Ar. Geog. Regio Mediterranea calidior. Deest in Gallia, in Italia media et boreali, et in Macaronesia.

T. VILLOSA (L., DC. Pr. iv. 202).

Mar. sept.—In monte Beni Hosmar prope Tetuan!

Mar. merid.—Ad radices Atlantis Majoris sat frequens. Ourika! Reraya! Sektana! et alibi visa!

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis. Gallia Mediterranea.

Specimen ex monte Beni Hosmar foliis radicalibus tantum donatum, sed ob nervorum dispositionem singularem satis certum!

ELÆOSELINUM (Koch in DC. Pr. iv. 215).

E. MEOIDES (Desf. Fl. Atl. i. 253, tab. 69, sub *Laserpitio*; Koch, l. c.), =Thapsia meoides (Guss. Prod. Fl. Sic. i. 370).

Mar. sept.—Tanger! (Salzm. Exsicc. sub "*Laserpitium meoides*")! (Ex herb. Schoushoe sub "*Ferula meoides*"), Coss. spec.! (Webb Herb.)!

Mar. merid.—Mesfoua! Mzouda! Aït Mesan circ. 1400 m.!

Ar. Geog. Regio Mediterranea occidentalis. Sardinia. Sicilia. Hispania. Africa borealis.

E. humile, nob.

Proximum *E. meoidi* et verisimiliter sicut subspecies enumerandum,

differt tamen notis insignibus. Statura multo minor; caules 6-12-pollicares; folia fere omnia radicalia subspithamæa, caulina ad vaginas amplas sæpissime reducta; umbella subexinvolucrata, foliolis 1-2 cito deciduis; involuella ex foliolis parvis setaceis deciduis. Styli florum fertilium, tam in hoc quam in *E. meoide*, reeti divergentes demum recurvi.

Mar. merid.—In aridis saxosis regionis inferioris. In prov. Shedma prope Aïn Oumast et alibi!

ELÆOSELINUM EXINVOLUCRATUM (*Coss. in Bull. Soc. Bot. Fr. xx. 247*). Sequitur descriptio Cossoniana:—

“Planta perennis caudice crassiusculo fruticescente descendente superne fibrillis petiolorum emarcidorum vestigiis *dense vestito*, caulem unicum emittente. Caulis ut tota planta glaber, glaucescens, gracilis, striatus, superne ramosus, paucifolius. *Folia* glauca, pleraque radicalia, ambitu triangularia, petiolata, petiolo inferne in vaginam margine membranaceam dilatato, *tripinnatisecta* partitionibus petiolulatis, *lobis* crassiusculis brevibus *ovatis*, *inferioribus* saltem *pinnatilobis lobulis abbreviatis obtusis*, caulina inferiora pauca subconformia petiolo in vaginam amplam dilatato, superiora sæpe ad vaginam ovatam reducta. *Umbellæ* longe pedunculatæ, 5-6-*radiatæ* radiis subæqualibus gracilibus teretibus lævibus, omnes fertiles in umbellula quaque permixtis floribus sterilibus pluribus. *Involucrum nullum. Involucella oligophylla, foliolis* linearibus inæqualibus *citissime deciduis*. Umbellulæ 7-9-floræ, pedicellis subæqualibus, demum flore longioribus. Calyx breviter 5-dentatus, dentibus latiusculis. *Petala lutea, integra, ovata, in lacinulam* linearem *introflexam acuminata*. Stylopodia crassa, depressa. *Styli* post anthesin divergentes, *latitudini stylopodiorum subæquilongi*. Ovarium læve, glabrum, campanulato-obconicum, jugis haud distinctis.”

Mar. merid.—Prope Keira, alt. circ. 1000 m., legit Balansa anno 1867. Caudicem unicum foliis radicalibus donatum legi supra Seksaoua, et specimen manens fructiferum ex rupibus præruptis avulsi!

Fructus maturus egregio auctori ineognitus major est quam in sp. affinis. Mericarpiæ sæpissime inæqualia, alterum alis minus evolutis præditum. Alæ perfectæ latæ purpurascens nitentes late expansæ, utrinque truncatæ, superne angulum obtusum, basi angulum acutum formantes. Stylopodium ex angulo obtuso inter alas exsertum; umbella primaria in spec. nostro decemradiata, laterales (floribus omnibus abortivis) 12-radiatæ.

Hæc species, cui proxima *E. Lagascæ*, Boiss. (*Thapsia tenuifolia*, Lag.), differt caudice fibrillis dense vestito, foliis glaucis crassioribus, umbellis pauciradiatis, involuero nullo, nec polyphyllis, involuellis oligophyllis cito deciduis, nec polyphyllis persistentibus, stylis denique brevioribus.

ELÆOSELINUM FÆTIDUM (*L., et DC. Pr. iv. 203, sub Thapsia; Boiss. Elench. 91*).

Mar. sept.—Tanger (*Webb Herb.*)! (*Salzm. Exsicc.*, sub “*Thapsia polygama*”)! (*Herb. Schousboe*, sub “*T. fætida*”), *Coss. spec.*! Prope Tetuan!

Ar. Geog. Hispania. Africa borealis.

In convalle Aït Mesan, alt. circ. 1400 m. legit ecl. J. D. Hooker folia *Umbelliferæ* ejusdam omnino incertæ, *Physospermum aquilegifolium*, Kochi referentia, sed ab hoc certe diversa. Planta ulterius inquirenda.

ARALIACEÆ (*Juss. Diet. ii. 348*).

HEDERA (*L. Gen. 283, ex parte*).

II. HELIX (*L., DC. Pr. iv. 261*).

Mar. sept.—Prope Tanger in monte Djebel Kebir! In monte Beni Hosmar prope Tetuan!

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Aït Mesan!

Ar. Geog. Europa media. Asia temperata. Japonia. Regio Mediterranea fere omnis. Var. *canariensis* in Macaronesia.

CAPRIFOLIACEÆ (*Juss. Gen. 110*).

SAMBUCUS (*L. Gen. 372*).

S. EBULUS (*L., DC. Pr. iv. 322; Schsb. Gew. Marok. 133*).

Mar. sept.—Tanger (*Schsb. l. c.*).

Ar. Geog. Europa temperata. Regio fere omnis Mediterranea, inclusa Madeira.

S. NIGRA (*L., DC. Pr. iv. 322*).

Mar. merid.—In convalle Aït Mesan a 1500 m. usque 1800 m. ! Visa etiam in convalle Amsmiz!

Ar. Geog. Europa media. Montes Europæ australis et Asiæ Mediterraneæ. Ex Africa non vidi. Deest in Macaronesia.

VIBURNUM (*L. Gen. 370*).

V. TINUS (*L., DC. Pr. iv. 324; Schousb. Gew. Marok. 130*).

Mar. merid.—In convalle Amsmiz circ. 1400 m. !

Ar. Geog. Regio Mediterranea fere omnis. In Oriente rarissime occurrit. In Macaronesia adest tantum sp. affinis *V. rugosum*, Pers.

— —. Var. corymbis laxioribus, floribus majoribus, foliis utrinque glabris superne lucidis. An idem ac *V. lucidum*, Mill.? Huc ducenda forma a Schousboe enumerata?

Mar. sept.—In monte Djebel Kebir, et alibi prope Tanger!

LONICERA (*L. Gen. 233*).

L. ETRUSCA (*Santi, Viagg. i. 113, tab. 1; DC. Pr. iv. 331*), = *L. caprifolium* (*Schousb. Gew. Marok. 73, non L.*).

Mar. sept.—“In sepibus prope Tingidem raro” (*Schsb.*).

Mar. merid.—In convalle Aït Mesan 1200–1400 m.!

In planta Atlantica corollæ tubus glandulis breviter stipitatis plus minus adpersus, folia præsertim juniora (etiam in pagina superiore) pilosa, caules subglaberrimi. Flores ex roseo lutescentes, odoratissimi.

Ar. Geog. Regio Mediterranea fere omnis. Floræ Maderensis civis, consideratur in ins. Canariensibus.

LONICERA IMPLEXA (*Ait. H. Kew.* i. 231; *DC. Pr.* iv. 331).

Mar. sept.—Tanger in monte Djebel Kebir et alibi!

Mar. occid.—Haud procul Mogador versus Aïn el Hadjar frustulum legit J. D. H.!

Mar. merid.—Prope Ourika ad radices Atlantis Majoris! Reraya (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Europa australis. Africa boreali-occidentalis. Bithynia (*sec. Boissier*). Legi pr. Smyrnam.

L. PERICLYMENUM (*L., DC. Pr.* iv. 331), var. *HISPANICA*, = *L. hispanica* (*Boiss. et Reut. Pug.* 52; *Walp. Ann.* v. 95), = *L. Periclymenum* (*Schousb. Gew. Marok.* 74).

Mar. sept.—Tanger in monte Djebel Kebir et alibi! (*Webb Herb.*)! (*Schsb. l. c.*). In sepibus prope Tetuan!

Ar. Geog. Species in Europa media et Scandinavia meridionali obvia. In Europa australi infrequens. Fertur in Græcia, sed ex Oriente spec. nulla vidit Boissier. In Africa boreali-occidentali rarius occurrit.

Notæ quibus distinguitur *L. hispanica* mihi videntur omnino fallaces. Indumentum in spec. nostris Tetuanensibus paræ et vix glandulosum. Folia variant latiora vel angustiora utrinque acuta, omnia petiolata vel suprema subsessilia. Dentes calycini aliquantulum longiores quam in spec. normalibus *L. Periclymeni*.

L. BIFLORA (*Desf. Fl. Atl.* i. 184, tab. 52), = *L. canescens* (*Schousb. Gew. Mar.* 74; *DC. Pr.* iv. 334).

Mar. occid.—Aïn el Hadjar haud procul Mogador! Prope Mogador (*Schsb. l. c.*). Pr. Sallé (*Broussonnet ex Cavan. Anal. Cienc. Nat.*).

Mar. merid.—Sheshiaoua! In umbrosis et hortis ad urbem Marocco! Reraya!

Ar. Geog. Species Maroecana et Algeriensis. Fertur etiam in Sicilia (*fide Gussone*).

Nomen Fontanesianum fere eodem tempore publici juris factum ac Schonsbocanum. Ipse Schousboeus tamen *L. bifloram* sicut synonymon *L. canescentis* citat, et prioritatem agnovit.

RUBIACEÆ (*Juss. Gen.* 196).

PUTORIA (*Pers. Syn.* i. 524).

P. CALABRICA (*L. fil., Suppl.* 120, sub *Asperula*; *Pers. l. c.*; *DC. Pr.* iv. 577).

Mar. sept.—Tetuan (1851)!

Mar. merid.—In convalle Aït Mesan 1200–1400 m.!

Ar. Geog. Regio Mediterranea calidior (exclusa Macaronesia). Ultra latit. gradum 40^{mum} vix se protendit nisi in litore Dalmatico et litoralis Austriaci, ubi usque Fiume obvia est.

Folia et corollæ tubus longitudine valde variabilia sunt. *P. Hispanica*, Boiss. Diagn. ser. 2, ii. 108, sistit formam foliis angustioribus longioribus, corollæ tubo elongato. Hujus specimina juxta Arundam (Ronda) et in montibus Sierra Nevada a me lecta videntur a typo satis diversa, sed formas intermedias vidi et legi prope Malaga. Specimina Oranensia ad var. *hispanicam* referenda sunt. Contra quæ legimus in Atlante Majore folia quam in typo breviora ostendunt.

CALLIPETIS (*Stev. Obs. Pl. Ross.* 69).

C. CUCULLARIA (L., sub *Valantia*; *DC. Pr.* iv. 613).

Mar. merid.—Prope Amsmiz! In convalle Amsmiz usque 1500 m.! Seksaoua!

Ar. Geog. Per regionem Mediterraneam calidiorem et siccioram ex Afghanistan usque territ. Maroccanum. In Europa nominis in Hispania centrali hucusque visa.

VAILLANTIA (*DC. Fl. Fr.* iv. 266), = *Valantia* (L., ex parte).

V. MURALIS (L., sub *Valantia*; *DC. l. c., Prod.* iv. 614).

Mar. sept.—In saxosis prope Tetuan! et in monte Beni Hosmar!

In spec. nostris caules et folia juniora ciliata.

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

RUBIA (*L. Gen.* 127).

R. TINCTORUM (L., *DC. Pr.* iv. 589; *Schousb. Gew. Marok.* 46).

Mar. merid.—In convalle Aït Mesan supra pagum Arround ultra 2000 m. s. m., verisimiliter introducta!

Sec. Schsb. l. c. frequens in sepibus et vepretis totius Imperii.

Species verisimiliter in regione Mediterranea indigena, diu culta et sæpe subspontanea facta. Imperii Maroccani dubia civis. A *R. peregrina*, ut mihi videtur, probe diversa.

R. PEREGRINA (L., *DC. Pr.* iv. 589), = *R. peregrina* β intermedia (*Gren. et Godr. Fl. Fr.* ii. 13).

Mar. sept.—Tanger! Inter Tanger et Tetuan! In rupestribus pr. Tetuan!

Mar. merid.—Ourika! In convalle Aït Mesan!

— —, var. *LUCIDA* (*Webb Phyt. Can.* ii. 187), = *R. lucida* (L., *DC. Pr.* iv. 590; *Schousb. Gew. Mar.* 46).

Mar. sept.—Prope Tanger in elivis maritimis!

Mar. occid.—Saffi (*Schsb. l. c.*).

Foliis in verticillo plerumque senis, quam in typo latioribus, apice rotundatis, caule lævi, videtur satis diversum; sed adsunt formæ inter-

mediæ, et duobus Cambessèdes, Webb, Grenier et Godron, hanc et sequentem ad varietates reduxi.

— —, var. *ANGUSTIFOLIA* (*Webb, l. c.*), = *R. angustifolia* (*L., DC. Pr. iv. 589*), = *R. longifolia* (*Poir. Suppl. ii. 703; DC. Pr. l. c.*), = *R. peregrina* (*Schousb. Gew. Mar. 46*), = *R. Requierii* (*Duby, Bot. Gall. i. 247*).

Mar. sept.—Prope Tetuan (*Webb Herb.*)!

Mar. occid.—Mogador (*Lowe*).

Mar. merid.—Shieshaoua! In convalle Aït Mesan circ. 1200 m.! Sektana (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Species Europam occidentalem ab Hibernia usque Fretum Herculis et regionem omnem Mediterraneam (inclusa Macaronesia) incolit. Fugit terras continentales. Varietates plurimæ per regionem Mediterraneam obviæ sunt.

R. fruticosa, species propria Canariensis fertur inter plantas Mogadorense a Broussonnet lectas, ex indicatione loci natalis erronea.

GALIUM (*L. Gen. 125*).

G. ELLIPTICUM (*Willd. Enum. Suppl. 1813; DC. Pr. iv. 599*).

Mar. sept.—? In monte Beni Hosmar prope Tetuan legimus frustula pl. nondum florentis, fere certe huc referenda!

Ar. Geog. Regio Mediterranea occidentalis (inclusa Macaronesia). Confer *Lowe, Man. Fl. Mad. pp. 388 et seq.*

G. POIRETIANUM, *nob.*, = *Rubia lævis* (*Poir. Voy. ii. 111; DC. Pr. iv. 589, non Thunb.*).

Mar. merid.—Ad radices Atlantis Majoris sat frequens. Mesfioua! Reraya! Aït Mesan! Mzouda!

Ar. Geog. Africa boreali-occidentalis.

Hanc speciem ad *Galium* referendam esse jam suspicatus est ill. A. P. De Candolle, nec video quo pacto a subgenere *Eugali*um, § *Leiogali*um divellenda sit. A *Rubiis* corolla quadrifida, fructus indumento aliquanto coriaceo, nec carnoso, recedit: *Leiogaliis* inflorescentia paniculata multiflora, foliis a basi sensim decreescentibus, et habitu toto convenit, *G. lævigato*, *L.*, proxime collocandum.

G. ERECTUM (*Huds. Fl. Angl. 68; DC. Pr. iv. 595*), var. *RIGIDUM*, = *G. rigidum* (*Vill. Dauph. ii. 319*).

Mar. merid.—Specimen unicum legi in rupibus supra Milhain!

Ar. Geog. Europa media. Regio Mediterranea fere omnis. Deest in ins. Canariensibus. Species affinis (*G. productum*, *Lowe*) in Madeira frequens.

G. CORRUDÆFOLIUM (*Vill. Dauph. ii. 320*), = *G. lucidum* (*All. Fl. Ped. No. 21, tab. 77. fig. 2*).

Mar. merid.—In monte Djebel Tezah circ. 2200 m. vix florens! Specimina florentia ex Atlante Majore benevole dedit cel. Cosson.

Ar. Geog. Species per Galliam australem, Hispaniam, valles calidiores Alpium, et totam Italiam vulgaris. Sec. Boissier deest in Oriente, sed adsunt plurimæ species affines. In hoc grege ut in eo *G. sylvestris* limites specierum vix notis stabilibus definiendi.

Galium aureum (*Vis. Ind. Ort. Padov.* 1842; *Boiss. Fl. Or.* iii. 61), = *G. melanantherum*, var. *latifolium* (*Boiss. Diagn. Pl. Or.* ser. 1, vi. 69). Mar. merid.—In rupibus supra Seksaoua!

Ar. Geog. In Oriente (sec. Boissier) late diffusum, et in Dalmatia obvium. Verisimiliter provenit in Algeria et alibi in Africa boreali.

Specimina majora nostra pl. Orientali omnino convenire videntur. Alia minora in monte supra Seksaoua lecta propiora videntur formæ nanæ *G. corrudæfolii*. Pace cel. Boiss. qui hanc speciem inter *Chromogalia* *G. vero* proxime collocavit, mihi videtur valde affinis *G. corrudæfolio*, et sicut hujus subspecies enumeranda.

G. SYLVESTRE (*Poll. Palat.* i. 151; *Koch, Syn. Fl. Germ.* i. 286), = *G. Bocconi* (*DC. Pr.* iv. 594, non *All.*), var. *ATLANTICUM*, nob., = *G. atlanticum* (*Ball in Journ. Bot.* 1873, 334).

Differt imprimis foliis surculorum steriliū et caulinis inferioribus abbreviatis obovato-acuminatis longiuscule mucronatis, interdum in pagina superiore scabris margine aculeolatis. Corolla parva gaudet segmentis acutis; antheræ ochroleucæ. Panicula cymiformis trichotomo-cymosa. Planta pumila, 3-4-pollicaris, laxe cæspitosa.

Mar. merid.—In regione superiore Atlantis Majoris. Legi in jugo Tagherot usque 2700 m.!

Exemplaria nostra fructu carent et ideo incerta sunt. Habui pro subspecie *G. sylvestris*; hodie non sine dubio sicut varietatem enumeravi.

Ar. Geog. Europa fere omnis, exclusa arctica. In montibus Hispaniæ et Italiæ. Spec. Africana non vidi.

G. ACUMINATUM (*Ball in Journ. Bot.* 1873, 334). (Plate XXI.)

Radix lignosa, e collo ramosissima; caules ad angulos ex setis brevibus patentibus vel subretrorsis scabri; folia sena, lineari-lanceolata, nitida, subcoriacea, nervo medio basin versus conspicuo, margine aculeata, in mucronem longum acuminata; panicula racemiformis, scil. rami laterales brevissimi trichotomi, interdum ad pedicellos 3 reducti, ex axillis foliorum superiorum 2-3 enati; corolla ochroleuca, segmentis obscure trinerviis, in vivo apiculatis; fructus (immaturus) demum granulatus.

Mar. merid.—In regione media et inferiore Atlantis Majoris. Distr. Reraya! In convalle Aït Mesan! a 1200 m. ad 2100 m. Habui a cel. Cosson spec. ab indigeno Ibrahim anno 1873 in monte Djebel Ouensa lecta!

Species ab omnibus mihi cognitis probe distincta. Spec. exsiccata *G. pulvinato*, Boiss., subsimilia sunt; sed inflorescentia et foliis longe mucronatis omnino diversa. Propius crit, ut videtur, *G. brunneum*.

Munby, nisi forsan vera affinitas in sectione *Euaparine*, DC., ejus species omnes annuæ, quærenda sit. In aprieis planta nostra fit contracta, foliis approximatis, caulibus 2-4-pollicaribus; in umbrosis occurrit habitu laxiore, caulibus usque pedem longis, foliis obverse lanceolatis, verticillis remotiusculis. Herba obscure virens, exsiccata nigrescit.

GALIUM CONCATENATUM (Coss. *Pl. Crit.* 38; *Walp. Ann.* iii. 901).

Mar. sept.—Tanger. Ex herb. Schousboe sub "*G. conglobatum*," spec. benevole comm. cel. Cosson!

Ar. Geog. Species hucusque nonnisi in utroque latere Freti Hereulis visa.

G. BOURGÆANUM (Coss. in *Bull. Soc. Bot. Fr.*), var. *MAROCCANUM* (Ball in *Journ. Bot.* 1873, 335). (Plate XXII.)

Radix lignosa, elongata, inter fissuras rupium serpens, caules numerosos pendentes vel prostratos, undique hirtos, emittens; folia inferiora obtusa, sequentia elliptico- vel obovato-lanceolata, in verticillo 6-9, superiora apice mucronata, omnia hirta, vel suprema glabrescentia tenuiter ciliolata; panicula multiflora, ramosa, pyramidata; corollæ segmenta albovirentia, in sicco nigrescentia insigniter trinervia, acuminata, extus pilosa; fructus glaber.

Mar. merid.—Sat frequens in rupibus prov. meridionalium Imper. Marocani, præsertim ad radices Atlantis Majoris. Prope Seksaoua! Milhaïn! Mtonga! Mskala!

Galium Bourgeanum, Coss., est species mihi non satis nota. Vidi tantum specimina "ex rupibus prope Tlemcen," in Bourgeau 'Pl. d'Algérie,' 1856, No. 107. Planta nostra differt corolla virescente, insigniter trinervi, nec alba subnervi.

A proximis *G. viridifloro*, Boiss., et *G. maritimo*, L., differt fructu glabro, nec hispido, cæterisque notis.

Ar. Geog. Africa boreali-occidentalis.

G. NOLI-TANGERE (Ball in *Journ. Bot.* 1873, 335).

Radix lignosa; caules herbaeci, cum foliis et pedicellis fragillimi, pilis rigidiusculis basi callosis vestiti; folia cruciata, ovalia, utrinque acuta, brevissime mucronata; inflorescentia depauperata; pedicelli pauci trichotomi terminales et axillares in verticillis superioribus; corolla alba; ovarium apice emarginatum papilloso-setosum.

Mar. merid.—In regione media Atlantis Majoris. In convalle Amsmiz a 1500 m. ad 1600 m.!

Proximum est sed probe distinctum *G. musciforme*, Boiss. *Diagn. Pl. Or.*; differt a nostro imprimis corolla rubra, pedunculis bracteatis, supremis solitariis, inferioribus a 3- ad 7-floris.

G. TUNETANUM (Lam. *Dict.* ii. 583).

Mar. merid.—Distr. Mesfoua versus Ourika! In convalle Aït Mesan eire. 1400 m.!

Ar. Geog. Africa boreali-occidentalis.

GALIUM GLOMERATUM (*Desf. Fl. Atl.* i. 128, tab 40; *DC. Pr.* iv. 606).
 Mar. sept.—Tanger (*Salzm. Exsicc.*)! Spec. ex herb. Schousboe comm.
 Cosson!

Mar. occid.—Casa Blanca!

Ar. Geog. Hispania australis. Africa boreali-occidentalis.

— — —, var. **CAMPESTRE**, *nob.*, = *G. campestre* (*Schousb. in Willd. Enum.* i. 152; *DC. Pr.* iv. 606).

Mar. sept.—Tanger in monte Djebel Kebir! In saxosis prope Tetuan!
Ar. Geog. Hispania australis. Africa boreali-occidentalis.

G. campestre (Duby, Bot. 248), = *G. corsicum* (Spreng. Syst. Veg. iv. pars 2, 39), est species aliena *G. rubro* proxime affinis.

Var. nostra a *G. glomerato* differt foliis inferioribus latioribus, sæpe quaternis, et corollæ lobis minus acutis; sed hæ notæ variabiles sunt, et specimina intermedia non desunt.

G. DIVARICATUM (*Lam. Dict.* ii. 580; *DC. Pr.* iv. 607).

Mar. sept.—In monte Djebel Kebir prope Tanger! Ex eodem loco spec. ex herb. Schsb. comm. Cosson!

Ar. Geog. Regio Mediterranea fere omnis. Gallia media. Adest in ins. Azoricis, sed hucusque in Macaronesia desideratur.

G. PARIENSE (*L., Sp. Pl.* 157; *Boiss. Fl. Or.* iii. 73), var. **LASIOCARPUM**, *Tausch*, = *G. litigiosum* (*DC. Fl. Fr.* iv. 263, et *Prod.* iv. 609).

Mar. merid.—Frequens ad radices Atlantis Majoris, unde per regionem inferiorem usque ad reg. subalpinam se protendit. Mesfioua! Ourika! Reraya! Seksaoua! Aïn Tarsil! In convalle Aït Mesan a 1200 m. usque 2200 m.! In monte Djebel Tezali usque 2500 m.! In convalle Amsmiz circ. 1600 m.!

Ar. Geog. Europa media. Regio omnis Mediterranea (inclusa Macaronesia).

Folia in spec. ex regione montana plerumque latiora.

G. SETACEUM (*Lam. Dict.* ii. 584; *DC. Pr.* iv. 609), = *G. capillare* (*Cav. Icon.* tab. 191, fig. 1), = *G. microcarpum* (*Vahl, Symb.* ii. 30), = *G. Urvillei* (*Req. in DC. Pr.* iv. 607), = *G. Sibthorpii* (*R. et Sch. Syst.* iii. 244).

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Sat frequens in saxosis calidioribus. Ex rupibus prope urbem Marocco! Tasseremout! Seksaoua! Mskala! Aïn Tarsil!

Ar. Geog. Regio Mediterranea fere tota (inclusa Macaronesia). Deest, ut videtur, in Lusitania et in Italia continentali.

Fructus ludit setis longiusculis patulis, vel paucis brevissimis, vel subnullis.

G. SACCHARATUM (*All. Fl. Ped.* No. 39; *DC. Pr.* iv. 607).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Prope Cap. Spartel! In monte Beni Hosmar prope Tetuan!

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Seksaoua! Tasseremout! Amsmiz!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia): in Oriente rarius occurrit. In Europa media hinc inde obvium, verisimiliter cum seminibus introductum, hodie spontaneum.

GALIMUM TRICORNE (*With. Brit. Pl.* ed. 2, i. 153; *DC. Pr.* iv. 608).

Mar. merid.—In convalle Aït Mesan circ. 2100 m.! In convalle Amsmiz alt. 1500–1600 m.!

Ar. Geog. Europa media. Regio Mediterranea imprimis orientalis. In ins. Madeira rarum, hucusque in ins. Canariensibus desideratur.

G. APARINE (*L., DC. Pr.* iv. 608; *Schousb. Gew. Marok.* 45).

Mar. occid.—In prov. Haha (verisimiliter pr. Mogador) *Schsb. l. c.*

Mar. merid.—In umbrosis ad urbem Marocco!

Ar. Geog. Europa fere omnis. Sibiria. Regio omnis Mediterranea (inclusa Macaronesia). In America introductum.

G. spurium (*L., DC. Pr.* iv. 608), var. *VAILLANTII* (*Gren. et Godr. Fl. Fr.* ii. 44), = *G. Aparine*, var. β (*Lam. Dict.* ii. 581; *Schousb. Gew. Mar.* 45), = *G. Aparine*, var. *Vaillantii* (*Koch, Syn. Fl. Germ.*), = *G. Vaillantii* (*DC. Pr. l. c.*).

Mar. occid.—“In aggeribus umbrosis prov. Haha” (*Schsb. l. c.*) verisimiliter in vicinitate urbis Mogador.

Mar. merid.—In regione media Atlantis Majoris. In convalle Aït Mesan a 1500 m. usque 2500 m.! In monte Djebel Tezah circ. 2200 m.!

Ar. Geog. Species habitat Europam mediam atque Asiam temperatam; in Scandinavia et in Europa australi rarius occurrit. In Oriente et in Africa boreali et in Macaronesia var. *Vaillantii* obvia est.

G. spurium, *L.*, *G. Aparini* arcte affine et nexu genetico conjunctum, sensu nostro pro subspecie habendum est; differt tamen notis satis stabilibus, et imprimis fructu quadruplo minore. Cultum et e seminibus reproductum vix variat.

Spec. nostra Atlantica referunt formam debilem—*G. spurium*, var. γ , *tenerum* (*Gren. et Godr. l. c.*), cujus synonymon est (sec. ecl. Boiss.) *G. aparinoides* (*C. Koch, Linn. xvii.* 38)—vix, ut mihi videtur, sicut varietas diversa distinguenda.

G. MURALE (*L.*, sub *Sherardia*; *All. Fl. Ped.* i. 8, tab. 77. fig. 1; *DC. Pr.* iv. 610).

Mar. sept.—Prope Cap. Spartel!

Mar. merid.—In prov. Shedna! In convalle Amsmiz circ. 1500 m.!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

ASPERULA (*L. Gen.* 121).

A. ARISTATA (*L., DC. Pr.* iv. 583).

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Ourika!

— —, var. *SCABRA*, = *A. scabra* (*Presl, Del. Prag.* 124; *DC. Pr.* iv. 584, non *Link*).

Mar. merid.—In convalle Amsmiz circ. 1500 m. ! In rupibus supra Seksaoua !

Ar. Geog. Verisimiliter per regionem Mediterraneam occidentalem diffusa (synonyma plurima valde dubia). Deest in Macaronesia.

Asperulæ hujus gregis quam maxime ludibundæ sunt, et characteres stabiles Botanici adhuc frustra quæsierunt. Duce cel. Visiani plurimas s. d. species sicut varietates *A. cyanchicæ*, L., enumerare mallet ; *A. aristata* aliæque nonnullæ forsân subspecies sistunt, sed res ardua est et studio acenrato eget.

ASPERULA HIRSUTA (*Desf. Fl. Atl.* i. 127 ; *DC. Pr.* iv. 583), = *A. repens* (*Brot. Phyt. Lus.* tab. 10 ; *Schousb. MSS. in Herb.*), = *A. algerica* (*Pers. Syn.* i. 124).

Mar. sept.—Tanger (*Schsb. Herb.*) ! Tetuan (1851) ! (*Webb Herb.*) !

Mar. occid.—In monte Djebel Hadid haud procul Mogador (*R. et Fr. spec. in Herb. Kew.*) !

Mar. merid.—Stirps speciosa, sat frequens ad radices Atlantis Majoris. Mesfioua ! Tasseremout ! Ourika ! Sektana ! Seksaoua ! Rarius occurrit in convallibus interioribus. In convalle Aït Mesan !

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

Hæc species valde ludibunda est ; variat scil. superne fere glabra plus minusve scabra, corollæ læte roseæ vel pallidæ vel sordidæ tubo longiore vel brevior, caulibus simplicibus vel ramosis, capitulis terminalibus multifloris seu panicifloris.

Asperula, sp. ?

In monte Beni Hosmar prope Tetuan legimus specimen immaturum nondum florens foliis omnibus longe et dense hirtis, verisimiliter ad *Asperulam* referendum, sed ab *A. hirsuta* fere certe diversum.

CRUCIANELLA (*L. Gen.* 126).

C. ANGUSTIFOLIA (*L., DC. Pr.* iv. 586).

Mar. sept.—Tetuan ! (*Webb Herb.*) !

Mar. occid.—Casa Blanca !

Mar. merid.—Prov. Shedma ! Ourika ! Seksaoua ! In convalle Aït Mesan circ. 1300 m. !

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia). In Gallia media sat frequens.

C. MARITIMA (*L., DC. Pr.* iv. 587 ; *Schousb. Gew. Mar.* 45).

Mar. sept.—Tanger ! (*Webb Herb.*) ! (*Schsb. Herb.*), *spec. comm. Cosson* ! (*Salzm. Exsicc.*) ! A Cap. Spartel meridiem versus !

Mar. occid.—Saffi !

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

SHERARDIA (*L. Gen.* 120).

S. ARVENSIS (*L., DC. Pr.* iv. 581; *Schousb. Gew. Marok.* 44).

Mar. sept.—Tanger! Cap. Spartel! Beni Hosmar pr. Tetuan!

Mar. merid.—In cultis non infrequens. Amsmiz! Mesfioua! In convalle Aït Mesan circ. 1300 m.!

Ar. Geog. Europa omnis (exclusa arctica). Regio omnis Mediterranea inclusa Macaronesia.

VALERIANEÆ (*DC. Fl. Fr.* iv. 237).VALERIANA (*L. Gen.* 44).

V. TUBEROSA (*L., DC. Pr.* iv. 637).

Mar. sept.—In monte Beni Hosmar prope Tetuan!

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia). Sibiria.

In Catalogo Plantarum prope Mogador lectarum a clar. Lowe adest *Valeriana Phu*. Hæc species certe non indigena est, sed, ut fit sæpius in Europa, e cultura forsitan evasit.

CENTRANTHUS (*DC. Fl. Fr.* iv. 238).

C. ANGUSTIFOLIUS (*DC. l. c., Prod.* iv. 632).

Mar. merid.—In convalle Aït Mesan a 1400 m. usque 1800 m. et nondum florens in jugo Tagherot circ. 2700 m.!

Ar. Geog. Species per montes Hispaniæ, Galliæ australis et Italiæ diffusa. Formæ arcte affines in Oriente.

Planta polymorpha, ejus limites ægre definiuntur. In speciminibus nostris caules e basi lignoso erecti simplices, nec adsunt rami secundarii ex axillis foliorum, ut sæpius fit in planta Europæa. Folia superiora lanceolata acuta fere æquidistantia. Characteres ex longitudine corollæ et calcaris mihi videntur fallaces.

C. CALCITRAPA (*L., et Schousb. Gew. Mar.* 11, sub *Valeriana*; *Dufr. Val.* 39; *DC. Pr.* iv. 632).

Mar. sept.—Tanger! (*Schsb. l. c.*) (*Salzm. Exsicc.*)! Prope Cap. Spartel! Prope Tetuan! In monte Beni Hosmar!

Mar. occid.—Casa Blanea, *J. D. H. spec.*! Prope Mogador (*Lowe, sp. in Herb. Kew.*)!

Mar. merid.—In convalle Aït Mesan usque 2200 m. In monte Djebel Tezah ad 2100 m.!

Specimen fructiferum a cl. Lowe prope Mogador lectum *C. macrosiphoni* (Boiss. Diagn. Pl. Or. ser. 1, iii. 57) simillimum est. Sed spec. nostra florentia, etsi foliis conformia, certe ad *C. Calcitrapam* genuinum referenda.

Ar. Geog. Regio Mediterranea fere omnis, imprimis occidentalis. In Oriente rarior, deest in ins. Canariensibus.

C. Calcitrapa, var. *orbiculata* (*DC. Pr. l. c.*), = *C. orbiculatus* (*Dufr. l. c.*),

est tantum lusus speciei foliis suborbiculatis subintegerrimis. Hue verisimiliter referendum spec. nostrum nondum florens in monte Beni Hosmar lectum.

FEDIA (*Mæench, Meth.* 486).

F. CORNUCOPILÆ (L., sub *Valeriana*; DC. *Fl. Fr.* iv. 250, et *Prod.* iv. 630).

Mar. sept.—Prope Tetuan! Inter Tetuan et Ceuta!

— —, Var. corollæ tubo ter longiore, = F. *graciliflora* (*Fisch. et Mey. Ind. Sem. Hort. Pet.* vi. 50; *Walp. Rep.* ii. 528). Hue etiam referendæ sunt *Valeriana Cornucopiæ* (*Schousb. Gew. Mar.* 11) et *C. scorpioides* (*Dufr. Val.* 55, tab. 1; DC. *Pr.* iv. 630).

Mar. sept.—Frequens circa Tanger! (*Schsb. l. c.*) (*Salzm. Exsicc.*, sub “*V. cornucopiæ*”)!

Ar. Geog. Species per regionem Mediterraneam calidiorem—peninsulam Ibericam australem, Italiam australem, insulas Sardiniam, Siciliam et Cretam diffusa, fit rarior in Oriente et deest in Macaronesia. Var. *graciliflora* per regionem litoralem Africæ borealis frequenter, et rarius in Hispania australi occurrit, sed forma typica non deest.

Nota diagnostica ex fructu desumpta, ut jam monuit cel. Boissier (*Fl. Or.* iii. 93), instabilis est et fallax, et *F. graciliflora* sine dubio ad varietatem reducenda est. *F. scorpioides*, *Dufr.*, est tantum lusus inflorescentia ex floribus subsessilibus scorpioidea.

VALERIANELLA (*Poll. Palat.* i. 29).

V. DISCOIDEA (*Willd. Sp. Pl.* i. 184, sub *Valeriana*; *Vahl, Enum.* ii. 21, sub *Fedia*; *Loisel. Not.* 148; *Gren. et Godr. Fl. Fr.* ii. 66), = V. coronata (DC. *Pr.* iv. 628, non *Fl. Fr.* iv. 241), = *Valeriana Locusta*, var. coronata (*Schousb. Gew. Mar.* 12, non L.).

Mar. sept.—In collibus et agris regionis Tingitanæ (*Schsb. l. c.*). In collibus inter Tanger et Tetuan (*J. D. H.*)!

Mar. occid.—Casa Blanca! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Sat frequens in regione inferiore et in convallibus Atlantis Majoris. Prov. Shedma! Mesfioua! Tasseremout! In convalle Aït Mesan 1200–1600 m.!

Ar. Geog. Regio fere omnis Mediterranea. In Macaronesia nonnisi in insula Madeira detecta.

V. AURICULA (DC. *Fl. Fr. Suppl.* 492, et *Prod.* iv. 627; *Gaud. Fl. Helv.* i. 84, sub *Fedia*).

Mar. sept.—Ex regione collina inter Tanger et Tetuan! Ex monte Beni Hosmar prope Tetuan!

Mar. merid.—In regione media Atlantis Majoris. In convalle Aït Mesan 1500–2300 m.! In monte Djebel Tezah 1800–2000 m.!

Ar. Geog. Europa media. Hinc inde occurrit in montanis Hispaniæ, Italiæ, et Rumiæ. In Asia Mediterranea nonnisi in Transcaucasia in-

dicata. Ex Africa non vidi. Specimina nostra ex ditione Tingitana immatura et ideo incerta.

VALERIANELLA CARINATA (*Loisel. Not.* 149; *DC. Pr.* iv. 629).

Mar. merid.—Distr. Reraya 1000–1200 m.! In monte Djebel Tezali 1700–2000 m.!

Ar. Geog. Europa media. Regio Mediterranea (exclusa Macaronesia). Fructus in spec. nostris puberulus.

V. MICROCARPA (*Loisel. Not.* 151), = V. mixta (*Vahl, Enum.* ii. 21, sub *Fedia*; *DC. Pr.* iv. 627), = V. puberula (*DC. Pr.* iv. 627).

Mar. sept.—Ex maritimis inter Tetuan et Ceuta!

Ar. Geog. Regio Mediterranea hinc inde: in peninsula Iberica et in Macaronesia nondum detecta. Forsan cum *V. Morisoni* conjungenda. Hæc per Europam fere omnem et regionem totam Mediterraneam diffusa est.

DIPSACEÆ (*Vaill. Mém. Ac. Par.* 1722).

DIPSACUS (*L. Gen.* 114).

D. SYLVESTRIS (*L., DC. Pr.* iv. 645; *Schousb. Gew. Mar.* 43).

Mar. sept.—Tanger!

Mar. occid.—In arvis “prope Saffi” (*Schsb. l. c.*).

Ar. Geog. Europa media. Regio Mediterranea omnis (inclusa Macaronesia).

SCABIOSA (*L. Gen.* 114).

S. MARITIMA (*L., DC. Pr.* iv. 657), = S. grandiflora (*Scop. Insubr.* iii. 29, tab. 14), = S. acutiflora (*Rchb. Cent.* iv. 326; *DC. Pr.* iv. 656), = S. ambigua (*Ten. Syll.* 63; *DC. Pr.* iv. 657).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Mar. occid.—Mazagan! Agadir (*Coll. Indig.* 1871)! Prov. Haha pr. Mogador!

Mar. merid.—Prope Mskala!

— —, var. MINOR. Capitulis fructiferis minoribus, involuclis minus pubescentibus.

Mar. occid.—Aïn el Hadjar haud procul Mogador! In monte Djebel Hadid (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—Ad radices Atlantis Majoris sat frequens. Seksaoua! Aïn Tarsil! Mtouga!

Ar. Geog. Species satis ludibunda per regionem omnem Mediterraneam (inclusa Macaronesia) late diffusa.

S. SEMIPAPPOSA (*Salzm. in DC. Pr.* iv. 658).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Tetuan!

Mar. occid.—Casa Blanca? (nondum fructifera)!

Ar. Geog. Africa boreali-occidentalis. Hispania australis.

SCABIOSA RUTÆFOLIA (*Vahl, Symb. ii. 26*), = *S. divaricata* (*Lam. Illustr. No. 1311*), = *S. urceolata* (*Desf. Fl. Atl. i. 122; DC. Pr. iv. 660*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Herb. Schousb.*), *Coss. spec.*!

Mar. occid.—Mazagan! Ain el Hadjar haud procul Mogador!

Ar. Geog. Peninsula Iberica. Italia. Africa borealis.

Ludit foliis pinnatifidis vel bipinnatifidis inter exemplaria eodem loco enata; planta fit interdum erecta parum ramosa vel ramis divaricatis e basi patentibus. Hæ formæ tamen veras varietates non sistunt.

S. STELLATA (*L., DC. Fl. Fr. iv. 231*), = *S. simplex* (*DC. Fl. Fr. l. c.*), = *S. monspeliensis* (*DC. Pr. iv. 655*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Mar. merid.—Frequens in regione inferiore ad radices Atlantis Majoris et in convallibus calidioribus. Mesfioua! Tasscremout! Ourika! Seksaoua! Amsmiz! In convalle Ait Mesan circ. 1600 m.!

— — —, var. *MONSPELIENSIS*, = *S. monspeliensis* (*Jacq. Misc. ii. 320*)?

Mar. occid.—Agadir (*Coll. Indig. 1871*)!

Mar. merid.—In prov. Shedma pluries visa et lecta!

Ar. Geog. Tam species quam varietas per regionem Mediterraneam occidentalem (exclusa Macaronesia) late diffusæ sunt.

Plurimi auctores species duas in regione Mediterranea occidentali admitterunt, sed has notis diversis distinxerunt. Ex multis exemplaribus accurate scrutatis discrimina ex corollis exterioribus longioribus vel omnibus subæqualibus, et corona membranacea præ setis calycinis paullo longiore vel brevior, fallacia videntur. Varietatem *monspeliensem* admisi, dum planta sub hoc nomine cognita minus ramosa est et caulis sæpius subsimplex; gaudet foliis pinnatisectis, segmentis sæpius dentatis, et setis ad basin involuelli longioribus foveolas 8 tubi calycini obtegentibus. *S. rotata* (*M. B. Fl. Taur. Cauc. iii. 102*), species affinis sed probe diversa, distinguitur imprimis squamis receptaculi setaceis, nec ut in *S. stellata* dilatatis scariosis.

Scabiosam quandam nondum florentem, foliis radicalibus tantum donatam ulterius inquirendam, legi in monte Beni Hosmar prope Tetuan!

PTEROCEPHALUS (*Vaill. Act. Ac. Par. 1722, 184*).

P. DEPRESSUS (*Coss. Bull. Soc. Bot. Fr. xx. 250*).

Sequitur descriptio Cossoniana:—"Planta *perennis subcaulis*, cæspitosa, humo adpressa, pulvinos latos densosque efformans, molliter breviterque villosa. Caudex crassus, lignosus, tortuosus, superne in caudiculos permultos ramosos divisus. Folia omnia *apice caudiculorum* in rosulas *congesta*, petiolata petiolo lato, *pinnatisecta*, 4-5-juga segmentis inferioribus linearibus vel linearis-spathulatis integris, cæteris obovato-cuneatis superne lobulatis lobulo exteriori profundiore lobulis obtusis, segmento superiore sæpius tripartito. Capitula e centro ro-

sularum foliorum enata, brevissime pedunculata. Involuerum uniseriale, foliolis circiter 12, extus molliter breviterque villosis, lanceolatis acutis, floribus subdimidio brevioribus. Receptaculum alveolatum, hirtulum. Corollæ extus serieeæ tubo violaceo, limbo inæqualiter 4-fido intus lutescente, exteriores laciniis oblongis obtusis paulo radiantibus, interiores laciniis lanceolatis. Stamina 4, corollam longe excedentia. Stylus exsertus, stigmate oblongo subbilabiato. Involucellum (immaturum tantum notum) dense villosum, foveolis obsoletis pilis occultatis, cylindraceo-turbinatum, corona brevi annulæformi. *Calyx coronam involucelli vix excedens, setis 14-18, involuella subquadruplo longioribus, dense plumosis, atro-violaceis.*"

Mar. merid.—Satis frequens, ut videtur, in regione media Atlantis Majoris. Primus Balansa legit in monte Sidi Fars pr. Keira. Legimus in convalle Aït Mesan usque in jugo Tagherot a 2000 m. ad 2700 m. ! In convalle Amsmiz 1500-1700 m. ! In monte Djebel Ouensa legit indigenus Ibrahim (*Coss. spec.*) !

Monente eel. auctore hic juxta *P. perennem*, Vaill., collocandus est, sed ab hoc et affinibus probe distinctus.

COMPOSITÆ (*Vaill. Act. Acad. Par. 1718*).

BELLIS (*L. Gen. 962*).

B. ANNUA (*L., DC. Pr. v. 304*).

Mar. sept.—Tanger ! (*Schousb. Herb.*). Inter Tanger et Tetuan !

Planta circa Tingidem sat frequens gaudet ligulis pulchre cæruleseentibus fere ut in specie Atlantica mox enumeranda. Cæteris notis nullis a typo differt.

— —, var. β , MINUTA (*DC. Pr. l. c.*), = *B. microcephala* (*Lange, Pug. 116*), ex *Boiss. Fl. Or. iii. 175*.

Mar. merid.—In Atlantis Majoris convalle Amsmiz, alt. 1600 m. !

Spec. nostris omnino conformia legit Balansa (*Pl. d'Algérie, 1852, No. 620*). Hæc statura minima, ligulis brevissimis involuerum parum excedentibus, foliis oblongo-spathulatis integerrimis a typo recedunt.

Ar. Geog. Per regionem Mediterraneam (inclusa Madeira) late diffusa, fugit regiones continentales a mari remotas. Varietas in Algeria, Hispania et (?) in agro Byzantino detecta est.

B. SYLVESTRIS (*Cyr. Plant. Rar. ii. 12, tab. 4* ; *DC. Pr. v. 305*), var. PAPPULOSA, = *B. pappulosa* (*Boiss. Voy. Esp. 303, tab. 91*).

Mar. sept.—Prope Tanger (*Webb Herb.*) ! In monte Djebel Kebir ! et juxta Cap. Spartel ! Inter Tanger et Tetuan !

Primo aspectu differt insigniter acheniis pappo setoso coronatis ; sed hæc nota, etsi gravis, eerte variabilis est. Inter specimina nostræ Tingitana adsunt nonnulla quæ statura majore, foliis latoribus nervis tribus prominentibus insignitis, pappo brevioris paucisetosi, ad *B. rotundifoliam* (*B. et R. Pug. 55*) proxime accedunt. Illius specimina typica ex ditone

nostra non vidi, sed vercor ne *Bellis rotundifolia* et affinis *B. atlantica* (B. et R. l. c.) inter varietates *B. sylvestris* reducendæ sint.

Ar. Geog. *B. sylvestris* per regionem Mediterraneam litoralem et sublitoralem late diffusa est; deest tamen in Macaronesia. Var. *pappulosa* nonnisi in Hispania australi et Africa boreali-occidentali finitima hucusque obvia fuit.

BELLIS CÆRULESCENS (*Coss. MSS.*).

Radix perennis (sæpius bi- vel triennis?) fibrillas longas filiformes emittens; folia omnia basilaria cochleariformia, limbo ovato-orbiculari interdum basi subcordato petiolo elongato suffulto, ex pube brevi subincana; scapi florigeri aphylli elongati tennes, capitula magnitudine fere *B. perennis* vel minora gerentia; ligulæ pulchre cærulescentes involucrium breviter pilosum duplo superantes; achenia compressa undique pilis patentibus brevissimis obsita, prorsus epapposa.

Planta delicatula loca umbrosa prædiligens a *B. rotundifolia* aspectu valde diversa. In hac achenia facie glabrescentia, margine setosa, apice pappo brevi semper coronata.

Mar. merid.—In regione media Atlantis Majoris sat frequens. In monte Djebel Sidi Fars ad 2000 m. primus detexit el. Balansa. Legimus in convalle Aït Mesan a 1800 m. usque 2400 m.! In monte Djebel Tezah 1800–2300 m.!

ERIGERON (*L. Gen.* 951).

E. LINIFOLIUM (*Willd. Sp. Pl.* iii. 1965; *Boiss. Fl. Or.* iii. 169), = *E. erispum* (*Pourr. Chlor. Narb.* 470), = *Conyza ambigua* (*DC. Fl. Fr. Suppl.* 468, et *Prod.* v. 381).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Mar. occid.—Mazagan! Mogador (*Lowe Cat.*).

Mar. merid.—Ad radices Atlantis Majoris prope Amsiniz!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). In Oriente nonnisi in terris continentalibus obvium.

NOLLETIA (*Cass. Dict.* xxxvii. 479).

N. CHRYSOCOMOIDES (*Desf. Fl. Atl.* ii. 269, sub *Conyza*; *Cass. loc. cit.*; *DC. Pr.* v. 366).

Mar. occid.—Mogador, præsertim in arenosis secus rivulum Oued Kseb! (*Brouss. ex Coss.*; *Lowe Cat.*, *Balansa spec.*)! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In glareosis ad radices Atlantis Majoris. Mesfiona! Ourika!

Ar. Geog. Africa boreali-occidentalis, usque ad fines Saharenses ditionis Constantinæ.

EVAX (*Gærtn. Fruct.* ii. 393).

E. PYGMÆA (*L.*, sub *Filagine*; *Pers. Syn.* ii. 422; *DC. Pr.* v. 458).

Mar. sept.—Tanger versus Cap. Spartel!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

See. eel. A. P. De Candolle exstat in herb. Broussonnet specimen *E. exigua* (Sibth., sub *Filagine*), forsā ex Mauritania allatum. Tam hæc quam *E. asterisciflora* in ditione nostra quærendæ sunt.

EVAX HELDREICHII (*Parl. Giorn. Tosc.* No. 2, *Pl. Nov.* p. 15), = *Filago atlantica* (Ball, *Journ. Bot.* 1873, 364).

Mar. merid.—In regione media Atlantis Majoris supra Arround in glareosis a 2200 m. ad 2400 m.!

Aspectu a planta Sicula omnino diversa, specimina nostra nulla nota gravi discrepant. Specimina quæ vidi ex Sicilia, ut jam monuit eel. Parlatore, ad glomerulum nanum nucis avellanæ magnitudine reducta. Planta Atlantica e collo radiceis ramos plurimos prostratos serpentes 4–6-pollicares emittit. Involueri pentagoni foliola nitida flaveseentia dorso viridia. Achenium sulcatum prismatieum subfusiforme.

Ar. Geog. Hæusque in Sicilia et Algeria orientali tantum detecta.

MICROPUS (*L. Gen.* 996).

M. SUPINUS (*L., DC. Pr.* v. 460).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Inter Tanger et Tetuan!

Mar. merid.—Ad radices Atlantis Majoris. Ourika! Reraya!

Ar. Geog. Peninsula Iberica. Africa borealis. Asia Mediterranea.

M. BOMBYCINUS (*Lag. Gen. et Sp.* 32; *DC. Pr.* v. 460).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. merid.—In regione inferiore Atlantis Majoris. Aït Mesan circ. 1500 m.! In convalle Amsmiz usque 1600 m.! Seksoua! et alibi visus!

Ar. Geog. Per regionem Mediterraneam late diffusus ultra Pyrenæos in Galliam anstralem se protendit. Deest in Macaronesia.

FILAGO (*L. Fl. Succ.* 302).

F. GERMANICA (*L., DC. Pr.* vi. 247).

Mar. sept.—Prope Tanger!

Mar. merid.—Misra ben Kara, haud procul urbe Marocco! Prov. Mtouga (*R. et Fr. spec. in Herb. Kew.*)!

Specimina nostra ad formam *F. canescentem*, Jord., referenda.

— —, var. γ , LANUGINOSA (*DC. l. c.*), = *F. lanuginosa*, *Req.*, = *F. eriocephala* (*Guss. Pl. Rar.* 344, tab. 69).

Mar. sept.—Tanger in monte Djebel Kebir!

Mar. merid.—In convalle Amsmiz circ. 1500 m.!

Ar. Geog. Species per Europam mediam et regionem Mediterraneam (inclusa Macaronesia) late diffusa. Var. *lanuginosa* in calidioribus magis frequens.

F. spathulata (*Presl, Del. Prag.* 93; *Boiss. Fl. Or.* iii. 246), = *F. germanica*, var. δ (*DC. Pr.* vi. 247).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)! Tetuan (1851)!

Mar. occid.—Casa Blanca! Mogador (*Brouss.*)? Confer *DC. l. c.*

Mar. merid.—Prov. Shedma! Prope Mskala!

Ar. Geog. Europa media. Regio omnis Mediterranea (inclusa Macaronesia).

Hanc formam *F. germanicæ* proximam haud sine dubio pro subspecie enumeravi.

FILAGO HETERANTHA (*Rafin. Giorn.* ii. fasc. xii. 170; *Guss. Fl. Sic.* ii. 864), = *F. Cupaniana* (*Parl. Spec. Nov.* 13).

Mar. merid.—In convalle Aït Mesan usque 2100 m.! In monte Djebel Tezah! Specimen incompletum et incertum in convalle Ourika!

Ar. Geog. Sicilia. Algeria. Marocco meridionalis.

Habui pro *F. Duriei* (*Coss. Pl. Crit.* 58) sed amicissimus Cosson certum me fecit de hac specie mihi prius non satis cognita.

F. GALLICA (*L., DC. Pr.* vi. 248).

Mar. sept.—Tanger! Cap. Spartel! Inter Tetuan et Ceuta!

Mar. occid.—Casa Blanca!

Mar. merid.—Frequens in regione inferiore Atlantis Majoris usque 2000 m. Mesfioua! Inter Tasseremout et Ourika (*G. Maw sp.*!) et alibi pluries visa!

Ar. Geog. Europa media. Regio omnis Mediterranea. Fertur in America australi; an indigena?

IFLOGA (*Cass. Dict.* xxxvii. 13).

I. SPICATA (*Forsk. Fl. Æg.-arab.* p. 73, sub *Chrysocoma*; *Vahl, Symb.* i. 70, sub *Gnaphalio*; *Sch. Bip. in Webb Phyt. Can.*), = *I. Fontanesii* (*Cass. l. c.*), = *Gnaphalium cauliflorum* (*Desf. Fl. Atl.* ii. 267), = *Trichogyne cauliflora* (*DC. Pr.* vi. 266), = *Gnaphalium ægyptiacum* (*Pers. Syn.* ii. 421), = *G. supracanum* (*Sibth. et Sm. Fl. Gr.* ix. p. 47, tab. 861).

Mar. occid.—Fertur in agro Mogadorensi ex spec. Broussonnetiano, sed a cæteris botanicis non visa et ideo incerta. Prope Agadir legerunt coll. indigeni Majo 1871!

Mar. merid.—Nascitur prope Misra Ben Kara! in planitie circa urbem Marocco, et inde versus radices Atlantis Majoris. Mesfioua! et alibi visa!

Ar. Geog. Per regionem Mediterraneam calidiorem et regiones subtropicales orbis veteris, ex India orientali usque insulas Canarienses, et in Africa australi late diffusa. Deest hæc species in Europa nisi in Hispania australi, ubi rarissima.

Gnaphalium Ruppellii (*Fres. Mus. Senck.* 79), quod idem est ac *G. Ammophila* (*Wall. Cat.* 2943), fertur in *DC. Prod. l. c.* inter synonyma hujus speciei polyonymæ. Sistit re vera varietatem vel subspeciem foliis subtus arancoso-lanatis. In planta Africana folia in pagina superiore araneoso-lanata, subtus nitida sunt.

PHAGNALON (*Cass. Bull. Phil.* 1819, p. 174).

P. SAXATILE (*L.*, sub *Conyza* ; *Cass. l. c.* ; *DC. Pr.* v. 396).

Mar. sept.—Tanger (*Schsb. Herb.*) ! Cap. Spartel ! Inter Tanger et Tetuan ! Prope Tetuan !

Mar. occid.—Casa Blanea !

Mar. merid.—Frequens per provincias meridionales Imperii et in regione inferiore Atlantis Majoris usque 2000 m. Legimus ad Tasseremout ! Milhaïn ! In convalle Aït Mesan ! In monte Djebel Tezah !

— —, var. *INTERMEDIUM* (*DC. Prod. l. c.*), = *Conyza intermedia* (*Lag. Elench.* 28), = *Phagnalon Lagascæ* (*Cass. loc. cit.*).

Mar. merid.—Valde frequens in regione calidiore prov. Haha ! Shedma pluribus locis ! Seksaoua ! In rupibus haud procul urbe Maroeco, *G. Maw* !

Ar. Geog. Species late diffusa per regionem Mediterraneam occidentalem (inclusa Macaronesia) ; nec ut mihi videtur satis diversa est species orientalis *P. græcum*, Boiss.

Optimo jure varietas nostra *P. intermedium* vocatur, dum inter formas vulgatiores *P. saxatilis* et *P. rupestris*, *DC.*, transitum præbet. Ut morem gererem magistris rei botanices *P. rupestre* sicut subspeciem enumeravi. Horum, re vera, formæ extremæ toto cœlo diversæ videntur ; sed, si seriem formarum intermediarum diligenter explorabis, vix invenies notam stabilem differentialem.

P. rupestre (*L. Mant.*, sub *Conyza* ; *DC. Pr.* v. 396).

Mar. sept.—Tanger (*Schsb. Herb.*) ! Cap. Spartel ! Tetuan ! (1851) !

Mar. occid.—Mogador (*Lowe Herb.*) !

Mar. merid.—Ad radices Atlantis Majoris pr. Seksaoua !

— —, var. *CALYGINUM*, *nob.*, = *P. calycinum* (*Cav.*, sub *Conyza* ; *DC. Pr.* v. 397), = *P. Bennettii*, *Lowe, MSS.*

Mar. occid.—Legimus prope Aïn el Hadjar ! et in monte Djebel Hadid ! et ibidem legerunt anno 1873 ell. Rein et Fritsch. Primus hanc formam legit beatus Broussonnet. Specimina inter hanc et typum *P. rupestris* ludentia legit in agro Mogadorensi *G. Maw* !

Ar. Geog. *P. rupestre* per regionem omnem Mediterraneam (inclusa Macaronesia) diffusum est. Var. *calycinum* in Madeira et in agro Mogadorensi obvia.

P. ATLANTICUM (*Ball, in Journ. Bot.* 1873, 364).

Suffruticosum, radice crassa lignosa, ramis numerosis, abbreviatis, adscendentibus, usque ad apicem foliaciis ; folia oblonga, obtusa, superne cinereo-viridia, subtus albo-tomentoso ; capitula terminalia, ex foliis supremis parum exserta ; involucri foliola eximie bicoloria, nempe pallide flava, macula curvata purpurea notata, exteriora acuta laxè imbricata basi fimbriata, media margine scarioso-dilatata, intima acuta cæteris longiora ; achenia ex pilis copiosis vitreo-luculentia.

Mar. merid.—In regione media Atlantis Majoris. In convalle Aït Mesan circ. 1600 m.!

P. purpurascens, Seh. Bip. in Webb Phyt. Can., habitu ad speciem nostram aliquid accedens, est *P. saxatili* propius, pedunculis elongatis filiformibus, terminalibus et axillaribus, involucri foliolis acutioribus.

GNAPHALIUM (*Don, Trans. Wern. Soc.* vi. 563).

G. LUTEO-ALBUM (*L., DC. Pr.* vi. 230).

Mar. merid.—Videtur haud infrequens, ex regione inferiore in zonam medianam Atlantis Majoris ascendens. Legimus pr. Ourika! et Aït Mesan usque 2000 m.! Specimina ex Oasi Akka benevole comm. cel. Cosson.

Ar. Geog. Per Europam medianam et regionem Mediterraneam omnem late diffusum, inde per regiones temperatas et calidiores totius orbis aufuga vel indigenum.

G. HELICHRYSOIDES (*Ball, in Journ. Bot.* 1873, 364). (Plate XXIII.)

Perenne, caespitosum, ex rhizomate crasso lignoso ramos plurimos brevissimos edens; folia lineari-oblonga, margine revoluta, subtus araneoso-lanata, in pagina superiore sparsim lanata, deinum glabrescentia viridia; scapi florigeri subbipollicares, erecti, herbacei, dense araneoso-lanati, squamis fuscis ebraeteolati; capitula solitaria, pro genere magna; involucri multiseriati foliola fusco-scariosa, oblonga, apice rotundata, exteriora latiuscula, interiora linearia subacuta; flosculi omnes tubulosi æquilongi, fœminei in seriebus 3–6 exterioribus, hermaphroditi numero variabiles interdum fœmineis subæquales, sæpius pauciores; antheræ longe caudatæ; receptaculum vix convexum, alveolorum margine demum læve.

Mar. merid.—In regione alpina Atlantis Majoris. In jugo Tagherot circ. 3000 m.!

Nuper mihi comm. cel. Cosson exemplaria in monte Afougneur ab indigeno Ibrahim lecta, quæ formam seu varietatem sp. nostræ referunt foliis angustioribus arete revolutis, in pagina superiore glaberrimis.

Planta anomala, quoad genus ægre definienda. Celebb. Benth. et Hooker in Gen. Plant. vol. ii. pp. 305–309 discrimen inter *Gnaphalium* et *Helichrysum* in ratione flosculorum ♀ et ♂ posuerunt, prioribus scilicet in *Gnaphalio* prævalentibus, in *Helichryso* deficientibus. Achenia in nostra pseudobiconata, ex pilis versus apicem achenii elongatis pappum exteriorem simulantibus. Hæc structura se ostendit in *Lucilia* (*Gnaphalii* grege Austro-americano) et interdum in *Gnaphalio* (*Omalotheca*, Cass.) *supino*, L. Affinitas proxima cum nostra stirpe inter *Gnaphalia* Andensia (§ *Merope*, Wedd.) quærenda.

Capitula facie ac struetura *Phagnalon* quodammodo referunt; sed invita natura cum hoc genere hucusque naturali plantam nostram iungeres.

Scapi herbacei squamigeri rationem vitalem longe diversam indicant.

HELICHRYSUM (*DC. Prod.* vi. 169).

H. STÆCHAS (*L.*, sub *Gnaphalio*; *DC. Pr.* vi. 182).

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

Species plurimæ affines per omnem regionem dispersæ.

LEYSSERA (*L. Sp. Pl.* 1249).

L. CAPILLIFOLIA (*Willd. Mag. Naturf. Berl.* 1811, 160; *DC. Pr.* vi. 279), = *Gnaphalium leysseroides*, *Dsf.*

Mar. merid.—Ad radices Atlantis Majoris. Mesfioua! Ourika!

Ar. Geog. Regio Mediterranea calidior (exclusa Macaronesia). In Europa nonnisi in Hispania austro-orientali detecta.

INULA (*L. Gen.* 956).

I. MONTANA (*L.*, *DC. Pr.* v. 468).

Mar. merid.—In Atlante Majore loc. dict. Tamarut specimen legerunt el. Rein et Fritsch! Specimen ab indigeno Ibrahim lectum benevole comm. eel. Cosson!

Ar. Geog. Montes Hispaniæ, Galliæ australis, Dalmatiæ et Italiæ. Loca calidiora Alpium Europæ. Specimina Algeriensia non vidi.

I. CRITHMOIDES (*L.*, *DC. Pr.* v. 470).

Mar. sept.—Tanger (*Webb Herb.*)! (*Schsb. Herb.*)!

Ar. Geog. Ad litora Mediterranea, et Europæ occidentalis. Deest in Macaronesia.

I. VISCOSA (*L.*, sub *Erigeron*; *Ait. H. Kew.* iii. 223).

Mar. occid.—Mogador (*Lowe Cat.*), a nobis etiam visa!

Mar. merid.—Legi mox floribundam prope Ourika! Specimina nondum florentia pluries vidi in regione inferiore!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

PULICARIA (*Gærtn. Fruct.* ii. 461).

P. ARABICA (*Cass. Dict.* xlv. 94; *DC. Pr.* v. 478; an *L.* sub *Inula*?).

Mar. sept.—Tanger (*Schsb. Herb.*). Ad lacus salsos prope Tanger (*Webb sp.*)!

Mar. occid.—Casa Blanca! Mazagan! Prope Mogador!

Mar. merid.—In planitie haud procul urbe Maroceo! Ex Ait Ibrahim spec. attulit indigenus Mardoehæus et benev. eomm. eel. Cosson!

Ar. Geog. Africa borealis. Asia Mediterranea. Peninsula Iberica.

Species valde ludibunda. Cel. Boissier (*Fl. Or.* iii. 205) nuper plantam Hispanicam a typo diversam judicavit et pro specie diversa (*P. hispa-*

nica) proposuit. Pace tanti viri notæ differentiales citati nimis inconstantes mihi videntur.

Pulicaria longifolia (Boiss. *Diagn. Pl. Or.* ser. 2, iii. 16; *Ball in Journ. Bot.* 1873, 364).

Differt a *P. arabica* radice bienni vel interdum perennante, cauli elato 1-3-pedali stricto, ramis paucis suberectis, foliis longioribus, basi semiamplexicaulibus, vix aut ne vix auriculatis. Hanc stirpem pro subspecie *P. arabicæ* anno 1873 proposui, sub eodem nomine triviali quo cel. Boissier stirpem suam orientalem jamdudum distinxit. Specimina Ægyptiaca a nobis nuper diligenter explorata notas nullas differentiales præbuerunt. Habitu a *P. arabica* longe recedit et facie *P. capensi* simillima est *P. longifolia*; sed involucria et achenia omnino *P. arabicæ*, et mihi videtur hujus sp. proles, facta biennis vel interdum perennans. Specimina a Balansa ad Oued Biskra lecta (*Plantes d'Algérie*, No. 774) sistunt formam intermediam.

Mar. merid.—Sat frequens in udis et secus aqueducta regionis inferioris. Legi prope urbem Marocco! In distr. Mesfioua! Prope Ourika! Prope Mzouda!

Ar. Geog. Africa borealis ex Ægypto usque Imp. Maroccanum. Peninsula Sinaïtica. Verisimiliter etiam in Africa tropica.

Inter plantas itineris Speke et Grant video *Pulicariam* cui scheda "*P. Grantii*, MSS.," secus Nilum lat. 16° N. lectam, quæ mihi vix a *P. longifolia* diversa videtur. Specimen alterum adest in herb. Kewensi a cel. Schweinfurth in Magna Oasi Lybica lectum cui scheda "*P. inuloides*, DC." Hæc a *P. longifolia* non differt nisi pedunculis crassioribus rigidioribus, et foliis subundulatis. Quid sit *P. inuloides* (DC. *Pr.* v. 480), ex specimine a b. Broussonnet "prope Mogador" lecto descripta, hucusque eruere non potui. Suspicio tamen hanc esse ipsissimæ *P. longifoliæ* lusum floribus radii irregulariter tubulosis, nec ligulatis.

P. ODORA (L., sub *Inula*; *Rchb. Fl. Exc.* ii. 1232; DC. *Pr.* v. 479).

Mar. sept.—Tanger (*Salzm. Exsicc.*, sub *P. dumetorum*)! (*Webb Herb.*)! Djebel Kebir! *Pr.* Cap. Spartel!

Mar. merid.—Ourika! Inter Tassereinout et Ourika (*Maw spec.*)!

— —, var. *MACROCEPHALA*, nob. Differt ligulis longioribus et capitulis fere duplo majoribus.

Mar. occid.—Ex monte Djebel Hadid haud procul Mogador attulerunt ell. Rein et Fritsch (*spec. in Herb. Kew.*)!

Ar. Geog. Species per regionem Mediterraneam præsertim occidentalem (exclusa Macaronesia) late diffusa. In Asia Mediterranea videtur rara.

P. MAURITANICA, Coss. in *Balansa Pl. Maroc.* 1867.

Species ab auctore nondum descripta, a cæteris hujus regionis omnino distincta. Nuper a cl. Warion in Algeria detecta.

Mar. merid.—Haud rara, ut videtur, in rupibus soli expositis prov. meri-

dionalium. Legimus prope Seksaoua! et in convalle Aït Mesan ad 1200 m.! In collibus pone urbem Marocco septentrionem versus (*G. Maw spec.*)! Ex provincia Shedma spec. attulerunt Rein et Fritsch! In rupibus prope Keira jam legit anno 1867 el. Balansa ex spec. in herb. nost.! Exstat spec. in herb. Schousboe sec. schedam MS. "prope Mogador" lectum. Verisimiliter a viro sedulo in itinere versus urbem Marocco detectum, nec in regione litorali, ubi nemini recentiorum obvium fuit.

Ar. Geog. Africa boreali-occidentalis regio continentalis.

PULICARIA SICULA (*L.*, sub *Erigeron*; *Moris, Fl. Sard.* ii. 363), = *Jasonia sicula* (*DC. Ann. Sc. Nat., et Prod.* v. 476), = *Inula elchrysocoides* (*Poir. Voy.* ii. 239; *Schousb. Herb., non Dsf. Fl. Atl.* ii. 275).

Mar. sept.—Tanger (*Schsb. Herb., comm. Cosson*)! (*Salzm. Exsicc.*)!

Ar. Geog. Europa Mediterranea. In Africa boreali infrequens. Deest in Macaronesia.

ODONTOSPERMUM (*Neck. Elem.* i. 20).

De hujus generis synonymia conf. Sch. Bip. in *Phyt. Canar.* ii. 232.

O. AQUATICUM (*L.*, sub *Bupthhalmo*; *Neck. l. c.*; *Sch. Bip. l. c.*), = *Asteriscus aquaticus* (*Mænoch, Meth.* 592; *DC. Pr.* v. 486, = *Nauplius aquaticus* (*Cass. Dict.* xxxiv. 273).

Mar. sept.—Tanger! Inter Tanger et Tetuan! Tetuan (1851)!

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In regione inferiore Atlantis Majoris. Reraya! Tassermout! In convalle Aït Mesan usque 1500 m.!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

O. MARITIMUM (*L.*, sub *Bupthhalmo*; *DC. Pr.* v. 486, sub *Asterisco*; *Sch. Bip. in Phyt. Canar.* ii. 233).

Mar. sept.—Prope Tetuan! Inter Tetuan et Tanger! In insulis Zafirinis (*Webb spec.*)!

— —, var. **MICROPHYLLA**? Plantam singularem nondum florentem foliis crebris minimis ulterius inquirendam legi semel Apr. 1871.

Mar. sept.—In declivibus maritimis prope Ceuta!

Ar. Geog. Species per zonam litoralem Europæ australis et Africæ borealis diffusa. In ins. Canariensibus olim reperta (sec. Buch) sed a recentioribus non visa?

O. ODORUM (*Schousb. Marok*, 185, tab. 7, sub *Bupthhalmo*; *DC. Pr.* v. 486, sub *Asterisco*; *Sch. Bip. in Phyt. Canar.* 236).

Mar. occid.—Sat frequens in regione litorali prope Mogador. Legimus pr. rivulum Oued Kseb! in via qua itur ad Castellum Shedma (*Maw spec.*)! et copiose prope Aïn el Hadjar! Habeo specimen a Balansa in sylva Arganiæ lectum anno 1867! Demum legi in maritimis ab urbe Saffi meridiem versus!

Ar. Geog. Species Maroccana in Canaria a Bourgeau (itinere secundo) detecta; in ins. Porto Santo erronee a beato Webb indicata. Confer Lowe, Man. Fl. Mad. i. 485.

Species nostra ab *O. graveolente* (Forsk. sub *Bupthhalmo*) differt indumento adpresse sericeo-velutino nec cinereo-hirtulo, foliis basi minus dilatatis vix cordatis, foliolis involucri et paleis receptaculi minus acutis. Species Forskohlæana tamen per totam regionem desertorum diffusa, et nuper ab indigeno Mardocheo in Oasi Akka detecta, non nihil variat, et *O. odorum* mihi videtur hujus proles litoralis et sient subspecies enumerandum. Planta Canariensis sistit var. *angustifolium*.

ODONTOSPERMUM IMBRICATUM (*Cav. Anal. Cienc. Nat.* 1801, iv. 94, sub *Bupthhalmo*; *DC. Pr.* v. 487, sub *Asterisco*), = *Bupththalmum elegans* (*Willd. Mag. Naturf. Berl.* 1807, 198), = *Bupththalmum chrysanthemoides* (*Poir. Dict. Suppl.* i. 747).

Mar. occid.—Occurrit rarissime in insula Mogador, specimen unicum detexi capitulo unico mox floribundo præditum!

Ar. Geog. Species Maroccana propria.

Quoad genus hæc mihi videtur dubia species, donec fructus (a Candolleo, ut videtur, non visus) characteres certiores suppeditet. Involucri foliola exteriora foliacea apice subspathulata dentata, interiora acuta, nervo medio prominulo mucronata.

PALLENIS (*Cass. Dict.* xxxvii. 275).

P. SPINOSA (*L.*, sub *Bupthhalmo*; *DC. Pr.* v. 487).

Mar. sept.—Tanger! (*Webb Herb.*)! (*Salzm. Exsicc.*, sub “*Bupththalmum aureum*”)! Inter Tanger et Tetuan (*J. D. H. spec.*)!

Mar. occid.—Casa Blanca!

Mar. merid.—Prov. Shedina pluribus locis! Prope urbem Marocco (*Maw spec.*)! Mesfioua!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

Stirps satis ludibunda; variat caule graciliore, habitu tenuiore, sed vix in varietates divellenda.

XANTHIUM (*L. Gen.* 1054).

X. STRUMARIUM (*L.*, *DC. Pr.* v. 522), var. *ANTIQUORUM* (*Wallr. Beytr.* 229).

Mar. sept.—Tanger (*Salzm. Exsicc.*, sub *X. sphærocephalo*)!

Ar. Geog. Species verisimiliter Europæ australis vel Asiæ Mediterraneæ indigena; inde per regiones temperatas calidiores totius orbis diffusa; in Africa verisimiliter introducta.

ANACYCLUS (*Pers. Syn.* ii. 464).

A. DEPRESSUS (*Ball in Journ. Bot.* 1873, 365). (Plate XXIV.)

Radix biennis vel perennis fusiformis verticalis, ramos plurimos pro-

stratos e collo emittens; rami apicem versus fureati, plerumque diephali, usque sub capitulo foliosi non inerassati; folia pinnata, pinnis alternis in lacinias lineares dissectis; involucri hemisphærici foliola herbacea, acutiuscula, rigida, margine non scariosa, regulariter imbricata; floseuli fœminei ligulati; ligulæ panæ breves erectæ, atrorubentes interdum albo marginatæ; receptaculi paleæ induratae, longiores quam in affini *A. Pyrethro*, apicem tantum herbaceæ, pilosæ; corollæ disci æqualiter quinquelobatae; achenia exteriora anguste marginata in dentes triangulares expansa, interiora non marginata, omnia cum tubo corollino compressa.

Mar. merid.—In regione superiore Atlantis Majoris. In convalle Aït Mesan! et in jugo Tagherot! a 2200 m. ad 2800 m.

Affinitas proxima cum *A. Pyrethro* (L., DC. Pr. vi. 15). Hic differt imprimis capitulis et ligulis duplo majoribus, ramis monoecephalis, paleis receptaculi dimidio brevioribus minus induratis, acheniis omnibus alatis. Facie proximus est *A. maroccano* nostro; hic autem facile distinguitur disci corollis biaristatis, pedunculis apicem inerassatis et acheniis anguste vel subexalatis.

ANACYCLUS CLAVATUS (*Desf. Fl. Atl.* ii. 287, sub *Anthemide*; *Pers. Syn.* ii. 465; *DC. Pr.* vi. 16), = *Anacyclus tomentosus* (*DC. Fl. Fr.* v. 481, et *Prod. l. c.*), = *Anthemis biaristata* (*DC. Fl. Fr.* iv. 204).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Inter Tetuan et Ceuta!

Mar. occid.—Mogador (*Lowe Cat.*).

Ar. Geog. Regio Mediterranea imprimis occidentalis (inclusa Macaronesia).

A. Maroccanus (*Ball in Journ. Bot.* 1873, 365).

Pumilus, prostratus; flosculorum exteriorum ligulæ breves, utrinque purpureæ, interdum albo-marginatæ, demum reflexæ; paleæ receptaculi apice setosæ. Involucrum et achenia fere ut in *A. clavata*.

Mar. merid.—In calidioribus Imperii Maroccani meridionalis. Prov. Shedma! ubi frequens, Mskala! Mesfious! Misra ben Kara! et prope Agadir (*Coll. Indig.* 1871)!

Facie ab *A. clavato* satis diversus. Forsan pro specie omnino diversa vindicandus.

A. RADIATUS (*Lois. Fl. Gall.* 583; *DC. Pr.* vi. 16), = *Anthemis valentina*, var. α (*L. Sp. Pl.* 1262).

Mar. sept.—Tanger! (*Webb Herb.*, sub "*A. Pyrethrum?*")! Tetuan prope ostium fluminis (1851)!

Mar. occid.—Casa Blanca! Mazagan! Saffi! In insula Mogador!

Mar. merid.—Prov. Shedma pr. Aïn Oumast! et alibi! Misra Ben Kara!

———, var. **OCHROLEUCUS**, *nob.* Ligulis pallide ochroleucis vel fere albis, foliis longioribus magis dissectis, et acheniis marginalibus angustius alatis. Facie *A. clavato* similior, sed hic differt involucri

phyllis apice ex appendiculatis et acheniis angustis alatis, internis subexalatis.

Mar. occid.—In arenosis maritimis prope Mogador ! *Lowe* (spec. in *Herb. Kew.*, sub "*A. clavatus*, var. β ") !

Hæc est forma quodammodo media inter *A. radiatum* et *A. clavatum*, priori tamen structura multo propior. Est verisimiliter (ex spec. supra citato) *A. clavatus*, var. β , *Lowe* in *Proc. Linn. Soc.* vol. v. p. 40.

Ar. Geog. Species per regionem Mediterraneam, præsertim occidentalem, late diffusa, fit rara in Græcia et Syria; in Macaronesia nondum detecta.

ANACYCLUS VALENTINUS (*L.*, ex parte, sub *Anthemide*; *DC. Pr.* vi. 16).

Mar. merid.—Satis frequens præsertim ad radices Atlantis Majoris. Legimus in prov. Shedma ! prope Sheshaoua ! prope urbem Marocco ! pr. Sektana ! Amsmiz circ. 1000 m. et iterum circ. 1500 m. ! Spec. ad Ourika et Mtouga lecta attulerunt ell. Rein et Fritsch !

Ar. Geog. Peninsula Iberica. Gallia australis. Africa boreali-occidentalis.

ACHILLEA (*L. Gen.* 971).

A. SANTOLINA (*DC. Pr.* vi. 31).

Mar. merid.—Ad radices Atlantis Majoris, distr. Reraya semel visa et lecta !

Ar. Geog. Species Orientalis et Algeriensis, præsertim in regione desertorum obvia, a Belutschistan usque ad Imp. Maroccanum. Species affinis (*A. santolinoides*, Lag.) in Hispania austro-orientali.

A. LIGUSTICA (*All. Fl. Bed.* no. 660, tab. 53, fig. 2; *DC. Pr.* vi. 26).

Mar. merid.—In regione inferiore Atlantis Majoris. Legimus in convalle Amsmiz circ. 1300 m. ! In rupibus supra Milhain ! Cel. Cosson benevole comm. spec. in montibus Ouensa et Afougueur ab indigeno Ibrahim lecta !

Spec. nostra a typo paululum diversa sunt, foliis magis hirtulis, rachi sæpe 2- vel 3-dentata.

Ar. Geog. Per Europam australem late diffusa. In Syria et in Africa boreali videtur rara.

—— —, var. *FOLIOSA*, *nob.* Differt foliis longioribus flaccidis versus apicem caulis parum diminutis. A typo aspectu valde diversa est, sed characteres graviores non video.

Mar. merid.—Legimus in convalle Ait Mesan a 1600 m. ad 1700 m. ! Taradut (*R. et Fr. spec. in Herb. Kew.*) !

In var. nostra folia inferiora ramorum abortivorum in axillis foliorum caulinorum accessoria videntur.

DIOTIS (*Desf. Fl. Atl.* ii. 261).

D. MARITIMA (*L.*, sub *Athanasia*; *Sm. Eng. Fl.* iii. 403), = *D. candidissima* (*Desf. l. c.*; *DC. Pr.* vi. 34).

Mar. sept.—Tanger! (*ex Herb. Schousboe spec.*)! (*Webb Herb.*)! In plaga Atlantica haud procul Cap. Spartel! Visa inter Tetuan et Ceuta!

Mar. occid.—In maritimis prope Saffi!

Ar. Geog. Secus oras Maris Interni fere totius. Ad litora Oceani ex Anglia usque iusulas Canarienses.

ANTHEMIS (*L. Gen.* 645).

A. TUBERCULATA (*Boiss. Fl. No.* 107; *Voy. Esp.* tab. 90).

Mar. sept.—In monte Djebel Dersa prope Tetuan legit Webb (*spec. in herb. nost.*)!

Mar. merid.—In regione superiore Atlantis Majoris, in monte Djebel Tezah a 2400 m. ad 2700 m.! Ex Djebel Ouensa spec. ab indigeno Ibrahim lectum benev. comm. ecl. Cosson!

Ar. Geog. Peninsula Iberica. Montes Africæ borealis.

A. FUSCATA (*Brot. Phyt. Lus.* 15; *Willd. Sp. Pl.* iii. 2182), = *Maruta fuseata* (*DC. Prod.* vi. 14), = *Perideræa fuscata* (*Webb, It. Hisp.* 37), = *Chamomilla fuseata* (*Gren. et God.* ii. 151).

Mar. sept.—Tanger! (*Webb spec.*)! Inter Tanger et Tetuan!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia). Fertur in Græcia, sed deest hæc species in Boiss. Fl. Orientali.

A. TENUISECTA (*Ball, Journ. Bot.* 1873, 368). (Plate XXV.).

Planta biennis (?), elegans, pedalis et ultra, erecto-ramosa; folia sparsa, bipinnatisecta, lobis tenuibus anguste linearibus, juniora eum ramis novellis velutina, demum viridia, pilosa; capitula terminalia, longiuscule pedunculata, pro genere parvula; involueri pauciseriati foliola obtusa, velutina; flosculi fœminei ligulis brevissimis aurantiacis aucti, hermaphroditi numerosi æquales; receptaculi paleæ flosculis æquilongæ, oblongæ, coneavæ, scariosæ, apice obtusiusculæ, subdentatæ; achenia curvata, exteriora profunde quinquecostata, in costis tuberculosa.

Mar. merid.—Haud procul urbe Marocco! In prov. Mesfoua!

Stirps bene distincta nescio cui proxima. Ob ligulas brevissimas recurvatas videtur discoidea. Ab ovibus crosa, et sub pedibus coneulcata specimina sæpius manea et indeora.

A. MIXTA (*L., Rehb. Ic.* xvi. tab. 110), = *Ormenis mixta* (*DC. Pr.* vi. 18).

Mar. sept.—Tanger! (*Webb Herb.*)! In monte Djebel Kebir! Cap. Spartel!

Mar. occid.—Aïn ei Hadjar haud procul Mogador!

Mar. merid.—Mesfioua! Mskala! Ourika (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio Mediterranea fere omnis (inclusa Macaronesia). Rarius occurrit in Europa occidentali.

Variat foliis majoribus pinnis superioribus dissectis. Ligulæ albæ basi plerumque ochroleucæ; in ditione nostra interdum flavæ et insigniter bicolores.

Prætergenus Linnæanum, et Cassinianum a Candolleo acceptum, auctores pro hac specie genera plurima—scil. *Marutam*, *Matricariam*, *Chamomillam*, et *Chamæmelum*—proposuerunt. Duce cel. Boissier genus *Ormenis* tenere maluisssem nisi obstitissent species plurimæ intermediæ, acheniis *O. mixtæ* propiores, sed habitu et facie *Anthemidibus* typicis simillimæ. Demum sententiam celeb. Benth. et Hook. (*Gen. Pl.* ii. 420) secutus sum, et nomen Linnæanum servavi.

ANTHEMIS HETEROPHYLLA (*Coss. MSS.*, sub *Ormenide*),= *Santolina scariosa* (*Ball in Journ. Bot.* 1873, 365).

Suffruticosa, caule lignoso, e basi ramosissimo, ramis strictis erectis monocephalis; folia infima pinnatipartita, segmentis laciniatis, media profunde dentata vel pinnatifida, segmentis oblongo-linearibus integris, suprema minima lineari-lanceolata denticulata; involucri foliola apice dilatata, fimbriato-scariosa; flores exteriores plerumque ligulati, ligulis flavis abbreviatis interdum deficientibus; disci corollæ aurantiacæ; receptaculi paleæ ovatæ, acutæ, mucronatæ, apice tantum pubescentes.

Mar. merid.—In regione inferiore et media Atlantis Majoris. Ourika! Aït Mesan! In convalle Amsmiz a 1000 m. ad 2000 m.!

Spec. altera nostris conformia, nisi ligulis longioribus, ex montibus Ouensa et Djebel Afougueur ab indigeno Ibrahim lecta misit cel. Cosson et me certiore fecit de identitate *A. heterophyllæ* suæ cum *Santolina scariosa* nostra. Species hæc gaudet re vera sæpissime floribus ligulatis et ideo vix in gen. *Santolina* retinenda. Auctoritatem cel. amici libenter agnosco; attamen vera affinitas plantæ nostræ cum *Santolina* quærenda mihi videtur. Facie proxima est *S. oblongifoliæ* (Boiss. *Diagn. Pl. Or.* ser. ii. fasc. 3, p. 18), sed involucri scariosis (fere *Anacycli radiati*, Lois.), paleis mucronatis, et floribus satis superque distincta est. Achenia immatura tantum vidi, omnino *Santolinis* conformia.

CLADANTHUS (*Cass. in Bull. Philom.* 1816).

C. ARABICUS (*L. Sp. Pl.* 1263, sub *Anthemide*; *Cass. l. c. et Dict.* ix. 342, tab. 87),= *Anthemis prolifera* (*Pers. Syn.* ii. 32),= *Cladanthus proliferus* (*DC. Pr.* vi. 18).

Mar. oecid.—Casa Blanca ! Djebel Hadid haud procul Mogador (*R. et Fr. spec. in Herb. Kew.*) !

Mar. merid.—Frequens in provinciis meridionalibus usque radices Atlantis Majoris. Prov. Haha ! Aïn Oumast ! Sheshaoua ! Mtouga ! Mesfoua ! Ourika (*Maw spec.*) !

Ar. Geog. Species Algeriæ occidentalis et Imperii Maroccani incola, nec in Arabia hucusque inventa.

Nomen a Persopio mutatum quia Patria “Arabia” erronee a Linnæo (*loc. cit.*) indicata. Sed J. Shaw, qui primus hanc stirpem descripsit et delineavit (*Spec. Phytog. Afric. No. 58*), nomen indigenum *Craffas* ab Arabis Mauritanis applicatum indicavit, et ideo latiore sensu nomen triviale Linnæanum haud improprium nec a posteris mutandum.

Flores interdum, nec semper, suaveolentes.

CHRYSANTHEMUM (*L. Gen. 966*).

C. SEGETUM (*L., DC. Pr. vi. 64*), = *Xanthophthalmum segetum* (*Sch. Bip. Tanac. 17*).

Mar. sept.—Tanger (*Webb spec.*) ! Mense Junio visum sed spec. nulla lecta !

Ar. Geog. Regio Mediterranea (exclusa Maearonesia). In Europa fere omni diffusum, sed fere semper in cultis et verisimiliter introductum.

C. CORONARIUM (*L., DC. Pr. vi. 64*), = *Pinardia coronaria* (*Less. Syn. 255*).

Mar. sept.—Tanger ! (*Webb Herb.*) ! (*Salzm. Exsicc.*) ! Tetuan !

Mar. occid.—Casa Blanca ! Agadir (*Coll. Indig. 1871*) !

Mar. merid.—Passim occurrit in regione inferiore. Legimus in prov. Shedma ! Prope Sheshaoua ! In ruderatis urbis Marocco !

Ar. Geog. Regio Mediterranea, imprimis ealidior (inclusa Maearonesia et ins. Azoricis). Deest in Gallia et in Italia superiore.

C. CARINATUM (*Schousb. Gew. Marok. 185, tab. 6 ; DC. Pr. vi. 64*), = *C. tricolor* (*Andr. Bot. ii. tab. 109*), = *Ismelia versicolor* (*Cass. Dict. xli. 40*).

Mar. occid.—Secus rivulum Oued Kseb prope Mogador, ubi primus legit b. Schousboe ! Vidimus mense Aprili florentem in atrio v. horto præfecti urbis Mogador.

Ar. Geog. Species propria Maroccana.

Ligulæ albæ basi flavescentes v. interdum purpurascens.

C. MYCONIS (*L., Boiss. Fl. Or. iii. 335*), = *Pyrethrum myconis* (*Mærch, Suppl. 287 ; DC. Pr. vi. 61*), = *Coleostephus myconis* (*Cass. Dict. xli. 43*).

Mar. sept.—Inter Tanger et Tetuan ! Prope Tetuan ! (*Webb Herb.*) !

Mar. occid.—Rabat (*R. et Fr. spec. in Herb. Kew.*) !

Ar. Geog. Regio Mediterranea omnis, imprimis in zona litorali (inclusa Macaronesia).

Affinitas hujus et sp. sequentis cum *C. segetum* et *C. coronario*, etsi acheniorum structura diversa.

Chrysanthemum macrotum (*Dur. in Duch. Rev. Bot.* i. 363, et *Expl. Alg.* tab. 56, sub *Coleostepho*), = *Myconia macrotus* (*Sch. Bip. in Walp. Ann.* i. 421), = *Glossopappus chrysanthemoides* (*Kze. in Flora*, 1846, 748).

Mar. occid.—Legi in monte Djebel Hadid haud procul Mogador!

Mar. merid.—Spec. unicum incompletum prope Milhaïn ad radices Atlantis Majoris legit J. D. Hooker!

Ar. Geog. Africa borealis. Hispania australis.

Priori valde affine et forsán tantum varietatem sistens. (Confer *Cosson, Notes Crit.* p. 38, et sententiam adversam *cl. Willkomm in Fl. Hisp.* ii. 106).

C. GLABRUM (*Poir. Voy.* ii. 241), = *C. paludosum* (*Desf. Fl. Atl.* ii. 283, tab. 238), = *Leucanthemum setabense* (*DC. Pr.* vi. 48), = *Hymenostemma Fontanesii* (*Willk. Fl. Hisp.* ii. 102).

Mar. sept.—Prope Tetuan florentem legi mense Aprili!

Ar. Geog. Peninsula Iberica. Africa borealis.

Confer Boissier et Reuter, 'Pugillus,' 57, qui plantam Africanam a *C. setabensi* diversam judicant. Ego non vidi nisi notas leves differentiales, et vix pro varietatibus diversis haberem.

C. WEBBIANUM (*Coss. ined.*, sub *Pyrethro*).

Mar. sept.—In monte Beni Hosmar pr. Tetuan legit Webb, unde spec. incompletum in herb. nost.! Legimus in eodem monte folia radicalia tantum, mense Aprili!

Mar. merid.—Specimina ex Atlante Majore loco d. Taradut lecta attulerunt *ell. Rein et Fritsch*, hodie in herb. Kew. conservata! Hæc, haud sine dubio, ad *C. Webbianum* retuli.

Ar. Geog. Species parum nota, forsán per Africam borealem diffusa.

Planta e grege *C. corymbosi*, L., et huic valde proxima. Nisi auctoritate tantorum virorum devinctus pro forma meridionali *C. corymbosi* habuerim. Huic nisi fallor referendum est "*Pyrethrum corymbosum*" ex monte Djebel Zaghouan (*Kralik, Pl. Tunetanæ*, No. 85).

C. GAYANUM (*Coss. et Dur. in Bull. Soc. Bot. Fr.* ii. 14, sub *Pyrethro*), = *C. Mawi* (*H. fil. in Bot. Mag.*).

Mar. merid.—Occurrit frequentissime in regione inferiore Atlantis Majoris ab 800 m. usque 1800 m. s. m. Legimus in convallibus Ourika! Aït Mesan! Reraya! Amsmiz! et in rupibus supra Seksaoua!

Ar. Geog. Species propria zonæ inferioris montium Africæ boreali-occidentalis.

C. Gayanum affine est *Chrysanthemis* (*Pyrethris* auct.) Hispanicis per Hisp. centralem et australem late diffusis, quorum plurima, scil. *Pyrethrum pulverulentum*, Lag., *Chrysanthemum radicans*, Willd., *C. aragonense*, Asso, *Pyrethrum sulphureum*, Boiss. et Reut., cl. Willkomm sub nomine "*P. hispanicum*" in Flora Hispanica, ii. 98, optimo jure conjunxit. *C. Gayanum* est pariter species valde ludibunda. Planta Maroccana habet involucri foliola minus acuta, intima apice longius scariosa quam specimina Algeriensia quæ vidi, et ligulas fere semper utrinque roseas vel pallide purpurascentes. Hæ mihi videntur notæ instabiles nec varietates definitas sistunt.

— —, var. *DEPRESSUM*, nob.

Differt a typo statura humili, ramis basi prostratis florigeris adscendentibus, herba tota viridi subglaberrima, involucri foliolis exterioribus latioribus, margine latiuscule scariosis, interioribus acutiuseculis ex nervo ad apicem usque non producto exapiculatis.

Mar. merid.—In Atlantis Majoris regione superiore a 2200 m. usque 3000 m.! Aït Mesan supra Arround! In jugo Tagherot! In monte Djebel Tezali!

Facie a typo satis diversa est planta nostra, nec transitum vidi; tamen pro varietate sine dubio habui.

CHRYSANTHEMUM MARESII (Coss. in *Bull. Soc. Bot. Fr.* ii. 16, sub *Pyrethro*), var. *HOSMARIENSE* (Ball in *Journ. Bot.* 1873, 366).

Mar. sept.—In parte superiore montis Beni Hosmar versus 1000 m.!

Planta nostra inter *C. Maresii* et *P. arundanum* (Boiss. Voy. Esp. 317), mihi nonnisi ex descriptione notum, ambigua videtur. *Pyrethro arundano* foliis et habitu valde affinis, differt tamen (ex fructu immaturo nimis incerto) acheniis 10-costatis, corona brevi inæquali subintegra donatis, nec 5-costatis, corona dentato-lacera.

Ar. Geog. Africa boreali-occidentalis.

C. ATLANTICUM (Ball in *Journ. Bot.* 1873, 366). (Plate XXVI.).

Pereune, subcæspitosum; rhizoma divisum; folia radicalia fasciculata, inæqualiter trisceta, lobo medio ternato, lateralibus a parte exteriori lobulo auctis, laciniis ovatis vel late oblongis, omnia pilosa, in pagina superiore demum calvescentia; pedunculi aphylli vel basin versus folio trisceto instructi; involucri foliola late oblonga, apice rotundata, pallide virentia nervo obscure viridi notata, margine lato fusco-scarioso; ligulæ latiusculæ ex albo rubellæ, subtus striis 4 saturatoribus notatæ; achenia 10-alata, alis diametrum achenii æquantibus in coronam brevem subæqualem productis. Stylus brevissime bifidus pseudoclavatus.

Mar. merid.—In regione superiore Atlantis Majoris. In jugo Tagherot! et in monte Djebel Tezali! a 2400 m. ad 3200 m.

Species, ut videtur, omnino distincta, et præter cæteras notas acheniis

insignibus a *C. Gayano* aliena. Structura fructus est tamen in *Chrysanthemis* valde variabilis.

CHRYSANTHEMUM CATANANCHE (*Ball in Journ. Bot.* 1873, 366; *II. fl. in Bot. Mag.* c. tab. 6107).

Perenne, subæspitosum; rhizoma lignosum, divisum; scapi adscendentes, infra medium foliacei; folia parva trifida, laciniis trisectis, dentatis simplicibusve, omnia in petiolum linearem limbo 3-4 longiorem attenuata; capitula magna solitaria; involueri foliola laxè imbricata, ovato-lanceolata, acutiuseula, albo-scariosa, lœcentia, dorso cartilagineo vel herbaeco parè pilosa; flosculi pro genere pauci, fœminei longè ligulati, hermaphroditi lutei antheris breviores; achenia (immatura) costato-alata, alis 4 vel 6, exteriora pappo auriculæformi ipsi achenio æquali, interiora (disci) pappo brevi inæquali subauriculato coronata; stylus bifidus ramis demum recurvis.

Mar. merid.—In regione superiore Atlantis Majoris. Ex convalle Aït Mesan usque in jugum Tagherot! In monte Djebel Tezah! a 2300 m. ad 3200 m.

Species insignis, decens florulæ alpinæ Atlantis. In planta spontanea ligulæ color ægre definienda ex ochroleuco purpurascens, basi atrorubens. In planta culta ligulæ fiunt flavescentes.

Ducibus celeb. Bentham et Hooker (*Gen. Plant.* ii. 424 et seq.) species Maroccanas e grege *Tanacetearum* in genera tria—*Chrysanthemum*, *Matricaria*, et *Tanacetum*—distribui, nec alium ordinem magis naturalem apud auctores summos invenire potui. Sed revera normæ ad genera naturalia stabilienda in hac serie prorsus desunt, dum characteres ex floribus et fructus structura desumpti cum ratione vitali et evolutionis ordine non conveniunt. Etiam auctores qui genera nova facillime admiserunt species toto cœlo alienas in eodem genere collocarunt, dum alias quam maxime affines nomine generico diverso salutarunt. Confer Willkomm in 'Flora Hispanica':—*Pyrethrum alpinum* et *P. corymbosum*; et contra, *Matricaria Chamomilla*, *Chamæmelum inodorum* et *Otospermum glabrum*, et sex species *Chrysanthemorum* ligulis aureis in genera quinque distributas.

MATRICARIA (*L. Gen.* 967).

M. CHAMOMILLA (*L., DC. Pr.* vi. 51).

Mar. sept.—Tanger (*Webb Herb.*)!

Ar. Geog. Species per Europam omnem (exclusa arctica), regionem omnem Mediterraneam (inclusa Macaronesia), et Sibiriam Uralensem dispersa.

M. GLABRA (*Lag. Elench. Hort. R. Matr.* 1805; *Gen. et Sp.* 30, 1816; et *DC. Pr.* vi. 55, sub *Pyrethro*), = *Pyrethrum arvense* (*Salzm. Exsicc.* 1825; *DC. Pr. loc. cit.*), = *Chamæmelum arvense* (*Rehb. fil.*), = *Otospermum glabrum* (*Willk. Fl. Hisp.* ii. 94), = *Pyrethrum Tingitanum* (*Bernh. Ind. Sem. Erf.* 1833).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! Cap. Spartel!
Ex collibus inter Tanger et Tetuan! Tetuan (1851)!

Ar. Geog. Peninsula Iberica australis et Africa Tingitana. Beatus Sch.
Bipontinus (Tanac. p. 49) veram affinitatem hujus et sequentis sp.
primus indicavit.

MATRICARIA FUSCATA (*Desf. Fl. Atl.* ii. 283, tab. 237, sub *Chrysanthemo*; *Willd. Sp. Pl.* iii. 2156, et *DC. Pr.* vi., sub *Pyrethro*).

Mar. sept.—? Tanger (*Webb Herb.*)!

Planta Webbiana a me non diligenter explorata incerta manet. Legi
prope Tingidem spec. nana *M. glabræ* e basi ramosa huic simillima,
etsi involucri et acheniorum structura satis diversa.

Ar. Geog. Hucusque nomen in Algeria meridionali et in agro Tuncetano
detecta.

Hæc species inter *Chrysanthemum* et *Matricariam* ambigua est sed ob
affinitatem cum priore huc referenda.

M. PUBESCENS (*Desf. Fl. Atl.* ii. 284, et *DC. Pr.* vi. 80, et *Benth. et Hook. Gen. Pl.*, sub *Cotula*; *Coss. et Kralik in Bull. Soc. Bot. Fr.* iv. 279, sub *Chlamydophora*).

Mar. merid.—Ex Oasi Akka spec. ab indigeno Mardochæo lecta benev.
comm. cel. Cosson!

Ar. Geog. Sahara Algeriensis et Marocana.

M. maroccana (*Ball in Journ. Bot.* 1873, 367).

Annua, pumila, e basi ramosissima; caules scapiformes, inferne foliosi;
folia subglaberrima, pinnatisecta, segmentis linearibus, sæpius alternis
integris dentatisve; involucri hemisphærici post anthesin incurvi foliola
subæqualia, scariosa, biseriata, paucis brevioribus basi adjectis; flosculi
fœminei ligulati, ligulis diametrum disci subæquantibus, hermaphro-
diti breves, steriles fertilibus intermixti; achenia sessilia, incurva, sub-
cylindrica, lucentia, sub lente minutissime striolata, pappo caduco, auri-
culæformi, versus apicem dilatato, tubum corollinum æquante vel
superante, coronata.

Mar. merid.—In prov. meridionalibus Imperii Maroccani sat frequens.
Shedma! Aïn Oumast! Mesfoua! Aïn Tarsil!

Affinitatem proximam cum *Cotula pubescente*, *Desf. loc. cit.* jam indi-
cavi; sed ex floribus fœmineis semper ligulatis, et illis disci minus nu-
merosis, herbæ glabritie, cæterisque notis minoris momenti, pro specie
diversa habui. Hodie ad subspeciem reducenda videtur. Pace celeb. Benthani et Hookeri hæc cum *Chlamydophora tridentata* (Ehrbg., *DC. Pr.* vi. 139) melius in genere *Matricaria* cum proxime affini *M. auriculata* (Boiss. *Diagn. Pl. Or.* ser. 1, xi. 23, sub *Chamæmelo*) collocanda est. *Chlamydophora tridentata* spontanea, ex spec. a me lectis, et ex descript. auctorum, semper discoidea est. In hac tamen, culta in Horto Paroliniano ex sem. Ægyptiacis, flores exteriores radiantes gaudent ligulis flavis.

TANACETUM (*L. Gen.* 944).

T. ANNUUM (*L., DC. Pr.* vi. 131).

Mar. sept.—Tanger! legi spec. nondum florens. Habui ex herb. Schousboe! (*Salzm. Exsicc.*)!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

ARTEMISIA (*L. Gen.* 945).

A. HERBA ALBA (*Asso, Syn.* No. 810, viii. tab. 1), = *A. arragonensis* (*Lam. Dict.* ii. 269; *DC. Pr.* vi. 101).

Mar. sept.—Tanger (*Schsb. spec.*)! Mers el Kebir (*Webb spec.*)!

Mar. merid.—Vidimus nondum florentem in prov. Shedma! et alibi in regione inferiore, et ad radices Atlantis Majoris prope Ourika!

Ar. Geog. In Hispania imprimis centrali et in convallibus calidioribus Pyrenæorum. Africa boreali-occidentalis. Ex Algeria spec. non vidi.

Legi in convalle Amsmiz ad circ. 1400 m. specimina nondum florifera sp. novæ (*A. maroccanæ*, Coss. MSS.) nondum publici juris factæ.

A. ABSINTHIUM (*L., DC. Pr.* vi. 125).

Mar. merid.—? Specimen nondum florens in prov. Mtouga lectum verisimiliter huc referendum!

Ar. Geog. Europa omnis et Asia temperata. Regio omnis Mediterranea.

SENECTIO (*L. Gen.* 1216).

S. VULGARIS (*L., DC. Pr.* vi. 341).

Mar. sept.—Tanger!

—— —, var. **TENERIFFÆ**, = *S. teneriffæ*, *Sch. Bip.*

Mar. merid.—In monte Djebel Tezah ad 2200 m.!

Ar. Geog. Species per regiones temperatas orbis veteris fere ubique diffusa. In America boreali verisimiliter introducta.

S. lividus (*L., DC. Pr.* vi. 343).

Mar. sept.—In monte Djebel Kebir prope Tanger! Cap. Spartel!

—— —, var. **PINGUIS**, *nob.* Caule humili crassiore; planta facie et odore *S. viscosum* simulans, ahenia tamen pubescentia.

Mar. merid.—In regione alpina Atlantis Majoris. Legi in jugo Tagherot ultra 3000 m.!

Ar. Geog. Regio Mediterranea, imprimis Europæ australis. Species deest in Asia Mediterranea, in Ægypto, et in Macaronesia.

Var. nostra forsitan ad *S. faniculaceum* (Ten., *DC. Prod. l. c.*) referenda est; hic certe inter formas *S. lividi* enumerandus.

S. LEUCANTHEMIFOLIUS (*Poir. Voy.* ii. 238; *DC. Pr.* vi. 344), = *S. humilis* (*Desf. Fl. Atl.* ii. 271; *DC. Pr. l. c.*), = *S. glaucus* (*DC. Pr.*

l. c., non *L.*), = *S. hyemalis* (*Salzm. Exsicc.*), = *S. vernus* (*Biv. Cent. i. 73*), = *S. trilobus* (*L., sec. Willk. et Lge.*).

Mar. sept.—Satis frequens prope Tanger (*Salzm. Exsicc.*)! Legimus in monte Djebel Kebir! et ad Cap. Spartel! Prope Tctuan!

Mar. occid.—Mogador! Aïn el Hadjar!

— —, var. *MAJOR*, *nob.* Differt statura majore, foliis basi auriculatis, ligulis longioribus sæpius expansis.

Mar. sept.—In monte Beni Hosmar prope Tetuan!

Specimina majora hujus var. nimis proxima sunt *S. petræo* (Boiss. et Reut. Pug. 50); hic, ut mihi videtur, inter formas *S. leucanthemifolii* enumerandus.

Ar. Geog. *S. leucanthemifolius* per regionem Mediterraneam australem satis frequens fit rarior in Oriente et in Peninsula Iberica. Deest in Macaronesia, ubi tamen nascitur subspecies mox enumeranda.

Hæc est species polymorpha cujus limites ægre definiendi. Proximus est, sed, ut videtur, satis diversus, *S. vernalis*, W. K., per regionem Mediterraneam orientalem diffusus, species pariter polymorpha, cujus varietates cum formis *S. leucanthemifolii* sæpe commutantur.

Senecio crassifolius (*Willd. Sp. Pl. iii. 1982; DC. Pr. vi. 344*).

Mar. occid.—Mogador (*Lowe Cat.*). In rupibus insulæ Mogador!

Ar. Geog. Nascitur hinc inde in maritimis per regionem omnem sp. præcedentis, cujus sistit subspeciem, seu forsan varietatem maritimam. In Macaronesia etiam obvia.

Ex capitulis crassioribus et foliis inerassatis subearnosis hæc stirps a præcedente facie diversa est, sed non video notas diagnosticas stabiles, et non desunt formæ intermediæ.

S. GALLICUS (*Chaix in Vill. Dauph. i. 331; DC. Pr. vi. 346*), = *S. squalidus* (*Willd. Sp. Pl. iii. 1991, an L.?*), = *S. coronopifolius* (*Willd. non Desf.*).

Mar. merid.—In planitie haud procul urbe Marocco spec. legit J. D. Hooker!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

— —, var. *SONCHIFOLIUS*, *nob.* Forsan est *S. auriculatus* (Desf. Fl. Atl. ii. 272, non Vahl). Differt foliis flaccidis irregulariter pinnatifidis basi in auriculam expansis. Plane apparet esse forma *S. gallici* locis humidis umbrosis enata.

Mar. merid.—In horto irrigato Ben Dreis urbis Marocco!

S. coronopifolius (*Desf. Fl. Atl. ii. 273; DC. Pr. vi. 344*).

Mar. merid.—Specimen unicum in convalle Aït Mesan ad 1100 m. alt. legit eel. Hooker! Frustulum in Oasi Akka ab indig. Mardochæo lectum benev. comm. Cosson!

Ar. Geog. Regio Mediterranea calidior a Persia et Arabia usque insulas Canarienses. Nascitur etiam in Græcia et in Asia temperata.

Hic mihi videtur *S. gallici* proles in regionibus siccis calidioribus enata.
Est fere certe *S. glaucus*, L. Sp. Pl. 1217; sed nomen ab auctoribus
confusum hodie melius negligendum.

SENECIO ERRATICUS (*Bert. Amœn. It.* 42; *DC. Pr.* vi. 349).

Mar. sept.—Tanger (*Webb Herb.*)! Legi nondum florentem et ideo incertum in monte Djebel Kebir!

Ar. Geog. Europa media. Regio Mediterranea (exclusa Macaronesia).

S. FOLIOSUS (*Salzm. Exsicc.* 1825; *DC. Pr.* vi. 351).

Mar. sept.—Tanger (*Salzm. Exsicc.*), in *Herb. Kew.*!

Ar. Geog. Ditio Tingitana. Hispania fere omnis, sed infrequens.

Mihi videtur nimis proximus *S. præalto*, Bert.

S. GIGANTEUS (*Desf. Fl. Atl.* ii. 273, tab. 234; *DC. Pr.* vi. 349).

Mar. merid.—Frequens secus rivulos ad radices Atlantis Majoris. In

Aït Mesan spec. nondum florentia legimus! et pluries alibi vidimus!

De identitate plantæ Maroccanæ non certissimus sum sed, habitus et folia conformia sunt. Canles a 6- usque 10-pedales et ultra; folia subtus leviter araneosa.

Ar. Geog. Africa boreali-occidentalis.

S. DORONICUM (L., *DC. Pr.* vi. 357), var. *HOSMARIENSE* (*Ball, Journ. Bot.* 1873, 367).

Differt a typo caule brevior, foliis caulinis paucis angustis inferioribus 2-3, limbo lato, in petiolum attenuatis, radicalibus late ovatis, basi truncata vel subcordata, in petiolum non decurrentibus.

Mar. sept.—In monte Beni Hosmar, prope Tetuan, altit. s. m. circ. 1000 m.!

Achenia immatura glabra.

Ar. Geog. *S. Doronicum* per montes excelsiores Europæ mediæ et australis a Carpathiis usque Apenninos et Pyrenæos diffusus est.

In regionibus confinibus adsunt formæ plurimæ arete affines, sicut subspecies vel varietates enumerandæ. In Europa orientali adest e. g.

S. transylvanicus, Boiss., in Gallia australi *S. Gerardi*, Gren. et God.,

in Hispania septentrionali *S. Lagascanus*, DC., in Hispania media

S. Carpetanus, Boiss. et Reut., demum in Hispania Bætica *S. eriopus*,

Willk. Hic novissimus pl. nostræ notis nonnullis accedit, at differt caule aphylo et foliis subtus albo lanuginosis.

S. PTERONEURA (*DC. Pr.* vi. 338, sub *Kleinia*).

Mar. occid.—Legimus mox floribundum in arenosis prope Mogador, ubi primus legit Broussonnet!

Ducibus celeb. Bentham et Hooker (*Gen. Pl.* ii. p. 447 et seq.) sub genere *Senecione* hanc plantam insignem enumeravi.

Ar. Geog. Siccies Maroccana.

CALENDULA (*L. Gen.* 990).

C. OFFICINALIS (*L., DC. Pr.* vi. 451).

Mar. sept.—In regione collina inter Tanger et Tetuan! In rupibus prope Tetuan!

Specimina nostra plantam spontaneam ostendunt quam ego nullibi in Europa vidi.

Ar. Geog. Sæpe recurrit per Europam calidiorem, sed vix aut ne vix spontanea videtur. Desideratur in Boissier, 'Fl. Orientali,' et spec. pl. spontaneæ non viderunt Willk. et Lange (cf. *Fl. Hisp.* ii. 126).

Species probabiliter ex Africa boreali in hortis Europæis olim allata.

C. ARVENSIS (*L., DC. Pr.* vi. 452).

Mar. sept.—Tanger! (*Webb Herb.*)!

Mar. occid.—Casa Blanca! Mogador! (*Lowe Cat.*).

Ar. Geog. Regio omnis Mediterranea, unde per Europam temperatam in cultis introducta fuit.

— —, var. *BICOLOR* (*DC. Pr. l. c.*), = *C. bicolor* (*Raf. Caratt.* p. 82; *Boiss. Fl. Or.* iii. 418).

Mar. sept.—Inter Tanger et Tetuan!

Ar. Geog. In Sicilia, Græcia, Ægypto et ins. Canariensibus indicata; verisimiliter per regionem fere omnem Mediterraneam dispersa.

— —, var. *ECHINATA*, *nob.*, = *C. echinata* (*DC. Pr.* vi. 453).

Mar. occid.—Prope Mogador legit et ill. A. P. De Candolle communicavit Broussonnet. Legimus in prov. Haha haud procul Mogador spec. descript. Candolleanæ conformia!

Mar. merid.—In prov. Shedma legit J. D. Hooker! Ad radices Atlantis Majoris legi prope Amsmiz!

Ar. Geog. Forma hujus sp. polymorphæ propria Imp. Maroecani.

— —, var. *MICRANTHA*, *nob.* Forma *C. parvifloræ* (*Raf. Caratt.* 83) valde proxima. Hæc tamen suadente eel. Cosson inter formas *C. stellatæ* enumeranda est; ex acheniorum structura spec. nostra ad *C. arvensis* retuli. Gaudet ramis suberectis, floribus quam in *C. arvensis* dimidio minoribus.

Mar. occid.—In maritimis haud procul Mogador!

Mar. merid.—In Oasi Sheshaoua! et in planitie ab urbe Maroceo meridiem versus!

C. STELLATA (*Cav. Ic.* i. 3, tab. 5; *DC. Pr.* vi. 454), = *C. sicula* (*Cyr., DC. Pr.* vi. 452).

Mar. merid.—In aridissimis pr. Aïn Oumast! Mskala! Seksaoua!

De hac specie confer Cosson et Kralik in *Bull. Soc. Bot. Fr.* iv. 282.

Præter synonyma a cll. auctoribus allata addenda erit forsitan *C. palæstina* (*Boiss. Diagn. Pl. Or. ser. 1, x.* 83). Specimina nostra omnia

locis aridis enata ex ligulis brevibus et foliis angustis ad *C. ægyptiacam*, Desf., accedunt, et notas diagnosticas stabiles vix inveniēs.

Ar. Geog. Species ludibunda per Africam borealem late diffusa, nascitur etiam in Sicilia et in Gallia australi et in Hispania austro-orientali, nec deest forma quam maxime affinis (*C. palestina*) in Oriente.

CALENDULA SUFFRUTICOSA (*Vahl, Symb.* ii. 94; *DC. Pr.* vi. 453; *Boiss. Voy. Esp.* tab. 99), = *C. marginata* (*Willd. Enum. Hort. Berol.* 935).

Mar. sept.—Tanger!

Mar. occid.—Casa Blanca!

Mar. merid.—In collinis prov. Haha!

Ar. Geog. Per Africam borealem late diffusa, habitat etiam in Hispania australi, in Lusitania et in ins. Canariensibus. Forma arcte affinis (subspecies vel varietas) *C. fulgida*, Raf., nascitur in Sicilia, nec a specie separanda est *C. Noëana*, Boiss., in agro Byzantino et in Rumelia inventa. Inter synonyma hujus speciei enumeranda est *C. lusitanica* (Boiss. *Diagn. Pl. Or. ser. 1, x.* 83) in rupibus Cintra prope Olyssiponem a cl. Guthnick et a me lecta. Achenia matura non vidi, nec habuit cel. auctor, sed a *C. suffruticosa* non mihi differre videtur nisi foliis latioribus apice rotundatis. Hæc vidi in var. mox enumeranda, nec desunt interdum in sp. affini *C. sinuata* (Boiss. et Gaill. *Diagn. Pl. Or. ser. 2, vi.* 109).

— — —, var. *TOMENTOSA*, = *C. tomentosa* (*Desf. Fl. Atl.* iii. 305, tab. 245), = *C. incana* (*Willd. Sp. Pl.* iii. 2341; *DC. Pr.* vi. 452).

Mar. sept.—Tetuan in rupibus (1851)! In monte Beni Hosmar! Ex Imp. Maroccano attulit Broussonnet, sed locus natalis haud indicatur. Forma a specie aspectu satis diversa, sed characteres a fructu desumpti nimis variabiles, et nullo jure pro specie diversa, in Willk. et Lange, *Fl. Hisp.* ii. 126, enumerata.

C. maroccana (*Ball in Journ. Bot.* 1873, 367).

Radix perennis; folia angusta elongata, inferiora lanceolata integra, vel in spec. macrioribus leviter repando-dentata, superiora linearia, omnia acuta; rami graciles erecti 6–24-pollicares; capitula quam in *C. suffruticosa* minora; ligulæ angustæ pallide flavæ vel citrinæ; achenia exteriora immarginata in rostrum tenue, involuerum ter superans, apice uncatum obtusum, producta, media margine cartilagineo pectinata, dorso cristulata, intima transverse rugosa.

Mar. merid.—Habitat in regione inferiore et in convallibus calidioribus Atlantis Majoris. Ansmiz! Mzouda! Seksaoua! necnon in rupes-tribus haud procul urbe Marocco, *G. Maw!* Lusum foliis latioribus legimus in convalle Aït Mesau! et in monte Djebel T'ezah!

Facie a *C. suffruticosa* valde, diversa, dignoscitur imprimis habitu gracilimo, et, præter alias notas, rostro acheniorum exteriorum longiore tenuiore, margine mediorum dilatato pectinato, et foliis semper basin

versus attenuatis margine non denticulatis. A *C. stellata* præter habitum singularem statim distinguitur radice perenni et rostro acheniorum obtuso, unicato. Pro specie propria habuerim, sed in hoc genere characteres stabiles omnino desiderantur.

ECHINOPS (*L. Gen.* 999).

E. SPINOSUS (*L., DC. Pr.* vi. 525).

Mar. merid.—Sat frequens in planitie et ad radices Atlantis Majoris usque 1500 m. alt. Legimus juxta Aïn Oumast! prope urbem Marocco! Seksaoua! In convalle Aït Mesan! et in prov. Mtonga, unde spec. attulerunt etiam cl. Rein et Fritsch!

Ar. Geog. Africa borealis. Fertur etiam in Syria et in insula Cypro, sed a recentioribus non visus.

Specimina nostra inter plantam Ægyptiacam et *E. Bovei* (Boiss. Diagn. Pl. Or. ser. 1, vi. 99) ambigua sunt. Hunc ad varietatem reducendum esse ipse ecl. Boissier (Fl. Or. iii. 429) suspicatus est. Antiqui, ut mihi videtur, formam intermediam cognoverunt et hanc pro typo speciei habuit Linnæus. Tabula citata Dodonæi (Pemptades, ed. i. 710, ed. ii. 722) potius ad plantam nostram quam ad Ægyptiacam referenda videtur. Eandem tabulam reproduxit Parkinson (Theatr. 977).

E. STRIGOSUS (*L., DC. Pr.* vi. 526).

Mar. merid.—Videtur infrequens in regione inferiore. Legimus inter Tasseremout et Ourika! et in distr. Reraya! Specimen prope Sektana lectum in Herb. Kew. coll. R. et Fr.!

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

XERANTHEMUM (*L. Gen.* 947).

X. MODESTUM (*Ball in Journ. Bot.* 1873, 368).

Affine *X. squarroso*, Boiss. Diagn. Pl. Or. ser. 1, fase. vi. p. 101, at differt insigniter hisce notis: involucri foliola exteriora angustiora, interiora anguste lanceolata mucronata (nec mutica), vix aut ne vix colorata, multo breviora (nec radiantia), paleas receptaculi integras lineares parum (nec triplo) superantia.

Mar. merid.—In regione inferiore et media Atlantis Majoris. Ludit erectum, 6-10-pollicare, prope Seksaoua! et nanum, caule e basi 2-3-furcato, foliis latioribus basi semiamplexicaulibus, in convalle Aït Mesan! et in monte Djebel Tezah! Altit. a 1200 m. ad 2000 m.

Achenia nigra, adpresse pilosa, striolata; pappi radii in setam scabram abrupte angustati, longiores paleis receptaculi æquilongi; folia (ut in *X. squarroso*) undique lanata mucrone nudo apiculata, mucro foliorum superiorum sæpe in apicem cartilagineum involucri foliolo similem productus. A *X. inaperto*, W., præter alias notas, differt receptaculi paleis integris.

CARLINA (*L. Gen.* 929).

C. LANATA (*L., DC. Pr.* vi. 546).

Mar. merid.—Seksoua! Mtouga? (spec. incertum)!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

C. RACEMOSA (*L., DC. Pr.* vi. 547), = *C. sulfurca* (*Desf. Fl. Atl.* ii. 251, tab. 224).

Mar. sept.—Tanger (*Salzm. Exsicc.*, sub *Atractylide gummifera*)! (*Schousb. Herb.*)!

Mar. occid.—Mazagan! Saffi!

Mar. merid.—In regione inferiore pluries visa. Mesfoua! Prov. Shedma (*J. D. II. spec.*)!

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis. Sardinia. Corsica.

C. INVOLUCRATA (*Poir. Voy.* ii. 234; *DC. Pr.* vi. 547).

Mar. occid.—In agro Mogadorensi (*Broussonet ex DC. loc. cit.*).

Ar. Geog. Africa borealis. Hispania. Oriens.

ATRACTYLIS (*L. Gen.* 939).

A. CANCELLATA (*L., DC. Pr.* vi. 550).

Mar. sept.—Tanger (*Salzm. Exsicc.*, sub "*A. canescens*")! Tetuan (1851)!

Mar. occid.—Aïn el Hadjar hand procul Mogador! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Satis frequens in regione inferiore et in convallibus calidioribus Atlantis Majoris!

Ar. Geog. Regio Mediterranea fere omnis (inclusa Macaronesia).

A. CÆSPIOSA (*Desf. Fl. Atl.* ii. 254, tab. 225; *DC. Pr.* vi. 550).

Mar. merid.—Ex prov. Mtouga spec. attulerunt Rein et Fritsch!

Ar. Geog. Africa boreali-occidentalis.

A. HUMILIS (*L., DC. Pr.* vi. 550).

Mar. merid.—Spec. exsiccata anni præcedentis legi in lapidosis aridissimis pr. Aïn Oumast!

Ar. Geog. In calidioribus Hispaniæ et Galliæ Narbonensis, et in Imp. Maroccano. Indicatur in Italia, sed nemini recentiorum, quoad sciam, obvia fuit.

A. FLAVA (*Desf. Fl. Atl.* ii. 254; *DC. Pr.* vi. 551).

Tabula Fontanesiana No. 262 a Candolleo citata deest in duobus a me scrutatis exemplaribus 'Floræ Atlanticæ,' nec in calce operis citatur.

Mar. merid.—? Legi specimen incertum in alveo torrentis Ourika, caule bicephalo exsiccato anni præcedentis, alteroque immaturo nondum florifero præditum!

Planta ulterius inquirenda.

Ar. Geog. Africa borealis. Arabia. Belutschistan.

ATRACTYLIS MACROPHYLLA (Desf. *Fl. Atl.* ii. 255, tab. 226; DC. *Pr.* vi. 548, sub *Carlina*).

Mar. merid.—Ad radices Atlantis Majoris et in convallibus calidioribus. Aït Mesan! Seksaoua! Pr. Aïn Tarsil!

Pulehra species, 2-3-pedalis, habitu *Kentrophylli arborei*, in 'Fl. Atlantica' sat bene delineata. Ex spec. incompletis immaturis, mihi videtur vera *Atractylis*, nec inter *Carlinas* enumeranda.

Ar. Geog. Africa boreali-occidentalis.

CARDUUS (*L. Gen.* 925, pro parte).

C. MACROCEPHALUS (Desf. *Fl. Atl.* ii. 245; DC. *Pr.* vi. 621).

Mar. sept.—Djebel Dersa pr. Tetuan (*spec. ex Herb. Webb*)!

Mar. merid.—Specimen unicum immaturum in convalle Aït Mesan legit J. D. Hooker! Specimen in monte Djebel Afougueur ab indig. Ibrahim lectum benev. eomm. cel. Cosson!

Ar. Geog. Naseitur in Africa boreali, in regno Neapolitano, et in Græcia.

Cel. Boissier (*Fl. Or.* iii. 516) hunc cum *C. nutante* conjunxit. Pauci viri mihi videtur forma tam constans ut pro subspecie habeatur. Diu hæsi an sp. sequentem etiam pro subspecie haberem. Characteres certos ex involuero et fructu desumptos non video, sed habitus et ratio vitalis nimis diversa videntur.

C. BALLII (*H. fil.*; *Ball in Journ. Bot.* 1873, 368).

Planta biennis, subacaulis, scapos 3-4 simplices crassos, sæpius abbreviatis, rarius pedales, monocephalos, paucifolios emittens; herba tota araneoso-canescens; folia radicalia pinnatifida, segmentis lobato-dentatis margine spinosis, superiora (in scapis pl. macrioris) angusta, linearia, spinoso-dentata; involucrium regulariter imbricatum, 7-9-seriatum, foliolis omnibus acutis, adpressis, nervo prominulo in mucronem producto munitis, interioribus subscariosis; pappus elongatus.

Mar. merid.—In clivo septentrionali Atlantis Majoris. Seksaoua! circa 1200 m. Aït Mesan! a 1700 m. ad 2400 m. Djebel Tezah! a 1600 m. ad 2000 m.

Facie *Carduo* nullo mihi cognito, nisi forte *C. Schimperii*, Hochst., sp. Abyssinica, comparandus.

Ex anthodii structura in grege *C. nutantis* collocandus, sed evolutionis ratione omnino diversus est. In illo et in *C. macrocephalo* caulis evolutus ramosus, in nostro etiam dum scapi fiunt elongati semper simplices crassi sunt, ex folii basi obsolete decurrente line inde spinulosi. Involucri foliola magis adpressa numerosiora in series 7-9, nec 5-6, disposita. Floret ineunte Majo, mense citius quam *C. macrocephalus*.

In convalle Aït Mesan legi specimen unicum incertum foliis valde spinosis, forsā hybridum inter *C. Ballii* et aliam speciem.

CARDUUS MYRIACANTHUS (*Salzm. Exsicc.*; *DC. Pr.* vi. 624), = *C. confertus* (*Mor. St. Sard. Fl.* ii. 5, non *Guss.*, nec *Ten.*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb spec.*)! Inter Tanger et Tetnan!

Mar. occid.—Casa Blanca! Agadir (*Coll. Indig.* 1871)! Mogador (*Broussounet, ex DC. Prod.*).

Mar. merid.—Prov. Shedma! Mesfioua! Mtouga!

Ar. Geog. Ditio Maroccana. Hispania australis (nonnisi in arenosis isthmi Gihraltarici sec. Willkomm). Sardinia.

An satis differt *C. Renterianus* (*Boiss. Diagn. Pl. Or.* ser. 2, iii. 43), = *C. Bourgæanus* (*Boiss. et Reut. Pug.* 62, non *Sch. Bip.*), = *C. myriacanthus* (*Boiss. Voy. Esp.*), quocum teste cl. Lange conjungendus est *C. pteracanthus* (*Durieu, Expl. Alg. tab.* 50), planta per Hispaniam et Algeriam latiuscule diffusa, dijudicare non audeo.

C. PYCNOCEPHALUS (*Jacq. Hort. Viud.* i. 17, tab. 44; *Boiss. Fl. Or.* iii. 520, an *L.?*).

Mar. occid.—Huc verisimiliter spec. a cl. Lowe prope Mogador lectum (*C. tenuiflorus*, *Lowe Cat.*).

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia), necnon in Britannia et Gallia occidentali. In Europa centrali forsā eum seminibus introductus.

— —, var. TENUIFLORUS, *nob.*, ramis latius alatis, = *C. tenuiflorus* (*Curt. Fl. Lond.* vi. tab. 55). Confer *Boiss. Fl. Or.* iii. 521.

Mar. merid.—Mtouga!

— —, var. SARDOÛS, *Boiss. l. c.*, = *C. Sardous* (*DC. Pr.* vi. 626).

Mar. merid.—Mesfioua!

Species polymorpha, cui, suadente cl. Boissier, conjungendi sunt *C. arabicus*, *Jacq.*, *C. albidus*, *M. B.*, et *C. cinereus*, *M. B.*

CNICUS (*L. Gen.* 926).

Ducibus celeb. Bentham et Hooker (*Gen. Pl.* ii. 468) genus a recentioribus male mulcatum ad normam Linnæanam restitui. Auctores recentiores gravissimi nomen Tournefortianum *Cirsium Cuico* Linnæano anteposuerunt, sed definitionem genericam Tournefortianam servare non potuerunt.

C. GIGANTEUS (*Willd. Sp. Pl.* iii. 1671; *Desf. Fl. Atl.* ii. 245, tab. 221, sub *Carduo*), = *Cirsium giganteum* (*Spreng. Syst.* iii. 375; *DC. Pr.* vi. 637), = *Carduus scaber* (*Poir. Voy.* ii. 231).

Mar. sept.—Vidi nondum florentem pr. Tanger!

Ar. Geog. Africa borealis. Italia meridionalis. Sicilia. Sardinia. Hispania australis prope Fretum Hereulis.

Nomen Poirertianum antiquius a Fontanesio immerito mutatum, sed a recentioribus prætermissum, in lucem revocare nolui.

CNICUS ECHINATUS (*Desf. Fl. Atl.* ii. 247, sub *Carduo*; *Willd. Sp. Pl.* iii. 1668), = *Cirsium echinatum* (*DC. Fl. Fr.* v. 465, et *Prod.* vi. 638).

Mar. merid.—? Frustulum spec. nondum florentis in convalle Aït Mesan circ. 1800 m. legit J. D. Hooker!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

C. ORNATUS (*Ball in Journ. Bot.* 1873, 368).

Caulis erectus, 5-pedalis et ultra; folia radicalia bipedalia, anguste lanceolata, profunde sinuata, caulina subintegerrima, basi semiamplexicaulia, decurrentia, suprema bracteiformia, omnia pannoso-lanata, in pagina inferiore nervosa, margine ad terminum nervorum divergentium spinis validis aureis ornata; involucri foliola regulariter imbricata, dorso herbacea, margine minute ciliolata, exteriora breviter mucronata, interiora linearia, apice scariosa colorata; achenia (nimis immatura) indeskripta.

Mar. merid.—In Atlantis Majoris, convalle Amsmiz! circ. 1300 m.

Pulchra species a *C. flavispina*, Boiss. (sub *Cirsio*), foliis et involuero satis superque distincta, et insuper spinis multo longioribus validioribusque. Proximus erit, ni fallor, species Abyssinica nondum descripta (?), "Coll. Schimper No. 51," quæ præter cæteras notas differt involucri foliolis cartilagineis, margine latiuscule scariosis denticulatis.

C. CHRYSACANTHUS (*Ball in Journ. Bot.* 1873, 369).

Caulis erectus, 3-pedalis et ultra; folia inferiora pinnatifida, segmentis dentato-lobatis, superiora angulato-dentata, suprema bracteiformia, omnia ex pube fusca araneosa obscure virentia, spinis aureis superne sensim crescentibus armata; spinæ bractearum capitula involucrantium ultra pollicem longæ; involucri foliola irregulariter imbricata, subæqualia, exteriora mucrone longo rigido munita, interiora colorata mucrone brevi aureo apiculata; achenia (nimis immatura)?

Mar. merid.—In regione media Atlantis Majoris. In convalle Aït Mesan! a 1700 m. ad 1900 m.

Ex spinis aureis *C. ornato* nostro subsimilis, sed involuero foliisque omnino diversus. Propior est *Cnicus polyacantha*, Hochst. (sub *Cirsio*), species Abyssinica, qui differt (cæteris neglectis) involucri foliolis utrinque dentes 2–4 spinosos gerentibus.

Flora montana Abyssinica Maroccanæ nullomodo arete affinis est. Mirum igitur quod inter *Carduineas* tres species novæ a nobis detectæ omnes proxime ad typos Abyssinicos accedunt!

C. SYRIACUS (*L.*, sub *Carduo*; *Willd. Sp. Pl.* iii. 1683), = *Notobasis syriaca* (*Cass. Dict.* xxv. 225; *DC. Pr.* vi. 660).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Mar. merid.—Prov. Mtouga!

Ar. Geog. Regio Mediterranea calidior (inclusa Macaronesia). Deest in Gallia australi et in Italia superiore.

De genere *Notobasi* confer Benth. et Hook. Gen. Pl. ii. 469.

CNICUS CASABONÆ (L., sub *Carduo*; Willd. Sp. Pl. iii. 1682), = *Chamaepeuce Casabonæ* (DC. Pr. vi. 658).

Mar. merid.—Legi nondum florentem in convalle Aït Mesan supra Arround alt. circ. 2200 m. ! et vidi spec. in Herb. Kew. fere eodem loco a ell. Rein et Fritsch lectum ! Demum spec. ex Atlante Majore ab indigeno Ibrahim lectum vidi in Herb. Kew. a ecl. Cosson receptum !

Ar. Geog. Hucusque tantum in insulis Italicis (Sardinia, Corsica, Ilva) et in *Stæchadibus* detectus.

Spec. meum habui pro *C. hispanico* (Lam., sub *Carduo*) donec spec. florentia citata vidi. Re vera folia in utraque planta non distinguenda. In stirpe nostra capitula in paniculam contractam subthyrsoideam disposita sunt; in *C. hispanico* capitula pauca corymbosa. An pro subspecie habenda ?

ONOPORDON (*L. Gen.* 927).

O. ILLYRICUM (L., Boiss. Fl. Or. iii. 560; Willk. et Lge., Fl. Hisp. ii. 178), = O. elongatum (Lam. Fl. Fr. ii. 6; DC. Pr. vi. 618).

Mar. merid.—In rudertis urbis Marocco ! Prov. Mzouda !

Ar. Geog. Europa australis a Lusitania ad Bosphorum. Asia Minor. Syria. Mesopotamia. Ex Africa boreali, nostris exceptis, spec. nulla vidi.

O. MACRACANTHUM (*Schousb. Gew. Mar.* 184, tab. 5 et 6; DC. Pr. vi. 618).

Mar. sept.—Legi prope Tetuan (1851) !

Mar. occid.—Casa Blanca ! Prope Mogador ! In prov. Ithia haud procul Mogador !

Ar. Geog. Species floræ Maroccanæ et Hispaniæ propria, nec alibi detecta.

Folia in spec. nostris bicoloria, superne glabrescentia viridia, in pagina inferiore albo-lanata, inferiora pinnatifida; involucri foliola exteriora tomentosa vix spinosa, media in spinas longas, subulatas, vix canaliculatas, patentes reflexasve, abrupte desinentia.

O. SIBTHORPIANUM (Boiss. et Held. in Fl. Or. iii. 561), = O. macracanthum (Fl. Gr. tab. 832, non Schousb.), var. VIRIDE, nob.

Foliis in pagina superiore glabris viridibus.

Mar. occid.—Mazagan ! et forsan hue frustulum incompletum prope Mogador lectum !

Ar. Geog. Species in Oriente latiuscule diffusa, ejus var. in Algeria interiore nascitur. Varietas nostra in litore Atlantico Imp. Marocani.

Nostræ plantæ valde proximum est "*O. ambiguum*," Balansa, Pl. d'Algérie, No. 794; sed hoc (monente eel. Boissier) ad *O. Sibthorpi-
num* referendum est.

Ab *O. macracantho* videtur toto cœlo diversa planta a nobis ad Mazagan lecta, gaudet spinis involueralibus dimidio brevioribus, squamis sensim nec abrupte acuminatis. Attamen hæ notæ videntur variabiles, et plurimæ species hujus generis incertæ mihi manent.

CYNARA (*L. Gen.* 928).

C. CARDUNCULUS (*L., DC. Pr.* vi. 620).

Mar. occid.—In parte superiore montis Djebel Hadid haud procul Mogador, loco a cultura remoto!

Nisi fallor, vidimus nondum florentem in regione inferiore Atlantis Majoris.

Ar. Geog. Per regionem fere omnem Mediterraneam (inclusa Macaronesia) dispersa sed sæpe introducta. Planta spontanea ubique rara.

C. HYSTRIX (*Ball in Journ. Bot.* 1873, 369).

Caulis erectus, gracilis, foliosus, 3-pedalis et ultra; folia inferiora longissima, pinnata, pinnis pinnatifidis laciniatis, rachi inter pinnas dentibus linearibus acutis herbaceis apice spinosis munita, superiora sensim decrescentia, omnia, præsertim in auriculis semiamplexicaulibus profunde laciniatis, spinis tenuibus acutissimis armata, in pagina inferiore, cum caule tereti gracili, albo-tomentosa; involucri foliola exteriora angusta adpressa, media cartilaginea e basi lata in mucronem (etiam in præfloreatione) patentem acuminata, interiora recta; achenia ignota.

Mar. merid.—Specimen unicum legi in collibus dittonis Reraya, haud procul Mulai Ibrahim!

Plantam nostram, etsi immaturam, inter *Cynaras* reponendam esse nihil dubito. A cæteris mihi cognitis omnino diversa est. Foliorum et pinnarum lacinia suprema integra linearis, insigniter elongata.

C. HUMILIS (*L., DC. Pr.* vi. 620), = *Bourgæa humilis* (*Coss. Pl. Crit.* 39).

Mar. sept.—Valde frequens in ditone Tingitana, ubi eam jam legerant Webb, Salzmann, cæterique botanici! Legi prope Tetuan (1851)!

Mar. occid.—Vidimus in monte Djebel Hadid haud procul Mogador!

Mar. merid.—Occurrit line inde in regione inferiore, sed non ubique.

Mesfoua! Specimen prope Ourika lectum in Herb. Kew. (*R. et Fr.*)!

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

SILYBUM (*Gærtn. Fruct.* ii. 378).

S. MARIANUM (*L., sub Carduo; Gærtn. l. c.; DC. Pr.* vi. 616).

Mar. sept.—Prope Tanger et Tetuan pluries visum!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). In Europa media et occidentali, verisimiliter introductum.

GALACTITES (*Mæneh, Meth.* 558).

G. TOMENTOSA (*Mæneh, l. c.*; *DC. Pr.* vi. 616), = *Centaurea Galactites, L.*

Mar. sept.—Tanger!

Ar. Geog. In Europa Mediterranea fere omni late dispersa. Africa borealis. Macaronesia.

STÆHELINA (*L. Gen.* 938).

S. DUBIA (*L., DC. Pr.* vi. 544).

Mar. sept.—In rupibus juxta Tetuan (1851)!

— —, var. capitulis (saltem virginicis) basi foliis cinctis. An forma inter *S. dubiam* et *S. bætican*, *DC.*, media?

Mar. merid.—Legimus mox floribundam in convalle Amsmiz, 1500–1700 m.!

Ar. Geog. *S. dubia*, per Europam austro-occidentalem a Dalmatia usque Lusitaniam dispersa, deest in Algeria et in Macaronesia.

CRUPINA (*Cass. Dict.* xii. 67).

C. VULGARIS (*Cass. Dict.* xlv. 39; *DC. Pr.* vi. 565), = *Centaurea crupina (L., Desf. Fl. Atl.* ii. 292).

Mar. merid.—Prope Seksaoua!

Ar. Geog. Species per Europam australem et Orientem dispersa, et inde per regionem Danubialem et Rossiam australem usque Soongariam. Ex Africa, præter spec. nostra, nunquam vidi.

VOLUTARELLA (*Cass. Bull. Philomath.* 1816; *Benth. et H. f. Gen. Pl.* ii. 476).

V. LIPPII (*L., sub Centaurea*; *Cass. Dict.* xlv. 39), = *Amberboa Lippii (DC. Pr.* vi. 559, et *auct. plur.*).

Mar. occid.—Prov. Haha haud procul Mogador! Aïn el Hadjar!

Mar. merid.—Shedina! In planitie et in rupibus arenaceis haud procul urbe Marocco!

Ar. Geog. Per regionem Mediterrancam calidiorem ex Arabia Petræa usque Macaronesiam diffusa. In Europa nunci in Hispania austro-orientali.

V. CRUPINOIDES (*Desf. Fl. Atl.* ii. 293, sub *Centaurea*; *DC. Pr.* vi. 559, sub *Amberboa*), = *Volutarella bicolor (Cass. Dict.* l. 250), = *Lacellia libyca (Viv. Fl. Lib.* 58, tab. 22).

Mar. occid.—Frustulum hujus speciei ex monte Djebel Hadid haud procul Mogador attulerunt Rein et Fritsch!

Mar. merid.—Legimus in rupibus supra Seksaoua ad radices Atlantis Majoris! Spec. ex Oasi Akka ab indigeno Mardocheo lectum benev. comm. cel. Cosson!

Ar. Geog. Africa borealis a Mari Rubro usque litora Oceani. Arabia. Palæstina.

Pace celeberr. Benth. et H. fil. hæc mihi videtur a priore probe distincta species. Video folia glabra *Crupinæ vulgaris*, involucri subglaberrimi (basi interdum arancoso-tomentosi) squamas lincis fuscis notatas, apice nigrescentes, sensim acutas, nec abrupte acuminatas cuspidato-mucronatas. Corollæ disci crocæ sunt, marginales læte purpureæ. Achenium, in utraque specie hilo laterali præditum, in *V. Lippii* villosum tenuiter rugosum vix foveolatum, in *V. crupinoide* minus villosum, inter costas insigniter foveolatum.

VOLUTARELLA LEUCANTHA (*Coss. Ind. Pl. Maroc.* 21, sub *Amberboa*). Mar. merid.—? Ex provincia Mtouga, spec. incompletum in Herb. Kew. asservatum attulerunt ell. Rein et Fritsch!

Species mihi parum cognita a cel. auctore nondum descripta. Monente cel. Cosson nascitur in Imp. Maroccano Transatlantico, in Oasi Akka, ubi eam legit indigenus Mardocheus.

V. MURICATA (*L.*, sub *Centaurea*; *DC. Pr.* vi. 559, sub *Amberboa*), = *Cyanopsis radiatissima* (*Cass. Dict.* xlv. 39).

Mar. merid.—Mesfioua! Prov. Mtouga ad fines prov. Shedma!

Ar. Geog. Hispania australis. Marocco meridionalis.

LEUZEIA (*DC. Fl. Fr.* iv. 109).

L. CONIFERA (*L.*, sub *Centaurea*; *DC. l. c., Prod.* vi. 665.).

Mar. sept.—In saxosis prope Tetuan (1851 et iterum 1871)! In monte Djebel Dersa prope Tetuan (*Webb, Herb.*)!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

CENTAUREA (*L. Gen.* 981; *Benth. et H. fil. Gen. Pl.* ii. 478).

C. TAGANA (*Brot. Fl. Lus.* i. 369, et *Phyt. Lus.* tab. 32; *DC. Pr.* vi. 566).

Mar. sept.—In monte Djebel Kebir et alibi in collibus prope Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! (*Schousb. Herb.*)!

Ar. Geog. Lusitania. Hispania australis. Sicilia (unde mihi spec. comm. cl. Todaro) Ager Tingitanus.

Hæc mihi videtur mera var. *C. africanæ* (*Lam. Dict.* i. 664), quæ vix differt nisi foliis inferioribus bipinnatifidis. In pl. nostra folia simplicia margine crenulato-serrata, vel basi grosse dentata, vel inferiora pinnatifida.

C. CHAMÆRHAPONTICUM, *nob.*, = *Cynara acaulis* (*L., Desf. Fl. Atl.* ii. 249, tab. 223), = *C. humilis* (*Juss. Gen.* 173, non *L.*), = *Rhaponticum acaule* (*DC. Pr.* vi. 664), = *Serratula acaulis* (*DC. Diss. Comp.* 31), = *Cestrinus carthamoides* (*Cass. Dict.* viii. 24).

Mar. occid.—? Folia radicalia fere certe huc referenda prope Agadir ab indigenis lecta anno 1871!

Mar. merid.—Occurrit haud raro in saxosis aridissimis planitici Maroccanæ. Prov. Shedma plurimis locis! Prope Seksaoua!

Ar. Geog. Africa borealis ad fines reg. desertæ. Nasceitur etiam in insula Cypro fide Sibthorp (Prod. Fl. Gr. ii. 158); sed a recentioribus non visa, nec enumeratur in Boiss. 'Fl. Orientali.'

Vidimus tantum sp. deflorata et subexsiccata. Floret verisimiliter hieme vel ineunte vere.

CENTAUREA COSSONIANA (Ball in Journ. Bot. 1873, 369), = *Rhaponiticum cauleseens* (Coss. in Bull. Soc. Bot. Fr. xx. 251).

Descriptionem Cossonianam proferre licet:

"Planta *perennis*, forma foliorum *Rhaponiticum acaule* exacte referens. Caudex crassus in radicem fusiformem abiens, superne basi petiolorum emareidorum squamæformium stipatus. *Caules* 1 vel 3–4, *folia plus minus superantes vel brevissimi*, simplices, erecti, teretes, striati, pube arachnoidea deteribili donati, demum glabriusculi, folia 1–2 gerentes. *Folia* supra pubescenti-floccosa, subtus cano-tomentosa; *radicalia* sæpius plurima, petiolata, *pinnatisecta segmentis* ambitu oblongo-vel ovato-cuneatis *inciso-pinnatifidis in rachim alato-folia-ceum* hinc inde dentatam *decurrentibus*; *caulina* 1–2, inferius segmentis sæpe integris, superius ad petiolum squamæformem reductum. Capitula magna, apice caudam solitaria, foliis floralibus haud involucreta. *Involucrum ovato-campanulatum, foliolis* pluriseriatis arete imbricatis, chartaceis, dorso superne subherbaceo striatis, *apice in appendicem scariosam ovatam vel oblongo-lanceolatam irregulariter lacera-*ram tenuissime ciliatam *dilatatis*, inferioribus multo minoribus lanceolatis dorso pubescenti-floccosis, mediis oblongo-lanceolatis, interioribus lanceolato-linearibus. Receptaculum piliferum pilis achæniis longioribus. *Flosculi omnes hermaphroditi, æquales*, pallide purpurascens? tubo longissimo, limbo ultra medium 5-fido laciniis angustis linearibus. *Filamenta papillosa. Antheræ basi sagittatæ*, auriculis contiguis connatis, caudis integris vel vix laceris, appendice terminali oblonga obtusa. Styli rami erecti, connati imo, apice obtusiusculo tantum liberi. *Achænia glabra, areola insertionis obliqua*, oblonga, *compressa*, obsolete pluristriata, superne paululum contracta margine prominulo brevissimo annulæformi denticulato coronata. *Pappi albi*, tubam flosculorum superantis *setæ* pluriseriatæ, *valide scabræ, fere barbellatæ*, sensim longiores, exteriores multo breviores, *interiores longiores conformes*."

Mar. occid.—Monente ecl. Cosson hæc planta nuper a mulione Ibrahim in monte Djebel Hadid, haud procul Mogador, lecta fuit.

Mar. merid.—Primus hanc stirpem legit Balansa in monte Djebel Orguis prope Imintenout. Legimus formam subacaulem in saxosis prope

Seksaoua! et formam caulescentem in monte supra Seksaoua ad 1600 m.!

Stirps insignis, quam nomine amicissimi Cosson, floræ Africæ borealis magistri, decorare proposui, fere eodem tempore quo sub *Rhapontico caulescente*, Coss. l. c., publici juris facta est. Quoad citatum "Coss. et Bal. in Balansa, Pl. Maroc. (1867)," hoc certe non valet, dum in itinere suo el. Balansa perpaucæ specimina legere potuit, et planta nostra in omnibus fere herbariis deest. Affinitatem proximam hujus eum *Rhapontico acauli* (quod duobus Benth. et H. fil. inter *Centaureas* enumeravi) nemo negabit, sed in hoc involueri structura valde diversa. In sp. nostra foliola involueri in seriis 8-10 æquidistantibus regulariter umbricata, exteriora lanceolata acuta margine chartacea exappendiculata, interiora appendice scarioso-fimbriata munita.

Achenia matura prismatica 5-costata, costis intermediis quinque subobsoletis. Pappi setæ demum purpurascentes scabræ, interiores plumosæ, plumulis deciduis.

Qui nobiseum animo haud æquo genus *Rhaponticum*, Lam., ad sectionem *Centaureæ* reductum a Bentham et Hooker (in Gen. Plant. ii. p. 479) observaverit, hanc speciem (a *Rhaponticis* nullo pacto dividendam) ad sententiam virorum illust. confirmandam agnoscere debet. Characterem technico est enim *Fornicio*, Cass., vel *Chartolepide*, DC., propior quam *Rhapontico*, a quo involuero et pappi setis interioribus plumosis longe distat, etiamsi facie simillima.

CENTAUREA CLEMENTEI (Boiss. ap. DC. Pr. vii. 303, et Voy. Esp. 346, tab. 102).

Mar. sept.—Prope Tetuan in monte Beni Hosmar, ubi jam legerat P. B. Webb!

Ar. Geog. In Hispaniæ australis montibus Sierra de Yunquera et in montibus Tetuanensibus.

Pulchra species a proxima *C. Ragusina*, L., satis superque distincta.

C. PULLATA (L., DC. Pr. vi. 577).

Mar. sept.—Tanger! (Webb Herb.)! Cap. Spartel! Inter Tanger et Tetuan! Prope Tetuan (1851)! Inter Tetuan et Ceuta!

Mar. occid.—Casa Blanca!

Mar. merid.—Milhaïn! Taradut (R. et Fr. spec. in Herb. Kew.)!

Ar. Geog. Regio Mediterranea imprimis occidentalis (exclusa Macaronesia).

— —, var. MINOR. Planta omnibus partibus minor, floeulis radiantibus et appendicibus foliolorum involueri brevioribus.

Mar. merid.—In saxosis apricis regionis inferioris haud infrequens. Prov. Shedma pluribus locis! Mzouda! Seksaoua!

C. INCANA (Desf. Fl. Atl. ii. 301, non Ten. nec Lag.), = C. pubescens (Willd. Sp. Pl. iii. 2322; DC. Pr. vi. 589).

Mar. merid.—Legimus nondum florentem in convalle Aït Mesan ad 1800 m. s. m. ! et prope Seksaoua !

—— —, var. *HOOKERIANA* (Ball in Journ. Bot. 1873, 370).

Primo intuitu a *C. incana* differt capitulis multo majoribus, flosculis aurantiaco-purpureis, involucri foliolorum spina media longiore, demum recurvata, utrinque spinulosa. Habui pro specie nova, donec specimina permulta hujus gregis diligenter explorata notas specificas lucusque indicatas fallaces esse me docuerunt.

Mar. merid.—In plautie Maroccana ad fines prov. Mtouga et Shedma !

Ar. Geog. *C. incana*, Desf., est species Africæ borealis propria ; species proxime affines per Orientem et Hispaniam diffusæ sunt.

Nomen *C. incana*, Desf., a Willdenovio ipso sub *C. pubescente* sua citatum, nescio quo pacto a recentioribus prætermisum, hic restitui.

Notæ differentiales quibus hæc et plurimæ species affines a *C. reflexa*, Lam., separantur mihi videntur nimis variabiles ; et sicut subspecies, sensu nostro, enumerandæ sunt *C. ornata*, W., *C. saxicola*, Lag., *C. omphalotricha*, Coss. et Dur., et *C. psilacantha*, Boiss. et Held. His forsân addenda est *C. persica*, Boiss. Hujus gregis varietas nostra regina est. Statura robusta 2-4-pedalis ; capitula magna colore ab aurantiaco ad atropurpureum variantia ; achenia basi copiose sericeo-pilosa ; pappus ut in *C. incana* typica longitudine medioeris, demum fuscus.

CENTAUREA CALCITRAPA (L., DC. Pr. vi. 597).

Mar. merid.—Prov. Shedma ! Prope Amsmiz !

Ar. Geog. Europa media. Regio Mediterranea fere omnis (inclusa Macaronesia). In ditione nostra videtur rara, et in agro Tingitano a botanicis plurimis perlustrata nondum detecta est.

C. MELITENSIS (L., DC. Pr. vi. 593), = *C. Apula* (Lam. Dict. i. 674 ; DC. Pr. l. c.).

Mar. sept. Prope Tetuan (1861) !

Mar. merid.—Mesfoua ! Mzouda !

Ar. Geog. Regio Mediterranea imprimis occidentalis (inclusa Macaronesia). Deest in Asia Mediterranea continentali et in Ægypto. Hæc species obvia est in Africa australi, in America australi extratropica (Chili, Bonaria, Monte Video, Brasilia), in California (Douglas), et in Insula Paschali (Chamisso). Dicitur introducta ; plantam tamen in Europa nullibi frequentem vix in tantis regionibus longinquis hominum opere allatam esse crediderim.

C. ERIOPHORA (L., DC. Pr. vi. 592).

Mar. occid.—Casa Blanca ! Agadir (Coll. Indig. 1871) !

Mar. merid.—Prov. Shedma !

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

CENTAUREA SULPHUREA (*Willd. Enum. No. 391; DC. Pr. vi. 593, non Lag.*).

Mar. merid.—Occurrit satis frequens ad radices Atlantis Majoris. Tassereimout! Mesfioua! Reraya! Amsmiz! Seksaoua!

Ar. Geog. Hispania. Africa borealis.

C. MAROCCANA (*Ball in Journ. Bot. 1873, 370*).

Planta humilis, dichotome ramosa; folia radicalia (et interdum caulina inferiora) pinnatifida, lobis angulato-dentatis; caulina lanceolato-vel linear-oblonga, integra acute denticulata; longe decurrentia; capitula terminalia inter folia suprema fulcrata; inflorescentia impia (scil. rami seniores abbreviati a junioribus superati); involucri ovato-conici elongati foliola nitida, cartilaginea, spina valida utrinque spinulosa coronata; foliolorum mediorum spinæ elongatæ, internorum abbreviatæ; achenia matura epapposa.

Mar. merid.—Sat frequens in regione arida subdeserta inter Mogador et urbem Marocco. Legimus pluribus locis in prov. Shedma, necnon ad radices Atlantis juxta Milhaïn!

Species fere media inter *C. eriophoram*, L., et *C. sulphuream*, W., ab utraque mihi videtur sat diversa; structura priori, facie tamen *C. sulphureæ* propior. A *C. eriophora* differt statura multo minore, capitulis minoribus dimidio angustioribus nitidis, nec basi arancoso-pubescentibus, spinis foliolorum involucri validioribus. *C. sulphurea* gaudet foliolis involucri omnino diversis, appendice spinis radiantibus coronata, spina media fusca quam cæteræ paulo longiore (fere ut in § *Seridia*), dum in *C. maroccana* spina media valida est potius subpinnatim spinulosa, spinulis utrinque sparsim dispositis. A *C. sicula*, L., structura et habitu longius distat. Nostra species est affinibus præcoecior; achenia jam matura vidimus dum *C. sulphurea* nondum florebat.

C. SERIDIS (L., *DC. Pr. vi. 599*), var. AURICULATA, = *C. auriculata* (*Balb. in Pers. Syn. ii. 91*).

Mar. occid.—In maritimis juxta Mazagan!

Ar. Geog. Hispania. Sardinia. Africa boreali-occidentalis.

Spec. nostra cum Sardois a cl. Mueller lectis bene conveniunt et inter *C. seridis* et *C. sonchifolia* ambigua sunt. Revera *C. sonchifolia* videtur esse tantum subspecies *C. seridis*. Forma vegetior foliis caulinis longe decurrentibus est *C. maritima* (L. Duf.; *DC. Pr. l. c.*). *C. seridis* typica, si bene hanc novi, nascitur in arenosis juxta S. Roque pr. Promont. Calpes.

C. POLYACANTHA (*Willd. Sp. Pl. iii. 2312; DC. Pr. vi. 600*).

Mar. sept.—In monte Djebel Kebir juxta Tanger! (*Salzm. Exsicc.*, sub *C. sphærocephala*)! Cap. Spartel! Inter Tanger et Tetuan (*Webb Herb.*)!

Ar. Geog. Species propria ditionis Tingitanæ et Hispaniæ australis juxta Fretum Herculis.

CENTAUREA ASPERA (L., *DC. Pr.* vi. 599).

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—In prov. Shedma, ut videtur, infrequens! Nuper habui a eel. Cosson spec. ab indigeno Mardocheo inter Mogador et Ilhir lectum!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

C. FRAGILIS (*Durieu in Rev. Bot.* ii. 429; *Walp. Ann.* i. 450).

Mar. sept.—Tanger (*Spec. ex Herb. Schousb.*, sub *C. sphærocephala*)! Cap. Spartel! In rupibus juxta Tetuan 1851 et 1871! In monte Djebel Dersa pr. Tetuan (*Webb Herb.*)!

— —, var. *INTEGRIFOLIA*, nob. Differt foliis etiam inferioribus subintegerrimis, caulinis obverse lanceolatis integerrimis basi semi-amplexicaulibus, supremis linearibus, ramis eximie angulatis subalatis. Valde similia sunt spec. a Balansa prope Aïn el Turk lecta.

Mar. occid.—Prope Mogador! In collinis prov. Iaha laud procul Mogador!

Ar. Geog. Africa boreali-occidentalis.

C. DILUTA (*Ait. Hort. Kew.* iii. 261; *DC. Pr.* vi. 591), = *C. elongata* (*Schousb. Gew. Marok*, p. 186, ex *Willd. Sp. Pl.* iii. 2300), = *C. raphanifolia* (*Salzm. Exsicc.* 1825). Verisimiliter eadem est *C. elata* (*Poir. Dict. Suppl.* ii. 150).

Mar. sept.—Legi florentem pr. Tanger 13 Jun.!(*Salzm. Exsicc.*)! (*Webb Herb.*)! Tetuan (1851)!

Ar. Geog. Species propria ditionis Tingitanæ.

C. SALMANTICA (L., *Desf. Fl. Atl.* ii. 303; *Benth. et H. fl. Gen. Pl.* ii. 482), = *Microlonchus salmanticus* (*DC. Pr.* vi. 563), = *Mantisalca elegans* (*Cass. Dict.* xxix. p. 31).

Planta polymorpha in species plurimas notis diagnosticis instabilibus definitas a recentioribus dilaniata. Characteres a pappo desumpti certe fallaces. Interdum achenia omnia papposa sunt, interdum exteriora calva; et hoc vidit etiam beatus J. Gay in spec. Gallicis *M. Duriæi*, Spach. Ex ditione nostra video varietates duas non semper facile distinguendas.

— —, var. *a. CLUSII*, = *Microlonchus Clusii* (*Spach in Ann. Sc. Nat.* 3^{me} ser. iv. 167; *Willk. et Lge. Fl. Hisp.* ii. 167).

Distinguitur capitulis majoribus, basin versus dilatatis, foliolis involucri apicem versus leviter pubescentibus mucrone brevi cito deciduo armatis.

Mar. sept.—Tetuan (1851)! (*Herb. Webb*)!

Mar. occid.—Casa Blanca!

Ar. Geog. Per regionem omnem Mediterraneam late dispersa. Fit rara in Oriente, adest in Madera sed in ins. Canariensibus nondum detecta.

— —, Var β . LEPTOLONCHUS, = *Microlophus leptolophus* (Spach, l. c.).

Planta gracilior gaudet capitulis angustioribus, foliolis involucri glaberrimis, mucrone subpersistente.

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Satis frequens in regione inferiore et ad radices Atlantis Majoris. Shedma! Sheshaoua! Prope urbem Marocco! Mesfioua! Aït Mesan! Amsniz! Seksaoua!

Ar. Geog. Hæc forma, quoad sciam, nonnisi in Africa boreali obvia est.

CARTHAMUS (*L. Gen.* 931; *Benth. et H. fil. Gen. Pl.* ii. 483).

C. LANATUS (*L., Boiss. Fl. Or.* iii. 706), = *Kentrophyllum lanatum* (*DC. Bot. Gall.* i. 293, et *Prod.* vi. 610), var. ELEGANS, nob. Folia omnia indivisa semiamplexicaulia, subtus eximie nervosa, nervis lateralibus intermediis usque ad apicem, exterioribus ultra mediam folii longitudinem, productis; involucri foliola exteriora foliacea ciliato-spinosa subpatentia, interioribus æquilonga nec longiora, intima angusta chartacea nervosa apicem versus ciliato-spinosa. Achenia immatura.

Mar. merid.—Spec. unicum prope Seksaoua legit J. D. H.! alterum legi in prov. Mtouga! et eandem plantam in prov. Mtouga legerunt Rein et Fritsch!

Pluribus notis planta nostra eum *Kentrophyllis bætico* (*Boiss. et Reut. Pug.* 65) convenit, et forsitan melius observata pro sp. distincta habenda erit.

Ar. Geog. Regio Mediterranea omnis (excl. Ins. Canariensibus). Occurrit etiam in calidioribus Europæ mediæ et orientalis.

C. CÆRULEUS (*L., Boiss. Fl. Or.* iii. 710; *DC. Pr.* vi. 615, sub *Carduncello*), var. TINGITANUS, = *C. tingitanus* (*L., Desf. Fl. Atl.* ii. 256).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)!

Mar. merid.—In convalle Aït Mesan a 1700 m. usque 2200 m.!

Ar. Geog. *C. cæruleus* per regionem omnem Mediterraneam (inclusa Macaronesia) dispersus est. Var. nostra, quæ differt tantum foliis pinnatifidis, fere ubique eum specie obvia.

CARDUNCELLUS (*Adans. Fam. Pl.* ii. 116).

C. PINNATUS (*Desf. Fl. Atl.* ii. 258, tab. 229, sub *Carthamo*; *DC. Pr.* vi. 614), = *Onobroma pinnatum* (*Spreng. Syst.* iii. 319).

Mar. sept.—Inter Tanger et Tetuan! (*Webb Herb.*)!

Ar. Geog. Africa borealis. Sicilia.

C. lucens (*Ball in Journ. Bot.* 1873, 370).

Acaulis; folia primaria integra ciliato-spinosa, sequentia pinnatipartita,

lobis eximie pectinato-spinosis; involucri foliolorum appendix late scarioso-pectinata, intimorum anguste marginata; achenia (immatura) videntur lævia.

Mar. merid.—In regione superiore Atlantis Majoris. In convalle Aït Mesan! a 2000 m. ad 2540 m.

Vivus a *C. pinnato* statim dignoscitur spinis longioribus albis lucentibus. Foliola involucri gaudent appendicibus longius pectinatis, et achenia videntur diversa. Melius cognitus pro specie omnino diversa facile accipiendus.

Nuper habui a cel. Cosson specimina in monte Djebel Ouensa ab indigeno Ibrahim lecta sub "*C. pinnatus*, Desf.," quæ certe ad subspeciem nostram spectant et appendicibus pectinatis gaudent.

SCOLYMUS (*L. Gen.* 922).

S. MACULATUS (*L., DC. Pr.* vii. 75).

Mar. sept.—Tanger! (*Webb Herb.*)!

Mar. merid.—In via qua itur a Shedma ad Mogador (*G. Maw sp.*)!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). In Nubia etiam obvia.

S. HISPANICUS (*L., DC. Pr.* vii. 76).

Mar. sept.—Tanger (*Webb Herb.*)! Tetuan (1851)!

Mar. occid.—Casa Blanca! Mogador! (*Lowe Cat.*).

Mar. merid.—Sat frequens in regione inferiore prov. Shedma! Prov. Mtouga! Ain Tarsil! Keira!

Ar. Geog. Regio Mediterranea fere omnis. In ins. Tencriffa obvia, deest in Madera.

CATANANCHE (*L. Gen.* 920).

C. CÆRULEA (*L., DC. Pr.* vii. 83).

Mar. merid.—Frequentissime occurrit ad radices et in convallibus calidioribus Atlantis Majoris. Aït Mesan! Reraya! Sektana! Mtouga (*R. et Fr. spec.*)!

Statura et foliorum forma valde ludibunda; folia fiunt integra, vel sinuato-dentata, vel grosse repando-dentata subpinnatifida.

— —, var. *TENUIS*. Differt statura minore, indumento argenteo-sericeo, foliis angustis linearibus integris, vel hinc inde lobis linearibus inæqualibus instructis.

Mar. merid.—In convalle Aït Mesan a 1500 m. ad 1700 m.! Supra Amsmiz! In rupibus supra Seksaoua!

Ar. Geog. *C. cærulea* per regionem Mediterraneam occidentalem (exclusa Macaronesia) dispersa est. Ex Galloprovincia usque in montosis Delphinatus se protendit.

C. CÆSPITOSUM (*Desf. Fl. Atl.* ii. 238, tab. 217; *DC. Pr.* vii. 83).

Mar. merid.—In monte Djebel Tezah a 2200 m. usque 2800 m.!

Ar. Geog. Africa borealis.

Ludit foliis obverse lanceolatis subintegerrimis, vel apicem versus cuneato-dilatatis 3-5-fidis, lobis acute dentatis.

CATANANCHE LUTEA (L., DC. Pr. vii. 83), = *Piptocephalum phaiolepis* (Sch. Bip. in *Bonplandia*, 1860, 359).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Specimen a Broussonnet in Herb. Kew.*)! Nos non vidimus.

Ar. Geog. Regio Mediterranea calidior ex Syria usque Hispaniam australem. Deest in Gallia, in Italia media et superiore, in Dalmatia, necnon in Macaronesia.

Cel. Schulz Bip. formam occidentalem, capitulis majoribus insignem, ab orientali (*P. luteum*, Sch. Bip.) distinxit, et invita natura ex his genus novum condidit.

C. MONTANA (Coss. et Dur. in *Bal. Pl. d'Algérie*)?

Mar. sept.—In monte Djebel Dersa prope Tetuan beatus P. B. Webb spec. legit quæ verisimiliter huc referenda sunt. *C. montana*, a ell. auctoribus nondum descripta, verisimiliter in ditione nostra alibi occurrit.

Ar. Geog. Africa boreali-occidentalis.

CICHORIUM (L. Gen. 921).

C. INTYBUS (L., DC. Pr. vii. 84).

Mar. merid.—Saksaoua!

Ar. Geog. Europa omnis (exclusa arctica). Sibiria. In regione Mediterranea occidentali minus frequens, et deest in Macaronesia.

Specimen nostrum incompletum et haud omnino certum.

C. pumilum (Jacq. Obs. iv. 3, tab. 80; Fl. Gr. tab. 822; DC. Pr. vii. 84), = *C. divaricatum* (Schousb. Gew. Marok, 184; Boiss. Fl. Or. iii. 716), = *C. Intybus*, var. *β. divaricatum* (DC. Pr. vii. 84).

Mar. sept.—Prope Tetuan ad ostium fluminis (1851)!

Mar. merid.—Pluries visum, legimus in prov. Shedma! Spec. ad Aït Ibrahim lectum ab indigeno Mardocheo benev. comm. cel. Cosson! Monente cel. Boissier stirps orientalis *C. pumilum* a *C. divaricato* occidentali non differt, et nomen antiquius prioritate gaudet. Hoc mihi videtur prioris subspecies facta annua vel interdum biennis.

Ar. Geog. Regio Mediterranea fere omnis, in Græcia et Dalmatia non indicatum. In Madera obvium, deest in Ins. Canariensibus.

HYOSERIS (L. Gen. 916, pro parte).

H. RADIATA (L., DC. Pr. vii. 79).

Mar. sept.—Tanger (*Webb Herb.*)! Tetuan (1851)! Inter Tanger et Tetuan!

Mar. merid.—Satis frequens ad radices Atlantis Majoris. Saksaoua! Milhaïn! et alibi pluries visa!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

TOLPIS (*Adans. Fam. Pl.* ii. 112).

T. BARBATA (*Gærtn. Fruct.* ii. 372, tab. 160; *DC. Pr.* vii. 86).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Tetuan (1851)!

Mar. occid.—Casa Blanca! Mogador! (*Lowe Cat.*).

Mar. sept.—In collibus prov. Haha!

— —, var. *GRANDIFLORA*, *nob.* Planta humilis, foliis quam in typo duplo latioribus exsiccato-dentatis, capitulis magnis, flosculis centralibus purpureis.

Mar. sept.—Cap. Spartel! Vidi hanc var. in altero litore Freti Herculis haud procul Algeciras.

Ar. Geog. *T. barbata* Hispaniæ et Africæ borealis civis est. In Macaronesia deest.

T. UMBELLATA (*Bert. Am. Ital.* 96; *DC. Pr.* vii. 86).

Mar. occid.—Mogador (*Lowe sp. in Herb. Kew.*, sub *T. crinita*)!

Mar. merid.—Haud infrequens in regione inferiore et in convallibus calidioribus Atlantis Majoris. Prov. Shedma! Aïn Oumast! Mesfiona! Ourika! Aït Mesan! Sektana! Seksaoua!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

Stirps valde ludibunda sæpius gaudet caule unico umbellatim ramoso.

Forma a basi ramosa simillima est *T. barbata*; ab hac differt foliis basin versus profunde dentatis subpinnatifidis et acheniis quadrisetosis.

In *T. barbata* achenia (an semper?) bisetosa.

RHAGADIOLUS (*Juss. Gen.* 168; *Benth. et H. fil.*

Gen. Pl. ii. 510).

R. STELLATUS (*L.*, sub *Lapsana*; *Willd. Sp. Pl.* iii. 1625; *DC. Pr.* vii. 77).

Mar. merid.—Videtur satis frequens in regione inferiore et in convallibus Atlantis Majoris. Misra Ben Kara! Tasseremout! Aït Mesan ad 1300 m. s. m. In convalle Amsniz circ. 1400 m.!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

R. HEDYPNOIS (*Coll. Fl. Ped.* No. 831; *Benth. et H. fil.* ii. 510). Inclusis *Hedypnoide cretica* (*DC. l. c. Pr.* vii. 81), *H. coronopifolia* (*DC. l. c.*), et *H. polymorpha* (*DC. l. c.*).

Mar. sept.—Tanger! Inter Tetuan et Ceuta!

Mar. occid.—Mogador! (*Lowe Cat.*).

Mar. merid.—Frequens in regione inferiore. Prov. Shedma! Prov. Mtouga! Mesfiona! et alibi obvius.

Stirps polymorpha, ejus formæ aspectu valde diversæ instabiles sunt.

Synonyma auctorum permulta sed plurima incerta.

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

R. pendulus (*Balb. in Pers. Syn.* iii. 369, sub *Hyoseride*; *DC. Pr.* vii. 82, sub *Hedypnoide*; *Benth. et H. fil. Gen. Pl.* ii. 510).

Mar. sept.—Tanger!

Mar. occid.—Casa Blanca! Circa Mogador frequentissime! In prov. Haha in reg. litorea! Ain el Hadjar haud procul Mogador! (*Lowe spec. in Herb. Kew.* prope Mogador lectum sub *II. arenaria*)!

Ar. Geog. Verisimiliter per regionem Mediterraneam fere omnem (exclusa Macaronesia) occurrit.

Cel. Boissier hanc stirpem ad varietatem prioris reduxit (conf. *Fl. Or.* iii. 719); ego tamen pro subspecie habui, dum characteres stabiliores præbet et formæ intermediæ vix inveniendæ. Pedunculi elongati graciles, sub anthesi sæpius (non semper) penduli, fructiferi parum incrassati. Ligulæ sæpius saturate luteæ, interdum aurantiacæ.

RHAGADIOLUS ARENARIUS (*Schousb. Gew. Marok*, 183, sub *Hyoseride*; *DC. Pr.* vii. 82, sub *Hedypnoide*; *Benth. et H. fil. Gen. Pl.* ii. 510).

Mar. sept.—Tanger (*Salzm. Exsicc.*, sub “*Hyoseris arenaria*”)!

Ar. Geog. Ditio Tingitana. Hispania australis in arenosis Gaditanis et alibi rarius in regione litorea.

Fertur in *Lowe Cat.* inter plantas Mogadorenses; sed species juxta Mogador perfrequens, et Schousboeo bene nota, ad *R. pendulum* certissime referenda est.

PICRIS (*L. Gen.* 909).

P. ALBIDA (*Ball in Journ. Bot.* 1873, 370).

Planta ramosa, decumbens; folia inferiora lyrato-pinnatifida, superiora angusta, basi semiamplexicaulia; pedunculi bracteati; capitula pro genere parva; involucri foliola exteriora dorso subindurata, interiora viridia pappum æquantia; ligulæ albidæ (exsiccatæ ochroleucæ), longiusculæ, exteriores dorso purpurascens; achenia omnia pappo plumoso coronata, rostro longiusculo ipso achenio subæquali.

Mar. merid.—Hinc inde in planitie Maroccana. Prope Sheshaoua! Mzouda! Seksaoua! et alibi.

Ex acheniorum structura ad § *Eupicridem* adjungenda, facie potius ad § *Spitzeliam*, et præsertim ad sp. sequentem accedit. Nostra species proxima est *P. campylocarpæ*, Boiss., et *P. longirostræ*, Sch. Bip. Prior differt habitu erecto, foliis sinuato-dentatis, indumento minus rigido.

In *P. albida* indumentum in herba tota biforme, ex pube brevi et setis rigidis apice non glochidiatis.

P. PILOSA (*Del. Fl. Eg.* 116, tab. 141. fig. 2; *DC. Pr.* vii. 130), = *Spitzelia ægyptiaca* (*Sch. Bip. in Flora*, 1833, 727), = *S. cupuligera* (*Dur. in Rev. Botan.* ii. 431).

Mar. occid.—Juxta Ain el Hadjar haud procul Mogador!

Mar. merid.—Frequens in regione inferiore et usque radices Atlantis Majoris. Prov. Haha! Shedma! Millaïn! Seksaoua! Amsmiz! Mskala! in rudcratis et locis umbrosis urbis Marocco!

Ar. Geog. Africa borealis ex Ægypto usque oras Oceani.

Aspectu variabilis, sæpius subpedalis, hæc planta in irrigatis fit luxurians 2-3-pedalis; sed structura involueri et fructus satis constans. Ducibus celeb. Benth. et H. fil. (Gen. Pl. ii. 512) et Boissier (Fl. Or. iii. 734) *Spitzeliam*, Sch. Bip. loc. cit. pro sectione *Picridis* habui. *S. cupulifera* a *P. pilosa*, Del., nec involuero nec fructu nec habitu differre videtur. Huic proximæ sunt *P. radicata*, Forsk. (= *P. lyrata*, Del.), et *P. atlantica*, Coss. (sub *Spitzelia*), sed in his pappus acheniorum exteriorum longiuseule fimbriatus est.

HELMINTHIA (Juss. Gen. 170).

II. ECHIOIDES (L., sub *Picride*; Gærtn. Fruct. ii. 398, tab. 169; DC. Pr. vii. 132).

Mar. sept.—Tanger!

Ar. Geog. Europa media. Regio omnis Mediterranea (inclusa Macaronesia).

Pace celeb. Benth. et Hooker fil., qui *Helminthiam* ad sectionem *Picridis* revocarunt, hoc mihi videtur genus omnino naturale, structura involueri propria bene definitum.

II. ACULEATA (Vahl, Symb. ii. 89, sub *Picride*; DC. Pr. vii. 132).

Mar. occid.—Legi prope Casa Blanca spec. unicum casu aliquo vel absconditum vel perditum!

Ar. Geog. Africa borealis. Sicilia.

CREPIS (L. Gen. 914).

C. TINGITANA (Salzm. Exsicc., sub *Hieracio*).

Mar. sept.—In monte Djebel Kebir et alibi in umbrosis prope Tanger! (Salzm. spec.)! (Webb Herb.)!

Mar. occid.—Casa Blanca!

Ar. Geog. Marocco. Hispania Gaditana a me prope Algeeciras, et a Bourgeo pr. Alcalá detecta, sec. Willk. et Lange.

Crepis spathulata (Guss. Pl. Cat. 1821; Spreng. Syst. iii. 651; et DC. Pr. vii. 153, sub *Barkhausia*) est planta aliena, quæcum cel. DC. l. c. speciem nostram perperam conjunxit. Stirps Sicula habet folia caulina pauca lineari-lanceolata, exauriculata, et achenia brevissime rostrata, nostra invicem gaudet foliis caulinis plurimis oblongis vel subspathulatis, sæpius dentatis, semper ad basin auriculatis, et acheniis longiuseule rostratis.

Habui ex Sicilia spec. *C. hyemalis*, Biv., pl. nostræ subsimilia, sed in hac foliola exteriora involueri latiora et laxiora. In *C. tingitana* capitula sat magna, ligulæ saturate flavæ, caules paulo supra basin ramosi, 1-2-pedales. *C. macrophylla* (Desf. Fl. Atl. ii. 231; DC. Pr. vii. 155, sub *Barkhausia*), ex descript. Fontanesiana, sp. nostræ valde proxima, est mihi species incerta, ejus exemplaria nulla vidi, nec vidit celeb. DC.

C. TARAXACIFOLIA (Thuill. Fl. Par. i. 409; DC. Pr. vii. 154, sub *Barkhausia*).

Mar. sept.—Tanger! (*Webb Herb.*)! Inter Tanger et Tetuan! Prope Tetuan!

Mar. occid.—Prov. Haha haud procul Mogador!

Mar. merid.—Mskala! Sheshaoua!

Planta polymorpha, cujus varietates extricatu nimis difficiles. Caule primario eroso vel coneulcato nascuntur e radice ramī accessoriī et planta fit biennis. Ex hac forma, in ditione Tingitana frequenti, nata est, nisi fallor, *C. taraxacoides* (Desf. *Fl. Atl.* ii. 231). Specimina in provinciis meridionalibus ad usum (*C. præcocem*, Balb.) referenda videntur. Formam sequentem distinguo.

— —, var. INTYBACEA (*DC. Prod. l. c.*), = *C. intybacea* (*Brot. Fl. Lus.* i. 321).

Mar. sept.—Cap. Spartel! Tanger (*Salzm. Exsicc.*)!

Planta Algeriensis a cel. Cosson pro *C. intybacea* determinata videtur diversa a pl. Candolleana, et a spec. Siculis quæ habui a cl. Tinco.

— —, var. STELLATA, *nob.* = *C. stellata* (*Ball in Journ. Bot.* 1873, 371). Differt a typo caule graciliore minus ramoso oligocephalo, foliis magis hispidis, involucri foliolis exterioribus angustioribus, valde inæqualibus magis adpressis, interioribus margine membranaceis, post anthesin concavis stellatim expansis, receptaculo minus piloso.

Mar. merid.—In provinciis meridionalibus Imperii Maroccani sat frequens. Haha! Tasseremout! Ourika! Seksaoua! Aït Mesan! usque 1400 m.

Forma Maroceana *C. taraxacifoliæ* habitu proprio potius quam charactere tecnico, distinguenda. Achenia foliolis involucri subæquilonga, exteriora foliolis adhærentia.

Ar. Geog. *C. taraxacifolia* est sp. polymorpha per Europæ et reg. Mediterraneæ partem occidentalem late diffusa.

CREPIS BULBOSA (*L.*, sub *Leontodonte*; *Tausch in Flora*, 1828; *Boiss. Fl. Or.* iii. 832), = *Ætheorhiza bulbosa* (*Cass. Dict.* xlviii. 425; *DC. Pr.* vii. 160).

Mar. sept.—Prope Tanger! Cap. Spartel! (*Webb Herb.*)!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

C. VIRENS (*L.*, *Boiss. Fl. Or.* iii. 848), = *C. polymorpha* (*Wallr. Sched. Crit.* 426; *DC. Pr.* vii. 162).

Mar. merid.—? Specimen unicum acheniis immaturis legi prope Milhain ad radices Atlantis Majoris!

Ar. Geog. Europa media. Regio Mediterranea fere omnis (inclusa Macaronesia) sed non ubique. In Oriente rara.

C. ΠΟΚΕΡΙΑΝΑ (*Ball in Journ. Bot.* 1873, 371; non *Clarke, Compos. Ind.* p. 255, anno 1876 edita).

Rhizoma perenne, erassum, sæpius furcatum; folia omnia radicalia, hirtula, lyrato-runcinata, lobis triangularibus acutis, sæpius denticulatis; scapus 4–8-pollicaris, adscendens, indivisus, medium versus bracteola

lineari instructus; involucri foliola exteriora angusta, interiora nigrescentia, apice ciliolata, lana brevi plus minusve adspersa; receptaculi foveolarum margo membranaceus lacero-ciliolatus; ligulæ flavæ dorso rubellæ; achenia elongata, non compressa, lævia vel oculo armato tenuissime asperula; pappi nivei setæ scabrinseulæ.

Mar. merid.—In regione superiore Atlantis Majoris, in monte Djebel Tezah! a 2400 m. ad 3000 m.!

Species probe distincta, affinis *C. oporinoidi* (Boiss. Voy. en Esp. ii. p. 388, tab. cxvii.). In hac seapus foliaceus, bifurcatus; folia pinnatifida, glabra, seu rarius parce lanata (nec hirtula), segmenta subæqualia, lineari-lanceolata, obtusiuscula, suprema elongata; margo foveolarum receptaculi brevissimus, vix membranaceus, sub lente minutissime ciliolatus; achenia costata. Celeb. Boissier *C. oporinoidi* sæpe pappum plumosum et achenia compressa tribuit. In speciminibus plurimis a me exploratis pappi setas simplices sub lente asperulas, et achenium maturum subcylindricum, nec compressum semper observavi. *C. cæspitosa*, G. G., Moris (sub *Barkhausia*), facie plantæ nostræ simillima, differt acheniis rostratis cæterisque notis.

PHLECASIUM (*Cass. Dict.* xxxix. 387).

P. PULCHRUM (*L., et DC. Pr.* vii. 160, sub *Crepide*; *Rchb. fl. Icon.* xix. tab. 80), = *P. lamsanoides* (*Cass. loc. cit.*).

Mar. merid.—In regione media et superiore Atlantis Majoris. In ascensu ad jugum Tagherot a 2500 m. ad 3000 m. ! In monte Djebel Tezah circ. 3000 m. !

Ar. Geog. Europa media. Montes regionis Mediterraneæ. In Oriente (ex Boissier *Fl. Or.*) nascitur etiam in regione calidiore. In Macaronesia non visa.

HIERACIUM (*L. Gen.* 913).

II. PILOSELLA (*L., DC. Pr.* vii. 199).

Mar. merid.—In regione media et superiore Atlantis Majoris. In jugo Tagherot 2400–2900 m. ! In monte Djebel Tezah ad 2400 m. !

Ar. Geog. Europa fere omnis (exclusa arctica). Asia temperata. Montes regionis Mediterraneæ (exclusa Macaronesia). In Asia Minore et in Græcia nonnisi in monte Olympo Bithynico detectum, sed adest forma proxime affinis *H. macranthum*, Ten.

Spec. nostra ex involucri pilis mollibus fuscis ad *H. pelleterianum* (DC. *Fl. Fr.* vi. 437, *Prod.* vii. 200) tendit. Hoc est nil nisi varietas sp. Linnæanæ, et formæ intermediæ nostris simillimæ in montibus Europa obviæ sunt.

ANDRYALA (*L. Gen.* 915).

A. RAGUSINA (*L., DC. Pr.* vii. 244).

Mar. occid.—In monte Djebel Hadid haud procul Mogador (*R. et Fr. spec. in Herb. Kw.*)!

Mar. merid.—Mesfioua ! Ourika !

Forma angustifolia, fol. inferioribus repando-dentatis, superioribus integerrimis. A formis similibus *A. integrifoliæ* distinguitur foveolarum receptaculi margine membranaceo laccro parce setoso, nec longe setoso.

— — —, var. LYRATA (DC. l. c.), = *A. lyrata* (Pourr. Act. Toul. iii. 308).

Mar. occid.—? Legi nondum florentem in collinis prov. Haha haud procul Mogador !

Folia in spec. nostris pinnatifida lobis oblongis obtusis. Planta incerta, ulterius inquirenda ; forsitan est *A. pinnatifida* (Ait. H. Kew. iii. 129 ; DC. Prod. l. c.), planta Canariensis et Maderensis ab auctoribus plurimis confusa. Hæc, etsi facie diversa, inter formas *A. integrifoliæ*, L., enumeranda videtur. Confer Lowe, Man. Fl. Mad. i. 564.

Ar. Geog. *A. ragusina* est imprimis sp. Hispanica, in Gallia australi et Corsica etiam obvia. Ex Dalmatia non vidi specimina, nec adest in Vis. Fl. Dalm. In Africa boreali vix ultra Imp. Maroccanum detecta.

ANDRYALA MOGADORENSIS (Coss. in Bull. Soc. Bot. Fr. xx. 252).

“Planta perennis, tota indumento molli cinereo-canescente demum rufescente tomentosa, caudice brevissimo sublignoso in radicem descendentem abeunte caules 1–2 vel plures emittente. Caulis erecti vel a basi arcuata ascendentes, indurati, superne ramosi, ramis patentibus. Folia integra vel obsolete subsinuata obtusa, infima obovato-oblonga in petiolum brevem attenuata, caulina approximata oblonga vel obovato-oblonga sessilia basi lata semiamplexicaulia. Capitula apice ramorum laxiuscule cymoso-corymbosa, breviter pedunculata, cymis corymbum generalem efformantibus. Involucrum lana densa sordide alba demum rufescente lanato-gossypinum pilis longioribus rigidioribus apice glanduligeris nigricantibus permixtis, foliolis linearibus acutis. Ligulæ aurantiacæ, involucrum superantes. Achænia fusco-purpurea, oblongo-cylindracea, inferne attenuata, sub-10-striato-costata, costis concoloribus.

“Caudice perenni, caulibus induratis, nonnullisque aliis notis ad *A. ragusinam*, L., collocanda, sed abunde distincta foliis omnibus obtusis integris vel obsolete subsinuatis, caulinis basi lata semiamplexicaulibus, pedunculis brevibus, involucri foliolis præter lanam pilis longioribus glanduliferis donatis, achæniis duplo brevioribus oblongo-cylindræis costis concoloribus, non lineari-cylindræis costis albidis.”

Mar. occid.—In rupibus maritimis insulæ Mogador ! ubi primus detexit el. Balansa. In maritimis ab urbe Saffi meridiem versus !

Achenia omnino matura fiunt atra costis concoloribus. Receptaculum ad foveolarum margines longe setosum.

Spec. quæ juxta Saffi legimus gaudent foliis angustioribus oblongis, subsinuatis.

ANDRYALA INTEGRIFOLIA (L., DC. *Pr.* vii. ; Willk. et Lge. *Fl. Hisp.* ii. 271).

Mar. sept.—Tanger! (*Webb Herb.*)! (*Salzm. Exsicc.*, sub *A. pinnatifida*) Tetuan (1851)!

Mar. occid.—Casa Blanca!

Mar. merid.—Amsmiz! Mtouga!

— — —, var. ARENARIA (*Boiss. Voy. Esp.* 394), = *A. arenaria* (*Boiss. et Reut. Pug.* 71).

Mar. sept.—Tetuan!

Mar. occid.—Mazagan! Mogador! In arenosis prov. Haha haud procul Mogador!

Mar. merid.—Shedma! Ourika!

Ar. Geog. Species regionis Mediterraneæ occidentalis.

Planta polymorpha in plurimas species falsas ab auctoribus dilaniata et synonymia igitur intricatissima. Forsan non satis differt *A. dentata*, Sibth. et Sm. ; = *A. tenuifolia*, DC., species in reg. Mediterranea orientali obvia.

A. LAXIFLORA (*Salzm. in DC. Pr.* vii. 247).

Mar. sept.—Tanger (*spec. ex Herb. Schousb.*)! (*Webb Herb.*)! Tetuan (1851)!

Mar. occid.—Casa Blanca (*spec. unicum legit J. D. II.*)!

Ar. Geog. Ditio Maroccana. Hispania australis.

HYPOCHLERIS (*L. Gen.* 918).

H. GLABRA (*L., DC. Pr.* vii. 90).

Mar. sept.—Inter Tanger et Tetuan!

Mar. occid.—Casa Blanca! Aïn el Hadjar haud procul Mogador!

Mar. merid.—Sektana! Seksaoua! Mskala!

— — —, var. foliis insigniter hispidis, pedunculis vix incrassatis, = *H. hispida* (*Brot. Fl. Lus.* i. 332)?

Mar. merid.—Prov. Shedma! In convalle Aït Mesan ad 1600 m. (?)!

Spec. ex Aït Mesan immaturum foliis runcinatis insigne incertum est.

— — —, var. SALZMANNIANA (*Coss. MSS.*), = *H. Salzmanniana* (*DC. Pr.* vii. 91), = *H. dimorpha* (*Salzm., non Brot. nec Ten.*).

Mar. sept.—Tanger (13 Jun.)! (*Salzm. Exsicc.*, sub “*H. dimorpha*”)!

Hæc est proles luxurians *H. glabræ*, pedunculis superne valde incrassatis, involucri foliolis paullo latioribus.

— — —, var. EROSTRIS (*Coss. et Germ. Fl. Par.*), = *H. arachnoidea* (*Poir. Dict.* v. 572 ; *DC. Pr.* vii. 90).

Mar. merid.—Mesfioua!

Ar. Geog. Species per Europam omuem (exclusa arctica) et regionem Mediterraneam late diffusa. In ins. Azoricis et in Madera obvia, deest, ut videtur, in ins. Canariensibus. Var. *Salzmanniana* est ditionis Tingitane et Hispaniæ Granatensis propria civis ; var. *erostri*s per regionem omnem imprimis calidiorem dispersa est.

HYPOCHÆRIS RADICATA (L., DC. *Pr.* vii. 91), var. **HETEROCARPA** (Moris, *Fl. Sard.* ii. 487), = *H. neapolitana* (DC. *Pr.* vii. 91), = *H. dimorpha* (Ten. *Syn. Fl. Neap.* 407, non Brot. nec Salzm.).

Mar. sept.—Inter Tanger et Tetuan (J. D. II., spec. unicum immaturum)! Tetuan (*Webb Herb.*)!

Mar. merid.—Sheshaoua! Shedma (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Per Europam omnem et regionem Mediterraneam fere omnem diffusa. Var. nostra frequentior in calidioribus. Species deest in ins. Canariensibus. In Madera adest tantum forma typica acheniis omnibus æqualiter rostratis.

Sectiones hujus generis *Arachnites* et *Porcellites*, a Candolleo in 'Pro-dromo' admissæ, et a Benth. et II. fil. (Gen. Pl. ii. 520) conservatæ, plane delendæ sunt; characteres enim quibus nituntur omnino variabiles nec ad species definiendum idoneæ.

II. ÆTHNENSIS (L. et DC. *Pr.* vii. 95, sub *Seriola*), = *Hypochæris serioloides* (Bert. *Misc.* ii. 236).

Mar. sept.—Tanger (*Webb Herb.*)! Tetuan (*Webb Herb.*)! In monte Beni Hosmar!

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

II. LÆVIGATA (L., et DC. *Pr.* vii. 96, sub *Seriola*), = *Piptopogon decipiens* (Cass. *Dict.* xlvi. 507).

Mar. sept.—Tanger (*spec. ex Webb Herb.*)! In monte Beni Hosmar pr. Tetuan! (*Webb Herb.*)!

Ar. Geog. Africa borealis. Sicilia.

H. lecntodontoides (Ball in *Journ. Bot.* 1873, 371).

Radix crassa, subverticalis, divisa, ad collum foliis emarceidis vestita; folia rosulata, obovato-spathulata, obtusa, vel obverse lanceolata, integra, acuta, repando-denticulata, subglaberrima, vel pilis crispis simplicibus vestita; scapi breves filiformes; involucrium paucifoliolatum, subregulariter imbricatum, extus pubescens, foliolis exterioribus interdum pilis paucis longioribus fuscis munitis; receptaculi paleæ hyalinæ, apice setiferæ, coloratæ, achenia (juniora) excedentes; pappus ex radiis 10 longioribus deciduo-plumosis basi parum expansis, cæteris (fere 10) dimidio brevioribus inter priores extus adjunctis; achenia nimis immatura ulterius observanda.

Mar. merid.—In regione superiore Atlantis Majoris. In convalle Aït Mesan! et in jugo Tagherot! a 2400 m. ad 3200 m.

Ab *H. lævigata* foliis integris rosulatis, scapis gracilibus nudis, involucri foliolis pubescentibus adpressis, videtur longe diversa. Cel. Cosson tamen certiore me fecit hanc esse priori proxime affinis. Achenia immatura videntur suberostrata illis *S. lævigatæ* dimidio breviora. Planta ulterius inquirenda pro subspecie haud sine dubio hic enumerata.

Pappus in pl. nostra radiis longioribus basi vix expansis ad confirmandam sententiam celeb. Benth. et II. fil., qui *Seriola* Linnæanam ad sectionem *Hypochæridis* reduxerunt, non parvi momenti est.

LEONTODON (*L. Gen.* 912; *Benth. et H. fil. Gen. Pl.* ii. 520).

L. AUTUMNALIS (*L., DC. Pr.* vii. 108, sub *Oporinia*; *Hoffm.*, sub *Apargia*), var. *ATLANTICUS*, *nob.* Planta nana glabra, foliis radicalibus runcinato-pinnatifidis, scapis nudis brevibus, involucri foliolis adpressis nitidis, acheniis immaturis apicem versus parum attenuatis, pappi radiis æqualibus ad basin vix incrassatis.

Mar. merid.—In convalle Aït Mesan circ. 2500 m.!

Var. nostra ad var. *minimam* (*DC. Pr. l. c.*) accedit. Hæc tamen differt capitulis albo-villosis.

Ar. Geog. Species per Europam septentrionalem et mediam, necnon in Sibiria late diffusa, fit rara in regione Mediterranea. Vidi tantum ex montibus Italiæ, Hispaniæ, Galliæ australis et Dalmatiæ.

L. ROTHII, *nob.*, = *Thrincia hispida* (*Roth, Catal. Bot.* i. 99; *DC. Pr.* vii. 100, non *Fl. Fr.*), = *Colobium hispidum* (*Roth in Rœm. Arch.* i. 38), = *Thrincia taraxacoides* (*Gaud. Fl. Helv.* v. 49, quoad var. *a.*), = *Leontodon nudicaulis* (*Solander MSS. in Herb. Mus. Brit.* pro parte, conf. *Lowe, Man. Fl. Mad.* i. 535), = *Hyoseris arenaria* (*Schousb. Gew. Marok*, 183), = *Thrincia mauritanica* (*Spreng. Syst.* iii. 666, pro parte), = *Leontodon saxatile* (*Lam. Dict.* iii. 531).

Mar. sept.—Frequens in dit. Tingitana. Legimus pluries prope Tanger! et juxta promont. Spartel! Inter Tanger et Tetuan! juxta urbem Tetuan! et formam foliis pinnatifidis inter Tetuan et Ceuta! Habui ex arenosis Tingitanis ex herb. Webb!

Mar. occid.—Casa Blanca! Aïn el Hadjar et Djebel Hadid haud procul Mogador!

Mar. merid.—Occurrit passim in regione inferiore et in convallibus calidioribus Atlantis Majoris. Prov. Shedma! Marocco in ruderatis! Seksaoua! Mtouga! Milhain! In convalle Aït Mesan usque 1600 m.!

— — —, var. *MAJOR* (*Boiss. Voy. Esp.* 378; *Willk. et Lge. Fl. Hisp.* ii. 214), = var. *microrhynchoides*, Ball, MSS. Folia sinuata vel repando-dentata, hispida, pilis (præsertim in nervo medio) apice glochidiatis; pappus 10-radiatus, radiis externis paucis brevissimis. Huc forsân *L. nudicaulis* (*Lowe, Prim. Fl. Mad.* 47, sub *Thrincia*) referenda.

Mar. occid.—Aïn el Hadjar haud procul Mogador!

Mar. merid.—In saxosis aridis reg. inferioris prov. Shedma! Misra Ben Kara haud procul urbe Marocco!

Ar. Geog. Species in regione Mediterranea occidentali (inclusa Macaronesia) satis frequens. Varietates plurimæ in Hispania et in Africa boreali.

Invito animo nomen novum proposui, dum nomen triviale *hispidus* ad aliam speciem Linnæanam pertinet; nomen *taraxacoides* improprium est, dum adsunt *L. taraxacum*, *L.*, *L. taraxaci*, *Lois.*, et *L. taraxacoides*, *Rœnsch.*, et *L. arenarius* ad formam *L. hirti* (*Thrincia hirtæ*

auct.) a plerisque auctoribus acceptum, *nudicaulis* denique ab ipso Solander eum *L. hirta* confusus.

LEONTODON MAROCCANUS (*Pers. Syn.* ii. 308, et *DC. Pr.* vii. 100, sub *Thrincia*), = *Thrincia mauritanica* (*Spreng. Syst.* iii. 666, pro parte), = *Hyoseris hispida* (*Schousb. Gew. Marok*, 183, sec *DC. Prod. l. c.*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb spec.*)! In monte Djebel Kebir! In collinis inter Tanger et Tetuan!

Mar. centr.—Prope Mequinez (*Schousb. l. c.*).

Mar. occid.—In montosis aridis ad Mogador (*Brouss. ex DC. Pr.*). A recentioribus non detectus.

Ar. Geog. Imp. Maroccanum. Hispania australis.

Stirps nostra, a Sprengelio et auct. recentioribus cum præcedente confusa, videtur probe diversa, capitulis majoribus, ligulis externis in dorso pilosis, ensulis fuscis, acheniis majoribus muriculatis, marginalibus pappo coroniformi fimbriato, interioribus rostro longissimo quam achenium ipsum quadruplo longiore donatis, receptaculo demum læviuseulo, margine foveolarum subobsoleto non piloso.

L. TINGITANUS (*Boiss. et Reut. Pug.* 68, sub *Thrincia*).

Mar. sept.—Prope Tanger (*Boiss. et Reut.*, spec. incompletum in *Herb. Kew.*)!

Ar. Geog. Nonnisi in ditione Tingitana et in Hispania Arundana hucusque detectus. Specimina completa Tingitana non vidi, sed exemplar Hispanicum anno 1851 prope Ronda lectum his omnino conforme est. Indumento ex pilis fureatis subvelutino insignis, verisimiliter probe distincta species.

Speciei hujus sectiones (*Thrinciæ*, auct.) extricatu difficillimæ sunt. Affinitas inter *L. hirtum* (*L.*, *DC. Pr.*, sub *Thrincia*) et *L. Rothii* nostro (*Thrincia hispida*, *Roth*) tam proxima ut exemplaria sæpe difficiliter dignoscuntur; et, pace tantorum virorum, sectio *Streckera*, a celeb. Benth. et H. fil. in *Gen. Pl.* loc. cit. laudata, a *Thrincia* separari nequit. Radix annua in *L. Rothii* (ut jam innotuit b. Lowe) in calidioribus sæpe perennans, et post annum primum fit præmorsa. Numerus radiorum pappi nullomodo constans ut voluerunt ell. Boiss. et Reuter, sed in *L. Rothii* sæpissime 10 et semper pauciores sunt quam in *L. hirta*. Pili non semper simplices.

L. HELMINTHOIDES (*Coss. et Dur. ined.*), var. = *L. hispanicus*, var. Ballii, *Benth.*

Seapi simplices breves, bracteis 1–2 linearibus instructi; involucri foliola pauciora quam in typo, minus regulariter imbricata—foliola interiora latiuscula; ligulæ extus rubellæ nec virentes; receptaculi foveolæ margine brevissime membranaceæ non ciliatæ; achenia apicem versus attenuata, vix, aut ne vix, rostrata, subtiliter corrugato-asperula (nec muriculata); pappi radii longiores plumosi, exteriores breviores setacei.

Mar. merid.—Legi in convalle Amsmiz circ. 1400 m. ! Spec. acheniis immaturis in monte Djebel Ouensa ab indigeno Ibrahim lecta benev. comm. cel. Cosson !

Ar. Geog. Species Algeriensis ; var. nostra Atlantis Majoris civis est.

L. helminthioides, *L. hispanico* (Mérat in Ann. Sc. Nat. xxii. 106 ; DC. Pr. vii. 127, sub *Asterothrici*) proximus est. Satis differre videtur acheniorum et pappi structura, et præsertim foveolis receptaculi sub-obsoletis margine non ciliatis. In hac grege tamen characteres valde incerti. Ne dicam de notis specificis, characteres etiam generis nullo modo stabiles sunt. Inter specimina *L. hispanici* ad Fuente de la Higuera lecta (Bourgeau, Pl. d'Espagne, No. 1600) video achenia exteriora incurva extus furfuracea, nec costato-striata, pappo brevissimo fimbrillifero coronata omnino ut in *Picride lyrata*, Del.

LEONTODON HISPIDULUS (Del. Fl. Eg. 117, tab. 42. sub *Crepide* ; DC. Pr. vii. 109, sub *Oporinia* ; Boiss. Fl. Or. iii. 727), = *Kalbfussia orientalis* (Jaub. et Spach, Ill. Pl. Or. iii. 117, tab. 283), = *Leontodon trivialis* (Ball in Journ. Bot. 1873, 372).

Mar. merid.—Legi prope Aïn Onnast, prov. Shedma ! et in prov. Mtouga ! Specimen unicum in ruderatis urbis Marocco !

Ar. Geog. Species Ægyptiaca et Maroccana, cujus var. (*Millina persica*, Boiss. Diagn. Pl. Or. ser. 1, vii. 3) per regionem desertorum Persiæ australis, Syriæ, Arabiæ diffusa est. Ex Algeria non vidi specimina. Planta ulterius inquirenda.

Planta glabriuscula vel pilis simplicibus adspersa ; scapus primarius erectus, posteriores plurimi adscendentes, omnes aphylli, sæpius furcati 1-3-cephali ; folia lanceolata repando-dentata ; involucrium glabrum vel tenuiter arancoso-pubescent ; receptaculum fimbriatum ; achenia in rostrum ipsis æquilongum attenuata, pappi radiis 10 plumosis, basin versus dilatata.

Stirps satis ludibunda. Spec. nostra aspectu diversa habui pro specie nova ; sed nullis notis gravioribus a planta Ægyptiaca separanda videntur. *Apargia annua* (Vis. Pl. Æg. 38, tab. 6), a cel. Boissier *l. c.* inter synonyma sp. nostræ citata, ob achenia pluries breviora, apice attenuata sed crostria, videtur satis distincta species.

Planta Maroccana sæpissime crosa vel sub pedibus conculcata videtur annua, radice indurata interdum (?) perennante.

L. SALZMANNI (Sch. Bip. in Flora, 1833, 723, et DC. Pr. vii. 101, sub *Kalbfussia*), = *Apargia verna* (Salzm. Exsicc. 1825).

Mar. sept.—Tanger ! (Salzm. Exsicc.) !

Ar. Geog. Africa borealis. Hispania. Var. vel subspecies (*Kalbfussia Mülleri*, Sch. Bip. *l. c.*) in Sardinia et Sicilia.

Nomen triviale Salzmannianum *vernus* prioritate gaudet, sed nolui mutare nomen ab aucteribus omnibus recentioribus acceptum.

LEONTODON TUBEROSUS (L., *Willd. Sp. Pl.* iii. 1549, sub *Apargia* DC. *Pr.* vii. 100, sub *Thrincia*).

Mar. sept.—Inter Tanger et Tetnan !

Ar. Geog. Regio Mediterranea fere omnis (inclusa Macaronesia).

Nomen generis *Leontodon* generis masculini a Linnæo pluribusque auctoribus ad gen. neutrum refertur.

TARAXACUM (*Hall, Stirp. Helv.* i. 23).

T. OFFICINALE (*Wigg. Prim. Fl. Hols.* 56), = T. Dens Leonis (*Desf. Fl. Atl.* ii. 228 ; DC. *Pr.* vii. 145).

Mar. merid.—Legi in irrigatis umbrosis juxta Sheshaoua ! Distr. Reraya ad 1000 m. ! et formam montanam in convalle Aït Mesan supra Arroumd circ. 2400 m. ! in jugo Tagherot usque 2800 m. ! et in monte Djebel Tezah !

Ar. Geog. Per hemisphæram borealem fere omnem diffusum, et inde in temperatis hemisphæræ australis introductum.

LACTUCA (*L. Gen.* 909).

L. VIMINEA (L., sub *Prenanthe* ; *Lam. Dict.* ii. 27, sub *Chondrilla* ; DC. *Pr.* vii. 176, sub *Phænopo* ; *Link, Enum. H. Ber.* ii. 281), = *Phoenixopus decurrens* (*Cass. Dict.* xxxix. 391).

Mar. merid.—Saksaoua ! Aït Mesan ! Spec. ex Djebel Ouensa ab indigeno Ibrahim lectum benevole comm. cel. Cosson.

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia). Occurrit etiam hinc inde in calidioribus Europæ mediæ.

Verisimiliter ad varietatem hujus sp. referendum est specimen nondum florens a cl. G. Maw inter Ourika et Reraya lectum, foliis (nisi imis) indivisis repando-denticulatis, lanceolatis acutis, basin versus in petiolum longum decurrentem sensim attenuatis.

L. TENERRIMA (*Pourr. Act. Toul.* iii. 322 ; DC. *Pr.* vii. 134).

Mar. merid.—In convallibus calidioribus et ad radices Atlantis Majoris haud infrequens. Aït Mesau usque 1000 m. ! In convalle Amsmiz 1200–1500 m. ! supra Seksaoua ! In rupibus supra Milhaïn circ. 1000 m. !

Ar. Geog. Hispania. Gallia australis. Italia subalpina. Ex Africa nisi nostra, spec. nulla vidi.

L. SALIGNA (L., DC. *Pr.* vii. 136).

Mar. sept.—Vidi nondum florentem prope Tanger ! spec. ex herb. Schonsboe dedit cel. Cosson !

Mar. merid.—? Legi nondum florentem in convalle Aït Mesan ! et folia in distr. Reraya ! Spec. incerta, verisimiliter hinc referenda.

Ar. Geog. Europa media. Regio Mediterranea fere omnis (exclusa Macaronesia).

PICRIDIMUM (*Desf. Fl. Atl.* ii. 220).

P. TINGITANUM (*L.*, sub *Scorzonera*; *Lam. Dict.* iii. 397, sub *Soucho*; *Desf. loc. cit.*; *DC. Pr.* vii. 182). Hue etiam *P. Hispanicum* (*Poir. Diet.* xl. 197); et sicut varietates enumeranda sunt *P. arabicum* (*Hochst. et Steud. in Schimp. Pl. Arab. Exsicc.*) et *P. orientale* (*Pers. Syn.* ii. 362).

Mar. sept.—Occurrit passim in hac regione. Tanger! (*Webb spec.*)! (*Salzm. Exsicc.*)! Cap. Spartel! In Tanger et Tetuan! Inter Tetuan et Ceuta!

Mar. occid.—Sali (*Brouss.*). Casa Blanca! Prope Saffi! Prov. Haha pr. Mogador! Aïn el Hadjar haud procul Mogador!

Mar. merid.—Valde frequens in reg. inferiore. Prov. Shedma! Prov. Mzouda! Mskala! Seksaoua! Mesfoua! Sektana (*R. et Fr. spec. in Herb. Kew.*)!

— —, var. *MARITIMUM*, = *P. ligulatum* (*Vent. Mahn., fide DC. Pr. loc. cit.* quoad plantam Mogadorenses). Hue etiam var. *diversifolium* (*DC. Pr. l. c.*).

Mar. occid.—Frequens in maritimis prope Mogador, præsertim in insula.

— —, var. *MAROCCANUM*, *nob.* Planta insignis, ulterius inquirenda, pedalis et ultra, e basi ramosa, ramis erectis æquilongis parve foliaceis; folia radicalia desunt, caulina semiamplexicaulia profunde pinnatifida lobis oblongo-euneatis 2-3 utrinque instructa, lobo terminali elongato; capitula aliquantulum minora, cætera ut in *P. tingitano*. Radix videtur annua?

Mar. merid.—Mesfoua!

Ar. Geog. Species per regionem omnem Mediterraneam calidiorem (inclusa Macaronesia) et inde per Indiam bor. occidentalem et Abyssiniam late diffusa. Deest in Gallia australi, in Italia continentali, et in Dalmatia.

Stirps polymorpha quoad formam foliorum et involucri valde ludibunda.

P. ligulatum auctorum vix pro varietate distinguendum est. Specimina hue a Candolle relata (*Prod. l. c.*) see. eel. Boissier potius formam maritimam *P. vulgaris* sistunt.

Achenia in *Picridiis* omnibus a me exploratis biformia sunt. Exteriorum series 3-4-, sæpius 4-sulcatæ (in eodem capitulo occurrunt ach. 5-sulcata), rugosa, tubereulata, interiora breviora subeonica, obsolete 4-sulcata, vix tubereulata, in *P. tingitano* integumento albescente crassiusculo munita (conf. Boiss. *Fl. Or.* iii. 327). Descriptio generis in *DC. Prod. l. c.* parum accurata.

P. VULGARE (*Desf. Fl. Atl.* ii. 221; *DC. Pr.* vii. 182), = *Scorzonera picroides*, *L.*

Mar. sept.—Tanger! (*Webb Herb.*)! Cap. Spartel (*Webb Herb.*)! Inter Tetuan et Ceuta!

Mar. occid.—Mogador (*Lowe Cat.*). Agadir (*Coll. Indig.* 1871)!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

Specimina nobis ex vicin. Agadir allata habent achenia longiora et centralia superne magis attenuata.

SONCHUS (*L. Gen.* 908).

S. OLERACEUS (*L. Sp. Pl.* 1116, excl. varr. γ et δ), = *S. ciliatus* (*Lam. Fl. Fr.* ii. 87; *DC. Pr.* vii. 185).

Mar. sept.—Tanger (*Webb Herb.*)! Cap. Spartel! Inter Tetuan et Ceuta!

Mar. oecid.—Mogador! (*Lowe Cat.*).

Mar. merid.—Sheshaoua! In horto irrigato Ben Dreis urbis Maroceo!
Spec. unicum in monte Djebel Tezah legit J. D. II.! Sektana (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Species eosmopolita per orbem terrarum fere omnem diffusa.

S. ASPER (*Vill. Dauph.* iii. 158), = *S. fallax* (*Wallr. Sched. Crit.* 432; *DC. Pr.* vii. 185).

Mar. sept.—Tanger (*Webb Herb.*)!

Mar. merid.—In convalle Aït Mesan 1200–1500 m.! Ourika (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Sicut *S. oleraceus* affinis *S. asper* per orbem fere totum diffusus est.

S. glaucescens (*Jord. Obs.* v. 75, tab. 5; *Boiss. Fl. Or.* iii. 796), = *S. græcus*, *Reut. MSS.*

Mar. merid.—Sheshaoua!

Ar. Geog. In Gallia australi, Hispania, Græcia, et in Oriente indicatus, verisimiliter per totam regionem Mediterraneam diffusa.

S. TENERRIMUS (*L., DC. Pr.* vii. 186), = *S. peetinatus* (*DC. loc. cit.*), forma perennis.

Mar. sept.—Tanger! (*Webb Herb.*)! In monte Beni Hosmar prope Tetuan (*Webb Herb.*)! Inter Tetuan et Ceuta!

Mar. oecid.—Mazagan! Mogador (*Lowe Cat.*).

Mar. merid.—Frequens in regione inferiore. Seksaoua! Aïn Tarsil! Milhaïn!

— —, var. *MARITIMUS*, *nob.* Forma insignis radice perenni rep-
tante, foliis in lacinias plurimas integras lineares, vel ovatas croso-
dentatas, v. pinnatifidas, retrorsum versas divisas; involuero nudo,
nec ad basin araneoso-floccoso.

Mar. sept.—In rupibus maritimis a promontorio Spartel meridiem
versus!

Mar. oecid.—Saffi! Prope Mogador in arenosis maritimis!

— —, var. *TUBERCULATUS*, *nob.* Priori subsimilis, sed major;
capitula majora; ligulæ aurantiacæ; folia minus divisa auriculis

magnis semiamplexicaulibus ; achenia insigniter rugoso-tuberculata, breviora et latiora quam in typo.

Mar. occid.—In rupibus insulæ Mogador !

— —, var. ARBORESCENS, = *S. arborescens* (*Salzm. Exsicc.*) !

Mar. sept.—Tanger (*Salzm. Exsicc.*) ! Inter Tanger et Tetuan ! Tetuan (*Webb spec.*) !

Planta elata 6–8-pedalis in sepibus et dumetis satis frequens. Caules farcti angulati ; folia interrupte pinnipartita, lobis subintegerrimis, terminali maximo triangulati-cordato vel trilobo. Ab hoc vix differt *S. pustulatus* (Willk. in *Fl. Hisp.* ii. 242).

Ar. Geog. *S. tenerrimus* per totam regionem Mediterraneam diffusus, et in Abyssinia obvius, in Imp. Maroccano vulgaris, deest, ut videtur, in Macaronesia.

Stirps in ditione nostra proteiformis variat statura, forma foliorum, radice annua vel perenni, pedunculis glabris vel glandulosis, sed ex his notis varietates satis stabiles ægre definiendæ.

SONCHUS FRAGILIS (*Ball, MSS. in herb.* 1851, in *Journ. Bot.* 1873, 372), = *S. tenerrimus*, *L.*, var. *pectinatus*, *Coss. in Bourgeau, Pl. d'Algérie* (1856, No. 84).

Radix perennis crassa, suberosa ; caules breves foliosi ; folia pinnata, segmentis subalternis, ovato-rotundis, acutiusculis ; pedunculi breves, apice non floccosi ; involucri foliola et ligulæ latiores quam in *S. pectinato* ; achenia vix, aut ne vix, compressa, costis 4–5 prominentibus, tam in costis quam in interstitiis lævissima.

Mar. sept.—In rupibus juxta urbem Tetuan ! et in monte Beni Hosmar Legi anno 1851 et iterum 1871.

Acheniis diversissimis præter habitum singularem a *S. tenerrimo* optime dignoscitur. Si ullo alio foret *S. maritimo*, *L.*, magis affine ; sed in hoc achenia 8-costata.

Ar. Geog. Algeria. Ditio Tetuanensis.

S. MARITIMUS (*L., DC. Pr.* vii. 187).

Mar. merid.—Secus rivulos irrigatorios urbis Marocco ! vidi etiam in prov. Mtonga !

Ar. Geog. Regio Mediterranea exclusa Macaronesia. Deest, ut videtur, in Græcia et in Asia Minore.

Folia in sp. Maroccanis integra margine minute denticulata.

S. ACIDUS (*Schousb. in Willd. Sp. Pl.* iii. 1511 ; *DC. Pr.* vii. 188).

Mar. occid.—Specimen unicum legi in arenosis maritimis pr. Mogador ! (*Broussonet*), ex *DC. Pr.*

Ar. Geog. Marocco occidentalis. In insula Lancerota sec. Bourgeau exstat planta unica forsitan casu aliquo a Mogador allata.

Species valde proxima *S. pinnato* Canariensi.

MICRORHYNCHUS (*Less. Syn. Compos.* 139).

M. NUDICAULIS (*L.*, sub *Chondrilla*; *Less. loc. cit.*; *DC. Pr.* vii. 180).

Mar. occid.—Casa Blanca! Prov. Italia prope Mogador! Agadir (*Coll. Indigeni* 1871)!

Mar. merid.—Frequens in saxosis aridissimis planitici, nec deest in regione inferiore Atlantis Majoris. Prov. Shedma prope Aïn Oumast! Mskala! Sheshaoua! Misra Ben Kara! Prope urbem Marocco! In convalle Aït Mesan usque 1500 m.! Milhaïn!

Spec. Maroccana sæpe ambigua sunt inter typum Candolleanum et var. *divaricatum*.

Ar. Geog. Africa borealis. Arabia. Hispania austro-orientalis. Ins. Canarienses. Deest in Madera.

M. SPINOSUS (*Forsk. Fl. Æg.-arab.* 144, sub *Prenanthe*; *Lam. Dict.* iii. 408, sub *Lactuca*; *DC. Pr.* vii. 189, sub *Soncho*; *Benth. et H. fl., Gen. Pl.* ii. 529), = *Zollikoferia spinosa* (*Boiss. Fl. Or.* iii. 826).

Mar. merid.—In saxosis aridis regionis inferioris et ad radices Atlantis Majoris. In collibus arenaceis prope urbem Marocco (*G. Maw*)! Supra Seksaoua! In convalle Aït Mesan! -

Ar. Geog. Regiones desertæ Arabiæ et Ægypti. Marocco meridionalis. Adest etiam in insulis Capitis Viridis et Canariensibus, et rarius in Hispania austro-orientali.

Cel. Boissier in 'Fl. Orientali' *Microhynchum* eum gen. Candolleanis *Zollikoferia* et *Brachyrampho*, et Cassinianis *Lomatolepide* et *Rhabdotheca* in unum genus optimo jure conjunxit; sed immerito nomen antiquius *Microhynchum* recentiori *Zollikoferiæ* postposuit.

TRAGOPOGON (*L. Gen.* 905).

T. PORRIFOLIUS (*L.*, *DC. Pr.* vii. 113).

Mar. merid.—Seksaoua! Ourika (*J. D. H.*)! Mesfioua (*J. D. H.*)! Mtonga (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio Mediterranea Europæ et Africæ borealis. Ins. Canarienses. In Europa media, verisimiliter introductus. Deest in Madera. In Asia Mediterranea adest forma proxime affinis (*T. longirostris*, *Bisch. in Webb Phyt. Can.* ii. 469).

Specimina ex Mtonga et Seksaoua formam hujus sp. (*T. australem*, *Jord. Cat. Dij.* 1848) sistunt. Hic mihi videtur fere medius inter sp. Linnæanam et *T. longirostrem*, quem, pæce eel. Boissier, pro varietate, vel ad summum pro subspecie habeo. Specimen a eel. Hooker ad Mesfioua lectum vix a *T. longirostri* differt.

T. GLABER (*L.*, et *DC. Pr.* vii. 111, sub *Geropogone*; *Benth. et H. fl. Gen. Pl.* ii. 531).

Mar. occid.—'In agro Mogadorensi (*Broussonnet*)!', *DC. Prod. l. c.*

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia), sed nullibi frequens.

UROSPERMUM (*Scop. Introd. No. 366*).

U. DALECHAMPII (L., sub *Tragopogone*; *Desf. Cat. II. Par. 90*; *DC. Pr. vii. 116*).

Mar. sept.—Inter Tanger et Tetuan (*J. D. II.*)! Vidi etiam in collibus pr. Tanger!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia). Fertur in Thracia et in insulis Archipelagi, sed a recentioribus non visum.

U. PICROIDES (L., sub *Tragopogone*; *Desf. Cat. II. Par. 90*; *DC. Pr. vii. 116*).

Mar. sept.—Tanger (*Webb Herb.*)! Cap. Spartel! Tetnan!

Mar. occid.—Casa Blanea! Mazagan! Djebel Hadid laud procul Mogador (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—Prov. Shedma!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

SCORZONERA (*L. Gen. 906*).

S. UNDULATA (*Vahl, Symb. ii. 86*; *DC. Pr. vii. 117*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Tetuan (*Webb Herb.*)!

Mar. merid.—Ad radices Atlantis Majoris. Mesfioua! Ourika! Tasse-remout! In convalle Aït Mesan!

Ar. Geog. Africa boreali-occidentalis. Fertur in Oriente, sed monente cel. Boissier planta orientalis ad speciem diversam, *S. Alexandrinam* suam, referenda est.

S. DELICIOSA (*Guss. Ind. Sem. II. Bocc. 1825*; *DC. Pr. vii. 118*), var. TETUANENSIS, = *S. Tetuanensis* (*Webb MSS. in Herb.*).

Mar. sept.—In regione collina inter Tanger et Tetuan! Prope Tetuan! (*Webb Herb.*)!

Planta Tetuanensis differt a typo Siculo foliis et involucri foliolis pluries (sæpe ter) latioribus. Specimina Algeriensia *S. deliciosæ* tamen intermedia sunt; et in hoc genere folia latitudine valde variabilia. Achenia matura non vidi.

Ar. Geog. *S. deliciosa* nonnisi in Sicilia et in Africa boreali-occidentali hinc inde detecta fuit.

S. PYGMÆA (*Sibth. et Sm. Fl. Gr. viii. 61, tab. 783*; *DC. Pr. vii. 123*).

Mar. merid.—In regione alpina Atlantis Majoris, in jugo Tagherot legi spec. nondum florentia quæ mihi incertæ sedis manebant donec cel. Cosson certum me fecit hæc ad *S. pygmæum*, *S. S.*, referenda esse, et spec. florentia, mense Junio 1874 in monte Djebel Ouensa ab indigeno Ibrahim lecta, benev. communicavit!

Ar. Geog. Distributio hujus sp. anomala est. Olim in monte Olympo Bithynico, nec alibi in Oriente, detecta, postea in summis caeminibus Atlantis Minoris Algeriæ (Djebel Che'liah, Djebel Tongour) reperta, et demum in Atlante Majori.

Cll. Rein et Fritsch in jugo Tagherot specimina incompleta *Scorzonera* sp. nanæ mox floribundæ legerunt, *S. pygmææ* certe affinis. In his folia subglaberrima, nec sericeo-lanata, et involuerum videtur diversum. Planta ulterius inquirenda.

SCORZONERA LACINIATA (*L., Boiss. Fl. Or.* iii. 757), = *Podospermum laciniatum* (*DC. Fl. Fr.* iv. 61, et *Prod.* vii. 111).

Mar. sept.—Inter Tanger et Tetuan!

Mar. merid.—In regione inferiore Atlantis Majoris haud infrequens. Aït Mesan usque 1600 m.! In convalle Amsmiz circ. 1400 m.! Distr. Reraya ad 1000 m.!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia). In Europa media calidiore.

Specimina nostra ambigua sunt inter typum et var. *calcitrapifoliam* (*Boiss. l. c.*), = *S. calcitrapifolia* (*Vahl, Symb.* ii. 87), formam foliorum segmentis latioribus.

LOBELIACEÆ (*Juss. Ann. Mus.* xviii. 1).

LAURENTIA (*Mich. Nov. Gen.* 18).

L. MICHELII (*A. DC. in Prod.* vii. 409), = *Lobelia Laurentia* (*L., Desf. Fl. Atl.* ii. 312; *Schsb. Gew. Marok*, 73).

Mar. sept.—In humentibus montis Djebel Kebir pr. Tanger (*Schsb. l. c.*).

Mar. occid.—Prope Aïn el Hadjar haud procul Mogador!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

CAMPANULACEÆ (*Juss. Gen.* 163).

JASTONE (*L. Hort. Cliff.* 426).

J. BLEPHARODON (*Boiss. et Rent. Pug.* 72).

Mar. occid.—Aïn el Hadjar haud procul Mogador!

Mar. merid.—Prov. Shedma juxta castellum gubernatoris

Ar. Geog. Peninsula Iberica. Marocco.

Verisimiliter ad subspeciem *J. montanæ*, *L.*, reducenda.

J. CORYMBOSA (*Poir. Dict. Suppl.* iii. 131; *DC. Pr.* vii. 415), = *J. montana* (*Schousb. Gew. Marok*, 70, non aliorum).

Mar. sept.—Tanger! (*Salzm. Exsicc.*, sub *J. arenaria*)! (*Schsb. l. c.*).
In monte Djebel Kebir!

Mar. occid.—Casa Blanca!

Ar. Geog. Africa boreali-occidentalis. Hispania Granatensis.

J. CORNUTA (*Ball in Journ. Bot.* 1873, 373).

Annua; caulis subsimplex in pedunculum nudum desinens; folia linearilanceolata, acuta, repando-dentata, superne vix decrescentia; involucri foliola 8–9, e basi lata ovata, longe acuminata, flores longitudine superantia, recurva, cum foliis et caule ciliato-pilosa; pedicelli cum calycibus glaberrimi; laciniae calycinae angustissimæ in setas capillaceas floribus æquilongas productæ; capitulum pro genere magnum.

Mar. occid.—In monte Djebel Hadid haud procul Mogador (*R. et Fr. spec. in Herb. Kew.*)! In collinis prov. Haha prope Mogador!

Mar. merid.—In prov. meridionalibus imper. Maroccani, sat frequens. Shedma! Ourika! Reraya! Mesfioua!

Species ex involucri foliolis cornuformibus insignis.

JASIONE ATLANTICA (*Ball in Journ. Bot.* 1873, 373).

Perennis; caules numerosi prostrati, ramis elongatis debilibus, sterilibus dense foliosis, floriferis superne nudis; folia obovato-linearia, obtusa, brevia; involucri foliola 12–13 ovata margine undulata, cum foliis et pedunculo glaberrima vel pube tenui leviter adspersa; pedicelli cum calycibus glaberrimi; laciniae calycinae breves vix quartam partem corollae longitudinis attingentes.

Mar. merid.—In regione superiore Atlantis Majoris. In convalle Aït Mesan! a 2000 m. ad 2500 m.

A. *J. amethystina*, Lag. et Rod., præter indumentum diversum, differt lacinii calycinis multo minoribus; sed forsân sicut hujus subspecies enumeranda.

CAMPANULA (*L. Gen.* 218).

C. DICHOTOMA (*L., DC. Pr.* vii. 462), = C. *afra* (*Cav. Anal. Cienc. Nat.* iii. 21).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Tetuan (1851)!

Mar. occid.—Casa Blanca! Salli (*Broussonnet, ex Cav. loc. cit.*).

Mar. merid.—Valde frequens in regione inferiore. Legimus in provinciis Haha! Shiedma! et Mtouga! et ad radices Atlantis Majoris ad Reraya! et Seksaoua! Ourika (*R. et Fr. spec. in Herb. Kew.*)!

— —, var. PARVIFLORA, = C. *Kremeri* (*Boiss. et Reut. Pug.* 75).

Mar. occid.—In monte Djebel Hadid! et juxta Aïn el Hadjar! haud procul Mogador.

Mar. merid.—In aridis juxta Mskala!

Inter specimina a nobis lecta nonnulla ambigua inter var. et formam normalem.

Ar. Geog. Regio Mediterranea occidentalis, imprimis calidior. Hispania austro-orientalis; Italia meridionalis. Africa borealis. In insulis Canariensibus obvia, deest in Madera. In Græcia erronee indicata. Var. nostra in Hispania et in Africa boreali obvia est.

C. MOLLIS (*L., DC. Pr.* vii. 463), = C. *velutina* (*Desf. Fl. Atl.* i. 180, tab. 51).

Mar. sept.—? Prope Tanger legit Broussonnet sec. Cavanilles (*Anal. Cienc. Nat.*).

Ar. Geog. Hispania australis. Imp. Maroccani dubia civis. Fertur in Creta, sed vix genuina.

A nemini recentiorum visa facile tamen in rupestribus Tingitanis invenienda. In promontorio Calpes hæc species erescit. Broussonnet

plantam genuinam ibi legit, et forsā eum plantis Tingitanis commiscuit.

Campanula maroccana, nobis.

C. molli proxima, differt tamen notis plurimis non parvi momenti. Indumentum ex pilis rigidis hispidulum, nec velutino-pubescens; caules ex pilis rigidiusculis præsertim in striis elevatis e basi foliorum decurrentibus setosi; folia viridia, sæpius erenato-denticulata, superiora cum calyce sæpe glabrescentia; laciniae calycinae magis oblongæ apice obtusiusculæ, appendicibus reflexis tubo calycino longioribus; corolla basin versus contracta, paullo longior, fere ad medium fissa lobis lanceolatis; calyx fructifer quinquenervius, nervis in pedunculo decurrentibus.

Mar. merid.—Valde frequens ad radices Atlantis Majoris et in convallibus inferioribus. Mesfioua! Ourika! Aït Mesan usque 1600 m.! Seksaoua! Amismiz! Milhaïn!

A *C. molli* ex forma calycis et corollæ diversa videtur probe distincta. An potius forma *C. flicaulis*, Coss. et Dur., sp. nondum descriptæ, cujus vidi spec. cultum in herb. Gay hodie Kewensi. Hæc tamen habet folia angustiora, et indumentum pareum videtur diversum.

C. ERINUS (L., DC. Pr. vii. 473).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Cap. Spartel! Tetuan (1851)

Mar. oecid.—Prope Mogador!

Mar. merid.—Ad radices Atlantis Majoris, et alibi in regione inferiore.

Mtougā! Aïn Tarsil! Visa etiam juxta Seksaoua! pr. Gurguri!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

C. RAPUNCULUS (L., DC. Pr. vii. 480).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Tetuan (1851)!

Mar. oecid.—In monte Djebel Hadid haud procul Mogador (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—Mesfioua! Ourika! In convalle Aït Mesan usque 1600 m.!

— —, var. CALYCE STRIGOSO, = *C. verruculosa* (*Link et Hffgg. Fl. Fort.* ii. 12, tab. 81).

Mar. sept.—Tanger! Djebel Dersa prope Tetuan (*Webb spec.*)!

Mar. oecid.—Casa Blanca! Mazagan!

Ar. Geog. Species per Europam omnem (excl. aretica), in Sibiria Uralensi, in Oriente et in Africa boreali diffusa. Var., vix enumeratū digna, hinc inde in reg. Mediterranea occurrit.

C. LÆFFLINGII (Brot. Fl. Lus., et Phyt. Lus. t. 18; DC. Pr. vii. 482),
= *C. erinoides* (*Cav. Anal. Cienc. Nat.* iii. 20; *DC. Pr.* vii. 480, an *L.* ?), = *C. Broussonnetiana* (*R. et Sch. Syst.* v. 104), = *C. Duriæi* (*Boiss. Voy. Esp.* 401). Huc etiam referendæ sunt velut varietates *C. Matritensis* (*A. DC. Mon. Camp.* 332; *DC. Pr.* vii. 481) et *C. decumbens* (*A. DC. Mon. Camp.* 334; *DC. Pr.* vii. 431).

Mar. sept.—Prope Tanger legit Broussonnet teste Cavanilles loc. cit. Tetuan (1851)!

Mar. occid.—Casa Blanca! Prov. Haha prope Mogador! Prope Mogador (*Broussonnet, teste A. DC.*).

Mar. merid.—In regione inferiore, præsertim ad radices Atlantis Majoris. Prov. Shedma! Prope Seksaoua! Mzouda! Formam debilem paucifloram (*A. decumbens, A. DC.*) legimus in convalle Aït Mesan circ. 1800 m.

Ar. Geog. Peninsula Iberica. Marocco.

Planta valde ludibunda. Variat habitu, caule erecto vel decumbente plus minus folioso, ramis paucis vel numerosis, patulis vel substrictis, corolla majore vel minore; et formæ diversæ haud raro simul cresunt. Spec. Maroccana sæpius majora quam Hispanica, pedalia et ultra, habitu *Speculariam Speculum* referentia.

Huic proximæ sunt *C. ramosissima*, Sibth. et Sm., *C. Kotschyana* (*A. DC. in Prod. vii. 481*) et *C. phrygia* (*Jaub. et Sp. Ill. Pl. Or. tab. 233*), quæ forsitan non satis diversæ sunt, et *C. specularioides* (*Coss. Pl. Crit. 41*).

SPECULARIA (*Heist. Syst. Pl. Gen. 8*).

S. FALCATA (*Ten. Prod. Fl. Neap. 16, sub Prismatocarpo; A. DC. Mon. Camp. 345, et Prod. vii. 489*).

Mar. merid.—Prope Amsniz! In monte Djebel Tezali 1700–1800 m.!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

Verisimiliter huic referenda est *Campanula Speculum* (*Schousb. Gew. Marok.*), dum nemini recentiorum hæc novissima obvia fuit.

TRACHELIUM (*L. Gen. 293*).

T. CÆRULEUM (*L., DC. Pr. vii. 491; Schousb. Gew. Marok. 71*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Brouss sec. Cav.*) Tetuan (1851)!

Mar. centr.—Prope Mequinez (*Schsb. l. c.*).

Mar. occid.—Prope Mogador (*Schsb. l. c.*).

Mar. merid.—Vidi plantam nondum florentem pr. Sheshaoua! et in distr. Mesfioua!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

T. ANGUSTIFOLIUM (*Schousb. Gew. Marok. 72; DC. Pr. vii. 491*).

Mar. centr.—Inter Mequinez et Fez (*Schsb. l. c.*).

Mar. merid.—Ad radices Atlantis Majoris et in convallibus calidioribus. Legimus in convalle Aït Mesan circ. 1400 m. In rupibus supra Seksaoua 1000–1200 m. et in rupibus Aïn Tarsil!

Ar. Geog. Species Maroccana.

Descriptio Schousbœana non accurata.

Radix perennis, crassa, lignosa, caules plurimos herbaceos erectos simplices foliaceos emittens; folia inferiora lineari-oblonga vel lineari-spathulata apicem versus denticulata, superiora angustiora, suprema

angustissima; corymbus contractus; calycis parvuli lacinia semimillimetrum longæ tubo subæquales; corolla demum valde elongata, tubo 10-12 millim. longo sæpius albido, laciniis $1\frac{1}{2}$ millim. longis acutis, cæruleis; stylus glaberrimus, minus quam in *T. cæruleo* exsertus; stigma clavatum, demum trifidum (nec bifidum); capsula immatura certe trilocularis; semina in quovis loculo 3-4). Ex errore Schonsbœi, qui se stigma bifidum vidisse credidit, cel. A. DeCandolle in Prodr. loc. cit. capsulam bilocularem descripsit.

ERICACEÆ (*Lindl. Nat. Syst.* 220).

CALLUNA (*Salisb. in Trans. Linn. Soc.* vi. 317).

C. VULGARIS (*L., et Schousb. Gew. Marok.* 174, sub *Erica*; *Salisb. l. c.*; *DC. Pr.* vii. 613).

Mar. sept.—In monte Djebel Kebir prope Tanger (*spec. ex herb. Schousb.*)! Tetuan!

Ar. Geog. Europa fere omnis a Scandinavia ad fretum Herculis. Asia Minor. Africa boreali-occidentalis. Versus orientem usque in Sibiriam Uralensem, et occidentem versus in insulas Islandiam et Terranovam se protendit.

ERICA (*L. Gen.* 192).

E. CILIARIS (*L., DC. Pr.* vii. 665).

Mar. sept.—Vidi folia in monte Djebel Kebir! *spec. florifera ex herb. Schousboc* prope Tanger lecta benev. comm. cel. Cosson!

Ar. Geog. Europa occidentalis ex Anglia usque fretum Herculis. Ditio Tingitana. Legi supra Algeciras.

E. AUSTRALIS (*L., DC. Pr.* vii. 666; *Schousb. Gew. Marok.* 175).

Mar. sept.—In monte Djebel Kebir prope Tanger! (*Schsb. l. c.*) (*Salzm. Exsicc.*)!

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

E. UMBELLATA (*L., DC. Pr.* vii. 666).

Mar. sept.—In monte Djebel Kebir pr. Tanger! (*Salzm. Exsicc.*, sub *E. umbellifera*)! Inter Tetuan et Ceuta!

Ar. Geog. Peninsula Iberica. Africa Tingitana.

E. MULTIFLORA (*L., DC. Pr.* vii. 667).

Mar. sept.—In monte Beni Hosmar prope Tetuan!

Ar. Geog. Regio Mediterranea ex Dalmatia usque fretum Herculis. Deest in Lusitania et in Græcia; conf. Boissier (*Fl. Or.* iii. 971). Forma proxime affinis in Anglia et in Gallia occidentali obvia est.

E. ARBOREA (*L., DC. Pr.* vii. 690; *Schousb. Gew. Marok.* 176).

Mar. sept.—In monte Djebel Kebir prope Tanger! (*Schsb. l. c.*). Tetuan!

Ar. Geog. Regio fere omnis Mediterranea (inclusa Macaronesia); in Oriente minus frequens.

ERICA SCOPARIA (L., DC. Pr. vii. 692).

Mar. sept.—In monte Djebel Kebir pr. Tanger! Cap. Spartel! Tetuan!
Ar. Geog. Regio Mediterranea occidentalis ex Dalmatia usque Macaronesiam. Gallia occidentalis a Pyrenæis usque Ligerim flumen.

ARBUTUS (L. Gen. 552).

A. UNEDO (L., DC. Pr. vii. 585).]

Mar. sept.—Vidi prope Tanger!

Mar. occid.—Legimus prope Aïn el Hadjar haud procul Mogador!

Mar. merid.—In convalle Aït Mesan circ. 1400 m. legit J. D. H.!

Ar. Geog. Regio Mediterranea fere omnis, imprimis occidentalis (exclusa Madera). Adest etiam in Gallia occidentali et in Hibernia australi.

PLUMBAGINEÆ.

STATICE (L. Gen. 388).

S. SINUATA (L., DC. Pr. xii. 635; Schousb. Gew. Marok.).

Mar. sept.—Tanger (*Schsb. l. c.*). Cap. Spartel! Inter Tetuan et Ceuta!

Mar. occid.—Mogador, *Lowe Cat.* (*Broussonnet, teste Cavanilles*). In provincia Haha haud procul Mogador!

Mar. merid.—Occurrit hinc inde in regione inferiore a mari remota. Prov. Shedma!

— —, var. INTEGRIFOLIA (var. β , L., *Boiss. in DC. Pr. l. c.*).

Mar. occid.—Casa Blanca! Specimina Maroccana in herb. DC. a Broussonnet lecta.

Ar. Geog. Species per regionem Mediterraneam (exclusa Macaronesia) late diffusa.

S. BEAUMIERIANA (*Coss. MSS.*)?

Mar. merid.—Legi in glareosis torrentis Ouel en Fisk prope Gurguri frustulum *Statice* e grege *S. sinuata*, ab hac tamen diversæ, verisimiliter ad sp. novam Cossonianam referendæ!

Ar. Geog. Species propria Imp. Maroccani meridionalis.

S. THOUINI (*Viv. Cat. H. di Negro*, 34; DC. Pr. xii. 636), = *S. sinuata*, var. γ (*L. Sp. Pl. i. 397*), = *S. alata* (*Willd. Enum. Hort. Berol. Suppl. 15*), = *S. Ægyptiaca* (*Pers. Syn., et Del. Fl. Eg. tab. 25*).

Mar. occid.—In arenosis a mari parum remotis prope Mogador! Prope Saffi! Agadir (*Coll. Indig. 1871*)!

Mar. merid.—Occurrit frequentissime in aridis saxosis regionis inferioris, imprimis in prov. Shedma! Ad radices Atlantis Majoris pr. Seksaoua!

Ar. Geog. Regio Mediterranea calidior. Africa borealis. Oriens. Græcia. Hispania australis. Fertur in Teneriffa (*fide spec. Broussonnetiano*) sed a recentioribus non visa.

STATICE MUCRONATA (*L. fil. Suppl.* 187; *DC. Pr.* xii. 639).

Mar. occid.—Sallé (*Broussonnet, ex Cav. Anal. Cienc. Nat.* iii. 38).

Mazagan! Saffi! Mogador in arenosis et in insula! ubi eam olim legit Broussonnet (*teste Cavan. l. c.*) et postea Lowe et Balansa.

Ar. Geog. Species propria reg. litoralis Atlantici Imp. Marocani.

Descriptio Boissieriana in *DC. Pr.* l. e. ex spec. incompletis haud omnino accurata. Folia radicalia late obovata vel sæpius suborbiculata in petiolum longum sensim attenuata. Ex planta in dumetis nascente (ut fere semper fit in ditione Mogadorensi) rami elati 2–3-pedales foliis carent, vel ex axillis squamarum fasciculos foliorum edunt (var. vel potius lusius a cel. Boissier memoratus); sed spec. humiliora quæ legi pr. Saffi et Mazagan in apricis enata gaudent ramis sterilibus foliaceis, foliis late orbicularibus in petiolum limbo æquilongum abrupte attenuatis. Alæ ramorum sæpissime, sed non semper, crispatae. Sp. nostræ proxima est *S. pectinata*, Ait., floræ Canariensis civis; videtur tamen satis distincta.

S. OVALIFOLIA (*Poir. Dict. Suppl.* v. 237; *DC. Pr.* xii. 646), = *S. auriculæfolia* (*Brot. et Vahl, Symb. pro parte*), = *S. mucosa* (*Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Spec. Maroceana legit Broussonnet quoad loc. natalem dubia.

Ar. Geog. Habitat in litore occidentali Galliæ, in peninsula Iberica, in litore Marocano, et in Macaronesia.

S. LYCHNIDIFOLIA (*Girard in Ann. Sc. Nat.* xvii. 18, tab. 3 A; *DC. Pr.* xii. 646).

Mar. sept.—Tetuan ad ostium fluminis (1851)! Tanger (*Salzm. Exsicc.*, sub *S. auriculæfolia*)!

Ar. Geog. Peninsula Iberica. Gallia occidentalis et australis. Africa Tingitana.

Spec. nostra ad var., seu potius formam, panienula confertiore (var. β , *corymbosa*, Boiss. in *Prod. loc. cit.*) referenda sunt.

S. OLEÆFOLIA (*Scop. Delic. Insubr.* i. 24, tab. 10), = *S. auriculæfolia* (*Vahl, Symb. pro parte*, et verisimiliter *Schousb. Gew. Marok.* 133), = *S. Scopoliana* (*Bertol. Fl. It.* iii. 528), = *S. densiflora* (*Guss. Prod. Fl. Sic. Suppl.*, non *Girard*), = *S. oxylepis* (*Boiss. in DC. Pr.* xii. 647).

Mar. sept.—? Tanger (*Schsb. l. c.*) (*Broussonnet, teste Cav. Anal. Cienc. Nat.* iii. 40).

Mar. occid.—? Mogador (*Schsb. l. c.*).

Ar. Geog. Italia continentalis et insulæ. Africa boreali-occidentalis.

Floræ Marocanæ dubia eivis, dum synonyma Schousbœi et Cavanillesii incerta, et nemini recentiorum in litore Tingitano species obvia fuit.

Cel. Boissier tamen spec. autoptica prope Oran leeta enumerat, et facile in territorio Marocano confini invenienda erit.

Cel. Boissier nomen triviale Scopolianum antiquius et satis aptum mu-

tavit, dum a Willdenovio, aliisque recentioribus, eum sp. alienis hoc confusum est. Pace tanti viri jus prioritatis illæsum manet.

STATICE OCCIDENTALIS (Lloyd, *Fl. Loire Inf.* 212; *DC. Pr.* xii. 648).

Mar. sept.—Tanger (*Durand, teste Boiss. in DC. Pr. l. c.*).

Ar. Geog. Litora Atlantica Europæ a Scotia usque Hispaniam borealem. Ditio Tingitana.

S. SPATHULATA (*Desf. Fl. Atl.* i. 275; *DC. Pr.* xii. 649), var. β , *EMARGINATA* (*Boiss. in Prod. l. c.*).

Mar. sept.—? Nondum florentem legi in maritimis inter Tetuan et Centa!

Ar. Geog. Species in litore Mediterraneo Africæ borealis; varietas ad Gibraltar et in ditone Tetuanensi obvia.

Ob spec. incompleta haud omnino certus sum de hac planta, quæ tamen spec. Calpensibus adamussim convenit.

S. GUMMIFERA (*Durieu in Boiss. et Reut. Pug.* 104), var. *CORYMBULOSA* (*Coss. Not. Crit.* 175; *Walp. Ann. Bot. Syst.* v. 716).

Mar. sept.—In insulis Zafarinis legit P. B. Webb, ex spec. in herb. nost.!

Ar. Geog. Species in Algeria prope Oran detecta; var. in Hispania austro-orientali et in ins. Maroccanis Zafarinis obvia.

Ex sect. *Limonio* adsunt *Statices* spec. duo mihi nimis incerta, nempe spec. incompletum nondum florens a me juxta Casa Blanca lectum, verisimiliter ad *S. delicatulam*, Girard referendum, alterum a ell. Rein et Fritsch prope Mazagan lectum, *S. lychnidifoliæ* subsimile, sed foliis apice non acuminatis diversum. Hujus gregis species plurimæ notis instabilibus dignoscendæ ad varietates vel subspecies reducendæ videntur.

S. ECHIOIDES (L., *DC. Pr.* xii. 665).

Mar. occid.—Frequens in saxosis prope Mogador a litore maris remotiusculis! Agadir (*Coll. Indig.* 1871)!

Ar. Geog. Regio Mediterranea imprimis litoralis, interdum in salsuginosis a mari remotis.

S. ORNATA, nob., = *S. læta* (*Ball in Journ. Bot.* 1875, 176, non *Moris, Fl. Sard.* iii. 42).

Planta perennis, ex caudice lignoso caules vel scapos plurimos, erectos, 2-4-pedales, subaphyllos emittens; folia radicalia glabra, lanceolato-spathulata, obtusissima, in petiolum sensim attenuata, caulina (non nisi in caulibus truncatis vel emorsis) radicalibus similia sed breviora, cætera ad squamas reducta; squamæ triangulares, parvæ, margine vix scariosæ; caules scapiformes teretes, in ramos strictos alternos dichotome divisi, subglaberrimi vel parce furfuracci; rami inferiores steriles abbreviati, superiores ramulosi, ramulis florigeris subarenatim patentibus; spiculæ unifloræ vel rarius bifloræ, rectiusculæ, in spicas laxas uni-

laterales dispositæ; spicularum bracteæ subæquales minutæ, margine scariosæ, acutiusculæ; calyx profunde 5-costatus, tubo basi tantum bractea suffulto, limbo brevissimo ex dentibus quinque coloratis cuspidatis, versus basin scarioso-marginatis; corolla parva, pulchre roseo-purpurea, tubo ex petalorum unguibus coalitis quam calyx sesquilongiore, limbo libero; filamenta tubo corollino ultra medium adnata; antheræ exsertæ; styli liberi.

Mar. merid.—Legimus ad ripas fl. Oued En Fys juxta Misra ben Kara! Species distinctissima in § *Polyarthrion*, Boiss., collocanda, cui proxima est *St. cæsia*, Girard, Ann. Sc. Nat. ser. 3, vol. ii. p. 325; Boiss. in DC. Prod. xii. 667; et melius a Cosson (Notes sur qu. Pl. d'Espagne, p. 176) definita. Ab hac nostra differt bracteis spicularum æqualibus, nec suprema subquadruplo longiore, calycis limbi parte scariosa ad marginem angustum versus basin dentium reducta, petalorum unguibus in tubum coalitis, nec medium versus liberis, staminibus amplius adnatis, antheris exsertis.

STATICE FERULACEA (*L.*, DC. Pr. xii. 668; *Schousb. Gew. Marok.* 134).

Mar. sept.—Tanger! (*Webb Herb.*)! (*Salzm. Exsicc.*)! (*Schsb. l. c.*).

Tetuan ad ostium fluminis (1851)!

Mar. oecid.—Casa Blanea (*J. D. H.*)! Mazagan (*R. et Fr. sp. in Herb. Kew.*)! Mogador (*Schsb. l. c.*).

Ar. Geog. Regio Mediterranea occidentalis (inclusa Maearonesia).

ARMERIA (*Willd. Enum. H. Berol.* 333).

A. GADITANA (*Boiss. in DC. Pr. xii. 675*), var. TINGITANA, nob., = A. tingitana (*Boiss. et Reut. Pug.* 102). Verisimiliter huc *Statice cephalotes* (*Schousb. Gew. Marok.* 132).

Mar. sept.—Legimus juxta Cap. Spartel, ubi jam legerant Boiss. et Reuter!

Ar. Geog. Peninsula Iberica austro-occidentalis. Var. nostra nonnisi in ditone Tingitana.

A. tingitana ab A. gaditana differt phyllis involueri latioribus, calycis calcaris brevioris, et pedicello paullo brevioris. Hæ notæ mihi videntur nimis inconstantes. Quoad characteres a forma foveolæ basilaris desumptos vereor ne hi fallaces sint. In spec. authenticis A. gaditanæ video foveolas quoad formam valde ludibundas.

A. MAURITANICA (*Wallr. Monog. Arm.* 217; DC. Pr. xii. 676).

Mar. sept.—Inter Tetuan et Ceuta! Formam foliis angustioribus pr. Tetuan legit P. B. Webb (spec. sub "*A. alliacea*," ex herb. Webb)!

Mar. oecid.—Casa Blanca!

Ar. Geog. Africa borealis. Varietas (*A. Welwitschii*, Boiss.) in Lusitania. Hæc est stirps variabilis. Involueri phylla interdum omnia mucronulata, interdum interiora mutica exteriora mucronulata, interdum (in spec. Algeriensibus a ecl. Cosson determinatis) phylla fere omnia mutica, pauca externa debiliter mucronulata. Vidi semel

erescentes formam foliis margine eiliolatis et formam f. glaberrimis donatam. Forsan ad formam angustifoliam referenda est *Statice alliacea* (Schousb. Gew. Marok. 131) in arenosis Tingitanis indicata.

ARMERIA PLANTAGINEA (*All. Fl. Ped.* ii. 90, sub *Statice*; *Willd. Enum. II. Berol.* 334; *DC. Pr.* xii. 683).

Mar. merid.—In regione media Atlantis Majoris. Legimus mox floribundam in monte Djebel Tezah a 2000 m. usque 2400 m.!

Ar. Geog. Europa media. Montes Hispaniæ, Lusitaniæ, Galliæ australis et Italiæ superioris.

Spec. nostra verisimiliter ad var. *leucantham* (Boiss. in *DC. Pr. l. c.*) referenda. Spec. altera nimis immatura, in convalle Aït Mesan eire. 2500 m. lecta, forsan potius ad *A. allioidem*, Boiss. tribuenda sunt.

LIMONIASTRUM (*Mœnch. Meth.* 423).

L. MONOPETALUM (L., et *Schousb. Gew. Marok.* 133, sub *Statice*; *Boiss. in DC. Pr.* xii. 689).

Mar. sept.—Tanger (*Schsb. l. c.*).

Mar. occid.—El Araïsch (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio Mediterranea, exclusa Græcia, Oriente et Macaronesia.

PLUMBAGO (*L. Gen.* 213).

P. EUROPÆA (L., *DC. Pr.* xii. 691), = P. lapathifolia (*Schousb. Gew. Marok.* 67, non *Willd.*).

Mar. occid.—Casa Blanca! Saffi (*Schsb. l. c.*). Agadir (*Coll. Indig.*)!

Mar. merid.—In regione inferiore haud infrequens. Amsmiz! Milhaïn!

Mtougā! Prov. Shedma (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

PRIMULACEÆ (*Vent. Tabl.* ii. 285).

CORIS (*L. Gen.* 243).

C. MONSPELIENSIS (L., *DC. Pr.* viii. 59; *Schousb. Gew. Marok.* 75).

Mar. sept.—Tetuan (1851)! Spec. a P. B. Webb lecta, verisimiliter etiam pr. Tetuan, in herb. Webb conservantur.

Mar. occid.—Prope Mogador! (*Balansa spec.* 1867)! In prov. Haha, verisim. haud procul Mogador (*Schsb. l. c.*). Prope Agadir (*Coll. Indig.* 1871)!

Ar. Geog. Regio Mediterranea occidentalis a Sicilia et Algeria usque litora Atlantica Marocana.

ASTEROLINUM (*Lk. et Hffgg. Fl. Port.* 333).

A. LINUM STELLATUM (L., sub *Lysimachia*; *Lk. et Hffgg. l. c.*; *DC. Pr.* viii. 68).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! In monte Djebel Kebir pr. Tanger!

Mar. merid.—In regione media et inferiore Atlantis Majoris. In convalle Amsmiz circ. 1400 m.! In monte Djebel Tezah usque 1800 m.!

Distr. Reraya eire. 1200 m.!

Ar. Geog. Regio omnis Mediterranea (exclusa Maearonesia).

ANAGALLIS (*L. Gen.* 206).

A. ARVENSIS (*L., DC. Pr.* viii. 69).

Mar. sept.—Tanger! (*Salzm. Exsicc.*, sub *A. parviflora*)! Tetuan (*Webb spec.*)!

Mar. oecid.—Mogador (*Lowe Cat.*).

— —, var. LATIFOLIA (*Lge. Pug.* 221; *Willk. et Lge. Fl. Hisp.* ii. 648), = *A. latifolia* (*L., DC. Pr.* viii. 70; *Schousb. Gew. Marok.* 66).

Mar. sept.—In monte Djebel Kebir pr. Tanger! Cap. Spartel! Tetuan (*Webb spec.*, sub *A. cærulea*)!

Ar. Geog. Species per orbem fere totum (reg. glacialibus exclusis) diffusa. Varietas per reg. Mediterraneam, Indiam Orientalem et Americam australem dispersa.

A. latifoliam, nulla nota constante dignoscendam, ad var. cl. Lange optimo consilio reduxit. Tam hæc quam var. typica ludunt floribus phœniceis vel cæruleis.

A. LINIFOLIA (*L., DC. Pr.* viii. 70; *Willk. et Lge. Fl. Hisp.* ii. 648), = *A. Monelli* (*Schousb. Gew. Marok.* et verisimiliter *L. Sp. Pl.* i. 211, excl. loc. nat. Verona).

Mar. sept.—Tetuan (*Webb Herb.*, sub *A. collina*)!

Mar. oecid.—In agris prov. Haha (*Schsb. l. c.*). Hue verisimiliter spec. incompletum a me lectum in monte Djebel Hadid hand procul Mogador!

Mar. merid.—In convalle Aït Mesan eire. 1400 m.!

— —, var. MICROPHYLLA, *nob.* In convalle Aït Mesan supra Arround ad 2400 m. s. m.!

Forma insignis multicaulis radice suffrutescente, caulibus adscendentibus, foliis minimis lanceolatis, corollis quam in typo minoribus.

— —, var. COLLINA, *nob.* = *A. collina* (*Schousb. Gew. Marok.* 64; *DC. Pr.* viii. 70). Non differt nisi statura sæpius majore, caulibus crassioribus basi frutescentibus, pedicellis fructiferis longioribus, corolla rosea vel purpurasecente, nec cærulea. In utraque forma foliola variant latiora vel angustiora.

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Spec. ex Herb. Schousb.*)!

Mar. oecid.—Casa Blanca! Mazagan! Aïn el Hadjar, hand procul Mogador! Prov. Haha pr. Mogador! (*Schsb. l. c.*). Agadir (*Coll. Indig.* 1871)!

Specimina Tingitana, nisi colore florum, a forma latifolia *A. linifoliae* Hispaniæ non distinguenda sunt. Confer *Willk. et Lge. Fl. Hisp.* ii. 649.

Ar. Geog. Tam *A. linifolia* quam *A. collina* peninsulæ Ibericæ et Africæ boreali-occidentalis cives sunt. *A. collina* in Sardinia a eel. Moris indicatur.

ANAGALLIS CRASSIFOLIA (*Thor. Chlor. Land.* 62; *DC. Pr.* viii. 70.)

Mar. sept.—In monte Djebel Kebir prope Tanger! (*Webb Herb.*)! (*Salzm. Exsicc.*)!

Ar. Geog. In Gallia austro-occidentali, in peninsula Iberica et in agro Tingitano, ubique rarius obvia.

A. TENELLA (*L., DC. Pr.* viii. 71).

Mar. sept.—Cel. Cosson benevole comm. spec. a Schousboeo lecta! Ego semel vidi folia, sed spec. nulla legi!

Ar. Geog. Europa occidentalis. Africa boreali-occidentalis.

Occurrit etiam (quod mirum) in insula Creta.

SAMOLUS (*L. Gen.* 222).

S. VALERANDI (*L., DC. Pr.* viii. 73; *Schousb. Gew. Marok.* 73).

Mar. sept.—Tetuan (1851)! "Ad fontes regionis Tingitanæ" (*Schousb. l. c.*).

Mar. occid.—Casa Blanca! Mogador! (*Lowe Cat.*). Juxta Aïn el Hadjar!

Ar. Geog. Species cosmopolita per orbem fere totum dispersa.

SAPOTACEÆ (*Juss. Gen.* 151).

ARGANIA (*R. et Sch. Syst.* iv. 502).

A. SIDEROXYLON (*R. et Sch. l. c.*; *DC. Pr.* viii. 187), = Sideroxylon spinosum (*L. Hort. Cliff.* 69; *ex A. DC. in Prod. l. c.*), = Elæodendron Argan (*Retz. Obs.* vi. 26; *Schousb. Gew. Marok.* 89).

Mar. occid.—Per zonam occidentalem Imperii Maroccani inter flumina Oumerbiali et Sous, et forsan, referentibus indigenis, usque fl. Oued. Noun. Prævalet in regione subltorca usque 20 chilometra ab Oceano remota, sed sparsim occurrit in terris interioribus usque 40 chilometra a mari longinqua. Legimus jam florentem decedente Aprili!

Arbor diametro demum 3-4-pedali, ramis patentibus, vetusta altitudine sua latior. Incolis oleum subamarum ad ciborum condimentum usitatissimum et lignum grave durissimum suppeditat. Fusius de hac stirpe Floræ Maroccanæ cive insigni in Schousboci opere citato et in Jackson.

Ar. Geog. Unicum *Sapotacearum* genus extratropicale, nisi in territorio Maroccano nullibi visum.

OLEACEÆ (*Lindl. Nat. Syst.* 307).

JASMINUM (*L. Gen.* 17).

J. FRUTICANS (*L., DC. Pr.* viii. 313; *Schousb. Gew. Marok.* 2).

Mar. occid.—Mogador (*Lowe Cat.*). In monte Djebel Hadid haud procul Mogador! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In saxosis reg. inferioris Atlantis Majoris. Tasseremout !
 Aït Mesan usque 1400 m. ! et alibi visum.
Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

FRAXINUS (*L. Gen.* 1160).

F. OXYPHYLLA (*M. B. Fl. Taurico-cauc.* ii. 450; *DC. Pr.* viii. 276), =
F. rostrata (*Guss. Pl. Rar.* 374, tab. 63; *DC. Pr. l. c.*), = *F. australis* (*auct. Gall.*; vix *F. excelsior* β , *australis*, *J. Gay in Camb. Enum. Bal.*).

Mar. sept.—Inter Tanger et Tetuan ! Tetuan (*Webb Herb.*) !

Mar. merid.—Pluries visa in regione inferiore et ad radices Atlantis Majoris. Ourika ! Reraya ! In convalle Aït Mesan usque 1700 m. !

Ar. Geog. Regio Mediterranea (exclusa Macaronesia). Regio Danubialis. Rossia australis.

Characteres quibus hæc a *F. excelsiore* separatur non graves; sed videntur satis constantes. Folia semper minora, subcoriacea, jugis paucioribus, pinnis acutioribus; samaræ angustiores, semper integræ, dum in *F. excelsiore* sæpius emarginatæ sunt. Pro subspecie enumeranda ?

F. DIMORPHA (*Coss. et Dur. in Bull. Soc. Bot. Fr.* ii. 367).

Mar. merid.—In convalle Ourika eire. 1000 m. ! In convalle Aït Mesan (spec. foliis tantum donatum legit J. D. H.) !

Planta singularis in statu dumetoso sterili a nobis visa pro *Fraxino* difficile habenda. Rami debiles 2–3-pedales ereeti, folia parvula, foliola variant ovata acuta vel suborbicularia obtusa.

PHILLYREA (*L. Gen.* 19).

P. LATIFOLIA (*L., DC. Pr.* viii. 292; *Schousb. Gew. Marok.* 3).

Mar. sept.—In monte Djebel Kebir pr. Tanger ! (*Schsb. l. c.*). In monte Beni Hosmar prope Tetuan (*J. D. H.*) !

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

P. MEDIA (*L., DC. Pr.* viii. 292; *Schousb. Gew. Marok.* 2).

Mar. occid.—Prope Mogador (*Lowe Cat.*), (*Schsb. l. c.*). In monte Djebel Hadid haud procul Mogador !

Mar. merid.—Pluries visa ad radices Atlantis Majoris. Ourika (*J. D. H.*) !

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

P. ANGUSTIFOLIA (*L., DC. Pr.* viii. 292).

Mar. sept.—In monte Djebel Kebir prope Tanger !

Hæc species a beato Lowe indicatur dubitative (signo interrogationis) prope Mogador; forsitan errore aliquo, dum nemini in hac ditione obvia fuit.

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

OLEA (*L. Gen.* 20).

O. EUROPEA (*L., DC. Pr.* viii. 284; *Schousb. Gew. Marok.* 3).

Mar. sept.—Frequens in ditione Tingitana; sæpius culta, sed in collibus saxosis facta spontanea et forsā indigena!

Mar. centr.—Circa Mequinez et Fez (*Schousb. l. c.*).

Mar. occid.—Frequens in regione inferiore. In monte Djebel Hadid prope Mogador videtur spontanea!

Mar. merid.—Colitur hinc inde in regione inferiore et in omnibus convallibus Atlantis Majoris usque 1500–1600 m. Interdum facta spontanea forma fruticosa spinosa in rupestribus obvia est!

Ar. Geog. Arbor diu culta, verisimiliter in Africa boreali et in Oriente indigena, hodie facta spontanea in calidioribus totius regionis Mediterraneæ.

APOCYNACEÆ (*Lindl. Nat. Syst.* 299).VINCA (*L. Gen.* 295).

V. MEDIA (*Lk. et Hffgg. Fl. Port.* tab. 90; *DC. Pr.* viii. 384), = V. major (*Schousb. Gew. Marok.* 106, non *L.*).

Mar. sept.—Frequens in sepibus et dumetis pr. Tanger! (*Salzm. Exsicc.*)! (*Schsb. l. c.*).

Ar. Geog. Peninsula Iberica. Italia meridionalis. Sardinia. Africa borealis.

NERIUM (*L. Gen.* 181).

N. OLEANDER (*L., DC. Pr.* viii. 420; *Schousb. Gew. Marok.* 106).

Mar. sept.—Ad ripas fluminum prope Tanger (*Schousb. l. c.*). Inter Tanger et Tetuan, et prope Tetuan a nobis visum!

Mar. occid.—Prope Mogador (*Lowe Cat.*). Prov. Haha haud procul Mogador!

Mar. merid.—Ad radices Atlantis Majoris et in convallibus inferioribus, fere semper ad ripas torrentium. Amsmiz! In convalle Aït Mesan et alibi usque 1500 m. observatum! Rarius occurrit in planitie. Secus rivulum ad fines prov. Haha et Shedma!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

ASCLEPIADEÆ (*R. Br. in Wern. Trans.* i. 12).PERIPLOCA (*L. Gen.* 300).

P. LÆVIGATA (*Ait. H. Kew.* i. 301; *DC. Pr.* viii. 498), = P. angustifolia (*Labill. Pl. Syr. Dec.* ii. 13, tab. 7; *Schousb. Gew. Marok.* 106).

Mar. occid.—Prope Mogador! (*Lowe Cat.*), (*Schsb. l. c.*). Magis frequens circa Aïn el Hadjar! et in parte inferiore montis Djebel Hadid!

Ar. Geog. Species late diffusa sed nullibi frequens. In Syria maritima semel legit Labillardière. In Lampedusa cæterisque insulis minoribus inter Siciliam et oras Africanas, et in Hispania austro-orientali rarius occurrit. Adest in Africa boreali, et in ins. Canariensibus; deest tamen in Madera.

GOMPHOCARPUS (*R. Br. in Wern. Trans. i. 37*).

G. FRUTICOSUS (*L., sub Asclepiade; R. Br. loc. cit.; DC. Pr. viii. 557*).

Mar. sept.—Prope Tanger spontaneus sed vix indigenus! Tetuan (*Webb spec.*)!

Ar. Geog. Fertur ex Arabia (ubi certe indigenus) per regionem Mediterraneam Europæ calidiorē et per Africam fere totam iter fecisse usque in insulas Canarienses et in Madagasear.

Inter plantas a beato Webb in Mauritania lectas video specimen ad *Gymnema subvolubile* (Decne. in *Ann. Sc. Nat. ix. 277*, et in *DC. Pr. viii. 621*) referendum. Hæc est species per Africam tropicam occidentalem late diffusa, sed a nemini in Imp. Maroccano observata, et vix indigena.

BOUCEROSIA (*Wight et Arn. in Contr. Bot. Ind. 34*).

B. MAROCCANA (*Hook. fil. in Bot. Mag. c. 6137*), = *Stapelia quadrangula* (*Schousb. Gew. Marok. 107, non Forsk.*).

“Ramis tetragonis marginibus angulato-sinuatis, foliis trulliformibus acutis, corollæ lobis nudis tubo intus hirsuto, coronæ stamineæ processibus quinque incurvis gynostegio ineumbentibus cum decem erectis geminatim collateralibus capitatis alternantibus.”

Mar. occid.—Legimus in insula Mogador! ubi olim legit Schousboe, et in regione litorali prov. Haha!

Mar. merid.—Videtur satis frequens in regione inferiore usque radices Atlantis Majoris. Onrika! Milhaïn! Mtouga! et alibi visa.

Proxime affinis *B. Gussonianæ* (Mikan in *N. Ac. Nat. Cur. xvii. 594*, tab. 41, et Desne. in *DC. Pr. viii. 649*, sub *Apteranthe*) quacum eam confudimus donec flores ex plantis in Angliam allatis vidit eel. J. D. Hooker et notas diagnosticas loc. cit. indicavit. Planta nostra differt ramorum costis in lobos triangulares elevatis, nec sinuato-undulatis, folio in apice ejusvis lobi trulliformi basin versus contracto, sæpius deflexo, nec sessili basi adnato adscendente. Flores pauciores longius pedicellati, corollæ lobis brevioribus minus reflexis, nudis nec, ut in *B. Gussoniana*, ciliatis. Coronæ stamineæ structura diversa. Lobi exteriores per paria cum interioribus quinque ineumbentibus alternantes in sp. nostra longitudine subæquales erecti, dum in *B. Gussoniana* ad globulos sessiles reducti sunt. Qui sententiam celeb. Benth. et II. fil. genus *Apteranthe* cum *Boucerosia* conjungendum esse ægre tulit, ex planta nostra edoctus facile consentiet.

GENTIANÆ (*Juss. Gen.* 141).MICROCALA (*Lk. et Hffgg. Fl. Port.* i. 359).

M. FILIFORMIS (*L.*, sub *Gentiana*; *Willd. Sp. Pl.* i. 633, et *Schousb. Gew. Marok.* 47, sub *Exaco*; *Lk. et Hffgg. l. c.*; *DC. Pr.* ix. 62).

Mar. sept.—Prope Tanger in subhumidis montis Djebel Kebir (*Schsb. l. c.*).

Ar. Geog. Europa occidentalis ex Lapponia usque fretum Hereulis. In Europa Mediterranea infrequens. In Græcia olim a Sibthorp lecta; prope Byzantium unico loco nascitur (*ex Boiss. Fl. Or.*).

CHLORA (*L. Gen.* 1258).

C. PERFOLIATA (*L.*, *DC. Pr.* ix. 69; *Schousb. Gew. Marok.* 172).

Mar. sept.—Rarius occurrit in ditone Tingitana (*Schsb. l. c.*). In monte Beni Hosmar nondum florens, forsân ad var. sequentem referenda !

Mar. merid.—Ad radices Atlantis Majoris. Prope Ourika ! Mesfioua (*J. D. H.*) ! Tasseremout (*J. D. H.*) ! Prov. Mtouga (*R. et Fr. spec. in Herb. Kew.*) !

Ar. Geog. Europa occidentalis. Regio omnis Mediterranea (exclusa Macaronesia).

— —, var. MASCARIENSIS (*Desf. Fl. Atl.* i. 327), = *C. grandiflora* (*Viv. Cors. App. Atl.* 4).

Mar. sept.—Prope Tetuan (1851) ! (*Webb Herb.*) !

Ar. Geog. Africa borealis. Corsica. Sardinia.

Hæc forma forsân sicut subspecies enumeranda est. Videtur satis constans, caulibus tenuioribus, cyma laxiore, corollæ duplo majoris segmentis basi transverse rugosis distinguenda.

C. IMPERFOLIATA (*L. fil. Suppl.* 218; *DC. Pr.* ix. 69).

Mar. sept.—Tanger (*Salzm. Exsicc.*, sub *C. crenulata*) ! (*spec. ex Herb. Schousboe*) !

Ar. Geog. Hispania. Gallia meridionalis. Ager Tingitanus. Ins. Canarienses.

ERYTHRÆA (*Rich. in Pers. Syn.* i. 283).

E. RAMOSISSIMA (*Vill. Dauph.* ii. 380, sub *Gentiana*; *Rich. in Pers. Syn.* i. 283; *DC. Pr.* ix. 57), = *Gentiana Centaurium*, var. β (*L. Sp. Pl.* i. 333), = *Chironia erythræa* (*Schousb. Gew. Marok.* 81).

Mar. sept.—Prope Tanger ! In dumetis montis Djebel Kebir (*Schsb. l. c.*). Prope Tetuan (1851) ! Ad ripas fluminis Tetuanensis (1851) !

Mar. occid.—Prope Mogador (*J. D. H.*) !

Mar. merid.—Mesfioua ! Inter Sektana et Frouga (*G. Maw*) ! Spec. ex Oasi Akka a Mardoehæo lectum benev. comm. cel. Cosson !

Ar. Geog. Europa occidentalis et centralis. Regio omnis Mediterranea

inclusa Madera, sed (?) exclusis ins. Canariensibus. Adest etiam in Asia temperata.

Species variabilis, quæ, nisi fallor, includit *E. pulchellam*, Fl. Dan., *E. latifoliam*, Sm., *E. tenuifloram*, Link, aliasque s. d. species Orientales. Synonymon Schousboeanum ob diagnosin maneam incertum.

ERYTHRÆA CENTAURIUM (*L.*, sub *Gentiana*; *Pers. Syn.* i. 283; *DC. Pr.* ix. 58), = *Chironia Centaurium* (*Schousb. Gew. Marok.* 81).

Mar. sept.—Juxta Cap. Spartel!

Mar. occid.—Casa Blanca! Prope Mazagan visa! Mogador (*Lowe Cat.*). Aïn el Hadjar! et in monte Djebel Hadid haud procul Mogador! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In prov. Shedma (*J. D. H.*)! Mesfioua! Milhaïn!

—— —, var. *SUFFRUTICOSA* (*Griseb. Mon. Gent.* 140, et in *DC. Pr.* ix. 70), = *Chironia suffruticosa* (*Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)! In monte Djebel Kebir! In monte Djebel Dersa prope Tetuan (*Webb spec.*)!

Ar. Geog. Species per Europam mediam et regionem omnem Mediterraneam (exclusa Madera) late diffusa; var. in Hispania et in Africa boreali.

Stirps quam maxime ludibunda, in plurimas species et varietates ab auctoribus dilaniata notis inconstantibus definitas. Inutile duco varietates ex florum magnitudine vel ex foliorum forma enumerare, dum formæ intermediæ passim obviæ sunt.

E. SPICATA (*L.*, sub *Gentiana*; *Pers. Syn.* i. 283; *DC. Pr.* ix. 60), = *Chironia spicata* (*Schousb. Gew. Marok.* 86).

Mar. sept.—Tanger (*Schsb. l. c. spec. comm. Coss.*)!

Mar. occid.—Mogador (*Lowe Cat.*).

Ar. Geog. Regio fere omnis Mediterranea; in Madera deest.

E. MARITIMA (*L.*, sub *Gentiana*; *Pers. Syn.* i. 283; *DC. Pr.* ix. 60), = *Chironia maritima* (*Willd. Sp. Pl.* i. 1069; *Schousb. Gew. Marok.* 85).

Mar. sept.—In aridis montis Djebel Kebir pr. Tanger (*Schsb. l. c. spec. comm. Cosson*)!

Ar. Geog. Habitat in maritimis fere omnis reg. Mediterraneæ (inclusa Macaronesia) et in Gallia occidentali et in Persia boreali.

CICENDIA (*Adans. Fam. Pl.* ii. 503).

C. PUSILLA (*Lam. Dict.* ii. 645, sub *Gentiana*; *DC. Fl. Fr.* iii. 663, sub *Exaco*; *Griseb. Mon. Gent.* 156, et in *DC. Pr.* ix. 61).

Mar. sept.—? Legi hanc sp. facile prætervisam in herbosis subhumidis prope S. Roque haud procul Gibraltar, et ex notula MS. (nescio quam auctoritate allata) ipsam in agro Tingitano obviam esse credo. Specimina Tingitana nulla vidi. Planta ulterius inquirenda.

Ar. Geog. Gallia media et occidentalis. Hispania borealis et centralis,

et verisimiliter in utroque litore freti Herculis. In Italia media et in Sardinia etiam obvia.

BORAGINÆ (*Juss. Gen.* 128)*.

HELIOTROPIUM (*L. Gen.* 179).

II. SUPINUM (*L., DC. Pr.* ix. 533).

Mar. sept.—Tanger (*Salzm. Exsicc.*), ex *DC. Pr. l. c.*

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia). Africa tropica et (?) australis. Formæ proxime affines in India orientali.

II. EUROPÆUM (*L., DC. Pr.* ix. 534; *Schousb. Gew. Marok.* 59).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. occid.—Casa Blanca (*R. et Fr. spec. in Herb. Kew.*)! Mazagan!

Mar. merid.—In prov. Shedina! et alibi visum loco non notato.

Ar. Geog. Per regionem omnem Mediterraneam diffusum. In Europa media, forsan introductum.

II. UNDULATUM (*Vahl, Symb.* i. 13; *DC. Pr.* ix. 536; *Schousb. Gew. Marok.* 60), = *H. crispum* (*Desf. Fl. Atl.* i. 151, tab. 41), = *II. maroccanum* (*Lehm. Asper.* 56; *DC. Pr. l. c.*).

Mar. occid.—In arenosis prope Mogador! ubi olim legerunt Broussonnet et Schousboe, et postea Balansa, ex spec. in herb. nost.

Ar. Geog. Africa borealis ex Mare Rubro ad Oceanum, sed potius infrequens. Africa tropica (Sennaar, Senegal). Palæstina.

H. maroccanum, *Lehm.*, videtur esse nil nisi var. levis *H. undulati* caulibus erectis et foliis aliquantulum latioribus. Sp. nostræ proxime affine, etsi in genere *Heliophyto* a Candolleo enumeratum, est *H. erosum* (*Lehm. Asper.* 54), stirps Canariensis et Senegalensis.

CYNOGLOSSUM (*L. Gen.* 183).

C. DIOSCORIDIS (*Vill. Dauph.* ii. 457; *DC. Pr.* x. 147), var. NEBRODENSE (*Guss. Prod. Fl. Sic.* i. 216; *DC. Pr.* x. 148).

Mar. merid.—In umbrosis juxta Sheshaoua! In convalle Ourika! In convalle Aït Mesan! Reraya (*R. et Fr. spec. in Herb. Kew.*).

Ar. Geog. Typus speciei Villarsianæ est forma septentrionalis in Gallia, Pedemontio et Hispania septentrionali potius rara. Varietas nostra per reg. Mediterraneam calidiorem late diffusa est, in Hispania, Sicilia, Africa boreali, et Oriente obvia.

Specimina nostra, in convalle Aït Mesan ad eire. 1500 m. lecta, sunt typo Gussoniano conformia, alia, quæ in eadem valle supra Arround eire. 2200 m. legimus, differunt statura pumila 8–12-pollicari, nuculis facie læviuseulis paræ glochidiato-aeuleatis.

* Normam Linnæanam scribendi—scil. *Borago* potius quam *Borrago*—secutus sum.

C. nebrodense, consentiente ecl. Boissier, nullis notis gravioribus a *C. Dioscoridis* distinguendum est.

CYNOGLOSSUM PICTUM (*Ait. Hort. Kew.* i. 179; *DC. Pr.* x. 147).

Mar. sept.—Videtur frequens in ditione Tingitana. Tanger (*Webb Herb.*)! In monte Djebel Kebir! Tetuan (1851)! Inter Tetuan et Ceuta!

Mar. occid.—? Folia fere certe huc referenda prope Agadir lecta attulerunt Coll. Indigeni (1871)!

Mar. merid.—Sheshaoua! Mesfioua (*J. D. H.*)!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

C. CLANDESTINUM (*Desf. Fl. Atl.* i. 159, tab. 42; *DC. Pr.* x. 148).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Inter Tanger et Tetuan!

Ar. Geog. Regio Mediterranea occidentalis a Sicilia usque fretum Herculis. Deest in Macaronesia et in Gallia australi.

In hac specie haud raro pedicelli plurimi bracteati. Ambigua igitur est inter sectiones a celeb. De Candolle stabilitas. Corolla extus villosa facile dignoscitur.

C. CHEIRIFOLIUM (*L., DC. Pr.* x. 154).

Mar. occid.—Prope Agadir (*Coll. Indig.* 1871).

Mar. merid.—Haud infrequens in saxosis aridis regionis inferioris et ad radices Atlantis Majoris. Prope Aïn Oumast! Mesfioua! Ourika!

Ar. Geog. Regio omnis Mediterranea occidentalis ab insula Melita usque Lusitaniam. Deest in Macaronesia et in Italia nisi meridionali.

— — —, var. HETEROCARPUM (*Kunze, Chlor. Austro-Hisp.* p. 49), = *C. arundanum* (*Coss. Not. Crit.* 41).

Mar. sept.—In monte Beni Hosmar prope Tetuan!

Ar. Geog. Hispania australis. Montes Tetuancenses.

Primo intuitu var. nostra satis differt tomento niveo crassiore et nuculis facie superiore læviuseculis centro parce cclinatis. Hæ notæ tamen variabiles sunt, nec desunt formæ intermediæ. Video nuculas *C. cheirifolii* in spec. nostris et Hispanicis valde ludibundas. Planta in aridissimis enata gaudet nuculis facie externa excavatis margine elevato, et cum spinis brevibus crassis basi conicis. In spec. alteris spinæ disci pauciores et tenuiores, marginales divergentes basi quasi confluentes. Stylus in utraque forma tenuis longitudine variabilis.

ECHINOSPERMUM (*Swartz in Lehm. Asper.* 113).

E. BARBATUM (*M. B. Fl. Taurico-cauc.* i. 421, sub *Myosotide*; *Lehm. Asper.* 128; *DC. Pr.* x. 137), var. *BREVISTYLUM*, nob.

Mar. merid.—Ad radices Atlantis Majoris in distr. Reraya haud procul Moulai Ibrahim!

Ar. Geog. Species orientalis ab India boreali-occidentali per Persiam et

Asiam Minorem usque Bulgariam, Rossiam australem et Sibiriam Uralensem dispersa. In Syria et Arabia etiam obvia. Spec. Africana præter nostra nulla vidi.

Spec. nostra varietati *E. cariensis*, Boiss., simillima, sed fructus structura ad typum Lehmannianum potius referenda; nec video discrimen nisi stylos in pl. nostra breviores.

ROCHELIA (*Rehb. in Flora*, 1824, p. 243).

R. STELLULATA (*Rehb. Fl. Excurs.* 346; *DC. Pr.* x. 176), = *Lithospermum dispernum* (*L. Sp. Pl.* 191).

Mar. merid.—In couvalle Aït Mesan circ. 1300 m. ! In couvalle Amsmiz ad 1400 m. !

Ar. Geog. Species imprimis orientalis ex India boreali-occidentali per Persiam et Asiam Minorem et Syriam usque in Europam austro-orientalem diffusa. In Hispania centrali et in Algeria interiore et in Imp. Maroecano hinc inde, sed rarius, obvia est.

BORAGO (*L. Gen.* 188).

B. OFFICINALIS (*L., DC. Pr.* x. 35).

Mar. sept.—Tanger (*Webb Herb.*) ! Tetuan (*Webb Herb.*) ! Vidi prope Tetuan sed spec. nulla servavi.

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). In Europa media haud infrequens sed verisimiliter introdueta.

ANCHUSA (*L. Gen.* 182).

A. UNDULATA (*L., DC. Pr.* x. 44).

Mar. merid.—Tetuan (1851) ! (*Webb spec.*) !

Ar. Geog. Per regionem Mediterraneam fere omnem (exclusa Macaronesia) sparsim occurrit.

A. MACROPHYLLA (*Lam. Ill.* no. 1816; *Desf. Fl. Atl.* i. 156; *DC. Pr.* x. 46).

Nascitur in Imperio Maroecano see. Desfontaines (loc. cit.), sed a nemini recentiorum visa.

A. ITALICA (*Retz. Obs.* i. 12; *DC. Pr.* x. 47; *Schousb. Gew. Marok.* 61).

Mar. sept.—Tanger ! (*Schsb. l. c.*). Vidi etiam prope Tetuan !

Mar. merid.—Rarius occurrit in arvis regionis inferioris, forsan introdueta. Mesfioua (*J. D. H.*) ! Reraya (*R. et Fr. spec. in Herb. Kew.*) !

Ar. Geog. Regio fere omnis Mediterranea (inclusa Macaronesia). In calidioribus Europæ mediæ etiam obvia.

A. ATLANTICA (*Ball in Journ. Bot.* 1873, 373).

Radix biennis; caulis erectus, simplex, foliosus; folia lanceolata, infe-

riora in petiolum latum attenuata, superiora sessilia, semiamplexicaulia; inflorescentia composita, ex racemis in axillis fol. superiorum ortis; corolla regularis, ochroleuca (in spec. nostris), hexamera; stylus inclusus; germen (immaturum) læve.

Mar. merid.—In regionis mediæ Atlantis Majoris convalle Aït Mesan a 1900 m. ad 2000 m.!

Ob fructum immaturum planta nostra quoad genus hand omnino certa; habitus tamen *Achusæ*, nullæ proxime affinis nisi forte spec. Abyssinicæ *A. affini*, R. Br. Planta tota molliter hispida; indumentum biforme ex pilis brevibus aliis longioribus patulis intermixtis.

NONNEA (*Medik. Phil. Bot.* i. 31).

N. NIGRICANS (*Lam. Dict.* iii. 627 (edit. Padov.), sub *Lycopside*; *DC. Fl. Fr.* iii. 626, et *Prod.* x. 31), = *Echioides nigricans* (*Dsf. Fl. Atl.* i. 163), = *Anchusa nigricans* (*Brot. Fl. Lus.* i. 298).

Mar. occid.—Casa Blanca!

Mar. merid.—Prope Gurguri (*J. D. H.*)! Prope Amsmiz!

Ar. Geog. Peninsula Iberica. Sicilia. Africa boreali-occidentalis.

Huc nimis proxima est *N. violacea* (*Dsf.* l. c., sub *Echioide*) quæ pro varietate tantum enumeranda videtur.

Speciem nostram, ut taceam de scriptoribus antiquis, Lamarek primus descripsit ex pl. in Hort. Paris. culta.

N. PHANERANTHERA (*Viv. Fl. Lib. Spec.* 9, tab. 1; *DC. Pr.* x. 33).

Mar. occid.—Mazagan! Saffi! Mogador! (*Brouss., sec. DC. Pr.* l. c.).

Ar. Geog. Africa borealis a Cyrenaica usque oras Atlanticas.

Huc verisimiliter referenda est *Lycopsis ciliata* (Schousb. *Gew. Marok.* 61) in arenosis pr. Mogador indicata. *L. ciliata* (Willd. *Sp. Pl.* i. 780) a Schousb. citata, monente eel. Boissier, est *Nounea lutea*, Rehb., species orientalis, nec in peninsula Iberica nec in Algeria (quoad sciam) hucusque detecta.

MYOSOTIS (*L. Gen.* 180).

M. LINGULATA (*Lehm. Asper.* 110; *Rehb. Ic. Fl. Germ.* xviii. tab. 120), = *M. caespitosa* (*Schultz, Fl. Starg. Suppl.* 11; *DC. Pr.* x. 105), var. *FOLIOSA*. Racemis elongatis insigniter foliaceis.

Mar. sept.—In uhginosis montis Djebel Kebir pr. Tanger meridiem versus!

Ar. Geog. Species per Europam omnem nisi arcticam, et regionem fere omnem Mediterraneam (excluis ins. Canariensibus) late diffusa, et usque in Asiam temperatam et Americam borealem provecta.

M. SYLVATICA (*Hoffm. Deutschl. Fl.* i. 85; *DC. Pr.* x. 107).

Mar. merid.—In monte Djebel Tezah a 2000 m. usque 3000 m.!

—— —, var. *ALPESTRIS* (*Koch, Syn. Fl. Germ.* 581), = *M. alpestris* (*Schmidt, Fl. Boh.* iii. 26).

Mar. merid.—In convalle Aït Mesan supra Arround, et in ascensu ad jugum Tagherot, a 2200 m. usque 3200 m.!

Ar. Geog. Species per Europam fere omnem et per montes regionis Mediterraneæ late diffusa, in Asia media et America boreali non deest. Forma montana (*M. alpestris*) in editioribus magis frequens.

Flores in spec. nostris minores quam in Europæis.

MYOSOTIS HISPIDA (*Schlecht. in Mag. Nat. Berl.* viii. 229; *DC. Pr.* x. 108).

Mar. sept.—Tanger! In monte Beni Hosmar prope Tetuan!

Mar. merid.—In regione media et superiore Atlantis Majoris. In convalle Aït Mesan et in jugo Tagherot a 2100 m. usque 3000 m.!

Ar. Geog. Europa fere omnis. Regio Mediterranea (inclusa Macaronesia).

Spec. ex Atlante variant pedicellis brevissimis, et interdum *M. uncatæ*, Boiss., simillima sunt.

M. VERSICOLOR (*Rchb. Fl. Excurs.* i. 341; *DC. Pr.* x. 109).

Mar. sept.—?Tanger (*Salzm. Exsicc.*). Vidi specimen Salzmannianum in herb. J. Gay; utrum ex ditione Tingitana haud certissimum sum.

Ar. Geog. In Europa occidentali frequens, fit rarior orientem versus. Naseitur in insulis Canariensibus et Azoricis, sed deest in reg. Mediterranea orientali.

M. STRICTA (*Link, Enum. H. Berol.* i. 161; *DC. Pr.* x. 109).

Mar. merid.—Legi in convalle Aït Mesan circ. 2100 m. In monte Djebel Tezah circ. 2200 m.!

Ar. Geog. Europa fere omnis, imprimis orientalis. In Oriente frequens, deest in Græcia et in Italia inferiore. In Hispania centrali obvia, desideratur in Lusitania et in Macaronesia. Spec. Africana (nostris exceptis) nondum vidi.

LITHOSPERMUM (*L. Gen.* 181).

L. ARVENSE (*L., DC. Pr.* x. 74).

Mar. sept.—Prope Tetuan!

Mar. merid.—In convalle Aït Mesan circ. 2100 m.!

Ar. Geog. In temperatis orbis veteris ex Hibernia usque Japoniam indigena species, in America introducta est. Adest in ins. Canariensibus.

L. INCRASSATUM (*Guss. Ind. Sem.* 1826, 6; *DC. Pr.* x. 74).

Mar. merid.—Ad radices et in regione superiore Atlantis Majoris. Tasseremout! In convalle Amsmiz circ. 1400 m. In convalle Aït Mesan supra 2200 m. In jugo Tagherot usque 3200 m. In monte Djebel Tezah circ. 3000 m.

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

Stirps variabilis, interdum ereeta ramis strictis, vel diffusis adscendentibus, vel in editioribus omnino prostrata ramis debilibus. Ut mihi videtur hæc est proles australis *L. arvensis*, a quo parum differt pedicellis incrassatis et nuculis paullo minus tuberculatis. Inter spec. nostra nonnulla valde accedunt ad *L. Splitbergeri*, Guss. Flores tamen semper pallide cærulei.

LITHOSPERMUM APULUM (*L.*, sub *Myosotide*; *Vahl, Symb.* ii. 32; *DC. Pr.* x. 75; *Schousb. Gew. Marok.* 60).

Mar. sept.—Tanger (*Webb Herb.*)! (*Schsb. l. c.*). In regione collina inter Tanger et Tetuan! Prope Tetuan!

Mar. merid.—Ad radices et in reg. inferiore Atlantis Majoris. Tassere-mout! Amsmiz! Seksaoua! In convalle Aït Mesan circ. 1400 m.!

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia).

L. MICROSPERMUM (*Boiss. Diagn. Pl. Or.* ser. 2, iii. 135), = *Rhytispermum apulum* (*Webb, Phyt. Can.* iii. 51, et in *Bourgeau, Plant. Canar.* No. 328, non *Link*), = *Lithospermum arvense*, var. *cærulescens* (*Bourgeau, Pl. Canar.* No. 318), = *L. Webbii* (*Coss. et Dur. MSS., Coss. Cat. Pl. Maroc. in Bull. Soc. Bot. Fr.* xxii. 23, sine descriptione).

Mar. occid.—In insula Mogador, ubi hanc sp. jam legerat cl. Balansa!

Ar. Geog. Insulæ Canarienses. Ditio Mogadorensis.

Hoc mihi videtur valde proximum *L. apulo*, et forsân sicut subspecies enumerandum.

Caules rigidi, interdum percunantes. Nuculæ fere ut in *L. apulo*.

L. FRUTICOSUM (*L.*, *DC. Pr.* x. 80).

Mar. sept.—Tanger, in monte Djebel Kebir! (*Salzm. Exsicc.*, sub *L. ericetorum*)!

Ar. Geog. Peninsula Iberica. Gallia occidentalis et australis. Ditio Tingitana. Species proxime affines, *L. hispidulum* (*Sibth. & Sm. Fl. Gr.* tab. 162), et *L. rosmarinifolium* (*Ten. Fl. Nap.* iii. 178), in Oriente et in Italia meridionali.

L. prostratum (*Lois. Fl. Gall.* i. 105; *DC. Pr.* x. 81) est mera forma, nec vera varietas, hujus sp. valde ludibundæ. Dignoscitur ramis prostratis et corolla extus pubescente. Re vera *L. fruticosum* ludit eaule ramoso, ramis strictis vel adscendentibus vel prostratis, et corolla extus glabra plus minusve pubescente. Specimina Tingitana plerumque ambigua sunt, habitu *L. fruticosi* sed corolla pubescente *L. prostrati*.

ECHIIUM (*L. Gen.* 191).

E. TUBERCVLATUM (*Lk. et Hffgg. Fl. Port.* i. 183; *DC. Pr.* x. 19), = *E. pustulatum* (*Lehm. Asper.* 448; *Rehb. Icones*, xviii. 1299; *Boiss. Fl. Or.* iv. 206; *vix Sibth. et Sm. Fl. Gr.* tab. 180).

Mar. sept.—Prope Tanger! Inter Tanger et Tetuan! Tetuan (1851)!

Ar. Geog. Per Europam australem, imprimis occidentalem hinc inde dispersum. Huc specimina ex Dalmatia (Unio, Itineraria, 1829) et teste Boiss. ex insula Zaeyntho. Adest in Africa boreali-occidentali, sed deest in Oriente et in Macaronesia.

Planta, quæ mihi videtur proles australis *E. vulgaris* et sicut subspecies enumeranda, satis variabilis est. Hujus varietas, nisi fallor, est *E. Gaditanum* (Boiss. Voy. Esp. 422). *E. pustulatum*, Fl. Gr. mihi videtur species diversa, propior *E. Rawolfii*, Delile, nuculis læviusculis lucentibus et indumento rigidior a pl. nostra dignoscenda.

ECHIMUM ANGUSTIFOLIUM (Lam. Ill. No. 1856; DC. Pr. x. 20)?

Mar. sept.—Prope Tanger legi florens et fructiferum die 13 Junii!

Ar. Geog. De hac specie et ejus distributione geographica nil affirmare audeo. Verisimiliter reg. Mediterranea occidentalis.

Planta nostra proxima est *E. angustifolio*, Coss. in Bourgeau, Pl. d'Espagne, No. 384, 989, et 1625, a cl. Willkomm ad *E. creticum*, L., relato; sed gaudet laciniis calycinis longioribus valde inæqualibus, qua nota ab *E. plantagineo* etiam distinguitur.

E. SERICEUM (Vahl, Symb. 35; DC. Pr. x. 21; Boiss. Fl. Or. iv. 207), var. *DIFFUSUM* (Boiss. l. c.), = *E. diffusum* (Sibth. et Sm. Fl. Gr. ii. 69, tab. 182; DC. Pr. x. 23).

Mar. merid.—Specimen unicum in itinere inter Shedma et Mogador lectum comm. cl. G. Maw!

Ar. Geog. Species regionis Mediterraneæ, imprimis orientalis civis, inde per African borealem omnem diffusa. In Europa nonnisi in Græcia et Italia australi detecta.

Spec. nostrum plantæ ex prov. Oran (Bourgeau, Pl. d'Algérie, 1856, No. 74) sat bene conveniens, differt, ut videtur, corollis paullo minoribus, staminibus parum exsertis.

E. sericeum est species variabilis, et formæ quas ecl. Boissier (loc. cit.) huc retulit aspectu valde diversæ. Huic valde, et forsan nimis, proximum est *E. setosum* (Vahl, Symb. ii. 32; DC. Pr. x. 21), quocum ecl. Cosson adunavit *E. humile* (Desf. Fl. Atl. i. 165), species perennis (vel biennis) caule inferne lignoso, staminibus longiuscule exsertis. *E. setosum* (Boiss. Fl. Or. iv. 209, et Del. Fl. Eg. tab. 17. fig. 2), est certe species diversa, annua, staminibus inclusis, stylo bifido lobis apice capitulatis; = *E. verecundum*, Viv., sec. Boissier, l. c.

E. PLANTAGINEUM (L., DC. Pr. x. 20), = *E. violaceum* (auct., an L.? Schousb. Gew. Marok. 61).

Mar. sept.—Tanger! (Salzm. Exsicc.)! (Schsb. l. c.). Prope Cap. Spartel!

Mar. occid.—Casa Blanca!

Mar. merid.—Distr. Mesfoua, haud procul urtem Marocco!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia). Gallia occidentalis.

— —, var. *GRANDIFLORUM* (*DC. Pr. l. c.*), = *E. grandiflorum* (*Desf. Fl. Atl. i. 168, tab. 45*).

Mar. sept.—Tanger in maritimis! *Pr. Cap. Spartel! Tetuan (Webb Herb.)!*

Ar. Geog. Occurrit hinc inde in reg. Mediterranea.

ECHIUUM LONGIFOLIUM (*Del. Fl. Eg. 40, tab. 16; DC. Pr. x. 23*).

Mar. merid.—? Frustulum inter Tasseremout et Ourika lectum comm. cl. G. Maw!

Ob spec. incompletum de hac specie hand certus sum. A typo vix differt nisi corollæ tubo rubro lobis violascentibus.

— —, var. *MAROCCANUM, nob.* Differt a typo indumento rigidiore, floribus minoribus versicoloribus. Planta annua (vel interdum bien- nis); caulis sæpius erectus vel e basi ramosus, ramis diffusis; indu- mentum strigosum; stamina subexserta; styli lobi apice capitulati, nueulæ majuseulæ, tetragono-conicæ, profunde rugoso-tuberculatæ. Forma aspectu inter typum et *E. Rauwolfii* (*Del. Fl. Eg. 51, tab. 19*) media. Hoc tamen differt nuculis nitidis obsolete tuberculatis.

Mar. occid.—Prope Mogador frustulum legi! Prope Agadir (*Coll. Indig. 1871*)!

Mar. merid.—Videtur satis frequens in aridis apricis regionis inferioris. Prov. Shedma pr. Aïn Oumast! Mesfoua! Amsniz! Milhain! Huc etiam spec. incompletum ex Mtouga (*R. et Fr. spec. in Herb. Kew.*)?

Ar. Geog. E. longifolium Ægypti, Nubiæ, et Arabiæ Petrææ civis est. Var. nostra nonnisi Maroceana videtur.

E. MODESTUM (*Ball in Journ. Bot. 1873, 373*).

Radix annua, tenuis; caulis erectus, tenuis, spithamæus, parce ramosus; folia lineari-oblonga, acutiuscula, pube brevi setis patulis longioribus intermixta vestita; inflorescentia scorpioidea subsimplex, rarius ramosa; bracteæ parvulæ calyce hirsuto breviores; corolla cærulea, pubescens, calyce duplo longior; antheræ parum exsertæ; stylus bifidus lobis apice capitulatis; nueulæ pyramidatae costatæ, parce et vix tuberculatæ.

Mar. occid.—Agadir (*Coll. Indig. 1871*)!

Mar. merid.—Frequens in aridis regionis inferioris. Aïn Oumast et alibi in pr. Shedma! Mesfoua! Mtouga! Seksoua!

Echio nullo nisi *E. maritimo* comparandum. Ab hoc differt statura humili, caule tenuiore, indumento patente magis hispido, staminibus exsertis (nec inclusis) et præsertim nueulis costatis nec dorso rotundatis. *E. maritimum* est sæpius bienne vel perenne et anno secundo ramos diffusos longiuseulos e basi caulis edit.

ECHIUM MARITIMUM (*Willd. Sp. Pl.* i. 788 ; *DC. Pr.* x. 23), = *E. micranthum* (*Schousb. Gew. Marok.* 62).

Mar. sept.—Tanger (*Webb Herb.*)! Tetuan (1851)! (*Webb spec.*)!

Mar. occid.—Casa Blanca! Prov. Haha pr. Mogador! Mogador (*Lowe Cat.*). In arvis prov. Haha rarissime (*Schsb. l. c.*).

Spec. nostra ex prov. Haha formam parvifloram hujus speciei sistunt, et procul dubio ad *E. micranthum*, *Schsb.*, referenda sunt.

Ar. Geog. Regio Mediterranea occidentalis ex Sicilia usque litora Atlantica. Deest in Macaronesia.

E. FLAVUM (*Desf. Fl. Atl.* i. 165, tab. 45 ; *Boiss. Voy. Esp.* 424), = *E. Fontanesii* (*DC. Pr.* x. 24).

Mar. sept.—Tetuan (*Webb spec.*)!

Ar. Geog. Hispania. Africa boreali-occidentalis.

E. POMPONIUM (*Boiss. Diagn. Pl. Or. ser.* i. 1, 93 ; *Voy. Esp.* tab. 124).

Mar. sept.—Tanger (*Salzm. Exsicc.*, sub *E. italico*)! Spec. ex herb. Schousboe comm. cel. Cosson!

Ar. Geog. Hispania. Africa boreali-occidentalis.

CERINTHE (*L. Gen.* 186).

C. MAJOR (*Lam. Dict.* iv. 67 ; *L.*, ex parte ; *Schousb. Gew. Marok.* 61), = *C. aspera* (*Roth, Catal. Bot.* i. 33 ; *DC. Pr.* x. 3).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Schsb. l. c.*). Cap. Spartel! Inter Tetuan et Ceuta!

Mar. occid.—Mogador (*Lowe Cat.*).

Ar. Geog. Regio Mediterranea, imprimis occidentalis (exclusa Macaronesia). In Asia Mediterranea nonnisi in insulis et in agro Byzantino obvia.

— —, var. *GYMNANDRA*, = *C. gymnandra* (*Gasparr. in Rend. Accad. Sc. Nat.* i. 72).

Mar. sept.—Ad ostium fluminis Tetuanensium (1851)! In monte Beni Hosmar pr. Tetuan!

Ar. Geog. Italia australis. Africa borealis.

CONVOLVULACEÆ (*Juss.* 132).

CALYSTEGIA (*R. Br. Prodr.* 483).

C. SOLDANELLA (*L.*, sub *Convolvulo* ; *R. Br. l. c.* ; *Choisy in DC. Pr.* ix. 433).

Mar. occid.—In arenosis prope Mogador legit Broussonnet teste Cavanilles (*Anal. Cienc. Nat.* iii. 17).

Ar. Geog. Species amphigea in litoribus Europæ occidentalis et regionis Mediterraneæ (incluso Mari Caspio) obvia, in Macaronesia tamen non indicata ; inde in litoribus temperatis Oceani Pacifici, scil. Australiæ, Novæ Zealandiæ, ins. Norfolk, Americæ australis et Californiæ dispersa. Ex Japonia spec. non vidi.

CALYSTEGIA SEPIUM (L., et Schousb. Gew. Mar., sub *Convolvulo*; R. Br. Prodr. 483; DC. Pr. ix. 433).

Mar. sept.—Tanger! Tetuan (1851)!

Ar. Geog. In temperatis orbis veteris late diffusa, inde in America boreali et australi et in Australasia dispersa. Spec. ex Macaronesia nulla vidi.

Spec. nostra intermedia sunt, sed propiora *C. sylvaticæ* (W. K. Pl. Rar. Hung. iii. 390, tab. 264, sub *Convolvulo*; Choisy in DC. Pr. ix. 433). Hæc tamen est mera varietas *C. sepium*; et formæ intermediæ præsertim in Italia sæpe occurrunt. Dignoscitur floribus majoribus, bracteis amplioribus magis patentibus obtusioribus. Folia variant basi cordata—auriculis rotundatis vel angulatis—vel sagittato-hastata.

CONVOLVULUS (L. Gen. 215).

C. CANTABRICA (L., DC. Pr. ix. 402).

Mar. merid.—Ad radices et in regione inferiore Atlantis Majoris. In convalle Amsmiz usque 1400 m.! In convalle Aït Mesan circ. 1300 m.! Prope Seksaoua!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia). Habitat etiam in calidioribus Europæ mediæ.

C. TRICOLOR (L., DC. Pr. ix. 405).

Mar. sept.—Tanger! Cap. Spartel! Tetuan (1851)!

Ar. Geog. Regio Mediterranea occidentalis a Sicilia usque Lusitaniam, et dit. Tingitanam. Deest in Macaronesia, necnon in Italia media et superiore.

C. UNDULATUS (Cav. Icon. iii. 39, tab. 277; DC. Pr. ix. 405), = *C. evoluloides* (Dsf. Fl. Atl. i. 176, tab. 49, et Sibth. et Sm. Fl. Gr. tab. 198).

Mar. sept.—Tanger (*Broussonnet ex Cavan. in Anal. de Cienc. Nat.*).

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In convalle Aït Mesan 1400–1500 m.!

Ar. Geog. Hispania centralis et australis. Liguria. Sicilia. Africa borealis.

Spec. ex Agadir allatum maximum, caules plurimos prostratos pedales et ultra e collo radiceis emittens; ex his prodeunt rami axillares, omnes cum caulibus flores plurimos subcapitato-congestos apice ferentes. Corolla extus glabra; capsula inmatura villosa, matura glabrescens; semina 1–2 in quovis loculo ex rugis prominentibus asperula.

C. ARVENSIS (L., DC. Pr. ix. 406; Schousb. Gew. Marok. 69).

Mar. sept.—Tanger! Tetuan (1851)!

Mar. merid.—In umbrosis irrigatis prope Sheshaoua!

Ar. Geog. Planta cosmopolita per totum orbem (exceptis terris arcticis) diffusa.

C. SABATIUS (Viv. Nov. Sp. Diagn. in calce Fl. Lib. Spec. 67; Rehb. Fl. Excurs. no. 2386; Bert. Fl. It. ii. 442; Moggr. Contrib. Fl. Ment. tab. 61), var. ATLANTICUS, nob.

Mar. merid.—In regione media Atlantis Majoris. In convalle Aït Mesan usque 2100 m.!

Ar. Geog. *C. sabatius* nonnisi in Liguria litorali rarius occurrit. Varietas vel subspecies *C. mauritanicus*, Boiss., in Algeria; var. nostra denique in Atlante Majore.

Pluribus notis planta nostra inter *C. sabatium* et *C. mauritanicum* (Boiss. Voy. Esp. 418, tab. 122) media est. Affinitatem proximam inter hos beatus Moggridge jampridem indicavit. Notæ quibus pl. nostra a typo recedit non gravis momenti. Ex radice lignosa procedunt caules plurimi tenues prostrati vel adscendentes, a spithama ad sesquipedem longi; folia parva ovalia vel subrotunda, basi truncata, breviter petiolata, cum caule pilis mollibus patentibus obsita; pedunculi axillares plerumque uniflori, rarius biflori, bracteis duabus linearilanceolatis medium versus instructi, dum biflori peduncululus alter nudus bracteolatus exit. Corolla pallide cærulea fauce oehroleuca, nec violacea ut in *C. sabatio*, nec in plicis rosea ut in *C. mauritanico*, sed in plicarum dorso pubescens; capsula glaberrima. Monente b. Moggridge planta spontanea in horto culta fit plus minusve scandens, et ex notula cl. G. Maw *C. mauritanicus* ex seminibus Algeriensibus enatus in horto ramos debiliter scandentes edit. In hoc segmenta calycina majora et magis foliacea quam in pl. spontanea. Cl. Choisy in DC. Pr. ix. 407, *C. sabatium* ad varietatem *C. pseudosiculi*, Cav., perperam reduxit. Hæc novissima est species omnino diversa, et meo sensu nil nisi varietas levis *C. siculi*, pedunculis bifloris, bracteis sæpe (non semper) a calyce remotioribus distinguenda. Hujus vidi specimina Algeriensia Canariensibus omnino conformia, et verisimiliter in ditione nostra invenienda sunt, sed hucusque nulla vidi.

CONVOLVULUS SICULUS (L., DC. Pr. ix. 407; Schousb. Gew. Marok. 70). Vulgaris in toto territorio Maroccano nobis noto.

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! Tetuan!

Mar. occid.—Prope Mogador (*Broussounet ex Carau. in Anal. Cienc. Nat.*) et in parum remoto monte Djebel Hadid! Prov. Ifaha haud procul Mogador! (*Schsb. l. c.*). Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In regione inferiore passim occurrit, et in Atlante Majore non deest. Shedna! Seksaoua! Mtouga! In monte Djebel Tezah usque 1800 m.! Alibi phryies visa!

Ar. Geog. Regio Mediterranea imprimis occidentalis (inclusa Macaronesia). Deest in Asia continentali.

C. ALTHÆOIDES (L., DC. Pr. ix. 409; Schousb. Gew. Marok. 69).

Mar. sept.—Tanger! (*Schsb. l. c.*).

Mar. occid.—In monte Djebel Hadid (*R. et Fr. spec. in Herb. Kew.*)! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Satis frequens in regione inferiore, etiam in saxosis aridissimis. Ain Oumast! Mesfona! In convalle Aït Mesan usque 1400 m.! Seksaoua! Sektana (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

CUSCUTA (*L. Gen.* 170).

C. PLANIFLORA (*Ten. Syll.* 128 ; *Fl. Nap.* tab. 220 ; *Engelm. Cusc. Sp.* 14), = *C. europæa*, var. *epithymum* (*Schousb. Gew. Marok.* 59).

Mar. sept.—Cap. Spartel (*Schsb. l. c.*).

Mar. occid.—? Agadir (*Coll. Indig.* 1871) !

Mar. merid.—? Prov. Shedma ! Mesfioua ! Misra Ben Kara ! et verisimiliter eadem species per provincias meridionales a nobis pereursas ubique communis.

Cuscutarum species et varietates in sicco extricare vix audco, sed opere Engelmänniano edoctus parum dubito quin *Cuscutæ* supra citatæ omnes ad *C. planifloram* (sensu latiore) referendæ sint. Specimina tamen prope Casa Blanca in foliis *Asphodeli* cujusdam lecta mihi videntur diversa, corollæ lobis longioribus acutioribus, et staminibus longioribus insignia. Monente Engelmann *Cuscutæ* indigenæ Canarienses—*C. epithymum*, *C. episonchum* et *C. epiplocamum*, Phyt. Canariensis omnes ad *C. planifloram* reducendæ sunt.

Ar. Geog. *C. planiflora* per regionem omnem Mediterraneam late diffusa est.

SOLANACEÆ (*Bartl. Ort. Nat.* 193).SOLANUM (*L. Gen.* 251).

S. NIGRUM (*L., DC. Pr.* xiii. pars 1, 50).

Mar. sept.—Tanger !

Mar. occid.—? Vidi prope Casa Blanca et Mazagan spec. verisimiliter ad typum referenda !

— —, var. *MINIATUM*, *M. K.*, = *S. miniatum* (*Bernh. in Willd. Enum.* 236 ; *Dunal in DC. Pr. l. c.* 56).

Mar. merid.—Sheshaoua !

— —, var. *HUMILE*, = *S. humile* (*Bernh. in Willd. l. c.* ; *Dun. in DC. Pr. l. c.*).

Mar. occid.—Mogador (*Lowe Cat.*).

— —, var. *SUFFRUTICOSUM*, = *S. suffruticosum* (*Schousb. in Willd. Enum.* 1236 ; *Dunal in DC. Pr. l. c.* 53).

Mar. occid.—In prov. Haha haud procul Mogador ! Planta a Schousbœo Willdenovio communicata verisimiliter ex agro Mogadorensi.

Ar. Geog. Species cosmopolita per totum orbem terrarum in temperatis et calidioribus, vestigia hominum secuta. Var. *suffruticosum* in subtropicis et tropicis orbis veteris occurrit.

S. villosum (*Lam. Dict.* iv. 289 ; *Dunal in DC. Pr.* xiii. pars 1, 58).

Mar. merid.—In ruderatis urbis Marocco !

Ar. Geog. Europa media et australis. Regio Mediterranea (inclusa Macaronesia). America australis.

S. nigro proxime affine. Pro subspecie licet dubitative enumeravi.

SOLANUM SODOMÆUM (L., *DC. Pr.* xiii. pars 1, 366).

Mar. sept.—Prope Tanger infrequens et forsā introductum !

Ar. Geog. In calidioribus regionis Mediterraneæ occidentalis. Deest in Gallia, in Italia superiore et in ins. Canariensibus. Varietas in Africa australi, ut videtur, indigena, neenon in Australia et in insula Mauritiî.

WITHANIA (*Pauq. Diss. de Bellad. ex Dunal in DC. Prod.*).

W. SOMNIFERA (L., sub *Physalide*; *Dunal in DC. Pr.* xiii. pars 1, 453).

Mar. merid.—In ruderatis et hortis urbis Marocco ! Inter Frouga et Sheshaoua legit G. Maw !

Ar. Geog. Regio Mediterranea calidior (inclusis ins. Canariensibus). India Orientalis. Africa orientalis et australis. Deest in Gallia, in Italia continentali et in Madera.

W. FRUTESCENS (L., et *Schousb. Gew. Marok.* 79, sub *Atropa*; *Pauq. Diss. de Bellad.*; *Dunal in DC. Pr.* xiii. pars 1, 457).

Mar. occid.—Casa Blanca ! In dumetis prope Mogador (*Schsb. l. c.*).

Mar. merid.—Frequens in regione inferiore et varius occurrit ad radices Atlantis Majoris. Prov. Haha ! Prov. Shedma ! Reraya usque 1000 m. ! Inter Frouga et Sheshaoua (G. Maw) !

Stirps aspectu variabilis. Interdum frutex 4–8-pedalis, foliis magnis suborbicularibus, interdum fruticulus foliis ovatis vix pollicem longis, et inter hos formæ intermediæ non desunt.

Ar. Geog. Hispania. Marocco. Insulæ Canarienses.

LYCIUM (L. *Gen.* 262).

L. VULGARE (*Dunal in DC. Pr.* xiii. pars 1, 509), = *L. barbarum* (L., *Sp. Pl.* ed. 2, nec ed. 1, sec. *Dunal*).

Mar. sept.—Vidi in sepibus prope Tanger, forsā introductum !

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—Pluries visum, sed semper in cultis vel prope pagos incolarum. Sheshaoua ! Ourika !

Ar. Geog. Verisimiliter ex Oriente in Europam et Africam borealem allatum. Frequens in sepibus Europæ mediæ et regionis Mediterraneæ, sed vix spontaneum.

L. AFRUM (L., *Dunal in DC. Pr.* xiii. pars 1, 521).

Mar. occid.—? Legi fructiferum prope Saffi 9 Junii 1871 ! Eandem speciem, nisi fallor, vidi prope Mazagan !

Ar. Geog. Sponte nascitur in Hispania (ubi rarum), in Oriente, in Africa australi, et (teste Rehb. fil.) in insulis Canariensibus.

Ex speciminibus incompletis planta nostra admodum incerta. Arbuscula humilis intricata, ramis brevibus, ex gemmis foliaceis in axillis foliorum delapsorum nodosis, baccis ovalibus solitariis pendulis.

LYCIUM MEDITERRANEUM (*Dunal in DC. Pr. xiii. pars 1, 523*); = *L. europæum* (*L., Schousb. Gew. Marok. 80*), nomen improprium a *Dunal* mutatum.

Mar. occid.—In insula Mogador! In collinis pr. Haha prope Mogador! (*Schsb. l. c.*).

Ar. Geog. Africa borealis. Oriens. Græcia. Macaronesia. Europæ australis dubia civis, sed sæpe introductum et hodie spontaneum.

Baccæ secundum anctores rarissime inveniuntur. In insula Mogador planta a locustis denudata baccas maturas oblongo-ovoideas saturate aurantiacas frequentes edit.

MANDRAGORA (*Juss. Gen. 125*).

M. OFFICINARUM (*L. Sp. Pl. ed. 1, 181*; *Dunal in DC. Pr. xiii. pars 1, 466*), = *Atropa Mandragora* (*L. Sp. Pl. ed. 2, 259*).

Mar. sept.—Prope Tanger folia visa! (*Salzm. Exsicc.*)! (*Blackmore, spec. in Herb. Kew.*)!

Ar. Geog. Regio Mediterranea calidior (exclusa Macaronesia). Deest in Gallia et in Italia superiore.

DATURA (*L. Gen. 246*).

D. STRAMONIUM (*L., Dunal in DC. Pr. xiii. pars 1, 540*).

Mar. sept.—Vidi in ruderatis circa Tanger!

Mar. occid.—Mogador (*Lowe Cat.*).

Ar. Geog. Ex America boreali dicitur oriunda; hodie per totam Europam calidiorem, regionem omnem Mediterraneam, et Americam australem facta est spontanea.

HYOSCYAMUS (*L. Gen. 247*).

H. ALBUS (*L., Dunal in DC. Pr. xiii. pars 1, 548*; *Schousb. Gew. Marok. 78*).

Mar. sept.—Tetuan (1851)!

Mar. merid.—Prov. Haha (*Schsb. l. c.*). In ruderatis urbis Marocco!
In convalle Aït Mesan circ. 2000 m.!

— — —, var. MAJOR (*L. Sp. Pl. ed. 2, 257*), = *H. majore* (*Mill. Dict.*; *Dunal in DC. Pr. xiii. pars 1, 548*).

Mar. occid.—Agadir (*Coll. Indig. 1871*)!

Ar. Geog. Species per regionem Mediterraneam diffusa. Var. in Oriente, Africa boreali et Macaronesia obvia.

SCROPHULARINEÆ (*R. Br. Prodr. 433*).

VERBASCUM (*L. Gen. 245*).

V. CALYCINUM (*Ball in Journ. Bot. 1875, 172*).

Planta tota albo-floccosa; caulis florifer fere bipedalis, subsimplex vel

breviter ramosus; folia inferiora obovato-acuta, in petiolum attenuata, caulina sessilia, superiora angustiora, floralia lanceolato-linearia, omnia, vel saltem inferiora, glomerulos florales superantia; flores fasciculati pedicellati, pedicellis calyceem subæquantibus; calycis segmenta oblongo-linearia, pro genere longissima, capsulam ovoideam sat magnam æquantia vel superantia; stamina inæqualia; filamenta purpureo-barbata. Ab affinibus (§ *Thapsoidea*, Benth. in DC. Prod.) calyce et foliorum forma bene distincta.

Mar. merid.—Ad radices Atlantis Majoris pr. Sektana! In convalle Aït Mesan circ. 1400 m.! Tamarat (*R. et Fr. spec. in Herb. Kew.*)!

VERBASCUM SINUATUM (*L., DC. Pr. x. 234; Schousb. Gew. Marok. 78*).

Mar. sept.—Tanger! (*Schsb. l. c.*).

Mar. occid.—Mogador (*Lowe Cat.*). Aïn el Hadjar haud procul Mogador!

Mar. merid.—Videtur haud infrequens in regione inferiore. In rudertis nr̄bis Marocco! prope Ourika! et alibi visum!

Ar. Geog. Per regionem omnem Mediterraneam (inclusa Macaronesia) late diffusum.

V. COSSONIANUM, *nobis*.

Elatum; inflorescentia racemoso-paniculata; folia crassa utrinque panoso-tomentosa, nervis in pagina inferiore prominulis, radicalia obverse oblongo-lanceolata in petiolum brevissimum sensim attenuata, margine leviter sinuata, caulina oblongo-lanceolata sessilia, basi subcordata, margine denticulata, superiora latiora abbreviata longiusecule acuminata; panicula magna (3-pedalis) subpyramidata; flores in fasciculis laxis pauci, 3–4, brevissime pedicellati; calycis segmenta oblonga breviuscula; corolla sat magna extus tomentosula; filamenta lanata; capsula?

Mar. merid.—In provincia Mtouga prope castellum gubernatoris florens 29 Maji!

In grege *V. speciosi*, Schrad., juxta *V. sinaiticum* (Benth. in DC. Pr. x. 236) collocandum. Ab hoc differt foliis crassioribus subtus insigniter nervosis, caulinis integris, floribus subsessilibus in fasciculis paucioribus, calycibus minoribus. Primo intuitu habui pro *V. granatensi*, Boiss. Hoc differt tamen foliis radicalibus et infimis profunde sinuato-crenatis, nervis subtus non prominentibus, floribus pedicellatis, calycis laciniis triangulari-ovatis. Pl. nostræ valde affine est frustulum fructiferum ab indigeno Mardochæo ad Tazeroualt lectum, a cel. Cosson benevole communicatum, sub "*V. pycnostachyum*, Coss. MSS." non Boiss. et Held. in Diagn. Pl. Or. Hoc verisimiliter ad sp. nostram referendum est.

V. LYCHNITIS (*Schousb. Gew. Marok. 77; an L.?*).

"In locis sterilibus siccis raro" (*Schsb. l. c.*). Præter Schousboe nemo

hane sp. in ditione nostra vidit. In Hispania australi a Willk. et Lange non indicatur, et floræ Maroceanæ dubia civis videtur.

Ar. Geog. Europa media. Regio Mediterranea præsertim orientalis. Deest in Macaronesia.

VERBASCUM HÆNSELERI (Boiss. *Voy. Esp.* 442; *Benth. in DC. Pr.* x. 239).

Mar. merid.—In distr. Reraya specimen unicum legit cel. J. D. Hooker!

Ar. Geog. Hispania centralis et australis. Atlas Major. Pl. nostra spec. Bourgæanis ex Sierra de Huescar et prope Riopar lectis bene congruit. Folia infima late ovata in petiolum abrupte angustata.

V. HOOKERIANUM, nob., = Verbaseum, nov. sp.? (*Ball in Journ. Bot.* 1875, 172).

Descriptio ex specimine unico a cel. J. D. Hooker lecto in regione inferiore Atlantis Majoris. Bienne (?), caulis bipedalis, ramoso-paniculatus; folia dense pannosa, tomento flavo obtecta, nervis in pagina inferiore valde prominulis, inferiora elliptico-lanceolata in petiolum longiuseulum attenuata, caulina sessilia ovata acutiuscula; flores parvi; calyceis segmenta abbreviata ex tomento deterrenti interdum virescentia, capsula ovato-conica obtusa dimidio breviora; stamina inæqualia.

Mar. merid.—In convalle Ourika!

Plantæ nostræ proximum est *V. Ternacha*, Hochst. in A. Richard Tent. Fl. Abyss. ii. 108. Hoc tamen ostendit stamina multo breviora subæqualia, et capsulam ovoideam et vix cum sp. Maroecana conjungendum videtur; sed in hoc genere ratio specierum valde incerta. Cel. Benth. in DC. Prod. vol. x., citat *V. Ternacha*, Hochst. (ex MS. Hoehstetteriano?), et dubitative—dum flores non vidit—sub *V. phlo-moide* collocauit.

CELSIA (*L. Gen.* 312).

C. CRETICA (*L. fil. Suppl.* 281; *DC. Pr.* x. 244), var. CAVANILLESII (*Willk. in Fl. Hisp.* ii. 545), = C. sinuata (*Cavan. in Anal. Cienc. Not.* iii. 68), = C. Cavanillesii (*Kunze, Chlor. Austro-Hisp.* No. 536).

Mar. sept.—Tanger (*Salzm. Exsicc.*, sub *C. cretica*)!

Mar. occid.—Videtnr satis frequens in regione sublitorali. Casa Blanca! Mazagan! Mogador! Agadir (*Coll. Indig.* 1871)!

Hæc planta, forsan melius pro subspecie sub *C. sinuata*, Cav., enumeranda, differt pluribus notis, et, nisi fallor, aream geographicam propriam habitat. *C. cretica* genuina insularum Cretæ, Sardinia, et Sicilia, necnon Algeriæ civis, gaudet caule simplici, floribus sessilibus in spicam laxam foliosam dispositis, foliis superioribus in bracteas floribus sublongiores abeuntibus. Hujus capsulam maturam non vidi. Stirps nostra Maroecana habet caulem ramis nonnullis brevibus

instructum, vel sæpe ex eaulē primario læso prodeunt rami plurimi erecti subæquales. Flores in racemis laxis magis distantes, pedicellis erecto-patulis, demum erassis, calyci subæqualibus, bracteis breves foliaceas superautibus fulerati. Folia primaria eito mareescentia, pinnatipartita, lobis subæqualibus rotundato-sinuatis, dum in *C. cretica* genuina sunt lyrato-pinnatifida. Capsula in pl. nostra sat magna ovato-conica vel ovato-globosa.

Hæc specimen omnia Hispanica et Maroccana et Canariensia quæ vidi sub "*C. cretica*." *C. laciniata* (Coss. in Balansa, 'Pl. d'Algérie,' no. 954) ex Biskra videtur etiam hæc referenda. Folia gaudent lobis ætiusculis; sed a foliis laciniato-pinnatifidis pl. Poiretiana diversæ sunt; cætera omnia ut in pl. nostra, nisi herba minus pubescenti-viscosa. Semina in spec. nostris superficie intertexta aulæiformia, majora quam in sp. affinis.

CELSIA BARNADESII (Vahl, *Symb.* ii. 39, sub *Verbasco*; Don, *Dict.* iv. 499; Willk. et Lge. *Fl. Hisp.* ii. 545), = *C. laciniata* (Poir. *Dict. Suppl.* ii. 147; Benth. in DC. *Pr.* x. 244), = *C. pinnatifida* (Boiss. et Reut. *Diagn. Pl. Hisp.* 22).

Mar. sept.—? Tanger (*Broussonnet in herb. Gouan*)?!

Ar. Geog. Species rara et paucis cognita. In Hispania centrali a Barnades et postea a Prolongo et Reuta detecta. Specimen in herb. Gouan a Broussonnet lectum originis incertæ in scheda manu Gouan "*ex Hispania*" notatur; sed in Hispania australi a nemini post Broussonnet visa; et fide Poiret Broussonnet plantam suam prope Tanger legit. Specimen incompletum ex herb. Webb, sub *C. betonicaefolia* serius citata, forsitan melius hæc referendum.

Ab affinis, præter alias notas, foliorum lobis acute incisus dentatisve, seminibus quam in *C. cretica* duplo minoribus, dignoscenda.

C. BETONICÆFOLIA (Desf. *Fl. Atl.* ii. 58; Benth. in DC. *Pr.* x. 245; Bot. Mag. xix. tab. 6066).

Mar. sept.—Tanger (*Salzm. Exsicc. teste Benth. l. c.*) (*Webb spec.*)!?

Ar. Geog. Algeria. Ins. Cap. Viridis. Ditio Tingitana. Hispaniæ dubia civis.

Hæc, nisi fallor, est stirps valde ludibunda, et forsitan melius cognita in subspecies dividenda. Semina ab auctoribus non descripta sæpe in herbariis desunt. Frustulum in herb. nost. a Webb prope Tanger lectum, mihi incertum videtur. Folium unicum a parte caulis inferiore *C. Barnadesii* magis quam huic conforme. Planta tota subglaberrima; folia floralia ad bracteis parvulas e basi lata acuminatas reducta; pedicelli erecto-patentes calyce ter quaterve longiores; calycis laciniae parvulae ovales integerrimae; corolla videtur lutea, quam in *C. betonicaefolia* genuina duplo major. Specimina ex insulis Capitis Viridis, aspectu satis diversa, habent folia caulina simpliciter serrata glabrescentia. Fide Lange exstat in herb. Schousboe specimen prope

Gades lectum, nemo tamen recentiorum hanc speciem in ditone Gadirana aut alibi in Hispania detexit. An error schedæ Schousboeanæ?

Celsia maroccana (*Ball in Journ. Bot.* 1875, 172).

Descr. emendata:—Planta biennis, vel perennis, tota molliter pilosa, vel rarius in aridis glabrescens, vix aut ne vix viscosa; folia radicalia petiolata lyrato-pinnatifida vel elliptico-spathulata, lobato-incisa, margine crenato-serrata, caulina fere omnia sessilia oblonga basi truncata inæqualiter dentata, floralia breviora apice acuminata, interdum in bracteis parvulas transeuntia; caulis subsimplex vel ramosus, e basi folii decurrente leviter angulatus; pedicelli patentes, fructiferi incrassati capsula matura duplo longiores; calycis laciniæ subfoliaceæ late ovatæ, grosse et inæqualiter serratæ, basin versus nervosæ, capsulam maturam subæquantes; corolla ampla ex aurantiao rubra; capsula ovato-conica vel ovato-globosa ex styli basi persistente mucronata; semina cylindrico-subconica utrinque truneata, transverse corrugata, pallidiora et profundius excavata quam in spec. Gorgonicis *C. betonicaefolia*.

Mar. occid.—In monte Djebel Hadid haud procul Mogador! Agadir (*Coll. Indig.* 1871)! Rabat (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—Haud infrequens in regione inferiore Atlantis Majoris, nec deest in saxosis aridis planitie Maroccanæ! Reraya (*J. D. H.*)! In convalle Aït Mesan usque 1400 m.! Seksaoua! Milhaïn! In prov. Shedma!

Melius explorata stirps nostra affinitatem proximam eum *C. betonicaefolia*, Desf. ostendit, et hodie non sine dubio pro subspecie habui; differt habitu, pedicellis brevioribus, calycis laciniis foliaceis, corollæ majoris colore diverso, capsula majore insigniter mucronata, et, nisi fallor, antheris anticis minus longe decurrentibus.

C. RAMOSISSIMA (*Benth. in DC. Pr.* x. 244).

Mar. occid.—Ex sylva Mamora inter flumina Sebou et Bouregreg a Durand allata.

Species tantum ex descript. Benthamiana l. c. cognita. Nemini recentiorum hanc sylvam leonibus et latronibus famosam penetrare licuit.

LINARIA (*Juss. Gen.* 120).

L. SPURIA (*L.*, sub *Antirrhino*; *Mill. Dict.* No. 15; *DC. Pr.* x. 263).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Tetuan (1851)!

Ar. Geog. Europa media. Regio omnis Mediterranea (inclusa Macaronesia).

L. GRÆCA (*Bory et Chaub. Expéd. Mor.* 175, tab. 21, sub *Antirrhino*; *Benth. in DC. Pr.* x. 263). Hujus var. levis est *L. commutata*

(Bernh. in Rehb. Ic. Crit. ix. tab. 815; Willk. et Lge. Fl. Hisp. ii. 559).

Mar. occid.—Casa Blanca! Prov. Haha prope Mogador! In monte Djebel Hadid (*R. et Fr. spec. in Herb. Kew.*)! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Satis frequens in regione inferiore. Prov. Shedma! Mesfiona! Amizmiz! In cultis urbis Marocco! Tamarat (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio Mediterranea imprimis occidentalis (inclusa Macaronesia). Adest in agro Byzantino, sed alibi in Asia Mediterranea vix detecta. In Gallia occidentali rarissime obvia.

LINARIA SAGITTATA (*Poir. Dict. Suppl.* iv. 19, sub *Antirrhino*; *H. fil. in Bot. Mag.* xcix. tab. 6060), = *L. heterophylla* (*Spreng. Syst.* ii. 790; *Chav. Mon. Antirr.* 112; *Benth. in DC. Pr.* x. 270), = *Antirrhinum heterophyllum* (*Schousb. Gew. Marok.* 181, tab. 3), = *L. Lancerothæ* (*Delile, Ind. Sem. H. Monsp.*), = *L. Webbiana* (*Vis. Ill. Ort. Bot. Pad.* 142).

Mar. occid.—Frequens in prov. Haha prope Mogador! (*Schsb. l. c.*) (*Lowe Cat.*). Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In prov. Shedma sat frequens! et hinc inde ad radices Atlantis Majoris. Ourika! Seksaoua!

Ar. Geog. Species Imp. Maroccani et insulæ Lancerothæ Canariensium propria.

Sæpius inter frutices late scandit, caulibus tenuibus flexuosis, interdum, sed rarius, convolutis. Vidimus etiam in saxosis formam pygmæam, ramis rigidis rectis, foliis præter infima omnibus linearibus.

L. LATIFOLIA (*Desf. Fl. Atl.* ii. 40, tab. 134; *Benth. in DC. Pr.* x. 271).

Mar. sept.—Prope Tanger (*Blackmore spec. in Herb. Kew.*)!

Ar. Geog. In Africa boreali-occidentali et in Hispania rarius occurrit.

L. TRIPHYLLA (*L.*, sub *Antirrhino*; *Mill. Dict.* No. 2; *Chav. Mon. Antirr.* 117).

Mar. sept.—Tanger!

Ar. Geog. Regio Mediterraneam fere omnis (exclusa Macaronesia).

L. VENTRICOSA (*Coss. in Bull. Soc. Bot. Fr.* xx. 252).

Descriptionem eel. auctoris citare malui:—

“*Perennis, elata*, glaberrima, glaucescens. Caules validi, sæpius 1 metr. elati, erecti, paniculato-ramosi. *Folia* approximata, alterna, sessilia, lanceolata, utrinque attenuata, integerrima, rigidula, penninervia, nervis primariis 3 prominulis e basi folii emergentibus, secundariis subobsoletis. Racemi terminales multiflori densiusculi, demum elongati. Pedicelli floriferi erecto-patentes, bractea et calyce breviores; fructiferi erecti. *Lacinia calycina* corolla multo breviores, capsulam subaequant, lanceolatae acutiusculæ. *Corolla magna* (fere ut in *L. vulgari*),

eum calcari *circiter 2 centim. longa*, pallide lutca venis croceo-fuscis notata, *tubo late campanulato-subventricoso*, calcari recto tenui *corolla subdimidio brevior*e, palato præminente subbilobo dense villosa villis intense luteis. Capsula didymo-subglobosa, loculis subæqualibus, utroque poro terminali 3-4-valvato dehiscente. *Semina parva, nigra, trigona, curvata, immarginata*, irregulariter striata *striis in tubercula decompositis*."

Mar. occid.—In sylva e *Callitri quadrivalvi* composita, ad orientem urbis *Mogador* cl. Balansa detexit. Prope Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Sat frequens in provincia Shedma! In convallibus interioribus et rarius ad radices Atlantis Majoris. In convalle Aït Mesan usque 1750 m. ! In convalle Amsmiz! Prope Keira (*J. D. H.*)!

Monente cel. Cosson *L. ventricosa* ad sectionem *Linariastrum* (Chav. et Benth. in DC. Prod. x. 271), § 2. *Speciosæ* (Benth. loc. cit. 274) pertinens, juxta *L. genistifoliam*, Mill., collocanda, a qua eximie differt laciniis calycinis minus acutis, corolla multo majore tubo ventricoso, calcari subdimidio brevior, etc.

LINARIA HETEROPHYLLA (*Dsf. Fl. Atl.* ii. 48, tab. 140; *II. fil. in Bot. Mag.* xcix. tab. 6041, non *Spreng.*, nec *Chav.*, nec *Benth. in DC. Prod.*), = *L. aparinoides* (*Willd. Sp. Pl.* iii. 247, sub *Antirrhino*; *Chav. Mon. Antirr.* 138; *Benth. in DC. Pr.* x. 275), = *Antirrhinum strictum* (*Sibth. et Sm. Fl. Gr.* tab. 594), = *Linaria stricta* (*Guss. Pl. Rar. Sic.* 250, et *Syn. Fl. Sic.* ii. 122 et 843; *Tenore, Syll. Fl. Neap.* 304), = *Antirrhinum multicaule* (*Tenore, Prod. Fl. Neap.* 36, non *L.*), = *Linaria reticulata* (*Rehb. Ic. Pl. Crit.* v. 19, tab. 431, non *Desf.*), = *Linaria sicula multicaulis molluginis folio* (*Bocc. Icon.* 38).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. occid.—Casa Blanca! Djebel Hadid haud procul Mogador (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—Sat frequens præsertim ad radices et in regione inferiore Atlantis Majoris. Mesfioua! Ourika! Amsmiz! Keira! Seksaua! In convalle Aït Mesan usque 1500 m. !

Ar. Geog. Species per Africam borealem late diffusa. Siciliæ et Italiæ meridionalis, necnon Cariæ et insulæ Cypri eivis indigena. Deest in Macaronesia.

Species quam maxime ludibunda ab auctoribus male muleata. Jam ante duo sæcula a Boceone memorata, et a Cupani, Robert, et Rivino delineata, a cel. Desfontaines nomine *L. heterophyllæ* publici juris facta est. Deinde Willdenovius speciei longe alienæ (*L. sagittatæ*, Poir.) nomen triviale *heterophylla* imposuit, et post eum plurimi auctores nomen antiquius Fontanesianum neglexerunt. Tabula Fontanesiana formam sp. nostræ surculorum foliis latioribus verticillato-senis exhibet. Hanc in Imp. Maroccano non vidimus, et spec. nostris

ex radicibus Atlantis propiora sunt quæ vidi ex Sicilia et Oriente, caulibus tenuioribus, foliis sureulorum linearibus confertis sæpius non verticillatis. In nostris tamen flores pallidiores ochroleuci, palato flavo nec aurantiaco donati.

— —, var. *TINGITANA*, nobis, = *L. tingitana* (*Boiss. et Reut. Pug.* 84).

Mar. sept.—Tanger (*Webb Herb.*)! In monte Djebel Kebir! Cap. Spartel! (*Boiss. et Reut. l. c.*).

Hæc mihi videtur mera varietas vegetior, foliis omnibus lanceolato-linearibus, nec anguste linearibus. Spec. nostra *L. heterophyllæ* prope Casa Blanca lecta formam intermediam sistunt. Secundum cll. auctores nascitur etiam in monte Santo prope Oran Algeriæ.

Linaria viscosa (*L.*, sub *Antirrhino*; *Dum. Cours Bot. Cult., ex Chav. Mon. Antirr.* 141; *Benth. in DC. Pr.* x. 276).

Mar. sept.—Prope Tetuan!

Mar. occid.—Casa Blanca!

Ar. Geog. Hispania australis. Africa boreali-occidentalis.

Differt a *L. heterophylla* statura multa minore, caule fere semper subsimplici, foliis sureulorum verticillato-ternis v. quaternis canlinis superioribus rarioribus, in apice caulis vel ramorum brevium carentibus, racemo abbreviato. Hæ notæ tamen in hac grege instabiles sunt, et hanc igitur sicut subspeciem enumeravi. Cll. Willkomm et Lange in *Fl. Hispanica* optimo consilio *L. Salzmanni* (*Boiss. Voy. Esp.* 456, tab. 128; *Benth. in DC. x.* 285), ad varietatem *L. viscosæ* reducerunt. Spec. nostra a Casa Blanca floribus pallide purpurascentibus forsanelius ad hanc varietatem referenda.

L. Maroccana (*H. fil. in Bot. Mag.* xeviii. tab. 5983).

Descriptio emendata:—Annuæ erectæ ramosa, superne viscoso-pubesceus, foliis in ramis inferioribus sureuliformibus sæpius lineari-lanceolatis, caulinis numerosissimis lineari-subulatis, racemis elongatis multifloris, pedicellis calyce longioribus, calyceis segmentis anguste linearibus obtusis adpressis, vel sub anthesi paululum recurvatis, calcaribus elongato recto corollam saturate coccineo-purpuream pedicellumque superante, labio superiore erecto bipartito lobis apice rotundatis, inferiore brevior truncato lobis lateralibus parvis rotundatis intermedio minimo, palato prominente emarginato aurco intus villosus, seminibus subcylindraceis transverse rugosis gibbosis.

Mar. merid.—In glareosis torrentium, in arvis et incultis saxosis ad radices Atlantis Majoris haud infrequens. Mesfioua! Ourika! Sektana!

Planta primo aspectu a *L. heterophylla*, in eadem regione obvia, omnino diversa. In hac rami plurimi stricti verticales e collo radice cum sureulis prodeunt, dum in *L. maroccana* caulis ramosus ramos nonnullos steriles sureuliformes et inde ramos floriferos adscendentes sæpius divisos edit, inflorescentia inde racemoso-paniculata. In

planta Hookeriana spontanea pedicelli duplo longiores, corolla major pulchre coccineo-purpurea, nec ut in tabula Fiteheana, e planta culta desumpta, læte purpurea. Semina tandem gibbosa quod vix aut ne vix occurrit in seminibus *L. heterophyllæ*. Ex observatione tamen plantæ e seminibus nostris in Horto Paroliniano pluribus annis cultæ et denique formarum *L. heterophyllæ* ineditarum in Herbario Algeriensi amicissimi Cosson, affinitas proxima nostræ plantæ et *L. heterophyllæ* mihi certior facta est. In planta culta folia latiora, anguste lineari-lanceolata, minus crebra, rami stricti subindivisi, racemus minus glandulosus subglaberrimus, corolla anno primo læte purpurea, fit demum pallidior et tota planta similior *L. tingitanæ* floribus purpureis quam *L. maroccanæ* nostræ. Pedicelli tamen et calycis lacinia longiores quam in omnibus *L. heterophyllæ* formis a me visis. Pro subspecie enumeravi donec res certior fiat.

LINARIA GALIOIDES (Ball in Journ. Bot. 1875, 173).

Radix perennis; caules floriferi plurimi erecti superne glabri, inferne cum sureulis sterilibus numerosis prostratis tenuiter pubescentes; folia caulina lineari-subulata, approximata pseudo-verticillata, superne pauciora, sureulorum verticillata, in quovis verticillo 5-6, latiora; pedicelli segmentis calycinis oblongo-linearibus parce glandulosi breviores; flores in racemo brevissimo conferti; calcar attenuatum corollæ flavæ labio inferiori aurantico æquilongum. Semina immatura angulosa, corrugata.

Mar. merid.—Regionis subalpinæ Atlantis Majoris incola in regionem inferiorem secus torrentes descendit. Legimus in convalle Aït Mesan supra Arround, a 2000 m. ad 2400 m. ! vidi specimen manens in glareis Oued en Fisk lectum, et alterum secus torrentem Ourika.

—— —, var. *PSEUDOSUPINA* (Ball, loc. cit.).

Caules omnes prostrati et fere omnes floriferi, folia subconformia breviora et latiora quam in specie. Facie *L. supinæ* similis, sed uisi fallor seminum structura aliena. Habitat in regione alpina in jugo Tagherot, a 2700 ad. 3200 m. !

Ob semina immatura de affinitate hujus plantæ haud omnino certus sum, sed verisimiliter in grege *L. heterophyllæ*, Desf., collocanda erit. Ab hac radice perenni, sureulis numerosis, calycis segmentis multo minoribus, racemo brevissimo, pedicellis pluries brevioribus, foliis caulinis pseudo-verticillatis, satis superque differt.

L. SPARTEA (L., et Cav. Ic. i. tab. 32, sub *Antirrhino*; Lk. et Hffgg. Fl. Port. 233, tab. 36; DC. Pr. x. 276), = *Antirrhinum junceum* (L. Herb.; Lam. Dict. iv. 332), = *Linaria juncea* (Desf. Fl. Atl. ii. 43).

In Imperio Maroccano legit Broussonnet, teste Desfontaines loc. cit. Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis. Ins. Canarienses.

A recentioribus in ditione nostra non visa, sed facile in territorio vix explorato invenienda.

LINARIA BIPARTITA (*Vent. Hort. Cels. tab. 83, sub Antirrhino; Willd. Enum. 640; DC. Pr. x. 277*), = *L. linogrisea* (*Lk. et Hffgg. Fl. Port. 240, tab. 41; DC. Pr. l. c.*), = *L. speciosa* (*Jacq. fil. Ecl. i. 141, tab. 95*), = *L. incarnata* (*Spreng. Syst. ii. 643*).

Mar. sept.—Tanger! (*Salzm. Exsicc., sub L. laxiflora*)!

Mar. occid.—Mazagan! Mogador! (*Broussonnet ex Benth. in DC. Prod.*). Aïn el Hadjar! In regione litorali pr. Haha frequens! Agadir (*Coll. Indig. 1871*)!

Mar. merid.—In zona *Arganiæ* prov. Shedma sat frequens; fit rarior in terris interioribus. Shedma pluribus locis! Mskala! Mtouga! Seksaoua! Tasseremout (*G. Maw spec.*)! Verisimiliter hue "*L. purpurea*" (*Lowe Cat.*) prope Mogador lecta. *L. purpurea* est sp. in reg. Mediterranea orientali indigena, in Hispania et in Africa boreali-occidentali omnino earens, et vix aut ne vix in agro Mogadorensi sponte provenit.

Ar. Geog. *L. bipartita* Africam boreali-occidentalem incolit et rarius in peninsula Iberica provenit.

Stirps satis variabilis, caulis plus minusve ramosus, ramis erectis interdum diffusis; pedicelli sæpius calyce ter quaterve longiores interdum vix sesquolongiores, calycis segmenta ovato-lanceolata variant margine membranaceo vel fere immarginata, subobtusata, acuta, vel acuminata. Inter spec. nostra nonnulla a *L. linogrisea* Lusitania nulla nota distinguenda, altera floribus minoribus, calycis segm. acuminatis, racemo pubescente, *L. sapphirinæ* (*Lk. et Hffgg. Fl. Port. 241, tab. 42*) simillima sunt. Hæc tamen (sec. Willk. et Lge. *Fl. Hisp. ii. 562*) habet semina semilunari-trigona angulata, nec ut in *L. bipartita* ovoideo-cylindrica transverse corrugata, et, uti jam suspicatus est eel. Benth., eadem est ac *L. delphinioides* (*J. Gay in Dur. Pl. Astur. Exsicc.*). Nomen Gayanum igitur recentius antiquiori postponendum. Specimina minora *L. bipartitæ* interdum *L. laxifloræ* (*Dsf. Fl. Atl. ii. 45, tab. 138; DC. Pr. x. 285*) valde similia sunt. Hæc novissima, nuper in territ. Maroccano detecta et mihi a eel. Cosson communicata, distinguitur ramis flexuosis nec strictis, floribus minoribus, et præsertim seminibus angularibus, testa reticulato-lacunosa nec transverse corrugata.

Pæc tanti viri, sectiones generis § 3. *Versicolores* et § 7. *Diffusæ* a eel. Benth. in DC. *Prod.* admissæ nimis affines sunt, et plurimæ species in sectionibus diversis enumeratæ ægre distinguuntur.

L. ARVENSIS (*L., sub Antirrhino; Desf. Fl. Atl. 45; Benth. in DC. Pr. x. 280*), var. β , *SIMPLEX* (*L., Chav. Mon. Antirr. 157*), = *L. simplex* (*Willd. Sp. Pl. iii. 213, sub Antirrhino; DC. Fl. Fr. iii. 588; Benth. in DC. Pr. x. 280*).

Mar. sept.—Tanger!

Mar. merid.—In convallibus et in regione superiore Atlantis Majoris. Supra Amsmiz! In convalle Aït Mesan supra Arround usque 2400 m.! In monte Djebel Tezah eire. 2500 m.!

Ar. Geog. Species per regionem Mediterraneam fere omnem (exclusa Macaronesia) diffusa. In calidioribus Europæ mediæ forsan eum seminibus introducta.

Specimina nostra subalpina a typo discrepant habitu diffuso, ramis brevibus basi supinis adscendentibus, foliis latioribus. Semina variant disco lævissimo vel minute tuberculato.

LINARIA MARGINATA (*Desf. Act. Soc. Par.* i. 36, tab. 7; *Benth. in DC. Pr.* x. 281).

Mar. sept.—In rupibus juxta Tetuan! In monte Beni Hosmar prope Tetuan (*Webb Herb.*)!

Mar. merid.—In rupibus supra Seksaoua circ. 1100 m.! In regione alpina Atlantis Majoris in jugo Tagherot circ. 3000 m. s. m.!

Ar. Geog. In Africa boreali-occidentalis hinc inde obvia; rarissime occurrit etiam in Hispania australi. Spec. Tetuanensibus omnino conformia bis inveni in rupibus circ. Ronda.

Specimina multa hujus gregis diligenter explorata mihi omnia ad speciem unicam collectivam *L. tristem* (*L.*, sub *Antirrhino*) reducenda videntur. Plurimæ tamen subspecies notis substabilibus definitæ enumerandæ videntur—scilicet *L. marginata*, *Desf.*, *L. lusitanica*, *Lk.* et *Hffgg.*, *L. melanantha*, *Boiss. et Reut.*, necnon *L. lurida*, nob. Nec forsan satis diversæ sunt *L. cæsia*, *Lag.*; *L. anticaria*, *Boiss. et Reut.* et *L. Rossmüssleri*, *Willk.* *L. marginata* distinguitur foliis quam in *L. tristi* latioribus, plerumque in verticillos subdistantes dispositis, herba viridi nec cæsia, corolla majuscula lutea, nec fusca vel fuscolutea variegata. Calcar variat rectum vel recurvum, rami prostrati vel adscendentes. In monte Beni Hosmar prope Tetuan specimina legimus foliis duplo latioribus, *L. anticariæ*, *B. & Reut.*, simillima, nisi floribus luteis nec albido-cærulescentibus.

Ex seminibus in jugo Tagherot lectis in Horto Paroliniano cultis enata sunt specimina Tetuanensibus simillima, quamvis planta spontanea *L. tristi* propius accedit foliis obtusioribus subcæsiis. Semina variant disco lævissimo vel tuberculato, et eodem modo ludit *L. tristis* genuina in rupibus Calpes satis frequens.

L. lurida (*Ball in Journ. Bot.* 1875, 173).

Perennis, e collo radices ramosissima, ramis decumbentibus surculis sterilibus numerosis; herba tota glaberrima, luride glauca; folia creberrima, crassiuscula, anguste linearia, pseudo-verticillata, subæqualia; flores in racemo brevi 5–8 majusculi; bractæ lineares, pedicellum, calyci subæqualem superantes; calycis segmenta interdum parce glandulosa, oblongo-spathulata, elongata; corolla sordide ochroleuca, calcar lutescente conico sensim attenuato subbreuior, galea apice bifida, labio inferiore macula obscura notato, valde convexo, fauce velutina. Semina satis matura non vidimus.

Mar. merid.—In regione alpina Atlantis Majoris legimus, a 3200 m. usque 3500 m., in jugo Tagherot!

Facie aliena sed structura valde proxima *L. marginatæ*, Desf., et præsertim *L. melananthæ*, B. et R. Png. Ab hac novissima corolla duplo majore et calcarī multo magis inflato, segmentis calycinis majoribus, foliis magis confertis, insigniter glauco-virentibus, diversa.

LINARIA AMETHYSTEÆ (*Lk. et Hffgg. Fl. Port. i. 253, tab. 47; Benth. in DC. Pr. x. 283*), = *Antirrhinum bipunctatum* (*Cavan. Ic. i. tab. 33, non L., ex Benth. l. c.*), = *A. polygonifolium* (*Poir. Herb., et verisimiliter, Dict. Suppl. iv. 26*), = *Linaria polygonifolia* (*Benth. in DC. Pr. x. 282, quoad plantam Gibraltariicam*).

Mar. occid.—? Legi in pr. Haha ab urbe Mogador meridiem verus!

Ar. Geog. *L. amethysteæ* genuina hucusque in peninsula Iberica nec alibi detecta.

Specimina nostra fructifera (corolla unica lilacina superstite munita), foliis angustioribus, racemo vix glanduloso et pedicellis brevioribus ad *L. Munbyanum* accedunt; sed semina insigniter concava margine subincrassato potius *L. amethysteæ*.

L. Broussonnetii (*Chav. Mon. Antirr. 169; Benth. in DC. Pr. x. 283*), quacum sicut varietas associanda est *L. multipunctata* (*Lk. et Hffgg. Fl. Port. i. 254, tab. 48*), et verisimiliter *L. ignescens* (*Kunze, Chlor. Hisp. 488*).

Mar. occid.—Spec. incompleta legi in arenosis prope Mogador! et prope Aïn el Hadjar! (Spec. prope Mogador lectum a Broussonnet in herb. Poiret descripsit cl. Chavaunes.)

Mar. merid.—Specimen prope Sektana lectum comm. cl. G. Maw!

Ar. Geog. Peninsula Iberica. Imp. Maroccanum. Uti jam suspicatus est cel. Benthām l. c., hæc est *L. amethysteæ* proxime affinis; differt etenim floribus majoribus flavis vel luteis palato sæpe croceo, bracteis inferioribus sæpe (non semper) reflexis. Racemus variat glandulosus vel subglaberrimus. Semina in spec. nostris disco tuberculata, in spec. Hispanicis sæpius lævia. Pro subspecie habui donec melius doceant qui plantam vivam observare possint.

L. Munbyana (*Boiss. et Reut. Pug. 89; Willk. et Lge. Fl. Hisp. ii. 567*).

Mar. occid.—In provincia Haha haud procul Mogador! et in arenosis juxta Aïn el Hadjar!

Mar. merid.—In convalle Amsmiz!

Ar. Geog. Africa boreali-occidentalis. Hispania australis.

Hæc planta, etsi aspectu valde diversa, est *L. amethysteæ* proxime affinis; nisi lusius tantos in hac grege vidissem pro specie probe diversa habuerim. Differt herba fere tota glaberrima (adsunt tamen glandulæ nonnullæ in calyce et pedicellis), foliis angustioribus linearibus, pedicellis calyce brevioribus, flore minore flavo, calcarī tenuiore, seminibus vix concavis anguste marginatis margine pallidiore vix incrassato. Hæc notæ tamen parum stabiles et hanc inter subspecies *L. amethysteæ*

animo fere certo enumeravi. Specimina *L. multicaulis*, Mill., a el. Balansa prope Mostaganem lecta (Pl. d'Algérie, 1851, No. 70) certe huc referenda et diagnosi Benthamianæ satis congrua sunt. Specimina authentica *L. multicaulis* non vidi, et el. Chavannes hanc ad varietatem *L. glaucæ* reduxit, facile tamen nomen hujus stirpis erit *L. multicaulis* (*L.*, sub *Antirrhino*). *Linaria atlantica* (Boiss. et Reut. Pug. 90) juxta Oran detecta, *L. Munbyanæ* simillima, distinguitur calcar corolla brevior, capsula minore, seminibus denique angulatis. Nuper habui a cel. Cosson spec. *L. atlanticæ* ad ostium fluminis Oued el Abid lecta calcar corollam æquante et capsula intermedia, seminibus ex eadem capsula alteris angulatis, alteris marginatis illis *L. Munbyanæ* simillima. An igitur *L. atlantica* ad var. heterospermam *L. Munbyanæ* reducenda?

LINARIA TOURNEFORTII (Poir. Dict. Suppl. iv. 24, sub *Antirrhino*; Lange, Ind. Sem. II. Hafu. 1859, 23), = *L. saxatilis* (Benth. in DC. Pr. x. 281), var. β . *GLABRESCENS* (Lange, l. c.), = *L. Perezii* (J. Gay in Durieu, Pl. Ast. Exsicc.). Synonymia auctorum valde dubia.

Mar. merid.—Specimen unicum in distr. Reraya legit J. D. H. !

— — —, var. γ , *MINOR* (Lange, l. c.), verisimiliter *L. glutinosa* (Lk. et Hffg. Fl. Port. tab. 39).

Mar. merid.—In regione media Atlantis Majoris. In convalle Aït Mesan supra Arround ! In monte Djebel Tezah ultra 2200 m. ! Tamarat (*R. et Fr. spec. in Herb. Kew.*) !

Ar. Geog. Species lincusque peninsulæ Ibericæ propria.

Species extricatu difficilis; formæ majores *L. verticillata*, Boiss., minores propius diversissimæ *L. Munbyanæ* accedunt. Specimen Hookerianum supra citatum, bienne vel perenne, satis bene cum *L. Perezii*, J. Gay, congruit, dum spec. nostra ex regione subalpina spec. Lusitanicis *L. glutinosæ* similia sunt. Nomen triviale *saxatilis* a el. Lange justa causa prætermissum dum *Antirrhinum saxatile* (*L. Sp. Pl.* 835) est sp. omnino incerta, a Linnæo *L. hirtæ*, sp. omnino alienæ, comparata. Quoad *L. saxatilem* (Lk. et Hffgg. Pl. Port. i. 238, tab. 40) res mihi videtur incerta. Auctores de seminibus siluerunt et tabula citata valde similis *L. verticillatæ*, Boiss. Vidi tamen in herb. J. Gay, hodie Kewensi, frustulum ex spec. in herb. Antoine de Jussieu a cel. Link communicatum nostræ sp. certe referendum. *L. glutinosa* (Lk. et Hffgg. Fl. Port. i. 238, tab. 40) a el. Lange omnino diversa habetur ob semina in tab. citata immarginata (an errore phytographi?). Vidi spec. ad Setnbal, loco classico ab auctoribus citato, ab equiti W. C. Trevelyan lecta tabulæ citatæ et spec. nostris Atlanticis conformia, seminibus distincte marginatis, margine angusto paululum incrassato.

L. RUBRIFOLIA (*Rob. et Cast. in DC. Fl. Fr. v. 410; Benth. in DC. Pr. x. 287*).

Mar. merid.—Legi in convalle Aït Mesan circ. 1700 m. !

Ar. Geog. Gallia australis. Hispaniæ montes. Sicilia. Sardinia. Atlas Major.

ANARRHINUM (*Desf. Fl. Atl.* ii. 51).

A. PEDATUM (*Desf. l. c.* tab. 141; *Benth. in DC. Pr.* x. 289).

Mar. sept.—Tetuan (*Webb spec.*)!

Mar. occid.—Casa Blanca!

Mar. merid.—In umbrosis prov. Shedma! et in aridissimis juxta Aïn Oumast! In convalle Aït Mesan usque 1350 m.! Ourika (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Africa borealis.

A. FRUTICOSUM (*Desf. Fl. Atl.* ii. 52, tab. 142; *Benth. in DC. Pr.* x. 289).

Mar. merid.—In convalle Amsmiz a 1300 m. usque 1500 m.!

Ar. Geog. Africa borealis. Forma valde proxima in Oriente et in Abyssinia.

Characteres quibus ccl. Bentham *A. orientale* suum ab *A. fruticoso* distinxit mihi videntur non satis constantes. Habitus in utraque sp. sæpe idem.

ANTIRRHINUM (*L. Gen.* 750).

A. ORONTIUM (*L., DC. Pr.* x. 290).

Mar. sept.—Tanger (*Webb Herb.*)! Prope Cap. Spartel! Tetuan! Inter Tetuan et Centa!

—— —, var. INDICUM (*Chav. Mon. Antirr.* 90), = *A. indicum* (*Royle Herb.*).

Mar. occid.—Mogador! Aïn el Hadjar! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Shedma! Seksaoua!

Var. hæc, a Bentham in *DC. Pr.* l. c. prætermissa, differt herba glaberrima, caule glabrescente, bracteis elongatis cum foliis angustis, corolla cum capsula minoribus.

—— —, var. GRANDIFLORUM (*Chav. Mon. Antirr. l. c.; Benth. in DC. Pr. l. c.*), = *A. calycinum* (*Lam. Dict.* iv. 365; *Salzm. Exsicc.*).

Mar. sept.—Prope Tanger frequens in arenosis maritimis! (*Salzm. Exsicc.*)!

Ar. Geog. Species in temperatis et subtropicis orbis veteris late diffusa, in America boreali verisimiliter introducta. Var. *indicum* in India boreali-occidentali, in montibus Nilghiri, in Abyssinia, in Macaronesia et in Africa boreali magis frequens. Var. *grandiflorum* in regione Mediterranea calidiore hinc inde occurrit.

A. MAJUS (*L., DC. Pr.* x. 291).

Mar. sept.—In rupibus montis Beni Hosmar! (*Webb Herb.*)!

Ar. Geog. In regione fere omni Mediterranea (exclusa Macaronesia) videtur spontaneum. In Europa media et in America extratropica obvium, sed verisimiliter introductum.

Antirrhinum tortuosum (*Bosc, Chav. Mon. Antirr.* 87; *Benth. in DC. Pr.* x. 291), var. CALYCE PUBESCENTE, = *A. majus*, var. angustifolium (*Willk. et Lge. Fl. Hisp.* ii. 583).

Mar. sept.—In rupibus juxta urbem Tetuan (1851)!

Planta nostra *A. tortuoso* genuino adamussim convenit nisi calyce et tubo corallino extus pubescentibus. Eandem plantam legi in rupibus Calpes calycibus interdum pubescentibus, sæpius glaberrimis. Differt ab *A. majore* ramis vel caulibus e rhizomate lignoso ortis, brevibus plus minusve tortuosis, cum foliis linearibus glaberrimis, floribus brevissime pedicellatis subfasciculatis, calycis segmentis angustioribus.

Ar. Geog. Per regionem Mediterraneam occidentalem (exclusa Macaronesia) hinc inde dispersum.

Ut morem gererem magistris *A. tortuosum* sicut subspeciem enumeravi. Mihi tamen nil nisi *A. majoris* forma rupestris videtur. Specimina tamen hujus in rupibus Beni Hosmar lecta omnino typica, et ab *A. tortuoso* aspectu diversissima.

A. INTRICATUM (*Ball in Journ. Bot.* 1875, 173).

Subspecies post *A. ramosissimum* (*Coss. in Bull. Soc. Bot.* ii. 254) enumeranda.

A. ramosissimo facie simillimum adeo ut sine floribus vix dignoscitur; sed eximie differt corollæ dimidio minoris extus pubescentis labio superiore emarginato vix bifido, non reflexo, labio inferiore brevissime trifido, lobis subæqualibus, fauce non clausa, segmentis calycinis brevibus acutis margine hirtulis (nec glaberrimis), capsula minore superne tenuiter glanduloso-pubescente. Semina compressa in utraque facie cristato-marginata. Testa ex alveolis inordinatis profunde rugosa. Embryo cylindrico-ovoidens vix curvatus. Fruticulus intricatus, sub anthesi aphyllus. Nisi certus fuissem stirpem nostram eum specie Cossoniana nexu genetico colligari vix in genere *Antirrhino* collocarem.

Mar. merid.—Legimus in glareosis Oued en Fisk prope Gurguri!

Ar. Geog. Species Africæ borealis civis; subspecies nostra Imper. Marocani propria.

SCROPHULARIA (*L. Gen.* 756).

S. ARGUTA (*Soland. in Ait. Hort. Kew.* ii. 342; *Benth. in DC. Pr.* x. 305).

Mar. merid.—Satis frequens in rupibus regionis inferioris præsertim ad radices Atlantis Majoris. Seksaoua! Milhaïn! Gurguri! Aïn Tarsil! Mskala! In collibus prope urbem Marocco (*G. Maw*)!

Ar. Geog. Insulæ Canarienses. Provinciæ meridionales Imper. Marocani. In regno Maseat (legit *Aucher*). In litore Africano Maris Rubri (legit *Lord*). In Hispaniæ Sierra de Gador unico loco (legit *Bourgeau*).

In spec. Canariensibus cymæ subsessiles, inflorescentia fere thyrsoidæa;

in Maroccanis cymæ semper pedunculo fulcratæ, sæpe ex axillis foliorum enato, rarius basi bracteato. De ramis subterraneis *S. argutæ*, confer Duricu in Bolle, Scroph. Canarienses, p. 12.

SCROPHULARIA SAMBUCIFOLIA (*L., Benth. in DC. Pr. x. 306*), = *S. mellifera* (*Ait. Hort. Kew. ii. 343; Desf. Fl. Atl. ii. 53, tab. 143; Salzmann. Exsicc.*), = *S. viridiflora* (*Poir. Voy. ii. 194*).

Mar. sept.—Tanger! (*Webb spec.*)! (*Schousb. spec.*)! (*Salzm. Exsicc.*)!
In monte Beni Hosmar pr. Tetuan!

Mar. occid.—Rabat (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Africa boreali-occidentalis. Peninsula Iberica.

Variat pedicellis dense glandulosis vel subglaberrimis, et segmentis calycinis late vel anguste marginatis.

S. SCORODONIA (*L., Benth. in DC. Pr. x. 307*). Var. foliis fere semper simpliciter nec duplicato-serratis, = *S. papillaris* (*Boiss. et Reut. Pug. 90*).

Mar. sept.—In fruticetis montis Djebel Kebir prope Tanger (*B. et R. l. c.*), (*Webb spec.*, sub *S. Scorodonia*, var. ?)!

Ar. Geog. Species occidentalis ex Anglia australi usque Macaronesium diffusa; in Hispania orientali rara vel omnino deficiens.

S. papillaris mihi videtur var. *S. Scorodonie* thyrso depauperato foliis simpliciter serratis. Folia inferiora in spec. Webbiano parce duplicato-serrata, serraturis quam in typo obtusioribus; thyrsos vix foliaceos.

S. AURICULATA (*L., Benth. in DC. Pr. x. 309*).

Mar. sept.—Tanger (*Webb spec.*)! In monte Djebel Kebir! Prope Tetuan!

— —, var. *PARVIFLORA*, nob. Foliis latioribus, caulinis inferioribus secus petiolum inæqualiter pinnatim lobatis, corolla et capsula minoribus. Facie *S. Smithii*, Horn., refert.

Mar. merid.—In irrigatis prope urbem Marocco! et in horto Ben Dreis!

Ar. Geog. Species in regione Mediterranea calidiore imprimis occidentali, ab insula Creta usque Lusitaniam late diffusa.

Affinitatem proximam hujus et *S. aquaticæ*, L., auctores indicarunt. Hæc forsitan pro subspecie enumeranda?

S. HISPIDA (*Desf. Fl. Atl. ii. 55*).

Mar. sept.—? Specimen vidi prope Tingidem lectum sed notulam frustra quæsi. Nascitur in Algeria occidentali ad fines Maroccanas, et procul dubio in ditone nostra indigena est.

Hanc speciem probe distinctam eel. Benth. in DC. Prod. l. c. cum præcedente immerito conjunxit. Differt etenim notis pluribus gravioribus, ut monuerunt eel. Boissier et Reuter in Pugillo cit. p. 91. Folia crassiora grosse dentato-lobata, dentibus semper acutis; thyrsus elongato-pyramidatus, nec interrupte spicatus; calycis segmenta an-

guste marginata; capsula minor ovato-conica, acuta; semina majora prismatico-cylindrica, truneata, longitudinaliter costata, costis granulatis.

Ar. Geog. Africa boreali-occidentalis.

— —, var. vel sp. nov.?

Mar. merid.—Prope Amsmiz legi frustulum plantæ insignis *S. hispida* pluribus notis aecedens, sed vix econjungendæ.

Planta 5-pedalis et ultra; caulis erectus, tetragonus, anguste alatus, ramis plurimis elongatis subbipedalibus instructus; folia caulina pinnatipartita lobis discretis petiolulatis omnibus lanceolatis acuminatis, acute duplicato-serrata; eymæ in ramis thyrsoidæ, superiores braetæa lineari suffultæ, approximatae, inferiores distantes basi foliaceæ; calycis segmenta orbicularia late scarioso-marginata; corolla parva oehroleuca, labio superiore bifido lobis rotundatis leviter sinuatis; staminodium parvum orbiculare; capsula ovato-conica acuta; semina *S. hispida*.

SCROPHULARIA LÆVIGATA (*Vahl, Symb.* ii. 67; *DC. Pr.* x. 309), = *S. trifoliata* (*Desf. Fl. Atl.* ii. 54; *Wyd. Mon. Scrof.* 34, non *L.*).

Mar. occid.—Mogador (*Lowe spec. in Herb. Kew.*)!

Ar. Geog. Africa borealis. Fertur in Oriente, sed spec. non vidi.

S. DURANDII (*Boiss. et Reut. Pug.* 91).

Mar. sept.—Specimen unicum in humidis prope Tingidem a Durand lectum exstat in herb. Pavon. hodie Boissieriano (*Boiss. e Reut. l. c.*).

Ex descriptione *S. auriculata* facie subsimilis, sed ex staminodio lineari in grege *S. canina*, *L.*, collocanda.

S. CANINA (*L., DC. Pr.* x. 315).

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—Sheshaoua! Shedma! Mtouga (*J. D. H.*)! Oned en Fisk (*R. et Fr. spec. in Herb. Kew.*)!

— —, var. *PINNATIFIDA* (*Boiss. Voy.* 446), = *S. pinnatifida* (*Brot. Fl. Lus.* i. 202; *Wyd. Mon. Scrof.* 43).

Mar. sept.—Tanger (*Webb Herb.*)!

Ar. Geog. Species in regione Mediterranea (exclusa Maenronesia) frequens. Oecurrit etiam in calidioribus Europæ mediæ.

— —, var. vel sp. nov.?

Mar. merid.—Legi in convalle Amsmiz spec. incompletum pluribus notis a *S. canina* diversum. Flores longius pedicellati; segmenta calycina minuta, corollæ vix sextam partem attingentia, orbiculata, anguste scarioso-marginata; corolla duplo major sordide oehroleuca, staminodio in duabus a me corollis exploratis nullo; semina nondum matura. Planta ulterius exploranda.

SCROPHULARIA FRUTESCENS (L., *Benth. in DC. Pr.* x. 316), = *S. pinatifida* (*Salzm. Exsicc., non aliorum*).

Mar. sept.—In arenosis maritimis juxta Tanger! (*Webb Herb.*)! (*Salzm. Exsicc.*)!

Ar. Geog. In litore Atlantico peninsulæ Ibericæ et Africæ Tingitanæ. Forma proxime affinis (var. vel subsp.) in Teneriffa.

DIGITALIS (*L. Gen.* 758).

D. LACINIATA (*Lindl. Bot. Reg.* tab. 1291; *DC. Pr.* x. 449).

Mar. sept.—Legimus nondum florentem in monte Beni Hosmar pr. Tetuan capsulis exsiccatis anni prioris adhuc onustam! In montosis Tetuanensibus (*Webb spec.*)!

Ar. Geog. Hispania Bætica. Montes Tetuanenses.

D. LUTEA (L., *DC. Pr.* x. 452), var. ATLANTICA, nob.

In hac forma flores minores (fere *D. australis*, Ten.), folia floralia subbracteiformia, flores superantia.

Mar. merid.—Legimus in convalle Aït Mesan circ. 2000 m.! Vidi spec. forsàn eodem loco lecta a ell. Rein et Fritsch!

Ar. Geog. Species in Europa media sat frequens; in Italia, Dalmatia et Hispania boreali obvia, alibi in regione Mediterranea rara vel omnino absens.

VERONICA (*L. Gen.* 25).

V. ANAGALLIS (L., *DC. Pr.* x. 467; *Schousb. Gew. Marok.* 4).

Mar. sept.—Tanger! (*Schsb. l. c.*).

Mar. occid.—Mogador! (*Lowe Cat.*).

Mar. merid.—In irrigatis urbis Marocco! Mesfioua! Mtouga! Amsmiz!

Ar. Geog. Species amphigea, in temperatis etiam subtropicis hemisphæræ borealis late diffusa. In Africa australi verisimiliter etiam indigena.

Spec. nostra ex prov. meridionalibus plurima ad *V. anagalloidem* (Guss. *Ic. Pl. Rar.* 5, tab. 3) accedunt. Hæc mihi videtur mera varietas formis intermediis conjuncta.

V. BECCABUNGA (L., *DC. Pr.* x. 468).

Mar. merid.—Legimus in uliginosis subalpinis convalis Aït Mesan in ascensu jugi Tagherot circ. 2600 m.! et secus torrentem prope pagum Amsmiz ad circ. 900 m.!

Ar. Geog. In temperatis hemisphæræ borealis præsertim frigidioribus late diffusa. In montibus excelsioribus regionis Mediterraneæ, Asiæ mediæ et Abyssiniæ. In insula Teneriffa olim lecta.

V. CUNEIFOLIA (*Don in Ann. Nat. Hist.* vii. 457; *Benth. in DC. Pr.* x. 472), = *V. cariensis* (*Boiss. Diagn. Pl. Or.* ser. 1, iv. 75), var. ATLANTICA, nob., = *V. atlantica* (*Ball in Journ. Bot.* 1875, 174).

Radix lignosa; caules plurimi, tenues, radicantes; folia petiolata, basi anguste cuneata, inciso-crenata, floralia braeteiformia integerrima, pedicello subæqualia vel longiora; segmenta calycina lineari-lanceolata inæqualia, quinto minore sæpe adjecto; corolla in racemis plurifloris majuseula, lobis staminibus æquilongis; capsula cordato-orbicularis, profunde emarginata; semina nimis immatura.

Mar. merid.—Habitat in regione subalpina et alpina Atlantis Majoris—in jugo Tagherot 2200 m. ad 3000 m. ! in monte Djebel Tezah 2200 m. ad 2800 m. !

Flores in var. nostra quam in planta Asiatica majores, eos *L. saxatilis*, L., æmulantes, folia præsertim inferiora angustiora sunt, et stamina longiora. Capsula videtur magis compressa; sed hæ notæ differentiales non gravis momenti. Olim pro subspecie habui; hodie pro varietate enumeravi.

Specimina in monte Djebel Tezah lecta flores paululum minores et folia pilosa subhispida ostendit.

Ar. Geog. Species hucusque nomini in montibus Asiæ Minoris detecta.

VERONICA ARVENSIS (L., DC. *Pr.* x. 483).

Mar. sept.—Tanger ! Cap. Spartel !

Mar. merid.—In regione subalpina Atlantis Majoris. In convalle Aït Mesan et in jugo Tagherot a 2000 m. usque 3000 m. In monte Djebel Tezah ultra 2300 m. !

Ar. Geog. In temperatis hemisphæræ borealis. In Macaronesia et in montibus Asiæ mediæ obvia.

Specimina in monte Djebel Tezah lecta ostendunt herbam dense pubescentem subvillosam, capsulas majores profundius emarginatas et pedicellos longiores calyci subæquilongos. Quæ tamen legi in convalle Aït Mesan inter hæc et typum Europæum fere intermedia sunt.

V. TRIPHYLLOS (L., DC. *Pr.* x. 486).

Mar. merid.—Specimen unicum legi in convalle Amsniz eire. 1400 m. !

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia). Europa media. Scandinavia australis.

V. TOURNEFORTII (Gmel. *Fl. Bad.* i. 39, anno 1805), = V. persica (Poir. *Dict.* viii. 542, anno 1808; Gr. et Godr. *Fl. Fr.* ii. 598; Willk. et Lge. *Fl. Hisp.* ii. 595), = V. Buxbaumii (Ten. *Fl. Nap.* i. 7, tab. 1; Benth. in DC. *Pr.* x. 487), = V. filiformis (DC. *Fl. Fr.* v. 388, non Sm.). Nomen triviale *persica* infaustum nullo jure prioritatis gaudet.

Mar. sept.—In maritimis prope Tanger !

Ar. Geog. Planta verisimiliter regionis Mediterraneæ totius (exclusa Macaronesia) indigena, in Europa media cultura introducta, hodie late diffusa. Adest etiam in montibus Himalaya. In America boreali forsitan introducta.

VERONICA AGRESTIS (*L., Benth. in DC. Pr. x. 487*).

Mar. merid.—In couvalle Aït Mesan supra Arround circ. 2100 m.!

Ar. Geog. In temperatis hemisphæræ borealis late diffusa. In Macaronesia non deest.

V. didyma (*Ten. Fl. Nap. Prod. 6*), = *V. polita* (*Fr. Nov. Fl. Suec. ed. 2, p. 1; Gr. et Godr. Fl. Fr. 598*).

Mar. sept.—In monte Beni Hosmar prope Tetuan!

Notæ plurimæ quibus hæc a simillima *V. agresti* dignoscitur certe variabiles sunt, et formæ intermediæ non desunt. Secundum ell. Grenier et Godron l. c. hæc differt seminibus in quovis loculo 8–10, stylo semper exserto, dum in *V. agresti* semina in quovis loculo 4–5, et stylus emarginaturam capsulæ non superat. Nescio an hæ notæ stabiles sint, et interea hanc sicut subspeciem enumeravi.

Ar. Geog. Distributio fere eadem ac *V. agrestis*.

V. CYMBALARIA (*Bodard, Diss.; Bertol. Amæn. Ital. 56; DC. Pr. x. 488*).

Mar. sept.—Tanger! In monte Beni Hosmar pr. Tetuan!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

V. HEDERIFOLIA (*L., DC. Pr. x. 488*).

Mar. merid.—In Atlante Majore videtur frequens. In couvalle Aït Mesan a 1400 m. usque 1600 m. et iterum supra Arround circ. 2200 m. et in jugo Tagherot usque 3000 m. In monte Djebel Tezah circ. 2000 m.!

Spec. in couvalle Aït Mesan circ. 1400 m. lecta iusigniter villosa.

Ar. Geog. Europa omnis (arctica exclusa). Montes regionis Mediterraneæ. Ex Macaronesia non vidi. In America boreali forsân introducta.

BARTSIA (*L. Gen. 730.*)

B. VISCOSA (*L., Eng. Bot. tab. 1045*), = *Eufragia viscosa* (*Benth. in DC. Pr. x. 543*), = *Trixago viscosa* (*Rehb. Fl. Excurs. 360*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Tetuan (*Webb Herb.*)!

Mar. occid.—Casa Blanca! Aïn el Hadjar haud procul Mogador (*J. D. H.*)!

Mar. merid.—Mesfoua! Mzouda!

Ar. Geog. Species imprimis occidentalis a Scotia et Hibernia usque Macaronesiam diffusa. In regione Mediterranea centrali et orientali minus frequens. In Brasilia australi verisimiliter introducta?

B. TRIXAGO (*L., Rebh. Ic. Fl. Germ. xx. tab. 103*), = *Trixago apula* (*Stev. Mem. Mosq. vi. 4; Benth. in DC. Pr. x. 543*), = *Bartsia veriscolor* (*Pers. Syn. ii. 151*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. occid.—Casa Blanca!

Mar. merid.—Tasseremout ad radices Atlantidis Majoris!

Ar. Geog. Fere eadem ac *B. viscosæ*, sed magis australis. Regio omnis Mediterranea. In Gallia occidentali rara. Abyssinia. In Africa australi et in Brasilia australi videtur spontanea.

BARTSIA ASPERA (*Brot. Fl. Lus.* i. 185, sub *Euphrasia*; *Willk. et Lge. Fl. Hisp.* ii. 614, in *Obs.*), = *Lasiopera aspera* (*Lk. et Hffgg. Fl. Port.* tab. 59), = *Odontites aspera* (*Boiss. Voy. Esp.* 473), = *Euphrasia asperrima* (*Spreng. Syst.* ii. 777).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Spec. ex herb. Schousboe benev. comm. eel. Cosson*)!

Ar. Geog. Lusitania. Ager Tingitanus. Nisi fallor, etiam in Hispania occurrit.

Specimina Hispanica vidi, sed locum natalem non notavi.

Huic arete affinis, et meo sensu sicut *B. asperæ* subspecies enumeranda est *B. spicata* (*Ram. in Bull. Philom.* xlii. 141, tab. 10; *Benth. in DC. Pr.* x. 545). Hæc est planta rupestris in Pyrenæis Gallicis rarissime obvia. Differt caule tenuiore flexuoso inferne nitido, nec rigido recto usque e basi aspero, parce ramoso, nec ramos plurimos abbreviatos floriferos edente, indumento totius plantæ molliore, floribus in spica laxiore dispositis, corollæ minus profunde divisæ tubo longiore, antheris minus exsertis, et capsula matura magis inflata.

B. PURPUREA (*Desf. Fl. Atl.* ii. 36, sub *Euphrasia*; *Don, Gen. Syst.* iv. 611, et *Benth. in DC. Pr.* x. 551, sub *Odontite*; non *Dub. Bot. Gall.* 354), = *Euphrasia squarrosa* (*Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! *Cel. Cosson* specimen ex herb. Schousboe sub "*Euphrasia linifolia*" benev. communicavit!

In scheda spec. Schousboeani *eel. Cosson* adiecit "flores lutei," et inde sententiam confirmavit hanc cum proxime affini *B. granatensi* (*Boiss. Elench.* sub *Odontite*) ad subspecies *B. Garideli*, nob. (*Euphrasia viscosæ*, L.) reducendas esse.

Ar. Geog. *B. purpurea* Hispaniæ australis et Afriæ borealis civis indigena est. Formæ arete affines in peninsula Iberica, Gallia meridionali, et Pedemontio obviæ sunt.

Auctoritatem celeb. *Bentham* et *Hooker* (*Gen. Plant.* ii. 977) libenter assecutus species ab auctoribus sub *Bartsia*, *Enfragia*, *Trixagine*, et *Odontite* sub genere Linnæano *Bartsia* enumeravi. Inde necessitate rei coactus nomen novum pro *Euphrasia viscosa*, L., = *Odontites viscosa* (*Rehb. et plur. auctt.*) proposui, et nomen auctoris qui primus hanc stirpem descripsit et delineavit prætuli. Confer *Garidel, Fl. Aix*, 351, tab. 80.

OROBANCHACEÆ (*Lindl. Introd. Nat. Syst.*).PHELIPÆA (*Desf. in Ann. Mus. Par. x. 298*).

P. CÆRULEA (*Vill. Dauph. ii. 406*, sub *Orobanche*; *C. A. M. Enum. Cauc. 104*; *Reut. in DC. Pr. xi. 5*), var.

Mar. merid.—Specimen unicum in convalle Aït Mesan legit J. D. II.!

Ar. Geog. Europa media et australis. Præter sp. nostrum ex Africa non vidi.

Specimen Hookerianum a typo differt calyce magis inflato, calycis dentibus angustis sinibus latis rotundatis divisus dente quinto brevissimo, corollæ labii inferioris lobis late ovatis vix aut ne vix acutis. Planta ulterius inquirenda.

P. SCHULTZII (*Mutel, Fl. Fr. ii. 352*, sub *Orobanche*; *Walp. Rep. Bot. Syst. iii. 463*; *F. Schultz, Arch. Fl. Fr. et All. 99 et seq.*), = *P. lavandulacea* (*Reut. in DC. Pr. xi. 7, pro parte*).

Mar. merid.—Ad radices Atlantis Majoris pr. Sektana legit J. D. II.!

Ar. Geog. Africa borealis. Hispania australis. Verisimiliter per regionem Mediterraneam diffusa, sed indicationes auctorum nimis incertæ.

P. MUTELI (*F. Schultz in Mutel Fl. Fr. ii. 353*; *Reut. in DC. Pr. xi. 9*; *Rchb. Ic. Fl. Germ. xx. tab. 150*).

Mar. sept.—Tanger (*Webb Herb.*)! Tetuan (*Webb spec.*)!

Mar. merid.—Millhian (*J. D. H.*)!

Ar. Geog. Per regionem Mediterraneam in radieibus Compositarum satis frequens. Ex Macaronesia non vidi.

— —, var. *NANA*, *Reut. l. c.*, = *P. nana* (*Rchb. fl.*; *Rchb. Icon. l. c. tab. 151*).

Mar. sept.—Legi prope Cap. Spartel!

Ar. Geog. In regione Mediterranea hinc inde.

P. LUTEA (*Desf. Fl. Atl. ii. 60, tab. 146*; *Reut. in DC. Pr. xi. 13*).

Mar. merid.—Legimus prope Misra ben Kara die deeedente, adeo ut insertionem fibrillarum explorare non licuit!

Ar. Geog. Africa borealis et rarissime occurrit in Hispania australi. In Arabia etiam indicatur.

Specimina nostra quinquepedalia, thyrso bipedali et ultra.

P. TINCTORIA (*Willd. Sp. Pl. iii. 352*, sub *Orobanche*; *Walp. Rep. Bot. Syst. iii. 462*; *Reut. in DC. Pr. xi. 13*), = *Lathræa Phelipæa* (*L., Brot. Fl. Lus. i. 184*), = *Phelipæa lusitanica* (*Coss. Not. Pl. Crit. 43*), = *P. lutea* (*Webb, It. Hisp. 21, et Salzm. Exsicc., non Desf.*).

Mar. sept.—Tanger in arenosis ad ostium fluminis! Eodem ipsissimo loco legit b. P. B. Webb, et mihi sub "*P. lutea*" communicavit! Tanger (*Salzm. et Schousboe, sec. Reuter, l. c.*).

Ar. Geog. In regione litorali Lusitaniæ, Hispaniæ Gaditanæ, et ditionis Tingitanæ hinc inde obvia.

Hæc et *P. senegalensis* (Reut. l. e.), *P. luteæ* valde proximæ, forsân pro subspeciebus habendæ.

OROBANCHE (*L. Gen.* 779).

O. RETICULATA (*Wallr. Gen. Orob. Diasc.* 42; *Reut. in DC. Pr.* xi. 16).

Mar. sept.—In monte Beni Hosmar prope Tetuan, in radicibus *Cistorum* parasita! Hinc etiâ specimen prope Tetuan lectum a P. W. Webb sub *O. fatida* (*Webb Herb.*)!

Ar. Geog. Hispania australis. Montes Tetuanenses.

O. FÆTIDA (*Desf. Fl. Atl.* ii. 59, tab. 144; *Reut. in DC. Pr.* xi. 18).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! In arenosis prope Tanger in radicibus *Medicaginum*! Tetuan (*Webb spec.*)!

Mar. occid.—Mogador (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—Prope Seksaoua!

— —, var. *COMOSA*, nob. Planta pedalis et ultra; spica densa crassa ex bracteis elongatis apice comosa; stigma truncatum biglobosum, stamina ut in typo ima basi pilosiuscula. Bracteæ insigniter cinereo-pubescentes.

Mar. merid.—Sektana ad radices Atlantis Majoris!

Ar. Geog. *O. fatida* peninsulam Ibericam et Africam borealem incolit.

Var. nostra insignis est et facie *O. crinitæ* (*Viv. Fl. Cors. Spec. Nov.* 11) simillima, sed vix a specie divellenda videtur. *O. fatida* est stirps valde ludibunda, et characteres stabiles quibus hanc ab *O. variegata*, Wallr., *O. densiflora*, Salzm., et *O. condensata*, Moris, distinguere vix inveniendi. Priores, cum *O. Hookeriana*, nob., sicut subspecies enumeravi, donec botanici qui eas in vivo sedule explorabunt meliora doceant.

O. variegata (*Wallr. Gen. Orob. Diasc.* 40; *Reuter in DC. Pr.* xi. 17).

Mar. sept.—Tanger (*Salzm. Exsicc.*, sub *O. strigosa*)!

Ar. Geog. Regio Mediterranea centralis. Europa media in calidioribus, sed nullibi frequens. Ex Hispania non vidi.

O. densiflora (*Salzm. Exsicc. MS.*; *Reut. in DC. Pr.* xi. 19), = *O. crinita* (*Rehb. Pl. Crit.* vii. fig. 922, non *Viv.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Ad ostium fluminis Tetuanensis (1851)! Specimen a G. Maw fere eodem loco lectum (1869) in *Herb. Kew.*!

Mar. occid.—Casa Blanca!

Ar. Geog. Hispania. Sardinia. Marocco.

Braectæ in spec. nostris floribus nondum explicatis longiores (et inde spica junior comosa), demum floribus æquilongæ.

Orobanche Hookeriana, *nob., nov. sp.*

Planta subglaberrima, pedalis et ultra; squamæ elongatæ longiuscule acuminatæ, laxæ; spica mediocriter densa; bractæ floribus explicatis dimidio breviores; calycis segmenta in lacinias breves subulatas divisa, in flore e basi spicæ utrinque adnata, in flore altero juniore tantum antice coalita; corolla atrorubens, laciniis calycinis duplo longior, insigniter bilabiata, labio superiore sinuato-denticulato, inferiore trilobo lobis rotundatis; stamina ima basi corollæ inserta, insigniter dilatata, basin versus pilosa; antheræ ad suturas glabræ; stigma capitato-bilobum.

Mar. merid.—In convalle Aït Mesan circ. 1300 m. legit J. D. II. ! Specimen ad eandem formam referendum legi pr. Seksaoua !

O. fœtidæ affinis, differt glabritie, bracteis et calycis laciniis brevioribus, staminibus ima basi tubi corollini nec paullo supra basin insertis, antheris glabris. Specimen prope Seksaoua lectum differt tamen antheris villosulis et staminibus circ. ad quartam partem supra basin tubi corollini insertis.

O. pruinosa (Lap. Suppl. 87; Reut. in DC. Pr. xi. 19).

Mar. sept.—Tanger (*spec. unicum legit J. D. II.*) !

Mar. occid.—Rabat (*R. et Fr. spec. in Herb. Kew.*) ! Casa Blanca !

Mar. merid.—Ad radices Atlantis Majoris. Sektana ! Reraya (*R. et Fr. spec. in Herb. Kew.*) !

— —, var. *SPECIOSA*, = *O. speciosa* (DC. Fl. Fr. Suppl. 395; Reut. in DC. Pr. xi. 19; non Walp. nec Dietr.).

Mar. merid.—Specimen unicum legit prope Sektana J. D. II. !

O. pruinosa et *O. speciosa* a Reuter l. c. pro speciebus distinctis descriptæ, in Fl. Hispanica, ii. 622, a Willkomm et Lange ad synonyma reducuntur. *O. speciosa* mihi videtur prioris var. insignis floribus majoribus et paucioribus, labiis minus papyraceis, superiore rubro-cærulescente, squamis insuper crebrioribus et majoribus distinguenda. Nomen *O. speciosa* antiquus. Hæc tamen forma rarius obvia, dum *O. pruinosa* genuina late diffusa est.

Quoad characteres a sepalis desumptos—scilicet *O. pruinosa* sep. bifidis laciniis subæqualibus, et *O. speciosa* sep. indivisis vel raro inæqualiter bifidis—video inter spec. simul crescentia, immo inter flores ex eadem spica desumptos, sepala omnino variabilia, habent nempe laciniis æqualibus et inæqualibus, tubo corollino brevioribus, æqualibus, vel longioribus. In spec. Maroccanis sepala indivisa nunquam vidi. Pari modo bractæ variant floribus breviores v. æquilongæ, et dum flores superiores nondum explcantur spica apice interdum comosa.

Ar. Geog. *O. pruinosa* per regionem Mediterraneam omnem (inclusa Macaronesia) late diffusa est. *O. speciosa* in Gallia meridionali et alibi indicatur, sed videtur satis rara.

OROBANCHE BARBATA (*Poir. Dict.* iv. 621 ; *Reut. in DC. Pr.* xi. 23, *non Richb.*) ?

Mar. sept.—Tanger in monte Djebel Kebir in radicibus *Cichoraceorum* parasita !

Mar. merid.—In regione inferiore Atlantis Majoris. Reraya ! In convalle Aït Mesan circ. 1300 m. !

Ar. Geog. Peninsula Iberica. Imp. Maroccanum.

Specimina Tingitana Algarbiana (Bourg. Pl. Port. No. 2011) simillima. Atlantica statura multo majora, caule debili sesquipedali, sed vix diversa.

Species mihi aliquantulum incerta, hinc *O. pruinosa*, illinc *O. minori*, Sutton, comparanda, illæ structura, huic facie propior. Stamina ostendit pilosa fere ad basin tubi corollini inserta inferne dilatata et incrassata. Bracteæ variant corolla breviores vel longiores, spica interdum, nec semper, apice comosa.

O. TETUANENSIS, *nob.* = *O. condensata* (*Ball MSS. olim, non Moris*).

Mar. sept.—In monte Beni Hosmar prope Tetuan die 12 Aprilis 1871 florifera lecta !

Planta subspithamæa; caulis cum squamis oblongo-linearibus subobtusis parce pilosus; spica densa; bracteæ lineares obtusiusculæ floribus sat magnis breviores; sepala lineari-lanceolata integra corolla dimidio breviora; corollæ sordide atropurpureæ labium superius subintegrum sinuato-denticulatum, inferius trilobum lobis subæqualibus; stamina ima basi corollæ inserta glaberrima; antheræ magnæ insigniter calcaratæ glaberrimæ; stylus sub apice parce glandulosus, stigma bilobum.

Primo aspectu *O. condensata* (Moris., St. Sard. El., Fl. Sard. iii. 247) subsimilis, sed sepalis angustis integris, bracteis brevioribus obtusis, staminibus et antheris glaberrimis videtur omnino distincta.

LENTIBULARIÆ (*Rich. Pl. Par.* i. 26).

PINGUICULA (*L. Gen.* 30).

P. LUSITANICA (*L., DC. Pr.* viii. 29).

Mar. sept.—Prope Tanger in monte Djebel Kebir (spec. ex herb. Schousboe benev. comm. ecl. Cosson) !

Ar. Geog. Species occidentalis ex Hibernia usque fretum Herculis diffusa.

ACANTHACEÆ (*R. Br. Prod.* 472).

ACANTHUS (*L. Gen.* 793).

A. MOLLIS (*L., DC. Pr.* xi. 270).

Mar. sept.—Tanger (*Webb Herb.*) ! In monte Djebel Kebir ! Prope Tetuan nondum florentem vidi !

Ar. Geog. Regio Mediterranea imprimis occidentalis (exclusa Macaronesia). In Asia Mediterranea rarissime obvius.

SELAGINEÆ (*Juss. Ann. Mus. Par.* vii. 71; *Lindl. Veg. Kingd.* 636, inclusis Globulariis).

GLOBULARIA (*L. Gen.* 112).

G. ALYPUM (*L., DC. Pr.* xii. 613; *Schousb. Gew. Mar.* 43).

Mar. sept.—Tanger (*Broussonet ex Cavan. Anal. Cienc. Nat.*). Tetuan!

Mar. occid.—In monte Djebel Hadid haud procul Mogador! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In provinciis australioribus (*Schsb. l. c.*). Ad radices Atlantis Majoris et usque 1800 m. s. m. Ourika! Amsmiz (*J. D. H.*)! Reraya! In monte Djebel Tezah! Mtouga! (*R. & Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio fere omnis Mediterranea. Deest in ins. Canariensibus, et Maderæ dubia civis est.

VERBENACEÆ (*Juss. Ann. Mus. Par.* vii. 63).

VERBENA (*L. Gen.* 32).

V. OFFICINALIS (*L., DC. Pr.* xi. 547).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Tetuan! (*Webb Herb.*)!

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—Sheshaoua! In ruderalis urbis Marocco!

Ar. Geog. Planta per orbem veterem (exclus. terris arcticis) vulgatissima. In America forsân introducta.

V. SUPINA (*L., DC. Pr.* xi. 548; *Schousb. Gew. Mar.* 4).

In ruderalis et arvis (loc. non indicato), *Schsb. l. c.*

Mar. sept.—Tetuan (*Webb spec.*)!

Mar. occid.—Mogador (*Lowe Cat.*).

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia).

VITEX (*L. Gen.* 790).

V. AGNUS CASTUS (*L., DC. Pr.* xi. 684).

Mar. sept.—Prope Tanger vidi nondum florentem!

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—Scens rivulum ad fines prov. Shedma et Haba! Alibi in regione inferiore vidimus, sed raro. Oued en Fisk (*R. et Fr. sp. in Hort. Kew.*)!

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

LABIATÆ (*Juss. Gen.* 110).

LAVANDULA (*L. Gen.* 711).

L. STÆCHAS (*L., DC. Pr.* xii. 144).

Mar. sept.—Tanger (*Webb Herb.*)! In monte Djebel Kebir! Inter Tanger et Tetuan! In rupibus prope Tetuan!

Mar. occid.—Mogador (*Balansa spec.*)! In monte Djebel Hadid! Agadir (*Coll. Indig.* 1871)!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

LAVANDULA PEDUNCULATA (*Cavan. Prælect.* 70; *DC. Pr.* xii. 144), =
L. Stœchas, var. β (*L. Sp. Pl.* 800).

Mar. merid.—Ad radices Atlantis Majoris. In convalle Ourika circ.
 1100 m.! Reraya ad 1000 m.!

Ar. Geog. Peninsula Iberica. Ins. Madera. Marocco meridionalis.
 Var. in Oriente.

L. DENTATA (*L., DC. Pr.* xii. 145).

Mar. sept.—Inter Tanger et Tetuan! In saxosis pr. Tetuan!

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Ourika!
 In convalle Aït Mesan usque 1400 m.!

— —, var. foliis arete revolutis fere ut in *L. santolinoide*, Jaub. et
 Spach.

Mar. merid.—In convalle Amsmiz circ. 1300 m.!

Ar. Geog. *L. dentata* per regionem Mediterraneam occidentalem calidi-
 orem ex Italia anstrali usque insulam Maderam dispersa est.

L. MULTIFIDA (*L., Benth. in DC. Pr.* xii. 147).

Mar. sept.—Inter Tanger et Tetuan!

Mar. occid.—Casa Blanca! Mazagan! Mogador (*Lowe Cat.*). Prov.
 Haha pr. Mogador! Agadir (*Coll. Indig.* 1871)!

— —, var. *INTERMEDIA*, nob.

Mar. occid.—In monte Djebel Hadid haud procul Mogador! Prov.
 Haha!

Mar. merid.—In saxosis aridissimis prope Aïn Oumast! In planitie
 distr. Mesfioua!

— —, var. *ABROTANOIDES*, nob., = *L. abrotanoides* (*Lam. Dict.*
 iii. 429; *Benth. in DC. Pr.* xii. 146).

Mar. merid.—In rupibus supra Seksaona! Aïn Tarsil.

Ar. Geog. *L. multifida* in regione Mediterranea calidiore imprimis occi-
 dentali (scil. in peninsula Iberica, Calabria et Africa boreali) diffusa,
 in Ægypto etiam obvia est. Var. *abrotanoides* est forma Canariensis
 et Maroccana.

Notæ diagnosticæ quibus *L. abrotanoides* a *L. multifida* distinguitur mihi
 videntur omnes instabiles, nec desunt formæ mediæ, var. *intermediam*
 sistentes. Secundum cel. Bentham loc. cit. *L. multifida* differt specieis
 brevioribus plerumque simplicibus, nec sæpius quinatis, foliorum
 lobis acutis, nec obtusis, foliis floralibus basi latis extus villosis, nec
 ovato-lanceolatis acuminatis, calyceibus demum valde tomentosus nec
 tenuiter puberulis. Spicas elongatas usque 3-pollicares vidi in spec.
 Algeriensibus et Hispanicis *L. multifidæ* genuinæ, et folia floralia in
 spec. Maroccanis angustiora quam in typo, fere eadem ac in *L. abrota-*
noidi Canariensi. In spec. omnibus Marroecanis (Tingitanis exexceptis)
 calyces tenuiter pubescentes dentium margine ciliati; foliorum lacinie
 indunt obtusæ vel acutæ.

Lavandula tenuisecta (Coss. MSS.), = *L. abrotanoides*, var. attenuata (Ball, in *Journ. Bot.* 1875, 174).

Rami tenues sulcati subglaberrimi superne aphylli; folia bipinnatifida laciniis obtusiusculis, ex pilis brevissimis rigidis scabrinseula; spica attenuata plerumque brevis solitaria; folia floralia (vel bractee) ovato-lanceolata acuminata basin versus pubescentia, calyce dimidio breviora; calyx eximie nervosus, tenuiter et vix puberulus, dentibus margine ciliolatis; corollæ tubus tenuior et paullo longior quam in *L. multifida*.

Mar. merid.—In regione inferiore et media Atlantis Majoris. Legimus in convalle Aït Mesan a 1400 m. usque 2200 m.! In monte Djebel Tezah usque 2100 m.! Specimina ex monte Djebel Ouensa ab indigeno Ibrahim lecta sub "*L. tenuisecta*, Coss." benev. communicavit eel. Cosson.

Characteres supra indicati in grege tam ludibundo non gravis momenti. Haud sine dubio hanc pro subspecie enumeravi. Pluribus notis *L. tenuisecta* *L. coronopifoliæ*, Poir., et *L. multifidæ* intermedia videtur, et specimina ex insulis Capitis Viridis in Herb. Kew. sub "*L. coronopifolia*" asservata nostris valde similia sunt.

MENTHA (*L. Gen.* 713).

M. SYLVESTRIS (*L., Benth. in DC. Pr.* xii. 166).

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—Sheshaoua (spec. mox floribundum legit J. D. II.)!

Ar. Geog. Regio omnis Mediterranea. Europa media. Asia temperata. In Africa australi forsitan indigena; in America verisimiliter introducta.

M. ROTUNDIFOLIA (*L., Benth. in DC. Pr.* xii. 167).

Mar. merid.—In ascensu ad jugum Tagherot nondum florentem legerunt Rein et Fritsch!

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia). Europa media. Asia temperata. Indigena, ut fertur, etiam in America boreali.

M. TIMIA, Coss. MSS.

Mar. merid.—Legimus nondum florentem in convalle Ourika! Specimina florentia ex monte Djebel Afougueur ab indigeno Ibrahim lecta comm. Cosson.

Descriptionem hujus, aliarumque plurimarum sp. novarum, viro amicissimo, Floræ Africæ borealis magistro, commendamus.

M. PULEGIUM (*L., Benth. in DC. Pr.* xii. 175).

Mar. sept.—Vidi prope Tanger nondum florentem! El Araïsch (*R. & Fr. spec. in Herb. Kew.*)!

Mar. merid.—In regione inferiore pluries visa. Legimus prope Misra ben Kara! et in distr. Mesfiona!

— —, var. *VILLOSA* (*Benth. loc. cit.*).

Mar. sept.—Prope Tetuan (1851)!

Ar. Geog. Species per regionem omnem Mediterraneam et Europam mediam, neenon in Abyssinia et in utraque America extratropica diffusa.

LYCOPUS (*L. Gen.* 33).

L. EUROPEUS (*L., Benth. in DC. Pr.* xii. 178), = *L. exaltatus* (*Schousb. Gew. Marok.* 4, non *L.*).

Mar. sept.—Tanger (*Salzm. spec., ex Benth. l. c.*), (*Schsb. l. c.*).

Mar. occid.—Mogador (*Broussonet, ex Benth. l. c.*).

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia). Europa media et borealis (exclusa aretica). Asia temperata.

Proenl dubio planta a Schousbæo indicata ad formam *L. europæi* foliis inferioribus pinnatifidis referenda est.

ORIGANUM (*L. Gen.* 726).

O. COMPACTUM (*Benth. Lab.* 334, et in *DC. Pr.* xii. 192), = *O. glandulosum* (*Salzm. Exsicc. non Desf.*), = *O. eretium* (*Schousb. Herb., non L., nec Sieber*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Schousb. spec.*)! Tetuan (1851)!

Mar. merid.—Mtonga (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Peninsula Iberica. Marocco.

THYMUS (*L. Gen.* 727).

T. SATUREIODES (*Coss. in Bull. Soc. Bot. Fr.* xx. 253).

Descriptionem eel. auctoris citare malui:—" Suffrutex 1-5 decim. altus, ramosissimus, dumosus, erectus vel diffusus, ramis vetulis tortuosus flexuosisve cortice rugoso rimoso cinereo-fuscescente, novellis breviter denseque pubescentibus. *Folia pleraque* in ramulos axillares *fasciculata* fasciculis ramorum vetulorum folio matrice orbatis, 4-9 millim. longa, *oblonga obtusa, inferne attenuata*, crassinsecula, *marginibus haud revolutis, nervis obsoletis*, ad basim utrinque pilis paucis rigidis ciliata, crebre glanduloso-punctata; *bractealia* herbaeca, *lanceolata* aenta, calycibus subæquilonga, pubescentia, ciliata, plana. Verticillastri 6-10-flori, apice ramorum in spicas breves subcapitatas approximati. Calyx in pedicello cernuus, extus pube brevissima obtectus, intus ad fauceem dense barbatus, subæqualiter bilabiatus, labii superioris ad tertiam partem tridentati dentibus ovato-lanceolatis lanceolatisve acutis, labii inferioris lineari-subulatis ciliatis. Corolla purpurea, tubo incluso. Stamina exserta, recta, distantia, didynama, inferioribus longioribus."

Mar. merid.—In regione inferiore Atlantis Majoris frequens, in regione media rara. In montibus prope Keira circ. 1000 m. detexit Balansa anno 1867 (*Coss. l. c.*). Legimus in convalle Aït Mesan usque 1400 m.! In convalle Amsmiz! Supra Seksaoua circ. 1100 m.! Ex provincia Mtonga (*R. et Fr. spec. in Herb. Kew.*)! In monte Djebel Tezah usque 1900 m.!

— —, var. *PSEUDOMASTICHINA*, nob. Foliis bractealibus et dentibus calycinis longe mucronatis corollæ subæquilongis, floribus læte roscis.

Mar. merid.—In convalle Ourika circ. 1100 m. ! in convalle Aït Mesan ! Monente cl. auctore :—*T. satureioides*, foliorum fabrica in genere insignis, habitu *Satureiam* refert, et inde nomen, sed ad *Thymum* pertinet calyce bilabiato fauce dense villosa, staminibus distantibus, non ascendentibus.”

In grege *T. vulgaris* collocandus.

THYMUS SERPYLLUM (L., DC. *Pr.* xii. 200), var. *ATLANTICUS*, nob. Differt a typo foliis elliptico-lanceolatis in petiolum brevissimum sensim attenuatis, apice obtusiusculis, eximie glanduloso-punctatis, florum verticillastris depauperatis in capitulum parvum laxum approximatis.

Mar. merid.—In regione alpina Atlantis Majoris, in jugo Tagherot, circ. 2900 m. !

Ar. Geog. Per Europam omnem (excepta peninsula Iberica australiore) a litore maris usque regionem glaciale summam Alpium diffusus. Adest in montibus Asiæ mediæ et Abyssiniæ.

T. LANCEOLATUS (Desf. *Fl. Atl.* ii. 30, tab. 128 ; *Benth. in DC. Pr.* xii. 203), = *T. Sibthorpii* (*Benth. Lab.* 345), var. *CRISPUS*, nob., = *T. pallidus*, Coss. MSS.

Planta nostra a typo differt foliis omnibus (floralibus exceptis) margine revolutis, floralibus (bractealibus) quam cætera paullo latioribus, floribus subminoribus, calycis labii inferioris dentibus eximie pectinato-ciliatis.

Mar. merid.—In regione inferiore Atlantis Majoris. Leginus in convalle Aït Mesan usque 1600 m. s. m. ! In convalle Amsmiz circ. 1400 m. ! Habui specimen ex monte Djebel Afougneur ab indigeno Ibrahim lectum, cui scheda “ *T. pallidus*, Coss. MSS.”

Ar. Geog. Africa borealis. Asia Mediterranea. Græcia.

Stirps variabilis, herba tota in spec. Orientalibus et in nostris pallide virens, in Algeriensibus plurimis quæ vidi minus pallida, et folia floralia cum calycibus subpurpurascentia. Calyx, ut monuit eel. Benth. l. c., et pariter fit in spec. nostris Maroccanis, variat labiorum dentibus longioribus brevioribusve, semper fauce eximie villosus. In tabula Fontanesiana folia floralia cæteris angustiora phytographus ostendit ; in spec. Algeriensibus quæ vidi æquilata sunt, in nostris paululum latiora. Var. nostra ludit herba tenuiter puberula vel pilis crebris hirtula.

T. CAPITATUS (Lk. et Hoffgg. *Fl. Port.* i. 123 ; *Benth. in DC. Pr.* xii. 204).

Mar. sept.—Tanger (*Broussonet, ex Benth. l. c.*). Vidi nondum florentem, et spec. ex Herb. Schousboe benev. comm. eel. Cosson ! Tetuan, nondum florens !

Ar. Geog. Regio Mediterranea calidior exclusa Macaronesia. Deest in Gallia et in Italia media ac boreali.

THYMUS BROUSSONNETII (Boiss. Voy. 492, tab. 141; Benth. in DC. Pr. xii. 205).

Mar. occid.—Mogador! (*Balansa spec.*)! (*Broussonnet ex Boiss. l. c.*) (*Lowe Cat.*). Prov. Haha ab urbe Mogador meridiem versus!

Mar. merid.—Prov. Shedma! Mskala! Mtouga! Seksaoua!

Fit rarior in terris interioribus a mari remotis. Descriptio Boissieriana ex planta culta, quam secutus est Benth., spec. nostris non admodum convenit. Canles video semper puberulos vel subtomentosos; folia subtus insigniter nervosa glabra, vel sæpius ex pube brevissima rigida scabrida; folia floralia ex basi lata ovata, calyce occultantia, semper hirtula et margine ciliata, demum læte colorata; calycis labium inferius superiore longius, dente medio majore; corollæ tubum elongatum extus pubescentem.

Species propria prov. austro-occidentalibus Imp. Marocani nullæ nisi sequenti proxima. Planta insectis invisæ in urbe Mogador ab incolis ad vestes conservandas adhibetur.

T. MAROCCANUS (*Ball in Jouru. Bot.* 1875, 174). (Plate XXVII.)

Planta ramosissima, ramis suberectis; folia sessilia, elliptico-lanceolata, basin versus attenuata, æqualia, subglaberrima vel hirtula, floralia conformia viridia, vel rarius aliquantulum purpurascentia; flores in capitulo brevi laxo approximati; calycis labii superioris tridentati dentibus subæqualibus, inferioris sublongioris dentibus anguste cuneatis tubo fere duplo longioribus; corollæ tubo extus hirtulo quam calyx sesquilingiore, labio inferiore trilobo lobis rotundatis, medio laterali-bus aliquid majore; stylus exsertus, stamina duo longiora subsuperans. Habitu ad *T. Broussonnetii*, Boiss., proxime accedit, sed præter alias notas foliis floralibus cæteris conformibus, et tubo corollæ multo brevior videtur probe distincta.

Mar. merid.—Ad radices Atlantis Majoris a 900 m. usque 1300 m. Tasseremout! Ourika! Reraya! In convalle Aït Mesan!

T. MUNBYANUS (*Boiss. et Reut. Pug.* 96), = *T. ciliatus*, var. *Munbyanus*, *De Noe*.

Mar. sept.—Inter Tanger et Tetuan! Frequens circa Tetuan, ubi legi 1851 et iterum 1871! et spec. habui ex herb. Webb.! In monte Beni Hosmar.

Proxime affinis *T. ciliato* (Desf. Fl. Atl. ii. 10, tab. 122, sub *Thymbra*; Benth. in DC. Pr. 205), a quo differt imprimis habitu, ramis sterilibus longis prostratis, floriferis erectis, foliis floralibus latioribus acuminatis eximie pectinato-ciliatis, calycis labio superiore profundius incisodentato. Pro subspecie enumeravi, dum formas intermedias non vidi. *Ar. Geog.* Algeria occidentalis. Ditio Tetuanensis.

Thymi specimen nondum floriferum legit in monte Djebel Tezah cel. J. D. Hooker *T. satureioidi* subsimile, sed foliis non fasciculatis diversum. Planta ulterius inquirenda.

SATUREJA (*L. Gen.* 707).

S. INODORA (*Salzm. in Benth. Lab.* 353; *Benth. in DC. Pr.* xii. 210),
= S. glomerata (*Schsb. MSS. in Herb.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (Spec. ex herb. Schousboe comm. cel. Cosson)!

Ar. Geog. In utroque latere freti Herculis.

In Flora Hispanica non indicata; legi tamen in collibus prope S. Roque

HYSSOPUS (*L. Gen.* 709).

II. OFFICINALIS (*L., DC. Pr.* xii. 251).

Mar. merid.—In regione alpina Atlantis Majoris. Legimus in jugo Tagherot a 3000 m. usque 3200 m.! et in vertice Djebel Tezah ultra 3300 m.! Spec. in monte Djebel Ouensa ab indig. Ibrahim lecta benev. comm. Cosson!

Ar. Geog. In montibus regionis Mediterraneæ a Sierra Nevada Hispaniæ usque Persiam. In Europæ mediæ rupestribus calidioribus et in montibus Asiæ mediæ. Spec. Africana præter nostra non vidi.

MICROMERIA (*Benth. in Bot. Reg.* xv. No. 1283).

M. MICROPHYLLA (*D'Urv. Enum. Pl. Arch. in Mém. Soc. Linn. Par.* 327, sub *Thymo*; *Benth. Lab.* 377, et in *DC. Pr.* xii. 219).

Mar. merid.—Frequens in regione inferiore Atlantis Majoris. Tassere-mont! Ourika! Aït Mesan! Aïn Tarsil! Sektana (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Africa borealis. Italia australis. Insulæ Sicilia et Melita. Planta ad Aïn Tarsil lecta sistit formam nanam dentibus calycinis brevissimis.

— — —, var. VILLOSA (*Benth. l. c.*).

Mar. merid.—In rupibus supra Seksaona! Keira (*Bolansa, spec. in Herb. Kew*, sub *M. microphylla*, var. *S. filiformis*, Desf., Coss.)

Spec. nostra diagnosi et tabulæ *Saturejæ filiformis* (Desf. *Fl. Atl.* ii. 8, tab. 121) omnino conformia. Præter indumentum a *M. microphylla* recedit calyceis subbilabiati dentibus longioribus subulatis, et cymarum pedunculis dimidio brevioribus.

CALAMINTHA (*Mæneh, Meth.* 408).

C. MENTHÆFOLIA (*Host, Fl. Austr.* ii. 129), = C. adscendens (*Jord. Obs. Fragm.* iv. 8, tab. 1). Var. BÆTICA, nob., = C. Bætica (*Boiss. et Rent. Pug.* 92).

Mar. sept.—Legi in monte Djebel Kebir prope Tanger! Tanger (*Salzm. Exsicc.*, sub *Thymus Calamintha*, testibus Boissier et Renter l. c.).

Ar. Geog. *C. menthæfolia* est species per Europam imprimis australem late diffusa. Var. nostra, quam legi in rupibus Calpes, videtur Hispaniæ, Lusitaniæ et Africæ Tingitanæ propria civis, nisi forte huc

referenda sit stirps Canariensis sub *C. officinali* β , *villosissima* in DC. Prod. l. e. memorata.

Consentientibus plerisque auctoribus *C. menthaefolia*, Host, est ipsissima *C. officinalis*, Benth. non Mœnch, dum *C. officinalis*, Mœnch, foret eadem ac *C. sylvatica* (Bromf. in Eng. Bot. Suppl., tab. 2897, Benth. in DC. Pr. l. e.). An hæ stirpes pro speciebus distinctis habendæ sint est mihi res adhuc incerta. Varietati nostræ valde proxima est *C. heterotricha* (Boiss. et Reut. Pug. 93), ad varietatem *C. menthaefoliae* forsan reducenda.

CALAMINTHIA ACINOS (L., sub *Thymo*; Benth. in DC. Pr. xii. 230), = *Acinos vulgaris* (Pers. Syn. ii. 131).

Mar. merid.—Ourika (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Europa media et australis. Regio Caucasio-Caspica.

C. GRAVEOLENS (*M. B. Fl. Taur.-Cauc.* ii. 60, sub *Thymo*; Benth. in DC. Pr. xii. 231).

Mar. merid.—In convalle Aït Mesan ad 1700 m. s. m.!

Ar. Geog. Regio fere omnis Mediterranea, imprimis orientalis. Adest etiam in Europa orientali. In Macaronesia nondum detecta.

C. ALPINA (L., sub *Thymo*; Benth. in DC. Pr. xii. 232), var. *PARVIFLORA*, *nob.*

Mar. merid.—In regione media et superiore Atlantis Majoris. In convalle Aït Mesan et in ascensu jugi Tagherot a 2200 m. usque 2800 m. In monte Djebel Tezah circ. 2500 m. In convalle Ourika spec. legit J. D. H.!

Radix perennis lignosa, caules tenues prostratos purpurasecentes emit-tens; folia petiolata integerrima vel obsolete denticulata; verticillastri pauciflori, ad apicem ramorum approximati; calyx ad quartam partem, interdum usque ad medium, fissus, labii inferioris longioris dentibus rectis v. subrecurvis subulatis, superioris dentibus ex basi triangulari acuminato-subulatis demum patentibus; corolla parva calyce vix duplo longior.

Ar. Geog. *C. alpina* per montes Europæ mediæ et australis late diffusa. In Asia Minore etiam indicata.

C. ATLANTICA (*Ball in Journ. Bot.* 1875, 175).

Inermis, valde ramosa, ramis adscendentibus prostratisve, interdum radicantibus; folia parvula longiuscule petiolata, late ovata, obtusa, subintegerrima, obscure crenulata; verticillastri subsexflori, interdum unicus solitarius terminalis, interdum 2, approximati; bracteæ subulatæ calyce breviores; calyx bilabiatus, usque ad medium vel ultra fissus, dentibus labii superioris ultra medium coalitis, lab. inferioris subulatis incurvatis; corollæ parvulæ tubo calyceis dentibus vix æquilongæ.

Mar. merid.—In Atlante Majore a 1200 m. usque 2400 m. Legimus in convalle Ourika! Aït Mesan supra Arround! In monte Djebel Tezah!

Statura multo minore, ramis debilibus fere prostratis, verticillis paucifloris et corollis parvis a *C. Clinopodio* insigniter differt; structura tamen vix satis distincta. Formam fere intermediam descripsit clar. Lange, *C. Clinopodium* β , *gracilis* (Lange, Pug. 177; Willk. et Lge. Fl. Hisp. ii. 416). Proxime affinis est etiam *C. Arundana* (Boiss. Voy. Esp. 498, tab. 145, sub *Melissa*; Benth. in DC. Pr. l. c.) species non satis nota, quæ ex descript. a nostra differt tantum foliis acutis dentatis et florum verticillis globosis. Interea planta nostra sicut subspecies enumeranda videtur, quæcum forsân adsocianda erit varietas Langeana supra memorata.

SALVIA (*L. Gen.* 39).

S. MAURORUM (*Ball in Journ. Bot.* 1875, 175). (Plate XXVIII.)

Proxima *S. Caudelabro* (Boiss. El. 72; Benth. in DC. Pr. xii. 263), cui sicut subspecies adsocianda, sed plurimis notis diversa. In nostra folia dimidio minora, rigidiora, superne lævia, margine integerrima; inflorescentia simplex, nec paniculata, ex cymis subtrifloris oppositis, sessilibus vel breviter pedunculatis, verticillos laxos paucifloros efformantibus; calycis breviter pedicellati, minute puberuli, vix aut ne vix glanduloso-viscosi dentes subæquales acuti mucronati; corolla gracilior, extus vix pilosa.

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Amsmiz circ. 1500 m.! In monte supra Seksaoua ad 1600 m.!

S. Blancoana (Webb et Held. in Blanco Pl. Jaen, No. 308), species dubia, a Willk. et Lge in Prod. Fl. Hisp. prætermissa, plantæ nostræ affinis est, et me iudice inter subspecies *S. Caudelabri* adnumeranda. Hæc gaudet panicula ramosa, floribus in cymis contractis pedunculatis, subsessilibus, calycis pilosi dentibus longioribus acutioribus. Hæc omnes a *S. officinali*, L., et *S. lavandulæfolia*, Vahl, præter alias notas, bracteis minimis ovali-acutis cito deciduis, nec late ovatis longiuscule acuminatis subpersistentibus, dignoscuntur.

S. INTERRUPTA (*Schousb. Gew. Marok.* 6, tab. 1; *Benth. in DC. Pr.* xii. 266).

Mar. merid.—In monte Beni Hosmar pr. Tetuan legit G. Maw, anno 1869, et spec. unicum nondum florens legit J. D. Hooker 12 Aprilis 1871!

Mar. occid.—Specimen ex monte Djebel Hadid haud procul Mogador ab indigeno Ibrahim lectum humanissime comm. cel. Cosson! In eodem monte verisimiliter legit h. Schousbœc semina in horto Hafniensi culta, unde specimen in opere citato delineatum.

Species Imp. Maroccani propria et verisimiliter rara. In regione inferiore Atlantis nullibi obvia.

[*S. viridis* (L., Benth. in DC. Pr. xii. 277), verisimiliter ex errore, schedæ Salzmännianæ, prope Tanger l. c. indicatur. Hæc species, a

neimine recentiorum in agro Tingitano visa, prope Malaga a Salzmann lecta fuit, et specimen Malacitanum facile sub rubrica "Tanger" evasit.]

[*S. tingitana* (Etl. Salv. No. 30; Benth. in DC. Pr. xii. 282) est planta Tunetana cui nomen falsum erronee attributum. Facile tamen in territorio Maroccoano invenienda erit, dum adest in agro Gaditano. Hæc est *S. fætida* (Lam. Ill. i. 69) et *S. coarctata* (Vahl, Enum. i. 277). Nomen triviale Lamarekianum aptum et Vahliano antiquius videtur hodie antefendum.]

SALVIA ARGENTEA (L., Benth. in DC. Pr. xii. 288).

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Videtur satis frequens in regione inferiore. Aïn Oumast! Shedma! Ourika (*R. et Fr. spec. in Herb. Kew.*)! Sektana (*R. et Fr. spec. in Herb. Kew.*)! Vidi etiam prope Seksaoua!

Ar. Geog. Regio Mediterranea ealidior (exclusa Macaronesia). Deest in Gallia et in Italia media æ superiore.

S. BICOLOR (Desf. Fl. Atl. i. 22, tab. 2; Bot. Mag. tab. 1774; Benth. in DC. Pr. xii. 288; Schousb. Gew. Marok. 5). Hujus lusus est *S. dichroa* (Hook. fil. in Bot. Mag. xeviii. tab. 6004).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)! (*Schsb. l. c.*). (*Broussonnet ex Benth. l. c.*).

Ar. Geog. Stirps insignis in regione litorali Africæ borealis rara, et in Hispania Bætica rarissima.

Circa Tingim sat frequens et valde ludibunda. Planta macrior 8-9-pedalis gaudet foliis circuitu late ovatis basi profunde cordatis, inferioribus pedem et amplius longis lobato incis, ex lobis inferioribus sese tegentibus pseudopeltatis, caulinis inferioribus petiolatis, pinatifido-lobatis, laciniato-dentatis, supremis subsessilibus. In specimenibus minoribus folia inferiora oblongo-lanceolata, basi truncata, parce dentata, interdum integerrima. Ex radice prope Tingim (nec ad radices Atlantis Majoris) lecta enata est *S. dichroa*, H. fil., lusus sp. nostræ, foliis basi truncatis subintegris, corollæ labio rosco.

Schousboe plantam jam defloratam mense Aprili prope Tingim vidit. Ego legi adhuc copiose florentem mense Junio 1871 in dumetis juxta urbem et in clivo montis Djebel Kebir.

S. algeriensis (Desf. Fl. Atl. i. 23, tab. 3; Benth. l. c.) foliis angustioribus (non semper ut in tabula citata integris) et floribus minoribus a *S. bicolori* distincta, forsan melius sicut subspecies enumeranda est.

S. OCHROLEUCA (Coss. in Bull. Soc. Bot. Fr. xx. 254).

Planta a me non visæ descriptionem auctoris citare licet:—

"Plantis *perennis*, caudice lignoso tortuoso. Caules circiter 20 centim. longi, erecti, herbaceo-subindurati, villosi pilis patentibus immixtis brevioribus glanduliferis. Folia bullato-rugosissima, utrinque, sed infra densius, pubescenti-velutina; radicalia plura rosulata, humo haud

adpressa, abrupte petiolata, ambitu *ovato-oblonga oblongave, plus minus incisopinnatifida*, lobis ovatis oblongisve obtusissimis inciso-erenatis; caulina 2-4, sessilia, subconformia; *floralia* sessilia, *ovata acuminata, calycibus paulo longiora vel breviora*, villosa. Racemi simplices, verticillastris 4-7-floris, sæpius caulem a tertia parte inferiore obtinentibus, inferioribus remotiusculis, superioribus approximatis. *Calyx* subsessilis, hirsuto-villosus pilis longis glanduliferis, campanulatus tubo brevi, *labiis subæquilongis distantibus, superiore oblongo summo apice 3-dentato dentibus ovato-acuminatis subæqualibus conniventibus, inferiore bifido laciniis lanceolatis in euspidem subspinescentem attenuatis*, tubo intus villosus. *Corolla ochroleuca* 10-15 millim. longa, calyce subduplo longior, *tubo infra faucem gibboso-ventricosus intus annulo piloso destituto, labio superiore arcuato galeato compresso apice rotundato emarginato extus pubescente, labii inferioris superiori subæquilongi lobis lateralibus oblongis suberectis medio patente concavo rotundato subemarginato*. *Connectiva inter se extremitate articulorum inferiorum connata, multo infra medium articulata, articulo superiore elongato lobo antheræ versatili oblongo donato, articulo inferiore crasso ad callum sterilem reducto*. Stylus apice bifidus, lobis subulatis inferiore multo longiore. Florifera Maio 1867 lecta."

Mar. merid.—In montibus ad austro-occidentem urbis Marocco, prope Keira, ad 1000 m. cl. Balansa detexit. Nos non vidimus.

"*S. ochroleuca* calyceis labio superiore summo apice tridentato dentibus conniventibus, corollæ tubo sub fauce gibboso-ventricosus intus exanulatus, labio superiore arcuato galeato compresso, labii inferioris lobis lateralibus oblongis suberectis medio patente rotundato concavo subemarginato, connectivis multo infra medium articulatis inter se extremitate articuli inferioris sterilibus ad callum reducti connatis ad sectionem *Plethiosphace* (Benth. in DC. Prod. xii. 288) pertinet et juxta *S. Austriacam*, L., colore corollæ collocanda, sed habitu, foliis multo angustioribus haud humo adpressis et forma dentium calycinorum eximie distincta."

SALVIA VERBENACA (L., Benth. in DC. Pr. xii. 294), var. *OBLONGIFOLIA* (Benth. l. c.), = *S. oblongata* (Vahl, Enum. i. 256), = *S. betonicæfolia* (Lam. Ill. i. 70).

Mar. occid.—Agadir (*Coll. Indig.* 1871)! In via qua itur a Mogador versus Shedma (*G. Maw spec.*)!

Mar. merid.—In prov. Shedma pluribus locis! In saxosis aridissimis prope Aïn Oumast!

— —, var. *PRÆCOX* (Lange, Pug. 179; Willk. et Lge. Fl. Hisp. i. 426), = *S. clandestina* (L., Benth. in DC. Pr. l. c.; Schousb. Ger. Marok. 5).

Mar. sept.—Tanger (*Schsb. l. c.*)!

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—In convalle Aït Mesan! a 1400 m. usque 2400 m.! Prope Seksaoua.

Ar. Geog. S. Verbenaca genuina per Europam fere omnem a Suecia usque litora Maris Interni diffusa est et prope Byzantium indicatur.

Var. *oblongifolia* in Hispania et Africa boreali obvia est. Var. *præcox* (*S. clandestina*) per regionem omnem Mediterraneam a Persia usque Macaronesium frequens occurrit.

SALVIA TARAXACIFOLIA (*Coss. in Bull. Soc. Bot. Fr. xx. 253; Bot. Mag. xeviii. tab. 5991*).

Sequitur descriptio Cossoniana:—

“Planta *perennis*, caudice lignoso tortuoso. Caules 5–25 centim. longi, erecti vel ascendentes, herbaeco-indurati, villosi pilis patentibus latis immixtis pilis brevioribus tenuioribusque glanduliferis. *Folia* radicalia plura rosulata, humo laud adpressa, pubescentia arachnoidea supra subcanescentia, *infra cano-tomentosa*, petiolata, *pinnatipartita vel pinnatisecta*, lobis 3–4-jugis ovatis oblongisve grosse dentatis rarius subintegris terminali majore; caulina pauca 2–4, subconformia, sæpe nulla; *floralia* persistentia, sessilia, *ovata in cuspidem spinescentem acuminata*, *calyces subæquantia vel breviora*, dorso lanata, margine ciliata. Racemi simplices, verticillastris 4–7-floris sæpius caulem a tertia parte inferiore obtinentibus, inferioribus remotis, superioribus subapproximatis. *Calyx* arachnoideo-lanatus pilis longioribus immixtis, tubuloso-campanulatus, coriaceus venulis anastomosantibus reticulato-venosus, *labio superiore truncato 3-dentato* inter dentes plano-subconcavo dente medio brevior, inferiore bifido, *dentibus omnibus patulis cuspidem subulata spinescente terminatis*, tubo intus sed *præsertim ad faucem dense barbato*. *Corolla* pallide purpurascens?, 2–3 centim. longa, calyce subtriplo longior, *tubo amplo intus ad tertiam partem inferiorem annulo pilorum donato*, labio superiore vix arcuato galeato compressiusculo extus pubescente, *labii inferioris superiori subæquilongi lobis lateralibus late ovato-rotundatis medio obovato-rotundato emarginato patente subconcavo*. *Connectiva inter se libera*, elongata, paulo *infra medium articulata articulo utroque lobum antheræ gerente*, articuli superioris lobo oblongo-lineari adnato, articuli inferioris crassioris lobo brevior oblongo. Stylus apice bifidus lobis subulatis inferiore paulo longiore. Nuculæ fusæ, nitidæ, obovato-subglobosæ.”

Mar. merid.—In glareosis torrentium ad radices Atlantis Majoris. Tasscremont! Onrika! Reraya (*R. et Fr. spec. in Herb. Kew.*)! In montibus ad meridiem urbis Maroc. prope Mulai-Ibrahim, ad 1000 m. cl. Balansa anno 1867 primus detexit; altero loco ignoto eorundem montium mulio Ibrahim auspice cl. Beaumier anno 1873 legit.

Monente ecl. auctore:—“*S. taraxacifolia* calyce campanulato dentibus

acutis patentibus maturatione haud mutatis, corollæ tubo amplo exserto intus piloso-annulato, labio superiore vix arcuato, labii inferioris lobis lateralibus patentibus medio late emarginato ad sectionem *Eusphace* (Benth. in DC. Prod. xii. 263) accedit, sed forsan melius ob connectivorum fabricam ad sectionem *Heterosphace* (Benth. in DC. Prod. xii. 350) referenda."

Species distinctissima prope *S. scabram*, Thunb., ut mihi videtur, collocanda. Corolla pallide, vel interdum saturate, purpurea.

SALVIA ÆGYPTIACA (L., Benth. in DC. Pr. xii. 355).

Mar. merid.—Legimus in alveo glareoso torrentis Oned en Fisk prope Gurguri! Spec. ab indigeno Mardoehæo in Oasi Akka lectum comm. cel. Cosson!

Ar. Geog. Regio Mediterranea calidior. In Oriente et in Africa boreali, nec in Europa occurrit.

NEPETA (L. Gen. 710).

N. MULTIBRACTEATA (Desf. Fl. Atl. ii. 11, tab. 123; Benth. in DC. Pr. xii. 374).

Mar. sept.—Prope Tetuan (*Maw spec. in Herb. Kew.*)!

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Shedma! Aït Mesan circ. 1300 m.! Prope Mzouda! Prov. Mtouga (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Africa boreali-occidentalis. Lusitania australis. Specimina Algeriensia a Maroccanis et Lusitanicis aliquid diversa sunt foliis basilioribus, caule simplici nec ut in nostris sæpe ramoso.

N. Apulei (*Ucria in Guss. Prod. Fl. Sic.* ii. 80; Benth. in DC. Pr. xii. 375), = *N. tuberosa* (Desf. Fl. Atl. ii. 10, non L.), = *N. rosea* (*Salzm. Exsicc.*), = *N. acerosa* (*Webb, It. Hisp.* 20).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb spec.*)!

Mar. occid.—Mogador (*Lowe Cat.*). Djebel Hadid (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—Ad radices Atlantis Majoris haud infrequens. Mesfioua! Tasseremout! Ourika! et alibi visa.

Ar. Geog. Africa boreali-occidentalis. Hispania australis. Sicilia.

N. Apulei a *N. multibracteata* differt foliis oblongo-lanceolatis, nec ovatis basi cordatis, caulinis fere omnibus sessilibus, spica tenuiore et bracteis oblongo-lanceolatis, nec lineari-subulatis. Hæ notæ tamen variabiles, et occurrunt spec. *N. multibracteata* a *N. Apulei* ægre distinguenda. Hanc ego pro subspecie enumeravi, et eodem pacto *N. reticulata* (Desf. Fl. Atl. ii. 11, tab. 124), in ditioe nostra nondum detecta, mihi videtur subspecies *N. tuberosæ* (L., Benth. l. c.).

N. ATLANTICA (*Ball in Journ. Bot.* 1875, 175).

Ex radice lignosa caules plurimos erectos, 1–2 pedales, simplices vel

breviter ramosos emittens; folia parva, breviter petiolata, ex basi truncate cuneato-ovata, in pagina inferiore insigniter nervosa, margine crenata, superiora sessilia, floralia minima, omnia tenuiter velutina; verticillastri remoti, cymis breviter pedunculatis sub 7-floris, bracteis minutis linear-lanceolatis; calyx ore obliquo, dentibus inæqualibus, sæpius coloratis, cuneatis, tubi tertiæ vel quartæ parti æquilongis; corolla pallide ochroleuca, tubo vix exserto.

Mar. merid.—*Leginus* in convalle Amsmiz 1200–1400 m.! Spec. ex monte Djebel Afougneur ab indigeno Ibrahim lecta comm. ecl. Cosson, sub *N. atlantica*, Coss. MSS.

Nostræ proxima est *N. marifolia*, Boiss. et Huet in Diagn. Pl. Or., ser. 2, fasc. iv. p. 24. In hac verticillastri florum approximati, calyx major, dentibus brevioribus et minus acutis, folia magis rugosa, bracteæ breviores et simul latiores, corolla denique gaudet tubo duplo longiore. *N. racemosa* (Lam., Benth. in DC. Pr. xii. 385), species etiam affinis, differt præter alias notas statura multo majore, corollæ tubo longiore, foliis et floribus duplo majoribus.

PRUNELLA (*L. Gen.* 735), interdum BRUNELLA.

P. vulgaris (*L. Sp. Pl.* 837), = *Brunella vulgaris* (*Benth. in DC. Pr.* xii. 410).

Mar. sept.—Prope Tanger (*Webb Herb.*)! (*Salzm. Exsicc.*)!

Ar. Geog. Species cosmopolita in temperatis totius fere Orbis late diffusa; in Asia usque Japoniam, in America boreali et in montibus reg. æquinoctialis, et in Australia.

CLEONIA (*L. Gen.* 736).

C. lusitanica (*L., Benth. in DC. Pr.* xii. 411).

Mar. sept.—Tanger (*spec. ex Herb. Schousboe*)!

Mar. merid.—Ad radices Atlantis Majoris. Tasseremout! Ourika (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Peninsula Iberica. Algeria. Marocco.

SIDERITIS (*L. Gen.* 712).

S. grandiflora (*Salzm. Pl. Exsicc. Ting.; Benth. in DC. Pr.* xii. 441).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! In monte Djebel Dersa pr. Tetuan (*Webb spec.*)!

Ar. Geog. Africa Tingitana et (rarissime) in Hispania australi.

S. scordioides (*L., Benth. in DC. Pr.* xii. 442), var. *angustifolia* (*Benth. l. c.*), = *S. hyssopifolia* (*L. Sp. Pl.* 803), = *S. hyssopifolia*, var. *angustifolia* (*Willk. et Lge. Fl. Hisp.* ii. 453).

Mar. occid.—? In monte Djebel Hadid haud procul Mogador (*R. et Fr.*

spec. in Hort. Kew.). Specimen incertum forsan melius sub var. *elongata* (Benth. l. c.) enumerandum.

Mar. merid.—? Spec. caule florifero anni præcedentis exsiccato legi in regione alpina montis Djebel Tezah !

Ar. Geog. Species per montes Hispaniæ, Italiæ, et Galliæ anstralis, necnon Alpes occidentales late diffusa. In Sicilia a Gussone indicata.

Specimina nostra nimis incompleta, forsan potius ad *S. incanam* referenda, rami tamen juniores cum foliis novellis subglaberrimi. Notæ quibus *S. hyssopifolia* a *S. scordioide* distinguitur mihi videntur nimis inconstantes, et formæ plurimæ intermediæ obviæ sunt.

SIDERITIS HIRSUTA (L., *Benth. in DC. Pr.* xii. 444).

Mar. merid.—Ad radices Atlantis Majoris prope Tasseremout !

Ar. Geog. Europa australi-occidentalis a Pedemontio usque Lusitaniam. Marocco. Var. in Algeria. Spec. typica Africana præter nostra non vidi.

S. ROMANA (L., *Benth. in DC. Pr.* xii. 445).

Mar. sept.—Legi juxta Tetnan anno 1851 !

Ar. Geog. Regio fere omnis Mediterranea (exclusa Macaronesia).

S. MONTANA (L., *Benth. in DC. Pr.* xii. 446), var. calycis tubo patule et molliter villosa.

Mar. merid.—In saxosis supra Seksaona !

Ar. Geog. Species in regione tota Mediterranea (exclusa Macaronesia) et in Europa orientali late diffusa.

S. VILLOSA (Coss. in *Bull. Soc. Bot. Fr.* xx. 255).

Sequitur descriptio Cossoniana :—

“Planta perennis caudice et parte inferiore caulium induratis, pilis patentibus molliter villosa. Caules plures, diffusi, 10–15 centim. longi, tetragoni angulis præminentibus obtusis, simplices vel inferne ramosi. Folia crenata, crenis etiam in superioribus muticis, inferiora petiolata ovata vel ovato-oblonga basi cordata, floralia subconformia minora in petiolum brevem attenuata calycibus longiora. Verticillastri 3–6-flori, remotiusculi vel superiores in spicam foliatam subapproximati. Calyx breviter pedicellatus, demum patenti-subreflexus, patule hispidus, campanulatus, tenuiter 10-nerviis, dentibus omnibus subæquilatis, ovato-triangularibus apice mucronatis, superiore longiore, tubi longitudine dimidia brevioribus, demum suberectis. Corolla 8–9 millim. longa, albida, sparse pilosa, tubo cylindrico, subrecto, calycem paulo excedente, intus piloso-annulato, labio superiore subascendente planiusculo integro, labio inferiore subæquilongo patente trilobo lobo medio latiore obovato integro. Stamina 4, tubo inclusa, superiora breviora antherarum lobis divaricatis, inferiora multo longiora antheris minutis crassis. Stylus inclusus, apice bifidus, lobo superiore tereti-truncato, inferiore dilatato basi superio-

rem amplectente. Nueulæ fuscæscientes albido-variegatæ, obovatæ, apice rotundatæ, glabræ, punctato-tuberculatæ."

Mar. merid.—Frequens in regione inferiore ac media Atlantis Majoris. In convalle Ourika! In convalle Aït Mesan usque in ascensu jugi Tagherot a 1400 m. ad 2700 m.! In monte Djebel Tezali circ. 1800 m.! In monte Sidi Fars pr. Moulai Ibrahim primus detexit el. Balansa.

Monente eel auctore, "ad sectionem *Hesiodia* (Benth. in DC. Prod. xii. 445) pertinens et juxta *S. montanam*, L., collocanda, sed ab illa et cæteris speciebus sectionis eaudice perenni distincta."

Specimina ex regione subalpina gaudent caulibus longioribus tenuioribus subprostratis, foliis majoribus minus villosis viridibus, caulinis latioribus inferioribus basi cordatis.

Hiæc stirps habitu et foliorum forma *Stachyem circinatam* aliquantulum refert. A charactere generico paululum recedit styli lobo inferiore minus dilatato superiorem vix amplectente.

SIDERITIS COSSONIANA, nob., = *S. Balansæ* (Coss. in Bull. Soc. Bot. Fr. xx. 256, non Boiss. Diagn. Pl. Or. ser. 2, iv. 35).

Dum nomen inventoris ad speciem alienam jam relatam est hanc stirpem novam cognomine eel. auctoris, floræ nostræ optime merentis, decorare juvat. Descriptionem auctoris citare malui:—

"Planta annua, pilis patentibus molliter villosa, a basi ramosa. Caules 12–15 centim. longi, tetragoni angulis præminentibus obtusis, laterales diffusi vel ascendentes, centralis erectus, simplices vel ramis paucis donati. Folia crenata crenis muticis vel in foliis superioribus crena terminali brevissime cuspidata, inferiora breviter petiolata ovata basi cuneata, floralia subconformia minora subsessilia calycibus sublongiora. Verticillastri 4–6-flori, inferiores sæpius remoti, superiores in spicam foliatam subapproximati. Calyx breviter pedicellatus, demum patenti-subdeflexus, patule hispidus, campanulatus, tenuiter 10-nerviis, dentibus omnibus subæquilatis subæquilongisque, ovato-lanceolatis apice in cuspidem longam subulatam attenuatis, tubi longitudine dimidia paulo longioribus, demum suberectis. Corolla 6–7 millim. longa, albida, sparse pilosa, tubo cylindrico, subrecto, calycis tubum excedente, dentes subæquante, intus obsolete piloso-annulato, labio superiore subascendente planiusculo integro, labio inferiore longiore patente trilobo lobo medio latiore obovato integro. Stamina 4, ad tertiam partem inferiorem tubi corollini inserta, tubo inclusa, superiora breviora antherarum lobis divaricatis, inferiora longiora antheris minutis cassis. Stylus inclusus, apice bifidus, lobo superiore tereti truncato, inferiore dilatato basi superiorem amplectente. Nueulæ fusæ, obovatæ apice rotundatæ, glabræ, sub lente punctato-tuberculatæ."

Mar. occid.—In arenosis haud procul Mogador et juxta Aïn el Hadjar

copiose crescentem floriferam et fructiferam post imbres mensis Maji legimus die 3 Junii!

Mar. merid.—In collinis prov. Haha ad 750 m. primus legit Balansa. In eadem provincia legimus circ. 200 m. ! et in saxosis prope Seksaoua ! Prov. Shedma (*Maw spec.*) !

A proxima *S. lanata* satis superque distincta foliis minoribus, floralibus distincte petiolatis, calycis dentibus subæquilongis demum erectis, nec inæqualibus reflexis, colore corollæ, stylo profundius bilobo, nuculis fusco-rufis nec nigrescentibus. A cæteris sectionis *Hesiodia*, Benth., plurimis notis longius distat.

MARRUBIUM (*L. Gen.* 721).

M. ALYSSON (*L., Benth. in DC. Pr.* xii. 448).

Mar. merid.—Mesfoua ! Seksaoua (*J. D. H.*) !

Ar. Geog. Per regionem Mediterraneam calidiorem dispersa. Deest in Græcia, in Italia media et superiore, in Gallia australia, in Lusitania et in Macaronesia.

M. VULGARE (*L., Benth. in DC. Pr.* xii. 453).

Mar. occid.—Agadir (*Coll. Indig.* 1871) !

Mar. merid.—Prov. Shedma ! In ruderatis urbis Marocco !

— —, var. LANATUM (*Benth. l. c.*), = M. Apulum (*Ten. Fl. Nap.* tab. 154).

Mar. occid.—Casa Blanca !

Mar. merid.—Ourika (*R. et Fr. spec. in Herb. Kew.*) !

Ar. Geog. Species cosmopolita per regiones temperatas et subtropicas orbis veteris neenon utriusque Americæ late diffusa. Non deest in Macaronesia.

M. ECHINATUM (*Ball in Journ. Bot.* 1875, 175).

Planta biennis (?) elata, 3–4-pedalis ; caules erecti, ramosi, hirti, quadranguli, sulcati ; folia petiolata, orbiculari-ovata, inæqualiter excisodentata, inferiora basi cordata, floralia basi cuneata, suprema subsessilia ; verticillastri remoti-20-30 flori, foliis breviores ; calyx infundibuliformis, profunde 10-sulcatus, dentibus 5 (rarins sex) instructus ; dentes calycini 3 superiores breves recti, apice mucronati, inferiores 2 (rarins 3) tubo calycis longiores, recti, apice hamati ; bractea calyce breviores longe villosæ ; corollæ labio superiore erecto bilobo, inferiore trilobo, lobo medio latiore rotundato emarginato ; labio utroque extus et intus velutino ; stamina brevia ; stylus brevissimus, bifidus ; nuculæ (immaturæ) ovoidæ læves.

Mar. merid.—Legimus in sepibus prope Amsmiz ! et specimen mancum attulerunt ell. Rein et Fritsch prope Sektana lectum !

Corollæ tubus intus nudus. Folia pro genere tenuia, superne viridia, subtus pallida nervosa non rugosa, utrinque breviter pubescentia. Species anomala, a charactere generico recedens, sed nullo pacto divellenda.

BETONICA (*L. Gen.* 718).

Genus naturale, charactere tecnico ægre definiendum, axe primario non evoluto, ramis floriferis axillaribus insigne.

B. OFFICINALIS (*L., Benth. in DC. Pr.* xii. 460), var. *ALGERIENSIS*, = *B. algeriensis* (*De Noé in Bull. Soc. Bot. Fr.* ii. 582).

Mar. sept.—Tanger (*Webb spec.*)!

Ar. Geog. Species per Europam omnem a Succia usque regionem Mediterraneam diffusa, rarior tamen in calidioribus. In Asia media et in prov. Caucasicis non deest. Var. *algeriensis* in Africa boreali et in Hispania Bætica. Legi prope San Roque!

B. algeriensis est forma australis sp. Linnæani, floribus minoribus et calycibus paullo majoribus diversa, sed nulla nota stabili definita.

STACHYS (*L. Gen.* 719).

S. GERMANICA (*L., Benth. in DC. Pr.* xii. 464).

Mar. sept.—Tanger (*Salzmann, sec. Benth. l. c.*). In monte Beni Hosmar pr. Tetuan (*Webb Herb.*)!

Specimina Schousbœana ex "Imperio Maroccano" a Benth. l. c. citata.

— — —, var. *CRETICA*, = *S. cretica* (*L., Willk. et Lge. Fl. Hisp.* ii. 440), = *S. cinerea* (*Salzm. Exsicc.*), = *S. heraclea* (*Webb Herb., non All.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)!

Differt a typo indumento molliter tomentoso nec lanato, calycibus villosis nec lanatis. Planta ulterius observanda inter *S. germanicam* et *S. heracleam*, All., pluribus notis media, haud sine dubio pro varietate enumerata.

Ar. Geog. Species per Europam mediam et regionem Mediterraneam fere omnem (exclusa Macaronesia) dispersa.

S. DURANDIANA (*Coss. in Bull. Soc. Bot. Fr.* xx. 256).

Descriptionem Cossonianam citare juvat.

"Perennis? tota pilis patulis aliis brevioribus glandulosis aliis elongatis dense hirsuta obscure virens. Caulis erectus, tetragonus angulis præminentibus obtusis, superne ramosus. Folia caulina petiolata, 3-4½ centim. longa, ovata vel ovato-triangularia, basi subcordata, dentata dentibus obtusiusculis vel acutiusculis, facie utraque hirsuta; floralia sessilia vel subsessilia, ovato-acuminata apice brevissime et molliter subspinescentia, dentata vel suprema ovato-oblonga integra, calycibus longiora. Verticillastri numerosi, subsexflori, inferiores tantum remotiusculi, cæteri in spicam foliatam approximati. Bracteolæ omnino obsoletæ. Calyx dense et patule hispidus, campanulato-tubulosus demum campanulatus, tenuiter 10-nervius, ore æquali, dentibus æqualibus triangularibus lanceolatisve, apice breviter spinulosus, tubi longitudine dimidia sæpius paulo brevioribus. Corolla 12-15 millim. longa, purpurea (sec. Durand), extus breviter hirta, tubo cylindrico subincurvo ad faucem haud ampliato calyce brevior ad tertiam par-

tem inferiorem *intus oblique annulato-piloso*, *labio superiore* subhorizontali subfornicato *integro vel subretuso*, *labio inferiore* multo longiore patente trilobo lobo medio maximo retuso. Stamina inferiora longiora demum ad latera faucis dejecta, antherarum lobis demum divaricatis. Nuculæ nigrescentes, obovatæ, apice rotundatæ, glabræ, læves.— Descriptio juxta specimina tria herbarii Durand nunc Bouchet."

Mar. centr.—Juxta Mequinez ab abbate Philippo Durand inventa.

"*S. Durandiana* canle, foliis, calycibusque hirsutis, verticillastris plerisque 6-floris, bracteis obsoletis, calyce laud bilabiato dentibus apice spinescentibus, corollæ purpureæ tubo incluso et intus piloso-annulato pertinet ad sectionem *Stachyotypus* (Dumort.; Benth. in DC. Prod. xii. 468). In speciminibus incompletis suppetentibus radix deficit. Si planta radice perenni gaudet, villositate, caulibus robustis superne ramosis, foliis caulinis ovato-triangularibus vel ovato-oblongis, corolla majuscula, ad gregem *Elutæ* (Benth. loc. cit. 468) referenda et juxta *S. sylvaticam*, L., collocanda. Si contra radicem annuam præbet referenda ad gregem *Agrestes* (Benth. loc. cit. 477) et juxta *S. marrubifoliam*, Viv., collocanda, a qua ut a *S. sylvatica* notis supra indicatis eximie differt."

STACHYS CIRCINATA (L'Hér. *Stirp.* i. 51, tab. 26; *Benth. in DC. Pr.* xii. 474), = *S. latifolia* (*Ait. Hort. Kew.* ii. 303), = *S. velutina* (*Willd. Enum. Suppl.* 41).

Mar. sept.—In monte Beni Hosmar pr. Tetuan (*Webb spec.*)!

Ar. Geog. Africa borealis. Hispania Bætica.

S. SAXICOLA (*Coss. in Bull. Soc. Bot. Fr.* xx. 257).

Descriptio Cossoniana ad calcem spec. a Balansa anno 1867 lectorum.

"*Perennis*, tota *densissime et mollissime niveo-lanata*. Caudex lignosus, tortuosus, caules plures emittens. *Caulis diffuso-procumbentes, obtuse tetragoni, lana copiosa oblecti*, simplices vel superne ramosi. *Folia caulina longe petiolata, ovato-suborbiculata basi late cordata*, crenata, crassinsecula, *utrinque niveo-tomentosa*; *floralia subconformia*, multo minora, sensim decrescuntia, *supremi calycibus breviora*. *Verticillastri* plures, subsexflori, *in spicam brevem densam approximati*. *Bracteolæ omnino obsoletæ*. *Calyx dense niveo-lanatus*, campanulatus, tenuiter 10-nervius, ore æquali, *dentibus æqualibus, ovatis apice abrupte in cuspidem brevem spinescentem acuminatis*, tubi longitudine dimidia brevioribus. *Corolla* (e fragmentis tantum nota) extus sparse et breviter hirta, *tubo cylindrico, subrecto, ad faucem vix ampliato*, calycem subæquante, ad mediam longitudinem intus *transversim annulato-piloso*, *labio superiore* subhorizontali subfornicato integro, *labio inferiore* multo longiore patente trilobo lobo medio maximo retuso. Stamina inferiora longiora, demum ad latera faucis dejecta, antherarum lobis demum divaricatis. Nuculæ nigrescentes, obovato-trigonæ, apice rotundatæ, glabræ, irregulariter reticulato-rugosæ."

Mar. merid.—In fissuris rupium prope Moulai Ibrahim primus detexit

Balansa. Spec. his conformia foliis utrinque albo-lanatis legimus prope Gurguri!

— —, var. *VILLOSISSIMA*, nob. Caulibus petiolisque longe villosolauatis, foliis subtus villosolauatis, in pagina superiore pilosis virescentibus.

Mar. merid.—Legimus copiose erescens in rupibus Aïn Tarsil!

Caudex crassus diametro pollicaris et ultra, ramos fragiles steriles diffusos floriferos adscendentes emittens. Corolla in spec. nostris pallide carnea gaudet tubo calycem paululum excedente labio superiore extus piloso, inferiore longiore trilobo, lobo medio emarginato, lateralibus parvis integris multo majore.

S. circinata affinis, sed facile dignoscenda foliis etiam floralibus petiolatis, verticillastris floralibus approximatis subspicatis, corolla minore, labii inferioris segmentis lateralibus parvis integris nec magnis subbilobis. Valde proxima est etiam *S. candida* (Bory et Chaub. Exped. Mor. 167, tab. 19; Benth. in DC. Pr. xii. 475). Hæc species, mihi non satis nota, differt sec. ecl. Cosson calycis dentibus lanceolatis in cuspidem attenuatis, nec ut in nostra ovatis apice abrupte in cuspidem brevem subspinescentem acuminatis. Video calycis dentes in sp. nostra et in affinibus valde variabiles, et notæ ex his desumptæ interdum fallaces sunt. Corolla tamen in *S. saxicola* longius exserta et, nisi fallor, forma labii inferioris diversa.

STACHYS MAWEANA, nob. Affinis *S. saxicolæ*, Coss., cui caudice perenni, indumento albo-lanato, caulibus adscendentibus, subsimilis; insigniter differt tamen foliis omnibus multo minoribus ovatis aentiusculis, basi late emneatis, in petiolum limbo æquilongum attenuatis crenato-dentatis, calycis minoris dentibus inæqualibus vix aut ne vix apice spinoso-mueronatis, corolla duplo majore, tubo parum exserto, labii inferioris lobo medio maximo subintegerrimo.

Mar. merid.—Frustula plantæ spontaneæ a cl. Maw inter Sektana et Frouga lecta nimis incompleta! ex radice ibidem lecta specimina florifera benev. communicavit cl. Maw!

Corolla, nisi fallor, pallide purpurea, pro grege magna.

S. ARVENSIS (L., Benth. in DC. Pr. xii. 477).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Mar. merid.—Mesfioua! Amsmiz (*J. D. H.*)! Sec. notulam MS. vidimus in convalle Amsmiz usque 2000 m.!

Ar. Geog. In temperatis totius orbis veteris late diffusa. In America tam tropicali quam extratropicali et in insula Sanctæ Helenæ verisimiliter cum seminibus introducta.

S. HIRTA (L., Benth. in DC. Pr. xii. 481).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Tetuan (1851 et 1871)!

Mar. occid.—In prov. Haha hand proenl Mogador!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

STACHYS ARENARIA (*Vahl, Symb.* ii. 64; *Desf. Fl. Atl.* ii. 21, tab. 126; *De Noë in Webb, Phyt. Canar.* iii. 98; *Benth. in DC. Pr.* xii. 485, ex parte).

Mar. sept.—Tetuan ad ostium fluminis (1851)! In montosis prope Tetuan (*Webb spec.*)!

Mar. occid.—Frequens in regione litorali, imprimis in arenosis, sed non deest in saxosis submontanis. Casa Blanca! Mazagan! Saffi! Mogador! In monte Djebel Hadid! et juxta Aïn el Hadjar!

Mar. merid.—Occurrit, sed rarius, in terris interioribus. Sektana! Seksaoua!

Ar. Geog. Africa borealis. Sicilia. Italia meridionalis. Ins. Canariensium dubia civis. Deest in peninsula Iberica.

Cel. Bentham loc. cit. *S. ibericam*, M. B., et *S. scordifoliam*, Willd., cum *S. arenaria* conjunxit, dum el. De Noë has omnes pro speciebus probe distinctis l. c. vindicavit. *S. scordifolia* est planta hortensis mihi non satis nota. *S. Iberica*, planta orientalis, est sane *S. arenariæ* valde proxima, differt tamen florum verticillis in spicam approximatis, dentibus calycinis inæqualibus, corollæ tubo annulo piloso destituto, labio superiore bifido, et, monente De Noë, cotyledonibus rotundatis nec ellipticis. In multis specim. *S. areuariæ* a me exploratis vidi semper dentes calycinos subæquales, corollæ tubum annulo piloso fere completo instructum, et labium superius integrum.

S. areuaria variat herba plus minusve pilosa, habet folia caulina late linear-oblonga pauciserrata plerumque obtusa, floralia integerrima apice cuspidato-mucronata. Nuculæ badiæ lævissimæ. Calyx intus glaberrimus nervis coloratis 18-22 (nec exacte 20) notatus.

S. mollis (*Willd. Herb.* no. 10877, ex *Benth. in DC. Pr.* xii. 475).

Mar. sept.—Prope Tetuan mense Junio 1851 legi florentem! eandem plantam nondum florentem legimus in monte Beni Hosmar mense Aprili 1871!

Ar. Geog. Species Marocana.

Descriptioni Benthamianæ l. c. adamussim convenit, nisi verticillastris supremis interdum (nec semper) approximatis. Procul dubio in grege *S. maritimæ* et *S. arenariæ* collocanda, huic proxima sed multis notis gravioribus diversa. Calyx multo brevior campanulatus dentibus brevibus apice mucronatis inæqualibus præditus; corollæ annulo piloso interno incompleto—scil. nonnisi inter insertionem staminum duorum inferiorum—labiis subæquilongis, superiore utrinque et præsertim extus villosa profunde bifido. Nuculæ oculo fortiter armato minute granulatæ. In *S. molli* herba tota albo-lanata; folia breviter crassa subrugosa, floralia integra apice calloso nec ut in *S. areuaria* mucronato-spinosa instructa. Willdenovius habuit hanc speciem inter plantas Maroccanas a Broussonnet lectas et (ut videtur) signo interrogationis “Mogador” pro loco natali adjecit.

LAMIUM (*L. Gen.* 716).

L. AMPLEXICAULE (*L., Benth. in DC. Pr.* xii. 508).

Mar. sept.—In monte Beni Hosmar prope Tetuan (*J. D. H.*)!

Mar. merid.—In Atlante Majore a 1500 m. usque 2900 m. In convalle Aït Mesan! et in ascensu ad jugum Tagherot! In monte Djebel Tezali!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). Europa fere omnis (exclusa arctica). Montes Himalaya. In America boreali forsitan advena, hodie spontaneum.

Inter spec. nostra Atlantica adest lusus herba tota (etiam in calycibus) glaberrima, calycis dentibus rigidioribus coloratis.

L. ALBUM (*L., Benth. in DC. Pr.* xii. 510), var. *PARVIFLORA*?

Mar. merid.—In convalle Aït Mesan circ. 1400 m.! Prope Amsmiz ad 1000 m. s. m.!

Planta mihi subdubia, a *L. albo* nisi corollis minoribus vix differt, sed facie aliquid diversa.

Ar. Geog. *L. album*, in temperatis frigidioribus Europæ et Asiæ mediæ frequens, rarius occurrit in montibus regionis Mediterraneæ. Deest in Hispania australi et in insulis Italicis; in Africa boreali a Fontanesio indicatum, sed a nemini recentiorum detectum.

L. FLEXUOSUM (*Ten. Fl. Nap.* ii. 19, tab. 52; *Benth. in DC. Pr.* xii. 511).

Mar. sept.—In monte Beni Hosmar prope Tetuan (*Webb Herb.*)! In eodem monte spec. mox floribundum legit *J. D. H.*!

Ar. Geog. Regio Mediterranea occidentalis a Sicilia usque fretum Herculis.

BALLOTA (*L. Gen.* 720).

B. HIRSUTA (*Benth. Lab.* 595, et in *DC. Pr.* xii. 518), = *B. orbicularis* (*Lag. Hort. R. Matrit.*), = *B. africana* (*Colm. Ap. Fl. D. Cast.* 121, non *Benth.*), = *Beringeria hispanica* (*Neck. Élém.* 312), = *Marrubium hispanicum* (*Cavan. et Desf., non L. Sp. Pl.* 816), = *Marrubium crispum* (*Desv. in Lam. Dict.* iii. 719, non *L.*).

Mar. occid.—Prope Mogador (*Broussonet ex Benth. l. c.*). Prope Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Satis frequens in dumetis et sepibus regionis inferioris. Prov. Haha! Prov. Shedma pluribus locis! Mesfioua! Mzouda! Sektana (*R. et Fr. spec. in Herb. Kew.*)!

Planta satis variabilis—folia sæpius late ovata utrinque rugosa superne subeineria, interdum latiora suborbicularia sat profunde cordata superne viridia; in spec. nostris folia omnia in pagina inferiore sordide albo-lanata eximie rugosa, floralia verticillos longitudine æquantia vel superantia.

Ar. Geog. Africa boreali-occidentalis. Hispania centralis et australis.

Calyx in spec. nostris descript. Benthamianæ non omnino conformis; limbus late expansus in lobos quinque ovatos acutos divisus, lobo singulo mucronato-denticulato, sæpius (sed non semper) dente majusculo inter lobos adjecto.

Specimen nostrum prope Mzouda lectum ad var. *hispidam* (Benth. l. c.) accedit, sed notæ differentiales indicatæ nimis variabiles.

BALLOTA NIGRA (*L., Benth. in DC. Pr. xii. 520*), lusus singularis vel forsân nova species?, = *B. bracteosa, nob. in Herb.*

Planta perennis? 2-3-pedalis, ramosa, ramis adscendentibus; folia late ovata, basi sæpius cordata, acuta, superiora longiuscule acuminata, grosse crenato-serrata, utrinque tomentoso-puberula, subtus cinerea; cymulæ laxiusculæ; flores bracteis obverse lanceolatis angustis integris, vel sæpius expansis foliaceis dentatis suffulti; calycis quinque-dentati limbo e basi infundibuliformi late expanso, in dentes integros acutos non mucronatos diviso; corolla parva calycem parum superans; nuculæ aterrimæ lucentes, nec ut in *B. nigra* atro-virides opacæ.

Mar. occid.—Legi floriferam et fructiferam die 5 Junii juxta urbem Mogador! Ex spec. immaturo in *Herb. Kew.* hæc est *B. nigra?* (*Lowe Cat.*).

Characteres enumerati ad speciem novam definiendam idonei, hanc tamen esse lusum singularem *B. nigræ*, sp. valde ludibundæ, valde suspicor. *Ar. Geog.* *B. nigra* per Europam mediam et regionem fere omnem Mediterraneam (exclusa Macaronesia) diffusa est.

PHLOMIS (*L. Gen. 723*).

P. CRINITA (*Cavan. Icon. iii. 25, tab. 247; Benth. in DC. Pr. xii. 538*), = *P. biloba (Desf. Fl. Atl. ii. 25, tab. 127)*, = *P. mauritanica (Munby, Fl. Alg. 60, tab. 3)*; sphalmate a Benth. l. c. citatur Moench, *Pl. Alg.*

Mar. sept.—Prope Tetuan (1851)! In monte Djebel Dersa prope Tetuan (*Webb Herb.*)!

Mar. merid.—In aridis pr. Aïn Onmast! Inter Tasseremont et Ourika (*G. Maw spec.*)! Sektana (*R. et Fr. spec. in Herb. Kew.*)!

Nomen Fontanesianum erroneum (ob corollæ labium superius in spec. suo casu aliquo fissum) et prætermittendum.

Ar. Geog. Africa boreali-occidentalis. Hispania.

P. HERBA VENTI (*L., Benth. in DC. Pr. xii. 542; Bot. Mag. li. tab. 2449*).

Mar. sept.—Tanger (*Webb spec.*)!

Mar. merid.—Inter Ourika et Reraya! Reraya! Sektana!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

PRASIUM (*L. Gen. 737*).

P. MAJUS (*L., Benth. in DC. Pr. xii. 556*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)! In monte Djebel Kebir! Inter Tanger et Tetuan! Tetuan (1851)!

Mar. occid.—Casa Blanca (*J. D. H.*)! Mogador (*Lowe Cat.*). Agadir (*Coll. Indig. 1871*)!

Mar. merid.—Vidimus ad radices Atlantis, sed specimina desunt! (spec. prope Aouina, prov. Shedma lecta benev. comm. Cosson)!

Ar. Geog. Regio fere omnis Mediterranea calidior. Deest in Gallia in Italia superiore et in insulis Canariensibus.

TEUCRIUM (*L. Gen.* 706).

T. FRUTICANS (*L., Benth. in DC. Pr.* xii. 575).

Mar. sept.—Tanger (*Salzmann, sec. Benth. l. c.*). In monte Beni Hosmar prope Tetuan!

Mar. occid.—In prov. Haha prope Mogador!

Mar. merid.—Haud infrequens ad radices Atlantis Majoris. Tassereimout! Ourika (*J. D. H.*)! Reraya! Sektana (*J. D. H.*)! Keira (*Balansa spec.*)!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia). In Gallia rarissimum.

T. BRACTEATUM (*Desf. Fl. Atl.* ii. 7, tab. 120, minus bona; *Benth. in DC. Pr.* xii. 582).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Secus viam ad Buhana (*spec. ex Herb. Schousb. comm. Cosson*)!

Ar. Geog. Africa boreali-occidentalis. Fertur in Hispania pr. Tarifa, sed a nemini recentiorum inventum.

T. COLLINUM (*Coss. in Bull. Soc. Bot. Fr.* xx. 258).

Sequitur descriptio Cossoniana:—

“Planta perennis, inferne frutescens, a basi ramosa, caudice lignoso in radicem fusiformem abeunte, a basi ramosa, tota velutino-villosa. Caules diffuso-adscendentes, vix tetragoni, subsimplices, sæpius ramulos nonnullos axillares abbreviatis emittentes, pilis patentibus mollibus longe villosi. Folia subapproximata, ovata basi cuneata in petiolum attenuata, crenata, nervis supra impressis infra præminentibus subbullata, supra villosa, subtus canescenti-subtomentosa, floralia subconformia in verticillos 4–6-folios disposita superiora oblonga calycees subæquantia. Verticillastri plures, distincti, subapproximati, 4–6-flori, superiores in spicam simplicem teretem foliatam conferti. Calyx breviter pedicellatus, late campanulatus tubo inferne valde gibbo, ore barbato, bilabiatus, dentibus labii superioris multo latioribus supremo transversim latiore subtruncato abrupte breviterque cuspidato, lateralibus ovatis apice rotundatis, labii inferioris dentibus lanceolatis in subulam brevem sensim attenuatis. Corolla purpurascens?, extus villosa, torsione tubi resupinata, labii superioris lobis cum labio inferiore in labium unicum 5-lobum utrinque lobis 2 ovato-rotundatis et lobo terminali suborbiculato multo majore concavo donatum confluentibus. Genitalia arcuata, longe exserta. Nuculæ fusæ, glabræ, grosse reticulato-rugosæ.”

Mar. occid.—Legimus in monte Djebel Hadid ubi hanc stirpem primus detexit Balansa! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Frequens in dumetis regionis inferioris. Legimus ac vidimus pluribus locis in provinciis Haha! et Shedma! et ad radices Atlantis Majoris. Amsmiz! Seksaoua! Mtouga (*J. D. H.*)! Prope Keira legit Balansa sec. Cosson l. c.

Monente cel. auctore, “*T. collinum* habitu, foliis floralibus verticillatis, calycis fabrica juxta *T. bracteatum*, Desf., ad sectionem *Stachyobotrys* a cl. Benthām (in DC. Prodr. xii. 581) refertum, collocandum.”

A *T. bracteato* satis superque differt foliis inferioribus basi semper cuneatis in petiolum sensim attenuatis, nec subcordatis, crenis foliorum inæqualibus acutioribus, floralibus brevissime nec longiuscule petiolatis fere omnibus serratis, nec integerrimis, calyce breviter nec longiuscule pedicellato, dente supremo latiore subtruncato breviter cuspidato nec longiuscule acuminato-cuspidato.

Planta satis variabilis, interdum erecta 2-pedalis et ultra, interdum humilis diffusa.

TEUCRIUM SCORODONIA (L., Benth. in DC. Pr. xii. 584).

Mar. sept.—In monte Djebel Kebir!

(In Imp. Maroccano legit Broussonnet sec. Desfontaines, Fl. Atl. 5.).

Ar. Geog. Europa fere omnis, imprimis occidentalis. Marocco. Madera. Specimina nostra ab Europæis nullo modo diversa.

T. PSEUDO-SCORODONIA (Desf. Fl. Atl. ii. 5, tab. 119; Benth. in DC. Pr. xii. 584).

Mar. sept.—Tanger (Salzm. Exsicc., spec. in Herb. Gay)!

— —, var. BÆTICUM, = *T. bæticum* (Boiss. et Reut. Pug. 98).

Mar. sept.—Tanger (Salzm. Exsicc.), sec. Boiss. et Reuter l. c. In monte Djebel Kebir!

Legi etiam in collibus juxta S. Roque.

Hæc est nil nisi var. levis *T. Pseudo-scorodonie*, foliis paullo minoribus et racemo tenuiter glanduloso-puberulo.

Ar. Geog. Species in Africa boreali-occidentali, in Italia meridionali et in Sicilia. Varietas in Hispania Bætica et in ditione Tingitana.

T. Pseudo-scorodonia, valde proximum, et *T. Scorodonie* notis non gravibus dignoscendum, forsā melius pro subspecie habendum.

T. RESUPINATUM (Desf. Fl. Atl. ii. 4, tab. 117; Benth. in DC. Pr. xii. 576).

Mar. sept.—Tanger (Salzm. Exsicc.)! (Webb spec.)! legi in herbosis uliginosis juxta pontem Romanum! et in clivo montis Djebel Kebir!

Ar. Geog. Africa boreali-occidentalis. Hispania.

T. decipiens (Coss. in Bull. Soc. Bot. Fr. xx. 258).

Sequitur descriptio Cossoniana:—“Planta annua, sæpius pusilla, villosa.

Caulis erectus, tetragonus, a basi divaricato-ramosus, pilis patentibus villosus. Folia præsertim superiora approximata, oblonga in petiolum attenuata, facie utraque viridia, præsertim infra villosa nervis vix prominulis, inferiora superne grosse paucidentata, floralia superiora subconformia angustiora integra flores excedentia. Verticillastri plurimi distincti, biflori, secundi, superiores in spicam foliatam foliis floralibus superioribus comosam approximati. Calyx pedicellatus, in pedicello nutanti-subdeflexus, campanulato-tubulosus tubo inferne gibbo, ore

barbato, obsolete bilabiatus, *dentibus subconformibus lanceolatis brevissime aristatis, supremo patenti-ascendente, cæteris erectis. Corolla* purpurascens, extus villosa, tubo haud torto *non resupinata* nempe labio inferiore deorsum spectante, labii superioris lobis cum labio inferiore in labium unicum 5-lobum utrinque lobis duobus et lobo terminali donatum confluentibus, lobis lateralibus superioribus triangulari-lanceolatis, inferioribus oblongis, terminali obovato-cuneato multo majore. Genitalia arcuata, longe exserta. Nuculæ fusco-nigrescentes, glabræ, reticulatæ."

Mar. occid.—In monte Djebel Hadid haud procul Mogador, ex spec. in herb nostro, primus detexit Balansa !

Mar. merid.—In regione inferiore hinc inde rarius occurrit. In provincia Shedma bis lectum ! Inter Ourika et Reraya (*Maw legit*) ! Sektana (*R. et Fr. spec. in Herb. Kew.*) !

A *T. resupinato* cui proximum distinguitur statura pumila 2–6-pollicari, verticillastris magis approximatis, corolla non resupinata. Ut mihi videtur, pro subspecie enumerandum. Dentes calycini in spec. ex Djebel Hadid quam in *T. resupinato* breviores, in spec. a Sektana æquilongi. Herba sæpius pube brevi tenui adspersa, interdum villosa.

T. mauritanicum (De Noë in Bull. Soc. Bot. Fr. ii. 585), mihi ex descriptione tantum nota, videtur *T. decipiente* valde proximum, nisi hoc foret idem ac *T. resupinatum* (Balansa, Pl. d'Algérie, No. 438, non Desf.), = *T. Balansæ* (nobis MSS. in herb. olim), a *T. resupinato*, ut mihi videtur, et certe a *T. decipiente*, diversum.

TEUCRIUM RUPESTRE (*Coss. in Bull. Soc. Bot. Fr. xx. 259*).

Descriptionem Cossonianam attuli :—" *Suffrutex* candice lignoso inerasato, multicaulis. Caules diffusi, subtetragoni, inferne et sæpius superne ramosi, præsertim in parte superiore pilis patentibus mollibus pubescentes vel demum glabrescentes. *Folia* caulina subdistantia, ovata vel ovato-subtriangularia obtusa basi truncata vel subcordata breviter petiolata, crenata, nervis supra impressis infra præminentibus bullata, supra glabrescentia, infra villosa-subtomentella, *floralia* infima subconformia, cætera oblonga integra calyces subæquantia. *Verticillastri* plures, 2-flori, in spicas apice caulis et ramorum conferti. *Calyx* pedicellatus, in pedicello nutans, campanulatus tubo inferne subgibbo, ore non barbato, bilabiatus, *dentibus labii superiori latioribus ovatis* supremo breviter acuminato, lateralibus brevioribus obtusis, *labii inferioris lanceolatis* apice vix mucronatis. *Corolla* lutescens, extus pubescens, tubo haud torto non resupinata, *labii superioris lobis cum labio inferiore in labium unicum 5-lobum* nempe utrinque lobis duobus superioribus lanceolatis elongatis, inferioribus oblongis et lobo terminali majore suborbiculato basi contracto concavo donatum confluentibus. Genitalia arcuata, longe exserta. Nuculæ (immaturæ tantum notæ) glabræ, rugosæ."

Mar. merid.—Prope Keira detexit Balansa anno 1867. Legimus in rupibus supra Seksoua copiose erescens ! In rupibus Aïn Tarsil !

Hæc stirps, ut jam monuit cel. Cosson, quoad affinitatem dubia, pluribus notis sectioni *Stachyobotrys* (Benth. in DC. Prod.) accedit, sed habitu et corollæ structura potius in § *Chamædrys* (Benth. l. c.) et juxta *T. fragile*, Boiss., collocanda mihi videtur.

TEUCRIUM GRANATENSE (Boiss. et Rent. Pug. 99), = *T. pyrenaicum*, var. *granatense* (Boiss. Elench. 78; Benth. in DC. Pr. xii. 591), var. *ATLANTICUM*, nobis.

Differt herba tota sordide albo-villosa; dentibus calycinis acutis, nec ut in typo Boissieriano acuminatis, cum foliis floralibus coloratis, corolla albida tubo quam in typo brevior.

Mar. merid.—In convalle Amsmiz circ. 1400 m.!

T. granatense, Boiss., hucusque tantum in Hispania australi detectum, locum medium tenet inter species facie prorsus diversas—*T. pyrenaicum*, L., et *T. cinereum*, Boiss.; et varietas nostra hinc et *T. granatensi* intermedia est. An igitur tres species enumerandæ, aut potius *T. cinereum* et *T. granatense* ad subspecies reducenda?

T. POLIUM (L., Benth. in DC. Pr. xii. 592), var. *VULGARE* (Benth. l. c.).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. occid.—Mogador (*Lowe Cat.*)? In prov. Italia prope Mogador!
—— —, var. *GNAPHALODES* (Benth. l. c.), = *T. gnaphalodes* (*Vahl, Symb.* i. 41).

Mar. occid.—Prope Mogador (*J. D. H.*)!

Mar. merid.—Sheshaoua! Misra ben Kara!

—— —, var. *ANGUSTIFOLIUM* (Benth. l. c.), = *T. capitatum* (*L. Sp. Pl.* 792; *Cavan. Icon.* ii. 17, tab. 119).

Floribus sæpius rosco-purpurascentibus in capitulis parvis tomentosis congestis, foliis oblongis vel oblongo-linearibus.

Mar. occid.—In monte Djebel Hadid laud procul Mogador (*R. et Fr. spec. in Herb. Kew.*)! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Forma in planitie et in regione inferiore Atlantis Majoris vulgatissima. Prov. Shedma! Tasscremout! Ourika! Reraya! In convalle Aït Mesan usque 1500 m.! Amsmiz! et alibi visum.

Hinc, nisi fallor, referenda est var. *purpurascens* (Benth. l. c.).

—— —, var. *ATLANTICUM*, nob.

Forma insignis forsân melius cognita pro specie diversa vindicanda, ramis prostratis floriferis indivisis, foliis profunde inciso-crenatis vel pinnatifidis, lobis oblongis apice rotundatis, breviter pilosis, obscure virentibus, floribus in spicam brevissimam confertis, corollis paullo majoribus purpurascentibus.

Mar. merid.—In convalle Aït Mesan circ. 1400 m.!

Ar. Geog. *T. Polium* per regionem omnem Mediterraneam (exclusa Macaronesia) late diffusum est.

Varietates emuneratæ (excepta forsân *atlantica*) nullis notis stabilibus definiendæ, etsi sæpe facie valde diversæ.

TEUCRIUM BULLATUM (Coss. in Bull. Soc. Bot. Fr. xx. 260).

Sequitur descriptio Cossoniana :—" *Suffrutex* caudice lignoso incrassato multicaulis. Caules diffusi vel ascendentes, vix tetragoni, inferne et sæpe superne ramosi pilis patentibus dense cinereo-villosi. *Folia sessilia*, caulina approximata, *oblongo-lanceolata* obtusa, a basi crenato, nervis supra impressis infra præminentibus *bullata*, marginibus revoluta, utrinque cinereo-villosa, floralia linearilanceolata integra calycibus paulo longiora spicas apice comantia. *Verticillastri in spicas oblongas* teretes apice caulis et ramorum *dense congesti*. *Calyx subsessilis*, erectus, *tubuloso-campanulatus*, ore non barbato, vix bilabiatus, *dentibus subæquilongis apice subulatis*, *labii superioris vix latioribus ovato-lanceolatis*, *labii inferioris lanceolatis*. *Corolla ochroleuca*, extus pubescens, tubo haud torto non resupinata, *labii superioris lobis amplis oblongis* divergentibus *subhorizontalibus genitalia excedentibus*, *labii inferioris trilobi lobis lateralibus oblongo-lanceolatis* margine inflexo plica superne ad basim lobulo auctis, lobo terminali majore suborbiculato basi contracto concavo. Genitalia areolata, lobis labii superioris breviora. Nuculæ nigrae, glabrae, rugosae."

Mar. merid.—In monte Aït-Ougourt juxta Keïra, ad austro-occidentem urbis Maroc, ad 1300 metr, legit Balansa anno 1867. Legimus in monte supra Seksaoua!

Monente cel. auctore :—" Caulibus suffrutescentibus, verticillastri in spicas oblongas teretes dense congestis, calyce tubuloso-campanulato, dentibus calycinis subæquilongis superioribus vix latioribus, nunculis rugosis ad sectionem *Polium* (Benth. in DC. Prodr. xii. 590) juxta *T. cinereum*, Boiss., colloecandum, sed abunde differt foliis sessilibus, dentibus calycinis longis, etc. Affine est *T. compacto* (Clement. in Lag. Nov. Gen. et Sp. 17; Benth. in DC. Prodr. xii. 586), a sectione *Scordium* (Benth. loc. cit. 585) removendo et quoque ad sectionem *Polium* referendo, sed eximie distinctum foliis sessilibus a basi crenatis, non petiolatis inferne cuneatis integris, dentibus calycinis longioribus, etc."

Radices lignosae in fissuris rupium aridissimarum serpentes caules plurimos 4-8 pollicares diffusos prostratos emittunt. Tota planta (in spec. nostris) pilis patentibus rigidiusculis cinereo-hirta; calyces tamen nisi dentibus margine ciliatis glabri. Species distinctissima juxta *T. compactum* et *T. verticillatum* (Cavan. Icon. ii. 77, tab. 198; Benth. in DC. Pr. xii. 593), ut mihi videtur, colloecanda.

AJUGA (L. Gen. 705).

A. IVA (L., sub *Teucro*; Schreb. *Unilab.* 25; Benth. in DC. Pr. xii. 600).

Mar. sept.—Tanger (*Webb Herb.*)! Cap. Spartel! Tetuan (1851)! Inter Tetuan et Ceuta!

Mar. occid.—Casa Blanca (*J. D. H.*)! Mogador! (*Lowe Cat.*). In monte Djebel Hadid haud procul Mogador (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—Frequens in arvis et saxosis e regione inferiore usque in

subalpinas Atlantis Majoris. In prov. Haha! Shedma, in aridis pr. Aït Oumast! Sheshaoua! Amsmiz! Seksaoua! In convalle Aït Mesan usque in ascensum jugi Tagherot 1400–2700 m. s. m.! In monte Djebel Tezah circ. 2000 m.!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

Corollæ indunt purpureæ, luteæ et rarius albidæ. Forma corollis luteis (*A. pseudo-Iva*, DC.) cum cæteris crescens nequaquam pro varietate habenda.

PLANTAGINEÆ (*R. Br. Prod.* 423).

PLANTAGO (*L. Gen.* 142).

P. MAJOR (*L., Dene. in DC. Pr.* xiii. pars 1, 694).

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—Sheshaoua!

Ar. Geog. In orbis veteris temperatis et calidioribus fere ubique diffusa.

In utraque America hodie spontanea, an introducta?

Planta Mogadorensis a b. Lowe ad var. *intermediam* (*Dene. l. c.*) relata; varietates tamen plurimæ descriptæ, formis intermediis inter se conjunctæ, vix certe dignoscendæ; alteræ potius lusus quam veræ varietates.

P. ALBICANS (*L., Dene. in DC. Pr.* xiii. pars 1, 705; *Schousb. Gew. Marok.* 49).

Mar. occid.—Casa Blanca! Mogador in insula!

Mar. merid.—Prov. Haha (ubi frequens)! (*Schsb. l. c.*). Prov. Shedma, pluries visa et lecta!

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

Folia et radices fere certissime ejusdem sp. prope Agadir legerunt coll. indigeni, 1871. Hæc et spec. nonnulla ex prov. Haha ad var. *latifoliam* (Willk. et Lge. *Fl. Hisp.* ii. 354) referenda videntur. Folia, in spec. nostris latitudine valde variabilia, video semper mucrone nudo cartilagineo apiculata.

— —, var. (vel forsán subspecies) *HUMILIS*, *nob.*

Folia breviora, pilosa, vix villosa, in pagina superiore cinereo-virentia nervosa; scapi breves 2–4-pollicares; spica brevior, oblongo-ovata, floribus 1–2 ad basin remotioribus. Facie potius *P. decumbenti* similis, sed calyx et fructus *P. albicantis*.

Mar. merid.—Legimus in convalle Amsmiz circ. 1500 m.!

P. DECUMBENS (*Forsk. Fl. Æg.-Arab.* 30).

Mar. merid.—In prov. Shedma in via qua itur versus Mogador (*G. Maw spec. in Herb. Kew.*)! Prope Misra ben Kara (spec. unicum defloratum)!

P. decumbens inter plantas Mogadorenses in Lowe Cat. enumeratur. Specimen tamen a beato Lowe ex Mogador in herb. Hooker conservatum, flore et fructu destitutum, mihi videtur nimis incertum, et forsán ad formam depauperatam *P. albicantis* referendum.

Ar. Geog. *P. decumbens* ex India boreali-occidentali per regiones aridas calidiores Asiæ, Mediterraneæ et Africæ borealis, usque in Imperium Maroccanum meridionale et in insulas Canarienses diffusa est. Species proxima et nimis affinis (*P. ovata*) in Hispania australi obvia est.

P. ovata (Forsk. Fl. Æg.-Arab. 31; Dene. in DC. Pr. xiii. pars 1, 706), nullis notis stabilibus a *P. decumbente* dignoscenda, mihi videtur tantum hujus varietas. Specimina Canariensia *P. decumbentis*, quibus nostra omnino conformia, intermedia sunt.

PLANTAGO LAGOPUS (L., Dene. in DC. Pr. xiii. pars 1, 716).

Mar. sept.—Tanger! Inter Tanger et Tetuan!

Mar. occid.—Mazagan! Mogador (satis frequens)! (*Lowe Cat.*). In provincia Haha prope Mogador! Aïn el Hadjar haud procul Mogador!

— —, var. LUSITANICA, = *P. lusitanica* (*Desf. Fl. Atl.* i. 135; Dene. in DC. Pr. l. c.; *Schousb. Gew. Marok.* 48).

Mar. sept.—Tanger (*Schsb. l. c.*).

Mar. occid.—Casa Blanca!

Mar. merid.—Mzouda!

Aspectu valde diversa, præsertim si specimina selecta in herbario vidimus.

Hæc est typo multo major, capitulis crassis cylindricis, caule basin versus elongato, adeo ut foliorum rosula sæpe a collo radicis remota; sed occurrunt formæ intermediæ, ut olim monuit b. Schousboe l. c., et hæc absque dubio sicut varietas enumeranda.

Ar. Geog. *P. Lagopus* per regionem Mediterraneam fere omnem (inclusa Macaronesia) diffusa est, prediligit regiones sublitorales et fit rarior in Oriente. Var. *lusitanica* occurrit hinc inde in zona calidiore.

P. AMPLEXICAULIS (*Cavan. Icon.* ii. 22, tab. 125; Dene. in DC. Pr. xiii. pars 1, 719; *Schousb. Gew. Marok.* 51).

Mar. merid.—Videtur satis rara in regione calidiore. Prope Seksaoua semel lecta! Schousboe non vidit sed semina habuit ex prov. meridionalibus.

Ar. Geog. Per Africam borealem, Hispaniam austro-orientalem, Græciam et Orientem dispersa, nullibi frequens. In insulis Canariensibus etiam obvia.

P. SERRARIA (L., Dene. in DC. Pr. xiii. pars 1, 731; *Schousb. Gew. Marok.* 50).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! “In ditione Tingitana rara” (*Schsb. l. c.*). Vidimus ac legimus copiose crescentem haud procul Cap. Spartel! et inter Tanger et Tetuan!

Mar. occid.—Prope Casa Blanca visa!

Ar. Geog. Regio Mediterranea occidentalis a Græcia usque insulas Canarienses. Deest in Gallia, in Hispania boreali et in Italia media et septentrionali.

P. MACRORHIZA (*Poir. Voy.* ii. 154; Dene. in DC. Pr. xiii. pars 1, 732; *Schousb. Gew. Marok.* 50).

Mar. sept.—Tanger (*Schsb. l. c.*). Cap. Spartel (*Schsb. l. c.*). Inter Tanger et Tetuan! Prope Tetuan!

Mar. occid.—Casa Blanca! Mazagan! Saffi! Mogador!

Mar. merid.—In prov. Shedma (*J. D. H.*)!

Ar. Geog. Per regionem Mediterraneam imprimis occidentalem dispersa.

Hispaniæ dubia civis. Deest in Macaronesia, nec vidi ex Asia Mediterranea.

Species imprimis maritima, rarius in terris a mari remotis obvia. Quoad flores et fructus *P. Coronopus* ab hac parum differt. An hujus subspecies facta perennis?

PLANTAGO CORONOPUS (L., *Dene. in DC. Pr.* xiii. pars 1, 732).

Mar. sept.—Tetuan! Nisi fallor, hanc etiam vidi prope Tanger!

Mar. occid.—Mazagan! Mogador (*Lowe Cat.*).

Mar. merid.—Mesfoua! Ansmiz! et alibi visa!

Specimina pumila tenuia forsán ad var. *simplex* (Dene. l. c.) referenda.

— —, var. MAROCCANA, nob.

Foliis brevibus latiusculis integris vel dentatis, spica villosa, bracteis flores excedentibus a typo differt.

Mar. occid.—In insula Mogador! et in arenosis prope urbem!

Mar. merid.—Juxta Mskala!

— —, var. CUPANI (Dene. l. c.), = *P. Cupani* (*Guss. Syn. Fl. Sic.* i. 199) lusus glabrescens.

Mar. merid.—Satis frequens ad radices Atlantis Majoris unde interdum in regionem subalpinam ascendit. Ourika! Aït Mesan usque in ascensum jugi Tagherot! Supra Ansmiz!

Quoad formam foliorum pinnatifidorum segmentis sæpius iterum profunde pinnatifido-dentatis falcatis, spec. nostra Siculis conformia sed folia nisi margine scabro subglaberrima.

Ar. Geog. *P. Coronopus* per regionem omnem Mediterraneam et Europam omnem (exc. arctica) late diffusa est; varietates plurimæ ægre limitandæ in reg. calidioribus hinc inde obviæ.

P. Cupani est forma minus variabilis et forsán sicut subspecies enumeranda.

P. MAURITANICA (*Boiss. et Reut. Pug.* 105; *Dene. in DC. Pr.* xiii. pars 1, 734).

Mar. merid.—In convalle Ansmiz a 1300 m. usque 1600 m.!

Ar. Geog. Africa boreali-occidentalis.

Sp. hanc cel. Decaisne foliis angustioribus a *P. Cynope* l. c. distinxit. Re vera folia variant angustiora, vel ut in sp. nostris breviora et latiora.

Præter cæteras notas *P. mauritanica* a *P. Cynope* optime dignoscitur foliolis calycinis subæqualibus, nec, ut in *P. Cynope*, insigniter inæqualibus. Forsán lapsu calami cel. Decaisne foliola calycina subæqualia *P. Cynopi* tribuit—testante ipsa natura, et consentientibus omnibus auctoribus, semper eximie inæqualia sunt.

P. PSYLLIUM (L., *Dene. in DC. Pr.* xiii. pars 1, 734; *Schonsb. Gew. Marok.* 54).

Mar. sept.—Tanger (Salzm. Exsicc. in Herb. Gay sub '*P. Psyllium*, L., non Fl. Gr.!) Inter Tanger et Tetuan! Tetuan!

Mar. occid.—Mogador! (*Lowe Cat.*).

Mar. merid.—Shedma (*J. D. H.*)! Mesfoua! Milhain! Seksaona!

Citatur inter synonyma hujus speciei *P. agrestis* (Salzm. Cat. Pl. Tingit.); opus citatum mihi incognitum est.

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). Etiam in Abyssinia inventa.

Plantago stricta (*Schousb. Gew. Marok.* 55; *Dene. in DC. Pr.* xiii. pars 1, 735).

Mar. occid.—Circa Mogador (*Schsb. l. c.*). In insula Mogador! Agadir (*Coll. Indig.* 1871)!

Differt a *P. Psyllio*, cui proxime affinis, indumento ex pilis papillosis brevissimis, ad nodos longioribus, plerumque retrorsis (non refractis ut habet cel. Decaisne), nec pubescenti-viscoso, foliis lineari-subulatis integerrimis, nec sæpius denticulatis, bracteis et foliolis calycinis apice teretibus obtusis, nec acuminatis. Hæ notæ videntur satis constantes sed non gravis momenti, et potius ad subspeciem definiendam idoneæ. Cel. Decaisne, nescio quomodo hallucinatus, bracteas inferiores acuminatas subaristatas l. c. descripsit.

Planta secundum auctores Imp. Maroccani propria. Cel. Decaisne tamen inter synonyma citat *P. rugosam* (Hochst. Pl. Adoen. Exsicc. No. 60), et ego vidi specimina in Herb. Kew. nostris conformia ex Afghanistan. Forsan enim aliis stirpibus per regionem desertorum usque provincias meridionales Imp. Maroccani dispersis enumeranda.

NYCTAGINEÆ (*Juss. Ann. Mus. Par.* ii. 269).

BOERHAVIA (*L. Hort. Cliff.* 17).

B. PLUMBAGINEA (*Cavan. Icon.* ii. 7, tab. 112; *Vahl, Enum.* i. 287), = *B. dichotoma* (*Vahl, Enum.* i. 290; *Chois. in DC. Pr.* xiii. pars 2, 454), = *B. pulverulenta* (*Dupuis in Vahl, Enum.* i. 291).

Mar. merid.—? Specimen nondum florens in prov. Shedma legit J. D. II.! Foliis et habitu spec. nostris Hispanicis et Africanis simillimum sed ex florum defectu omnino incertum.

Ar. Geog. Africa tropica et subtropica. Arabia Felix. In Hispania australi rarissima.

B. MAROCCANA (*Ball in Journ. Bot.* 1875, 176).

Basi lignosa, truncus scil. diametro pollicaris et ultra caules plurimos herbaceos vel inferne lignosos emittens; rami tenues adscendentes, dichotome divisi, sub lente hirtuli; folia petiolata, lanecolato-ovata, acuta, inferiora latiora obtusa, basi cuneata (nunquam subcordata), margine integra subundulata, insigniter bicoloria, in pagina superiore viridia nitida, inferne glauca; inflorescentia paniculato-cymosa, eymis contractis (pseudumbellis) ex floribus numerosis (quorum plurimi abortientes cito decidui) basi 2–3-bracteolatis; perigonium extus (cum apice ramulorum) glanduloso-viscidum, parte libera ex segmentis 5 inflexis post anthesin cum andrœcio et stylo simul decidua, tubo fructu aduato, quinquecostato, pilis nonnullis longioribus et glandulis subsessilibus vestito; antheræ tres; stigma capitatum.

Mar. merid.—In rupestribus collinis prope urbem Marocco primus detexit

el. G. Maw! In prov. Shedma anno 1873 specimen (in Herb. Kew. conservatum) legerunt ell. Rein et Fritsch!

B. elegans, Choisy, foliis ovatis bicoloribus plantæ nostræ aliquantulum accedit, sed inflorescentia diversa, perigonio non viseido, aliisque notis satis diversa. A *B. plumbaginea*, Cav., longius distat *B. maroccanu* floribus ter minoribus, et ab omnibus mihi notis perigonii segmentis inflexis, genitalia includentibus.

ILLECEBREÆ (*R. Br. Prod.* 413).

Ordo naturalis a celeb. R. Brown anno 1810 definitus, jure prioritatis et nomine satis apto ab auctore imposito, posteris commendatur.

IIERNIARIA (*L. Gen.* 308).

II. ANNUA (*Lag. Gen. et Sp.* 12), = II. cinerea (*DC. Fl. Fr.* v. 375, et *Prod.* iii. 367).

Mar. merid.—Satis frequens in regione inferiore et ad radices Atlantis Majoris. Prov. Shedma! Misra ben Kara! Mesfioua! Amsmiz! Mzouda! Ourika (*R. et Fr. spec. in Herb. Kew.*)!

— —, var. VIRESCENS, = II. virescens (*Salzm. Exsicc.*; *DC. Pr.* iii. 367).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Cap. Spartel! Tetuan (1851)!

Mar. occid.—Mazagan!

Ar. Geog. Regio Mediterranea occidentalis (inclusa Macaronesia).

H. virescens non differt nisi indumento parco, et meram varietatem sistit.

Spec. prope Mazagan lecta typo propiora fere intermedia sunt.

II. HIRSUTA (*L., DC. Pr.* iii. 367).

Mar. sept.—Prope ostium fluminis Tetuanensis (1851)!

Ar. Geog. Regio omnis Mediterranea. Europa media. Asia temperata.

Nescio an huc referenda sit *H. hirsuta* (Schousb. *Gew. Marok.* 107) in agro Tingitano inventa. Suspicio hanc ad *H. annuam* referendam esse.

II. POLYGONOIDES (*Cavan. Icon.* ii. 27, tab. 137; *DC. Pr.* iii. 368), = II. erecta (*Desf. Fl. Atl.* i. 214), = Ilceebrium suffruticosum (*L. Sp. Pl.* 298), = Paronychia suffruticosa (*Lam. Fl. Fr.* iii. 230).

In regno Maroccoano legit Broussonnet, fide Cavan. in *An. Cienc. Nat.* iii. 29.

Ar. Geog. Hispania australis. Africa boreali-occidentalis.

CORRIGIOLA (*L. Gen.* 378).

C. TELEPHIFOLIA (*Pourr. Chlor. Narb.* 20; *DC. Pr.* iii. 367), = C. litoralis (*Schousb. Gew. Marok.* 131, non *L.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Cap. Spartel!

Ar. Geog. Verisimiliter per regionem fere omnem Mediterraneam dispersa. Deest in Macaronesia. *C. litoralis*, *L.*, proxime affinis, etiam in Europa occidentali, neenon austro-orientali (Hungaria, Rossia media, et ect.), et secus litora Maris Caspii obvia est.

Hæc est *C. litorali* omnibus partibus major, radix sæpius (non semper) perennis; notæ diagnosticæ ex foliolis calycinis a Grenier et Godron in Fl. Fr. i. 614, indicatæ mihi non constantes videntur. Meo sensu pro subspecie *C. litoralis* enumeranda.

ILLECEBRUM (*L. Gen.* 290).

I. VERTICILLATUM (*L., DC. Pr.* iii. 370; *Schousb. Gew. Marok.* 103).

Mar. sept.—Tanger (*Webb Herb.*)! In aridis monte Djebel Kebir (*Schousb. l. c.*).

Mar. occid.—Mogador! (*Broussonnet ex Cavan. An. Cienc. Nat.* iii. 29).

Ar. Geog. Regio Mediterranea occidentalis (inclusa Macaronesia). Europa occidentalis. Occurrit etiam in Dania, in Rossic insula Osilia, et in insula Coreyra.

PARONYCHIA (*Juss. Mém. Mus. Par.* i. 388).

P. CYMOSA (*L., et Schousb. Gew. Marok.* 103, sub *Illecebro*; *DC. in Dict. Encyc.* v. 26, et *Prod.* iii. 370).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! In monte Djebel Kebir (*spec. ex Herb. Schousb.*)!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia), a Gallia australi usque Lusitaniam; ex synonymo falso in Creta erronee indicatur.

P. ECHINATA (*Desf. Fl. Atl.* i. 204, sub *Illecebro*; *Lam. Fl. Fr.* iii. 232; *DC. Pr.* iii. 370), = *Illecebrum cymosum* (*Fl. Gr.* tab. 245, non *L.*).

Mar. sept.—Frequens in ditione Tingitana. Tanger (*Salzm. Exsicc.*)! In monte Djebel Kebir! Cap. Spartel! Inter Tanger et Tetuan!

Ar. Geog. Regio Mediterranea imprimis occidentalis (inclusa Macaronesia). In Oriente nonnisi in insulis Creta et Melo huensque detecta.

P. ARGENTEA (*Lam. Fl. Fr.* iii. 230; *DC. Pr.* iii. 371), = *Illecebrum Paronychia* (*L., Schousb. Gew. Marok.* 103).

Species polymorpha per Imperium Maroccanum nobis notum a litore Oceani usque regionem superiorem vulgatissima. Varietates ægre definiendas maxima pro parte prætermisi.

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Schsb. l. c.*)! Tetuan (1851)!

Mar. occid.—Mazagan! Mogador! (*Lowe Cat.*). Prov. Haha prope Mogador! Aïn el Hadjar! Agadir (*Coll. Indig.* 1871)

Mar. merid.—Reraya! Aïn Tarsil! Aït Mesan! In ascensu jugi Tagherot usque 2900 m.! In convalle Amsmiz visa!

Ar. Geog. Regio omnis Mediterranea. In insulis Canariensibus rara.

— —, var. MAURITANICA (*DC. Pr. l. c.*), = *Illecebrum mauritanicum*, *Willd.*; differt foliis latioribus capitulis et remotioribus.

Mar. sept.—Cap. Spartel! Inter Tetuan et Ceuta!

Ma. occid.—Agadir (*Coll. Indig.* 1871)!

— —, var. *sepalis glabris vix encullatis.*

Mar. occid.—Mogador!

— —, var. *VELUTINA*, foliis et caulibus pube brevi cinerea vestitis.

Caulēs sæpe tantum latere superiore velutini.

Mar. merid.—In calidioribus regionis inferioris. Misra ben Kara!

Milhain! Amsuiz!

Folia aqua decocta instar Theæ ab incolis utuntur.

PARONYCHIA CAPITATA (L., sub *Illecebro*; *Lam. Fl. Fr.* iii. 229; *DC. Pr.* iii. 371), var. *NIVEA*, = *P. nivea* (*DC. in Dict. Encyc.* v. 25, et *Prod. l. c.*).

Mar. merid.—In convalle Aït Mesan circ. 1400 m.!

— —, var. *ATLANTICA*, *nob.* Forma a typo aspectu valde diversa sed vix notis stabilibus distincta. Folia parva numerosa obtusa recurvata pubescentia, stipulis lanceolato-acuminatis longiora; flores in capitula parva sæpius terminalia glomerati; bracteæ floribus vix longiores late obtusæ cuspidatæ. Facie *P. aretioidi*, DC., similis, differt foliis patulis denum recurvis, sepalis magis inæqualibus, acutioribus, ovato-lanceolatis.

Mar. merid.—Satis frequens in convalle Aït Mesan usque in ascensum jugi Tagherot a 2000 m. usque 2900 m.!

Ar. Geog. *P. capitata* per montes Europæ mediæ et regionis fere omnis Mediterraneæ (exclusa Macaronesia) diffusa est.

Notæ diagnosticæ quibus *P. nivea* a *P. capitata* distinguitur mihi videntur instabiles. Specimina Gallica a ell. Grenier et Godron (*Fl. Fr.* i. 611) imprimis sepalis in *P. capitata* æqualibus obtusis, in *P. nivea* valde inæqualibus acutis, distinguuntur; sec. el. Willkomm et Lange (*Fl. Hisp.* iii. 151) sepala *P. niveæ* apice angustata, obtusa. In spec. permultis a me exploratis sepala video variabilia et ambigua, interdum subæqualia, sæpius tamen plus minusve inæqualia, semper nervosa præsertim in facie interiore. In specie homologa *P. argentea* stipulæ et bracteæ ludunt minores vel majores, latiores vel angustiores, unde mihi incertum videtur an *P. aretioides* pro specie probe diversa habenda sit. Varietas nostra *atlantica*, stipulis et bracteis in hoc grege minimis, et *P. aretioides*, DC., forsan melius sicut subspecies enumerandæ.

P. MACROSEPALA (*Boiss. Diagn. Pl. Or.* ser. 1, iii. 11, et *Fl. Or.* i. 745; *Ball in Journ. Bot.* 1875, 204).

Mar. merid.—Videtur satis frequens in aridis regionis inferioris et ad radicis Atlantis Majoris. Aïn Onnast! Mesfioua! Seksaoua! In rupestribus collinis prope urbem Marocco (*G. Maw*)! In convalle Aït Mesan circ. 1300 m.!

Stirpem hanc pro specie nova descripsi eodem nomine quo eam jampridem salutavit cel. Boissier. Frustula ex Attica quæ debeo humanitati cel. amici habent folia majora plana, et descriptioni Boissierianæ haud omnino conformia, tamen notis gravioribus omnino conveniunt. Adjeci descriptionem stirpis nostræ, unde differentię minores facilius emineant.

Planta perennis; caules numerosi prostrati; folia parva oblongo-ovata acutiuscula, dense pubescentia, demum patulo-recurva; stipulæ lanceolatae, longiuscule acuminatae; bracteae ovato-lanceolatae, abbreviatae, sepalis vix æquilongae; sepala subincrassata, vix aut ne vix nervosa, oblongo-linearia, obtusa, mutica, valde inæqualia, apice plus minusve recurva, etiam in præfloratione basi discreta; petala (vel staminodia?) minima, linearia; stylus brevis, profunde bifidus, ramis stigmatosis divaricatis; stamina brevissima, ima basi disco calycino adnata; antherae breves.

A grege *P. capitata* differt imprimis sepalis (v. calycis segmentis) majoribus crassioribus subnerviis, valde inæqualibus, adeo ut sep. maximum minimo duplo longius est, fructiferis recurvis, nec clausis, petalis (v. staminodiis?) linearibus, nec nullis, antheris fere dimidio minoribus, stylo denique brevior.

— —, var. *QUERIODES* (*Ball in Journ. Bot. l. c.*). Bracteis brevioribus, foliis velutinis rigidiusculis approximatis, insigniter recurvis a typo recedit, et facie *Queriam hispanicam* simulat.

Mar. merid.—Specimen unicum legi in convalle Aït Mesan circ. 1400 m.!

Ad typum *P. macrosepala* relata fere ut var. nostra *atlantica*, *P. capitata*.

Ar. Geog. *P. macrosepala* hucusque in Græcia, Asia Minore, et in insulis adjacentibus, necnon in ditione Maroccana meridionali detecta, forsan alibi in regione Mediterranea calidiore invenienda.

SCLERANTHUS (*L. Gen.* 562).

S. ANNUUS (*L., DC. Pr.* iii. 378), var. *VERTICILLATUS* (*Fenzl, in Ledeb. Fl. Ross.* ii. 156), = *S. verticillatus* (*Tausch in Flora*, 1829.)

Mar. merid.—In monte Djebel Tezali circ. 2100 m.!

Ar. Geog. *S. annuus* per Europam omnem (exclusa arctica), Asiam mediam et regionem fere omnem Mediterraneam diffusa est. Adest in Abyssinia, sed ex Macaronesia non vidi.

AMARANTACEÆ (*R. Br. Prod.* 413).AMARANTUS (*L. Gen.* 1068).

A. CHLOROSTACHYS (*Willd. Amarant.* 34, tab. 10, fig. 19; *Moq. in DC. Pr.* xiii. pars 2, 250).

Mar. merid.—Prov. Shedma juxta castellum Præfecti!

Ar. Geog. Regio fere omnis Mediterranea. In utraque America obvi-
us, an indigenus?

A. BLITUM (*L., Moq. in DC. Pr.* xiii. pars 2, 263).

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—In prov. Shedma!

— — —, var. *GRÆCIZANS* (*Moq. l. c.*), = *A. græcizans*, *L.*

Mar. occid.—Mazagan! Saffi!

Ar. Geog. Species variabilis per orbem terrarum (exceptis terris Aus-
tralisicis) in temperatis calidioribus et subtropicis latissime diffusa.

A. ALBUS (*L., Moq. in DC. Pr.* xiii. pars 2, 264).

Mar. occid.—Mazagan!

Ar. Geog. Secundum auctores ex America boreali migrata, hodie spon-
tanea in regione Mediterranea occidentali (exclusa Macaronesia).

A. DEFLEXUS (*L., Willd. Amarant.* 10, tab. 10, fig. 20), = *A. prostratus*
(*Balb. Misc. Bot.* 44, tab. 10), = *Euxolus deflexus* (*Rafin. Fl. Tell.*
42; *Moq. in DC. Pr.* xiii. pars 2, 275).

Mar. sept.—Tanger! (*spec. ex Herb. Schousboe*)!

Ar. Geog. Species per regionem Mediterraneam occidentalem, et Euro-
pam austro-occidentalem diffusa. In America meridionali advena?
Deest in Macaronesia.

ACHYRANTHES (*L. Gen.* 113).

A. ARGENTEA (*Lam. Dict.* i. 545; *Moq. in DC. Pr.* xiii. pars 2, 315;
Schousb. Gew. Marok. 101).

Mar. sept.—Tanger! In fruticetis regionis Tingitanæ (*Schsb. l. c.*).
Tetuan (1851)!

Mar. occid.—Salli (*Broussonnet ex Cavan. in Anal. Cienc. Nat.*).

Ar. Geog. Africa borealis. Arabia Felix. Sicilia. Sardinia. Italia
meridionalis. In Hispania nonnisi ad Gibraltar. Macaronesia. Se-
negambia. Africa australis.

POLYCNEMUM (*L. Gen.* 53).

P. FONTANESII (*Durieu et Moq. in DC. Pr.* xiii. pars 2, 335).

Mar. occid.—Casa Blanca (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—Ad radices Atlantis Majoris haud infrequens. Tassere-
mout! Ourika! Ait Mesan circ. 1300 m.!

Ar. Geog. Africa boreali-occidentalis.

In spec. nostris folia nunquam subcurva.

CHENOPODIACEÆ (*Lindl. Nat. Syst.* ed. 2, 203).BETA (*L. Gen.* 310).

B. VULGARIS (*L., Moq. in DC. Pr.* xiii. pars 2, 55), var. *MARITIMA* (*Moq. l. c.* 56), = *B. maritima* (*L., Willk. et Lge. Fl. Hisp.* i. 274).

Mar. sept.—Tanger (*J. D. H.*)! Tetuan (1851)!

Mar. occid.—Casa Blanca! Mogador! (*Lowe Cat.*). Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Oecurrit hinc inde in regione inferiore. Sheshaoua! In ruderatis urbis Marocco! Seksaoua!

Ar. Geog. *B. vulgaris* verisimiliter sponte nascitur in regione omni Mediterranea, in Europa media et occidentali, et in India orientali.

Beta Cicla (*Willd. Sp. Pl.* i. 1309), a Schousboe (*Gew. Marok.* 108) prope Tingium indicata, est forma *B. vulgaris* eulta vel ex cultura evasa.

CHENOPODIUM (*L. Gen.* 309).

C. OLIDUM (*Curt. Fl. Lond.* v. tab. 20; *Eng. Bot.* tab. 1034), = *C. fœtidum* (*Lam. Fl. Fr.* iii. 244, non *Schrad.*), = *C. Vulvaria* (*L., Moq. in DC. Pr.* xiii. pars 2, 64).

Mar. occid.—Agadir (*Coll. Indig.* 1871)!

Mar. merid.—In ruderatis urbis Marocco!

Ar. Geog. Europa media et australis. Africa borealis.

C. OPULIFOLIUM (*Schrad. in DC. Fl. Fr.* v. 352; *Moq. in DC. Pr.* xiii. pars 2, 67).

Mar. sept.—Vidi nondum florens prope Tanger!

Ar. Geog. Europa media. Regio Mediterranea (exclusa Macaronesia). In America boreali obvium sed verisimiliter advena.

C. MURALE (*L., Moq. in DC. Pr.* xiii. pars 2, 69; *Schousb. Gew. Marok.* 108).

Mar. sept.—Prope Cap. Spartel!

Mar. occid.—Casa Blanca! Mogador (*Lowe Cat.*). Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Sheshaoua! In ruderatis urbis Marocco! Seksaoua!

Ar. Geog. Herba amphigea per Europam temperatam, regionem omnem Mediterraneam, Indiam orientalem, et Africam australem diffusa; etiam in America australi et in Australia obvia.

C. ALBUM (*L., Moq. in DC. Pr.* xiii. pars 2, 70).

Mar. occid.—Saffi! Mogador! (*Lowe Cat.*).

Ar. Geog. Regio omnis Mediterranea. Europa media et borealis (exclusa arctica). Asia media et calidior. Obvium etiam in America boreali, nec non in insulis Sandwich.

C. AMBROSIODES (*L., Moq. in DC. Pr.* xiii. pars 2, 72; *Schousb. Gew. Marok.* 108).

Mar. oceid.—Mogador (*Lowe Cat.*).

Mar. merid.—*Legimus* nondum florens in ruderatis urbis Marocco! et ad radices Atlantis Majoris prope Gurguri!

“ In locis arenosis depressis udis raro occurrit ” (*Schsb. l. c.*).

Ar. Geog. Verisimiliter ex zona tropica in Europam et alio introdueta, hodie herba eozmopolita per totum orbem in temperatis calidioribus et in tropicis diffusa. Rarius occurrit in Europa media vestigia hominum secuta.

ATRIPLEX (*L. Gen.* 1153).

A. LACINIATA (*L., Moq. in DC. Pr.* xiii. pars 2, 93).

Mar. sept.—Tanger (*Salzm. Exsicc.* in herb. Gay, sub *A. laciniata* et *A. glauca*)!

Mar. merid.—In regione inferiore solo gypsaceo. Sheshaoua! Misra Ben Kara! In planitie prope urbem Marocco (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. In litoribus Europæ occidentalis et regionis omnis Mediterraneæ (inclusa Macaronesia). Occurrit etiam in America septentrionali.

A. laciniata et *A. rosea*, *L.*, sunt species arete affines et valde ludibundæ, ab auctoribus sæpe confusæ. Loci natales igitur non semper certi.

A. PATULA (*L., Moq. in DC. Pr.* xiii. pars 2, 95).

Mar. sept.—Prope Tanger nondum florentem vidi!

Ar. Geog. Europa fere omnis (excepta arctica). Regio Mediterranea fere omnis (exclusa Macaronesia).

A. PARVIFOLIA (*Lowe, Prim. Fl. Mad.* 16; *Moq. in DC. Pr.* xiii. pars 2, 99, non *Kunth*).

Mar. occid.—Saffi in maritimis! In rupibus insulæ Mogador

Ar. Geog. Macaronesia. Marocco occidentalis.

A. HALIMUS (*L., Moq. in DC. Pr.* xiii. pars 2, 100).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. merid.—Sheshaoua! Mtouga (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio Mediterranea omnis, imprimis litoralis. Africa australis. Indicatur etiam in utraque America, an indigena?

OBIONE (*Gærtn. Fruct.* ii. 198).

O. PORTULACOIDES (*L., sub Atriplici; Moq. Chenop. Enum.* 75, et in *DC. Pr.* xiii. pars 2, 112).

Mar. sept.—Vidi nondum florentem prope Tanger! (*Spec. ex Herb. Schousboe comm. Cosson*)!

Ar. Geog. Litora Atlantica Europæ occidentalis et regionis fere omnis Mediterraneæ (exclusa Macaronesia). Occurrit etiam in Africa australi et in America boreali.

CHENOLEA (*Thunb. Nov. Gen. Pl.* 9).

C. CANARIENSIS (*Moq. Chenop. Enum.* 96; et in *Webb Phyt. Canar.* iii. 205, tab. 200), = C. lanata (*Moq. in DC. Pr.* xiii. pars 2, 129), = *Salsola lanata* (*Masson in Buch. Besch. Can.* 141, non *Gmel.*, nec *Pall.*, nec *Vahl*, nec *Poll.*).

Mar. occid.—In rupibus insulæ Mogador!

Ar. Geog. Stirps monotypica Macaronesiæ et litoralis Maroccanæ propria.

Nomen triviale *lanata Salsolis* plurimis alienis olim applicata nulla prioritate gaudet, et nomen *canariensis* jamdudum a cel. auctore evulgatum non mutandum videtur.

SALICORNIA (*L. Gen.* 10).

S. FRUTICOSA (*L. Sp. Pl.* ed. 2, 51, pro parte; *Ungern-Sternberg in Atti Congr. Bot. Fir.* 294, et verisimiliter *Schousb. Gew. Marok.* p. 1), = *Arthrocnemum fruticosum* (*Moq. Chenop. Enum.* iii., et in *DC. Pr.* xiii. pars 2, 151, pro parte).

In arenosis maritimis totius imperii (*Schsb. l. c.*).

Mar. merid.—Nondum florentem legimus in saxosis solo gypsaceo prope Sheshaoua!

Synonymon *Schousbœanum* forsitan ex parte ad *Arthrocnemum glaucum* referendum.

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia). Europa occidentalis. Africa australis. America fere omnis. In litore maris vel rarius in solo salso vel gypsaceo a mari longinquo.

ARTHIROCNEMUM (*Moq. Chenop. Enum.* 111).

A. GLAUCUM (*Del. Fl. Eg.* 69; *Ungern-Sternberg in Atti Congr. Bot. Fir.* 283), = A. fruticosum (*Moq. l. c.*, et in *DC. Pr.* xiii. pars 2, 151, pro parte), = *Salicornia macrostachya* (*Moric. Fl. Ven.* i. 2; *Guss. Prod. Fl. Sic.* i. 2).

Mar. sept.—? Fere certe in arenosis pro Tanger visum!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia) et in insulis Capitis Viridis. Americæ borealis dubia civis.

Clar. Ungern-Sternberg loc. cit. gregem *Salicorniearum* eximie tractavit, et observationibus accuratis illustravit.

Stirpes supra enumeratæ aspectu ægre dignoscendæ, ab auctoribus sub *Salicornia fruticosa* vel *Arthrocnemo fruticoso* semel confusæ, nunc in generibus diversis criterio securo separandæ sunt. Structura seminis in *Salicornia* albumine carentis, embryo conduplicato, in *Arthrocnemo* albumine præditi, embr. virgulæformi, immo ratio vitalis totius plantæ omnino diversa. Cæterum diagnosin harum specierum ell. Willkomm et Lange (*Fl. Hisp.* i. 263, 264) jampridem recte indicarunt.

SUÆDA (*Forsk. Fl. Æg.-Arab.* 62).

S. FRUTICOSA (*Forsk. Fl. Æg.-Arab.* 70; *Moq. in DC. Pr.* xiii. pars 2, 156), = *Salsola fruticosa* (*L., Eng. Bot.* tab. 635; *Schousb. Gew. Marok.* 109).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. occid.—Saffi! Mogador! (*Schsb. l. c.*).

Mar. merid.—Nondum florens sed fere certa in solo gypsaceo prope Sheshaoua! et Misra Ben Kara!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). Europa occidentalis. Senegal. Indicatur etiam in California.

S. MARITIMA (*L., sub Chenopodio; Dumort. Fl. Belg.* 22), = *Chenopodia maritima* (*Moq. in DC. Pr.* xiii. pars 2, 164).

Mar. merid.—Nondum florens sed fere certa in solo gypsaceo prope Sheshaoua!

Ar. Geog. Regio omnis Mediterranea. Europa occidentalis et media. Fertur etiam in insula Cuba et in ins. Sanctæ Helenæ.

CAROXYLON (*Thunb. Gen. Nov. Pl.* 37).

C. ARTICULATUM (*Moq. in DC. Pr.* xiii. pars 2, 175), = *Salsola articulata* (*Cav. Icon.* iii. 43, tab. 284, non *Forsk.*), = *Halogeton articulatum* (*Bunge in Lehm. Rel.*).

Mar. merid.—Legimus prope Sheshaoua solo gypsaceo!

Ar. Geog. Hispania australis. Ægyptus. Marocco meridionalis.

SALSOLA (*L. Gen.* 311).

S. VERMICULATA (*L., Moq. in DC. Pr.* xiii. pars 2, 181), var. *a*, FLAVESCENS (*Moq. l. c.*), = *S. flavescens* (*Cavan. Icon.* iii. 45, tab. 288).

Mar. sept.—Tanger (*Salzmann sec. Moq. l. c.*).

— — —, var. MICROPHYLLA (*Moq. l. c.*), = *S. microphylla* (*Cavan. Icon.* iii. 45, tab. 287), = *S. brevifolia* (*Desf. Fl. Atl.* i. 218; *Salzm. Exsicc.*)!

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. merid.—? Sheshaoua (nondum florens et subincerta)!

Ar. Geog. Regio Mediterranea calidior (inclusa Macaronesia). Asia media. Deest in Gallia et in Italia continentali. Var. in regione Mediterranea imprimis occidentali.

S. LONGIFOLIA (*Forsk. Fl. Æg.-Arab.* 55; *Moq. in DC. Pr.* xiii. 182, non *Lam.*).

Mar. occid.—Legimus in insula Mogador et in arenosis prope urbem!

— — —, var. VERTICILLATA, = *S. verticillata* (*Schousb. Gew. Marok.* 109).

Mar. occid.—In insula Mogador (*Schsb. l. c.*).

Mar. merid.—Hinc inde in regione inferiore obvia. Seksaoua! In gla-

reosis Oued en Fisk prope Gurguri! Ourika (*R. et Fr. spec. in Herb. Kew.*)! Inter Sektana et Frouga (*G. Maw*)!

Differt a typo foliis brevioribus tenuioribus subverticillatis.

Ar. Geog. Species per Africam borealem dispersa, etiam in Hispania austro-orientali et in Macaronesia obvia. Var. *verticillata* nonnisi in Marocco meridionali et in insulis Canariensibus detecta.

SALSOLA KALI (*L., Ten. Syll. Fl. Nap.* 124; *Moq. in DC. Pr.* xiii. pars 2, 187).

Mar. occid.—Casa Blanca!

Spec. nostra ad var. *α, hirtam* (*Moq. l. c.*) referenda.

Ar. Geog. In litoribus fere omnibus Europæ, Asiæ, Africae borealis et Americae septentrionalis (exceptis terris arcticis) diffusa.

Inter *Chenopodiaceas* plurimas in solo gypsaceo enatas a nobis prope Seksaoua lectas adsunt specimina nondum florentia prorsus incerta, ex foliis et habitu ad *Camphorosma* speciem forsitan referenda.

Cll. Rein et Fritsch prope Mogador legerunt specimen nondum florigerum verisimiliter ad *Anabasin articulatam* (*Moq. in DC. Pr.* xiii. pars 2, 214) referendum. Planta ulterius inquirenda.

THELYGONUM (*L. Gen.* 1098).

T. CYNOCRAMBE (*L., Koch, Syn. Fl. Germ.* 703), = *Cynoerambe prostrata* (*Gærtn. Fruct.* i. 362, tab. 75; *A. DC. in Prod.* xvii. 33).

Mar. sept.—In monte Beni Hosmar prope Tetuan! Exstat in herb. Webb verisimiliter prope Tetuan lectum!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

Præcuntibus celeb. Lindley et G. D. Koch, Alph. De Caudolle loco suo vero in serie naturali *Salsolis* proximo hanc plantam dubiæ affinitatis collocavit. Minus fauste nomen genericum Linnæanum ab auctoribus fere omnibus recentioribus receptum mutare statuit.

POLYGONEÆ (*Juss. Gen.* 82).

EMEX (*Neck. Elem.* ii. 214).

E. SPINOSA (*L., et Schousb. Gew. Marok.* 169, sub *Rumici*; *Campd. Rum.* 58, tab. 1; *Meisn. in DC. Pr.* xiv. 40).

Mar. sept.—Tanger! (*Schsb. l. c.*).

Mar. occid.—Casa Blanca! Mogador! (*Lowe Cat.*). Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Pluries visa in prov. Shedma et alibi!

Ar. Geog. Regio omnis Mediterranea, imprimis litoralis. In Brasilia introdueta.

RUMEX (*L. Gen.* 357).

R. CRISPUS (*L., Meisn. in DC. Pr.* xiv. 44).

Mar. merid.—In hortis et ruderalis urbis Marocco!

Ar. Geog. Species cosmopolita ex temperatis frigidioribus usque in zonam æquinoctialem. fere ubique diffusa, rarior tamen in calidioribus. Ex Macaronesia non vidi.

RUMEX CONGLOMERATUS (*Murr. Prod. St. Gott.* 52; *Meisn. in DC. Pr.* xiv. 49), = *R. acutus* (*Sm. Fl. Brit. et E. Bot.* tab. 724).

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—Mzonda! Prope Amsmiz visa!

Ar. Geog. Europa fere omnis (exclusa arctica). Oriens. Teneriffa, Africa borealis et australis. America borealis.

R. PULCHER (*L., Meisn. in DC. Pr.* xiv. 58).

Mar. sept.—Tetuan (1851)!

Mar. occid.—Casa Blanca!

Mar. merid.—Sheshaoua! In glareis Oued en Fisk pr. Gurguri (*J. D. H.*)!

Ar. Geog. Regio omnis Mediterranea. Europa media. Africa australis. In Brasilia, forsân introducta.

R. BUCEPHALOPHORUS (*L., Schousb. Gew. Marok.* 168; *Meisn. in DC. Pr.* xiv. 62).

Mar. sept.—Species in ditione Tingitana vulgatissima. Tanger! (*Schsb. l. c.*). In monte Djebel Kebir! Cap. Spartel! Inter Tanger et Tetuan! Tetuan (1851)! In monte Beni Hosmar! Inter Tetuan et Ceuta!

Mar. occid.—Mazagan! Mogador! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Ocenrrit hinc inde, rarius tamen quam in regione sublitorali. Seksaoua (*J. D. H.*)! Mtouga!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

R. INTERMEDIUS (*DC. Fl. Fr.* vi. 369; *Meisn. in DC. Pr.* xiv. 65).

Mar. sept.—Legi mox floribundum prope urbem Tetuan et ad ostium fluminis mense Junio, 1851!

Ar. Geog. Europa austro-occidentalis ex Sicilia usque fretum Herculis. Spec. Africana præter nostra non vidi.

Radix in spec. nostris et in plurimis Europæis in Herb. Kew. conservatis horizontalis vel oblique descendens. Spec. prope urbem Tetuan lectum habet indumentum ex pilis brevioribus papillaribus ut in *R. papillari* (*Boiss. et Reut. Pug.* 107), quæ mihi omnino var. *R. Acetosæ* videtur.

R. THYRSOIDES (*Desf. Fl. Atl.* i. 321; *Meisn. in DC. Pr.* xiv. 66).

Mar. sept.—Legi inter Tetuan et Centa!

Ar. Geog. Africa borealis. Insulæ Italicæ majores (Sicilia, Sardinia, Corsica). In Hispania nonnisi prope Gibraltar.

R. TINGITANUS (*L., Meisn. in DC. Pr.* xiv. 67; *Schousb. Gew. Marok.* 169).

Mar. sept.—In arenosis maritimis pr. Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! (*Schousb. l. c.*).

Zanoni primus hanc speciem publici juris fecit et icone satis bona illustravit anno 1675. Confer 'Istoria Botannica,' p. 14, fig. 5 (nec 6 ut habet Meisner l. c.). Semina et spec. exsiccata Tingitana Zanoni habuit anno 1664 ab Alexandro Balam mercatore Anglico, qui serius semina et specimina Morisonio præbuit.

Ar. Geog. Species per regionem Mediterraneam imprimis occidentalem dispersa. Fertur in insulis Græciæ, sed deest in Oriente continentali. Forma arete affinis (*R. madereusis*, Lowe) in Macaronesia.

Rumex pictus (*Forsk. Fl. Æg.-Arab.* 77), = *R. lacerus* (*Balb. Miscell.* 19; *Meisn. in DC. Pr.* xiv. 69), = *R. bipinnatus* (*L. fl. Suppl.* 211; *Schousb. Gew. Marok.* 169).

Verisimiliter est etiam *R. roseus* (*Desf. Fl. Atl.* i. 320, et *Poir. Diet.* v. 65, non L.).

Mar. occid.—Frequens et satis ludibundus occurrit in arenosis juxta urbem Mogador! et in proxima prov. Ithala meridiem versus! (*Schsb. l. c.*).

Stirps nostra ludit foliis simpliciter pinnatifidis ut in spec. orientalibus, vel bipinnatifidis ut in descript. Schousbœana. Flores vidi maseulos et hermaphroditos in eodem racemo, nec, ut vult cel. Meisner l. c., dioicos. Hæc nota tamen in grege *R. Acetosæ* certe variabilis.

R. pictus est procul dubio ipsissimus *R. lacerus*, Balb., in arenosis Ægypti inferioris ad Rosetta alibique obvia, nomen igitur antiquius prioritate gaudet.

R. pictus a *R. tingitano* differt statura humiliore, caulibus e collo radicis valde ramosis, foliis profundius divisis, valvis subcallosis. In *R. tingitano* nervus medius valvarum nec incrassatum nec (nisi rarius ad imam basin) prominens, dum in *R. picto* nervus ad basin valvarum prominens sæpius in callum parvum lanceolatum incrassatum. Pro subspecie *R. tingitani* enumerandus.

Ar. Geog. *R. pictus* in arenosis Afriæ borealis et Arabiæ et Syriæ obvius est.

R. SCUTATUS (L., *Meisn. in DC. Pr.* xiv. 69), var. *INDURATUS*, nob., = *R. induratus* (*Boiss. et Reut. Pug.* 107).

Mar. merid.—In convalle Aït Mesan a 1500 m. usque 2000 m.! In jugo Tagherot (*R. et Fr. spec. in Herb. Kew.*)! Spec. ex monte Djebel Ouensa ab indigeno Ibrahim lecta benev. comm. cel. Cosson!

R. induratus, Boiss. et Reut., pluries a me in Hispania lectus, mihi videtur mera varietas *R. scutati*. Spec. nostra Atlantica fere intermedia sunt sed stirpi Boissierianæ propiora.

Ar. Geog. *R. scutatus* per Europam fere omnem (exclusa aretica) et in montibus regionis Mediterraneæ late diffusa est. Deest in Macaronesia et in Insulis Britannicis. Varietas *induratus* in peninsula Iberica et in Atlante Majore.

Rumex Papilio (Coss. in *Ball. Soc. Bot. Fr.* xx. 260).

Sequitur descriptio Cossoniana :—

“*Planta perennis*, pluricaulis, glaberrima, caudice brevissimo verticali in radicem fusiformem sublignosam abeunte. Caules erecti, adscendentes vel diffusi, superne ramosi, ramis divergentibus. *Folia* omnia petiolata, inferiora ovato-oblonga sæpe in parte inferiore latere altero angulo præminente donata, media *ovato-triangularia* acuta basi *hastata* auriculis divergentibus vel divaricatis, superiora linearia, ocreis ovato-oblongis margine integris. *Flores polygamo-monoici*, in racemos paucos laxos aphyllous paniculam generalem efformantes dispositi, fasciculati, fasciculis 2–5-floris remotis, pedicellis tenuiter filiformibus arcuato-deflexis. Calycis laciniæ exteriores oblongo-lineares primum erectæ, dein reflexæ; *valvæ accrescentes, demum diaphano-membranaceæ, tenuiter venosæ* venis vix prominulis, margine denticulatæ, ambitu orbiculato-subrhombæ, *supra medium utrinque profunde incisæ et inde quadrilobæ* lobis inferioribus ovato-suborbiculatis, superioribus minoribus oblongis divergentibus, in disco pilis paucis papilliformibus donatæ, basi callo destitutæ vel una alteraque callo obsoleto prædita.

R. Papilio, e forma valvarum papilionem referentium denominatus, in sectione *Acetosa* (Meisn. in DC. *Prod.* xiv. 64) juxta *R. scutatum*, L., collocandus, sed ab illo et cæteris speciebus affinibus valvis quadrilobis eximie distinctus.”

Mar. merid.—In fascibus annis Ghaghaia ad meridiem urbis Marocco primus legit Balansa sec. Cosson l. c. Specimen unicum in convalle Aït Mesan legit cel. J. D. Hooker!

Nisi valvarum forma planta hæc a *R. scutato* ægre distinguenda. Paucioris auctoris amicissimi hæc mihi videtur subspecies, vel forsitan lusus singularis *R. scutati*.

In spec. Hookeriano desunt pili papilliformes in disco ab auctore indicati.

R. VESICARIUS (L., Meisn. in DC. *Pr.* xiv. 70), var. *RHODOPHYSA* (Ball in *Journ. Bot.* 1875, 205).

Differt a typo radice bienni, anno primo caules plurimos breves ramosos emittens. In his racemus abbreviatus pauciflorus, pedunculi florum geminorum brevissimi paulo infra medium articulati, valvæ maximæ, pulchre rosæ, alis reflexe plicatis adhærentibus, interdum callosis, callo inter alas latente, nervo marginali nullo. Anno secundo caulis fit altior alterne ramosus.

Mar. merid.—Legimus fructiferum eire. 1500 m. s. m. in monte prope Seksaoua!

Pedicellis brevissimis, et radice multicauli *R. roseus* (L., Meisn. l. c. 71) plantæ nostræ simillimus, differt tamen valvis margine nervo continuo spinuloso-ciliato cinctis.

Ar. Geog. *R. vesicarius* per regiones tropicas orbis veteris, Asiam Mediterraneam calidiorem et Africam borealem (inclusis ins. Canariensibus) late dispersa est, nullibi, ut videtur, communis. In Europa nonnisi in Sicilia et Græcia hucusque detectus. In insula St.-Helenæ indicata et (quod mirum) in insula Cuba.

POLYGONUM (*L. Gen.* 495).

P. EQUISETIFORME (*Sibth. et Sm. Prod. Fl. Gr.* i. 266; *Meisn. in DC. Pr.* xiv. 85), var. *TRIGYNUM* (*Boiss. Voy. Esp.* 554), = *P. suffruticosum* (*Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Inter Tetuan et Ceuta in salsis uliginosis nondum florens!

Specimina omnia quæ vidi ad var. *trigynum* referenda. Auctores *Fl. Græcæ* specimina ex insula Creta fl. digynis descripserunt, an errore aliquo vel forsitan ex lusu abnormali?

Ar. Geog. Regio Mediterranea calidior (exclusa Macaronesia). In Europa nonnisi in Hispania et Græcia obvia.

P. MARITIMUM (*L., Meisn. in DC. Pr.* xiv. 88; *Schousb. Gew. Marok.* 178).

Mar. sept.—"In arena mobili juxta Tingidem" (*Schsb. l. c.*). In arenosis maritimis a Promontorio Spartel meridiem versus!

Mar. occid.—Mazagan! Mogador! (*Lowe Cat.*).

Ar. Geog. In litoribus Atlanticis Europæ et Africæ boreali-occidentalis ex Hibernia usque insulas Canarienses; inde per litora omnia maris Mediterraneæ et Ponti Euxini diffusa. In litore Atlantico Americæ borealis etiam obvium.

P. AVICULARE (*L., Meisn. in DC. Pr.* xiv. 97; *Schousb. Gew. Marok.* 178).

Mar. sept.—Tanger (*Schousb. l. c.*).

Mar. merid.—In ruderatis urbis Marocco! In convalle Ait Mesan supra Arround, circ. 2100 m.!

Ar. Geog. Species cosmopolita per orbem fere omnem e regionibus arcticis usque zonam torridam, et in insulis Oceanicis diffusa.

P. litorale (*Link in Schrad. Journ.* i. 54; *Meisn. in DC. Pr.* xiv. 98).

Mar. occid.—In maritimis prope Mazagan!

Ar. Geog. In litore Atlantico Europæ occidentalis et Imp. Marocani. Occurrit etiam hinc inde in litore maris interni. Indicatur etiam in Sibiria et in America boreali. Spec. Africana præter nostra non vidi.

Proles *P. avicularis* habitu proprio sed notis nullis gravioribus distincta, pro subspecie enumeranda.

P. SERRULATUM (*Lag. Gen. et Sp.* 14; *Meisn. in DC. Pr.* xiv. 110),

var. *SALICIFOLIUM*, = *P. salicifolium* (*Delile, Fl. Eg.* 12; *Meisn. l. c.*),
= *P. Pseudohydropiper* (*Salzm. Exsicc.*).

Mar. sept.—Prope Tanger in maritimis! (*Salzm. Exsicc.*)!

Ar. Geog. Species per regionem omnem Mediterraneam diffusa, occurrit etiam in Africa tropica et australi, in Australasia et in insula Java. Varietas nostra hinc inde in regione Mediterranea calidiore.

P. salicifolium distinguitur ocreis apice integerrimis, vel breviter ciliatis, nec ciliis longis rigidiusculis munitis. Forsan pro subspecie enumerandum?

THYMELEÆ (*Adans. Fam. Pl.* ii. 14).

DAPHNE (*L. Gen.* 485).

D. GNIDIUM (*L., Meisn. in DC. Pr.* xiv. 538; *Schousb. Gew. Marok.* 175).

Mar. sept.—Tanger! (*Schsb. l. c.*).

Mar. occid.—Casa Blanea! (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—Ad radices Atlantis Majoris, unde in convalles interiores se protendit. Aït Mesan usque 2000 m.! Ourika (*R. et Fr. spec. in Herb. Kew.*)! In convalle Amsmiz visa!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

D. LAUREOLA (*L., Meisn. in DC. Pr.* xiv. 539).

Mar. merid.—In convalle Aït Mesan a 1600 m. usque 1800 m.! Spec. ex monte Djebel Ouensa ab indigeno Ibrahim lecta benevole comm. eel. Cosson.

— —, var. ? *HOSMARIENSIS*, *nob.* Specimina incompleta die 12 Aprilis 1871 lecta, fructu unico brevissime pedicellato, a typo foliis tenuioribus subtus glaucescentibus diversa sunt. Planta ulterius inquirenda.

Mar. sept.—In monte Beni Hosmar prope Tetuan!

Ar. Geog. D. Laureola per Europam mediam et in montosis Europæ australis et Asiæ Minoris diffusa est. Adest in ins. Azoricis. Specimina Africana præter nostra non vidi.

PASSERINA (*L. Gen.* 487).

P. ARVENSIS (*Lam. Fl. Fr.* iii. 218, et *Meisn. in DC. Pr.* xiv. 551, sub *Thymelæa*), = *P. annua* (*Wikstr. Thymel. in Act. Holm.* 320; *Spreng. Syst.* ii. 239; *Gren. et Godr. Fl. Fr.* iii. 60), = *P. Stellera* (*Coss. et Germ. Fl. Par.* 478), = *Thymelæa Passerina* (*Lange in Willk. et Lge. Fl. Hisp.* i. 298), = *Stellera Passerina* (*L. Sp. Pl.* 512; *Rehb. Ic. Fl. Germ.* No. 1167).

Mar. merid.—Occurrit hinc inde in regione inferiore. Mzonda! Mesfioua (*G. Maw*)! In collibus saxosis pr. urbem Marocco (*G. Maw*)! Reraya (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Europa media. Sibiria Altaica. In regione Mediterranea fere omni (exclusa Macaronesia).

PASSERINA VIRGATA (*Desf. Fl. Atl.* i. 331, tab. 95), = *Thymelæa virgata* (*Meisn. in DC. Pr.* xiv. 552).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Mar. merid.—Ourika! (*R. et Fr. spec. in Herb. Kew.*)!

— — —, var. *BROUSSONNETII*, *nob.* Foliis inferioribus et mediis viridibus glaberrimis, nec, ut in typo, sericeo-villosis incanis.

Vidi spec. ex Imp. Maroecano a Broussonnet lectum in herb. Gouan hodie Kewensi, et simile habuit Desfontaines in 'Fl. Atlantica' l. c. memoratum.

Mar. merid.—In regione inferiore Atlantis Majoris legimus in convalle Aït Mesan circ. 1400 m.! et in convalle Amsmiz!

Ar. Geog. Africa boreali-occidentalis. Spec. Hispanica ex loco non indicato vidit Lange.

P. VILLOSA (*L.*, sub *Daphne*; *Wikstr. Thymel. in Act. Holm.* 1818, 332), = *Thymelæa villosa* (*Endl. Gen. Suppl.*; *Meisn. in DC. Pr.* xiv.), = *Passerina tingitana* (*Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salm. Exsicc.*)! In monte Djebel Kebir!

Ar. Geog. Peninsula Iberica. Ditio Tingitana.

A b. Meisner, l. c., indicatur in Teneriffa fide Bory St. Vincent. In catalogo tamen (in opere 'Iles Fortunées') deest hæc species, nec ab aliis scriptoribus memoratur. Eodem modo planta dubitative "ex Oriente" (*Herb. DC.*) enumerata fere certe erronea.

P. CANESCENS (*Schousb. Gew. Marok.* 176; *Willd. Sp. Pl.* ii. 432), = *Thymelæa canescens* (*Meisn. in DC. Pr.* xiv. 556).

Mar. sept.—"Passim in regione Tingitana" (*Schsb. l. c.*).

Legi in altero latere freti Hereulis prope San Roque, sed in ditioe Tingitana a nobis non visa.

Ar. Geog. Hispania australis. Ditio Tingitana.

P. HIRSUTA (*L.*, *Schousb. Gew. Marok.* 176), = *Thymelæa hirsuta* (*Endl. Gen. Suppl.*; *Meisn. in DC. Pr.* xiv. 557).

Mar. sept.—Tanger (*Schsb. l. c.*). Prope lacum salsum (*Webb Herb.*)!

Mar. centr.—Mequinez (*Schsb. l. c.*).

Mar. occid.—Casa Blanca (*R. et Fr. spec. in Herb. Kew.*)! Aïn el Hadjar haud procul Mogador!

Mar. merid.—Videtur infrequens. Legimus semel ad radices Atlantis Majoris prope Seksaoua!

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

Auctores recentiores duce eel. Endlicher genus Linnæanum *Passerina* ad species duas Capenses limitarunt, et species regionis Mediterraneæ sub genere *Thymelæa* reducerunt (inclusa *Stellera Passerina*, *L.*). Auctor

generis *Thymelæa* dicitur Tournefort in Instit. R. Herbariæ, p. 594. Tournefort tamen genus nullo criterio definivit, et species omnes hodie ad *Daphnem*, *Passerinam*, *Gnidiam* aliaque genera affinia relatas sub *Thymelæa* semel conjunxit, eodem pacto quo Cupani cæterique auctores antiquiores vocem *Thymelæa* adhibuerunt. Ducibus Grenier et Godron ego nomen genericum Linnæanum pro hac grege servavi.

LAURINEÆ (*DC. Théor. Elém.* 247).

LAURUS (*L. Gen.* 503).

L. NOBILIS (*L., DC. Pr.* xv. pars 1, 233; *Schousb. Gew. Marok.* 179).

Mar. sept.—Tanger in dumetis et sylvis (*Schsb. l. c.*). Vidimus in monte Djebel Kebir non certe spontanea! Prope Tetuan ad radices montis Beni Hosmar verisimiliter indigena!

Ar. Geog. In regione fere omni Mediterranea (exclusa Macaronesia) hodie spontanea. Secundum auctores ex Asia Mediterranea in Europam et Africam allata; sententia, ut videtur, valde incerta.

SANTALACEÆ (*R. Br. Prod.* 350).

OSYRIS (*L. Gen.* 1101).

O. ALBA (*L., DC. Pr.* xiv. 633).

Mar. sept.—Inter Tanger et Tetuan! Prope Tetuan!

Mar. merid.—Supra Amsmiz! Ourika (*J. D. H.*)!

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia).

O. LANCEOLATA (*Hochst. et Steud. in Pl. Exsicc. Un. Itin.* 1832; *A. DC. in DC. Pr.* xiv. 633), = *O. quadrifida* (*Salzm. Exsicc.* 1825), = *O. quadripartita* (*Dcne. in Ann. Sc. Nat. sér.* 3, vi. 65, tab. 6).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! In insulis Zafarinis (*Webb spec.*)!

Mar. merid.—Occurrit rarius ad radices Atlantis Majoris. Prope Gurguri! Inter Sektana et Frouga (*G. Maw*)! Ourika (*R. et Fr. spec. in Herb. Kew.*)!

Nomen triviale Decaisneanum primum typis evulgatum, nomen Salzmannianum antiquius; ambo tamen impropria, flores enim, ut jam monuit eel. A. DC. l. c., sæpius trimeri etsi interdum tetrameri.

Ar. Geog. Africa borealis. Peninsula Iberica.

THESIUM (*L. Gen.* 292).

T. HUMILE (*Vahl, Symb.* iii. 43; *DC. Pr.* xiv. 651).

Mar. merid.—In monte supra Seksaoua circ. 1400 m.!

Ar. Geog. Regio Mediterranea calidior (inclusa Macaronesia). In Austria prope Wiener Neustadt (*fide Koch, Syn. Fl. Germ.* ed. 3, 539); an casu aliquo cum seminibus introductum?

ARISTOLOCHIEÆ (*Endl. Gen. Plant.* 114).ARISTOLOCHIA (*L. Gen.* 1022).

A. PISTOLOCHIA (*L., DC. Pr.* xv. pars 1, 485).

Mar. merid.—In convalle Aït Mesan circ. 1600 m. !

Ar. Geog. Régio Mediterranea occidentalis, ex litorali Nieæensi usque Lusitaniam. Oocurrit in Imp. Maroccano et in Sardinia. Ex Algeria non vidi, et deest in Macaronesia.

A. LONGA (*L., Duchartre in DC. Pr.* xv. pars 1, 486), = A. pallida (*Salzm. Exsicc., non Willd.*).

Mar. sept.—Tanger! (*Webb spec.*)! (*Salzm. Exsicc.*)! In monte Djebel Kebir! Inter Tanger et Tetuan!

Ar. Geog. Regio Mediterranea occidentalis (inelusa Macaronesia). Oocurrit etiam rarissime in Gallia occidentali.

Characteres quibus *A. Fontanesii* (Boiss. et Reut. Pug. 108) ab *A. longa* distinguitur mihi videntur omnino fallaces (confer Duchartre, l. c. p. 487). Sistit varietatem perigonio (vel calyce) majore.

A. BÆTICA (*L., Duchartre in DC. Pr.* xv. pars 1, 490), = A. glauca (*Desf. Fl. Atl.* ii. 324, tab. 250).

Mar. sept.—Tanger! Tetuan (*Webb Herb.*)!

Mar. occid.—Mogador (*Lowe spec. in Herb. Kew.*)! Agadir (*Coll. Indig.* 1871)!

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

CYTINEÆ (*A. Brogn. in Ann. Sc. Nat.* i. 29).CYTINUS (*L. Gen.* 1232).

C. HYPOCISTIS (*L. Gen.* 1232).

Mar. sept.—In monte Djebel Kebir prope Tanger in radicibus *Cistorum*!

Mar. merid.—Taradut (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio Mediterranea fere omnis (inclusa Macaronesia). Spec. ex Asia Mediterranea continentali non vidi.

EUPHORBIA CÆÆ (*R. Br. in Flind. Voy., Rem.* 23).EUPHORBIA (*L. Gen.* 609).

E. PEPLIS (*L., DC. Pr.* xv. pars 2, 27).

Mar. sept.—Tanger, folia visa! (*spec. ex Herb. Schousboe comm. Coss.*)!

Mar. occid.—Mazagan!

Ar. Geog. Species in litoribus fere omnibus arenosis regionis Mediterraneæ et Europæ occidentalis obvia.

E. CHAMÆSYCE (*L., Boiss. in DC. Pr.* xv. pars 2, 34).

Mar. occid.—Mazagan!

—— —, var. *CANESCENS* (*Boiss. l. c.*), = *E. canescens* (*L., Cavan. Icon. tab. 63*), = *E. massiliensis* (*Salzm. Exsicc.*)

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

Varietas *canescens* in typum formis intermediis transit.

EUPHORBIA AKENOCARPA (*Guss. Cat. Hort. Boccadif., et Prod. Fl. Sicc. i. 540; Boiss. in DC. Pr. xv. pars 2, 119*), = *E. sphærococca* (*Salzm. Exsicc.*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Ar. Geog. Hispania australis. Ager Tingitanus. Italia meridionalis.

E. RUPICOLA (*Boiss. Elench. No. 174, Voy. Esp. 566, tab. 161, et in DC. Pr. xv. pars 2, 131*), = *L. diffusa* (*L. Duf. in Bull. Soc. Bot. Fr. viii. 445*), = *E. dumetorum* (*Coss. et Dur. in Balansa Pl. d'Algérie, ex Boiss. l. c.*), = *E. squamigera* (*Lois. Fl. Gall. i. 345*).

Mar. sept.—Prope Tanger! et prope Tetuan! legi floriferam nondum fructiferam. Frustulum ex *Herb. Webb.* in monte Djebel Dersa prope Tetuan lectum certe huc referendum!

Ar. Geog. Rarius occurrit in Hispania australi (ad Gibraltar, ubi ipse legi) et alibi, et in Algarbiis et in Africa boreali-occidentali. Semel lecta prope Telonem “casu aliquo adventitia” ex *Boiss. l. c.* Nomen *Loiseleurianum* omnino improprium et negligendum.

Cel. J. D. Hooker specimen nimis immaturum prope Casa Blanca legit *E. rupicolæ* affine, sed vix huic referendum. Planta ulterius inquirenda.

E. PUBESCENS (*Vahl, Symb. ii. 55; Boiss. in DC. Pr. xv. pars 2, 134*).

Mar. sept.—Frequens in humidis et umbrosis. Tanger! Cap. Spartel! Ad lacum salum (*G. Maw*)! Tetuan (1851)!

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—Prov. Shedma! In irrigatis urbis Marocco! Amsmiz!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

E. PTEROCOCCA (*Brot. Fl. Lus. ii. 312; Boiss. in DC. Pr. xv. pars 2, 136*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Cap. Spartel! Inter Tanger et Tetuan!

Ar. Geog. Per regionem Mediterraneam calidiorem imprimis occidentalem (ex insula Zaeyntho usque Macaronesiam) dispersa, sed infrequens. Deest in Gallia et in Italia continentali.

E. HELIOSCOPIA (*L., DC. Pr. xv. pars 2, 136*).

Mar. sept.—Tanger! Cap. Spartel! Tetuan

Mar. merid.—Videtur rara. Specimen unicum in ditione Mesfioua legit J. D. H.!

Ar. Geog. Europa omnis a Lapponia usque fretum Herculis et Bos-

phorum. Asia media usque Japoniam. In regione Mediterranea fere omni, sed fors in calidioribus introducta. In montibus Himalaya. Cum seminibus in America boreali, Africa australi et alibi introducta.

EUPHORBIA EXIGUA (*L., DC. Pr. xv. pars 2, 139*).

Mar. sept.—Frequens in dit. Tingitana necnon in Tetuanensi, sed, ut videtur, omnino deficiens in provinciis meridionalibus. Tanger! (*Webb Herb.*)! Cap. Spartel! Inter Tanger et Tetuan! Tetuan (1851)!

Ar. Geog. Regio omnis Mediterranea. Europa media. Succia australis.

E. INCONSPICUA (*Ball in Journ. Bot. 1875, 204*).

Herbacea, nana, erectiuscula; caulis subsimplex, vel e basi ramosus; umbella pauci- (2-3-)radiata; folia omnia breviter petiolata, inferiora late cuneato-obcordata, basi truncata, apice mucronata, utrinque denticulata, sinu lato emarginata, superiora sensim angustiora, demum oblongo-linearia, acuta vel sæpius apice emarginata tridentata, floralia conformia; glandularum cornua longiuscula; capsula longiuscule pedicellata, lævis, trisulcata; styli tantum ima basi coaliti, divergentes; semina tuberculis acutis exasperata; caruncula substipitata, agoriciformis.

Mar. merid.—Legimus prope Ourika! et spec. unicum prope Seksaoua!

Specimina fructifera ex oasi Akka ad meridiem urbis Tarudant ab indigeno Mardochæo lecta sub "*E. glebulosa*, Coss. et Dur.," benevole communicavit cel. Cosson!

E. glebulosæ (Coss. et Dur. in Bull. Soc. Bot. Fr. iv. 493) vel (forsan errore typographi) *E. globulosæ* (Boiss. in DC. Pr. xv. pars 2, 139) nostra planta proxime affinis est, et post eam fors in sicut subspecies enumeranda, differt tamen notis haud levibus. *E. glebulosa* gaudet enim umbella 3-5-radiata, foliis inferioribus angustis, superioribus latioribus ovato-rhombeis, capsula brevissime nec longiuscule pedicellata. Semina in specim. Algeriensibus *E. glebulosæ* ego vidi semper minute tuberculata, multo minus aspera quam in planta nostra; hæc nota tamen, monente cel. Cosson, variabilis est.

Radix in *E. glebulosa* interdum perennans.

E. FALCATA (*L., Boiss. in DC. Pr. xv. pars 2, 140*).

Mar. occid.—In prov. Haha prope Mogador! In monte Djebel Hadid haud procul Mogador!

Mar. merid.—In prov. Shedma! Mesfioua!

— —, var. RUBRA, *Boiss. l. c.*, = *E. rubra* (*Cavan. Icon. i. tab. 34*).

Mar. occid.—Casa Blanca!

Mar. merid.—Seksoua!

Ut jam movit cel. Boissier, hæc est potius lusus quam vera varietas.

Ar. Geog. Regio Mediterranea omnis (exclusa Macaronesia). Hinc inde occurrit in calidioribus Europæ mediæ et in India boreali-occidentali.

EUPHORBIA PEPLUS (L., *Boiss. in DC. Pr.* xv. pars 2, 141).

Mar. sept.—Tetuan (1851)!

Mar. merid.—In ruderatis urbis Marocco!

Ar. Geog. Europa omnis (exclusa arctica). Regio omnis Mediterranea (inclusa Macaronesia). In America boreali et in Africa australi introducta.

E. peploides (*Gouan, Fl. Monsp.* 174; *Boiss. in DC. Pr.* xv. pars 2, 141).

Mar. merid.—Misra ben Kara! legi etiam ad Gibraltar et prope Alge-ciras.

Valde proxima præcedenti et sicut subspecies enumeranda.

Ar. Geog. Per regionem Mediterraneam hinc inde occurrit. Ex Maca-ronesia non vidi.

E. MEDICAGINEA (*Boiss. Elench.* No. 176, *Voy. Esp.* 569, tab. 162, et in *DC. Pr.* xv. pars 2, 144), = *E. italica* (*Salzm. Exsicc. Tingit.*, non *Lam.*), = *E. latifolia* (*Salzm. Exsicc. Malacitana*, non *C. A. M.*).

Mar. sept.—Satis frequens in ditiorie Tingitana. Tanger! (*Salzm. Exsicc.*)! Inter Tanger et Tetuan! Tetuan (1851)! Inter Tetuan et Centa!

— —, var. *OBLONGIFOLIA*. Folia caulina lineari-oblonga vel oblongo-ovalia, apice truncata, sinu lato emarginata.

Mar. merid.—Legi in provincia Shedma prope castellum præfecti

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

E. PINEA (L., *Boiss. in DC. Pr.* xv. pars 2, 145).

Mar. merid.—Ad radices Atlantis Majoris supra Milhain!

Ar. Geog. Species in maritimis regionis Mediterraneæ late diffusa. In Madera indicata, deest, ut videtur, in ins. Canariensibus.

E. pineæ proxime affines sunt *E. segetalis*, L., regionis Mediterraneæ occidentalis et Europæ centralis civis, et *E. Portlandica* in maritimis Atlanticis Europæ ex Hibernia et Scotia usque Lusitaniam obvia.

Specimina nostra prope Milhain lecta, fere media inter *E. pineam* et *E. segetalem*, priori adduxi, dum radix perennis, vel potius perennans. Planta enim floret anno primo et ex radice interdum (nec semper) pereunante nascitur caulis florifer anno sequente. *E. pinea*, ut mihi videtur, est var. maritima perennans *E. segetalis*, et ab hac sententia cel. Boissier non alienus foret. *E. Portlandica*, pace viri celeb., ab *E. segetali* longius distat radice perenni longævo, caulibus numerosis diffusis, et forma foliorum, et meo sensu subspeciem sistit. Tam *E. pinea* quam *E. segetalis* in insula Madera inveniuntur, quod facile credendum si has pro formis unius ejusdemque speciei agnoscimus.

E. MEGALATLANTICA (*Ball in Journ. Bot.* 1875, 205).

Descriptio ex spec. unico manco :—Perennis (?) e basi ramosus (?) ramis adscendentibus; folia sessilia, integerrima, obverse lanceolata, apice mucronata, reflexa (in spec. nost.); umbella 5-radiata, radiis bis bifidis;

folia floralia rhombeo-orbicularia; cornua glandularum involucri ipsarum latitudinem subæquantia, subelavata; capsula longiuscule pedicellata, in eocceæ dorso papilloso-verrucosa; styli longiusculi, subcoaliti, vix apice fureati; semina nigra, tuberculis minutis in lineas longitudinales dispositis asperula. Pluribus notis inter *E. segtalem* et *E. nicæensem* media, ab utraque capsula, seminibus, catervisque notis diversa. Forsan propior erit *E. cerebrina*, Hochst., mihi ex descript. tantum cognita, sed in hac glandulæ et semina omnino diversa.

Mar. merid.—Ad radices Atlantis Majoris prope Ourika!

EUPHORBIA RIMARUM (*Coss. in Bull. Soc. Bot. Fr.* xx. 261).

Sequitur descriptio Cossoniana:—"Planta perennis, pumila, pallide virens, caudice verticali in radicem fusiformem abeunte. *Caules* numerosi, tenues, *inferne suffruticosi* superne herbacei, basi ramosi, superne simplices vel parce ramosi, diffusi vel ascendentes, puberuli vel glabri. *Folia* minuta, oblongo-obovata vel suborbiculata, *integra*, apice obtusa vel subretusa mucronata, glabra vel puberula, *breviter petiolata*, patula, *floralia subconformia* latiora interdum basi truncata. *Umbellæ sæpius triradiatæ*, radiis longiusculis simplicibus vel bifidis. *Involucri* turbinati lobi ovati emarginati denticulato-ciliati, *glandulæ* transverse oblongæ truncate *cornutæ cornibus tenuibus* glandulæ latitudini subæquilongis parallelis. *Capsula* oblonga, trisuleata, *coccis lævibus* dorso subcarinatis *non alatis*, stylis ima basi tantum coalitis. *Semina* oblongo-obovata subtrigona, nigrescentia, *undique et crebre tuberculato-globulosa* tuberculis primum albidis, caruncula brevi obconico-subglobosa."

Mar. merid.—Occurrit frequentissime in rupibus præsertim in regione inferiore Atlantis Majoris a 500 m. usque 1600 m. s. m. Primus legit Balansa anno 1867 in provincia Haha et in rupibus prope Moulai Ibrahim. Legimus in convalle Ait Mesan! in convalle Amsiniz! supra Seksaoua! Milhaïn! et prope Mskala ad fines prov. Shedma!

Species affines, scil. *E. herniariæfolia*, Willd., *E. deflexa*, Sibth. et Sm., *E. kanaorica*, Boiss., *E. bounophila*, Boiss., et *E. caudiculosa*, Boiss., gaudent omnes seminibus varie foveolatis et insculptis, nec, ut in *E. rimarum*, tuberculatis. Hæ omnes sunt species alpinæ, dum *E. rimarum* est planta rupestris regionis calidioris incola.

E. TERRACINA (L., Boiss. in DC. Pr. xv. pars 2, 157), = *E. heterophylla* (Desf. Fl. Atl. i. 382, tab. 102), = *E. seticornis* (Poir. Voy. ii. 173), = *E. leiosperma* (Sibth. et Sm. Fl. Gr. v. 51, tab. 465), = *E. purpurascens* (Salzm. Exsicc. Tingit.), = *E. leiosperma* (Salzm. Exsicc. Tingit.), forma ab *E. leiosperma*, Fl. Gr. diversa, = *E. verna* (Salzm. Exsicc. Malacitana).

Synonyma alia quindecim a eel. Boissier l. c. citata prætermisi. Species proteiformis in omni territorio Maroccano nobis noto frequens, præsertim in zona litorali.

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! Cap. Spartel! Tetuan! (1851 et 1871)! Inter Tanger et Tetuan! Inter Tetuan et Ceuta

Mar. occid.—Casa Blanca! Mogador! (*Lowe Cat.*). In monte Djebel Hadid! et juxta Aïn el Hadjar haud procul Mogador! Agadir (*Coll. Indig.* 1871)!

Mar. merid.—Mzouda! In prov. Haha! Prope urbem Marocco (*R. et Fr. spec. in Herb. Kew.*)! Pluries visa locis non notatis.

Inter cæteras formas a nobis lectas adsunt spec. foliis crebris lineari-lanceolatis a forma typica valde diversa. Adsunt tamen tot tantæque formæ intermediæ ut varietates distinctæ vix enumerandæ videntur.

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

EUPHORBIA PARALIAS (*L., Boiss. in DC. Pr.* xv. pars 2, 167).

Mar. sept.—Vidi nondum florentem prope Tanger!

Mar. occid.—Mogador! vidi etiam in maritimis prope Saffi!

Ar. Geog. In litoribus marinis totius regionis Mediterraneæ, necnon Europæ occidentalis ab Anglia et Hibernia usque Lusitaniam.

E. CHARACIAS (*L., Boiss. in DC. Pr.* xv. pars 2, 172).

Mar. sept.—In monte Beni Hosmar!

Ar. Geog. Europa austro-occidentalis a Sicilia usque Lusitaniam. Marocco septentrionalis. Occurrit etiam in insula Creta.

Euphorbias cactoides Maroccanas eximie tractavit ecl. Cosson (in *Bull. Soc. Bot. Fr.* xxi. pp. 162 et seq.). Annuente ecl. auctore descriptiones et observationes infra transcripsi, verbo nullo mutato nisi locis natalibus nonnullis additis vel emendatis.

“*De Euphorbiis cactoides Maroccanis.*

“EUPHORBIA sect. DIACANTHIUM (quoad species Maroccanas).

“*Plantæ* perennes, habitu *Cactum referentes*, inferne *indurato-frutescentes* cortice cinereo, *ramis caruosis 3- polygonis* virenti-subglaucescentibus angulis haud undulato-lobatis. *Spinæ stipulares* 2, breves vel elongatæ, *inferne in scutellum confluentes* scutellis in angulis caulibus decurrentibus vel non decurrentibus. *Folia ad tuberculum minutum* cum scutello confluentia et *demum obsoletum reducta*. *Puncta depressa* (puncta gemmipara) *massula cellulosa*, gemmam ramalem vel floriferam emissura *repleta*, ad scutellum sita vel ab illo plus minus remota. Involucra saltem lateralia bracteis binis oppositis suffulta. Bracteolæ inter flores masculos fimbriatæ vel lacerae, sæpe basi connatæ. Glandulæ carnosæ. Capsulæ pericarpium crassum induratum. Semen subglobosum, *ecarunculatum*.

“E. RESINIFERA (*Berg. in Berg. et C.-F. Schmidt, Offiz. Gewächse*, t. 34 d, f, m-x; *Coss. in Bull. Soc. Roy. Belg.* x. 5-12, descriptio manca), = E. officinarum (*Jackson, Acc. Marocc.* ed. 3, p. 134, ex parte, t. 6. fig. sinistra).

“*Planta a basi ramosa*, a 25 centim. ad 1 metr. et ultra alta. *Caules*

paree ramosi, *tetragoni*, rarius *trigoni*, *faciebus subconcavis*, angulis obtusis. *Spinæ stipulares breves*, patentcs, inferne in *scutellum subtriangulanti-ovatum confluentes*, *scutellis* remotiusculis in angulis caulis non decurrentibus. *Puncta gemmipara paulo infra medium spatii interstipularis sita*. Cymæ pedunculatæ, pedunculo crasso ramis cymæ subæquilongo, sæpius 3-floræ, involuero calyciformi medio subsessili, lateralibus pedunculatis. Involucrum calyciforme campanulato-eyathiforme *glandulis carnosis majusculis* transverse oblongo-subrhombeis vel late obovato-euneatis, *aureo-flavis*. *Capsula* majuscula, *depressa profunde tricocca*, *coccis lævibus a latere compressis* sed valde convexis dorso *acute carinatis*. Semen subpapillosum.

“*E. resinifera* inter omnes species sectionis *Diacanthium* *scutellis* hand confluentibus donatas caulibus tri- vel tetragonis insignis.”

Mar. merid.—Ad radices Atlantis Majoris ad austro-orientem urbis Marocco. In ditione Mesfioua vidimus plantam vivam ex collinis vicinis in hortum allatam! Plantam in provincia Demenet satis frequentem ab indigenis usque Mogador allatam in Angliam misit b. Carstensen, et serius spec. altera in Horto Kewensi culta ex montosis prov. Ntifa obtinuit. Specimina in Horto Parisiensi culta ex Ouanyna in ditione Mesfioua attulit indigenus Ibrahim aspice b. Beaumier.

EUPHORBIA BEAUMIERANA (*Hook. f. et Coss. ap. Coss. in Bull. Soc. Bot.* xix. 59 sine descriptione), = *E. officinarum* (*Jackson, Acc. Marocc.* ed. 3, p. 134, ex parte, t. 6. fig. dextra), = *E. officinarum* (*Boiss. in DC. Pr.* xv. pars 2, 84, ex parte).

“Planta a 25 centim. ad 2 metr. alta. *Caulis primarius* oblongo-obovatus demum superne valde incrassatus obpiriformis in longitudinem non excrecens ibique ramos numerosos inæquales ipsum longissime superantes emittens. *Rami* sæpius 9-10goni, sinibus inter costas profundis. *Spinæ stipulares breves* patentcs haud deflexæ, inferne in *scutellum oblongo-lineare confluentes*, *scutellis linealiter in costa decurrentibus*. *Puncta gemmipara ad quintam partem inferiorem spatii interstipularis sita*. Cymæ, flores et fructus ignoti. *Involucri glandulæ intense rubræ* (see. Jackson).”

Mar. merid.—In regione litorali ad meridiem urbis Mogador usque ad Oued Noun; in provinciis Haha et Agadir (*Jackson*); specimina viva in regione litorali inter Mogador et Agadir lecta attulerunt indigeni mense Junio 1871, hodie in Hort. Kew. culta; ad Agadir, ubi montes contegit et sæpe specimina maxima occurrunt, mulio Ibrahim sub auspiciis el. Beaumier legit, unde specimina in Horto Parisiensi culta; in ditione Ksima inter Agadir et Oued Noun legit rabbinus Mardochée.

“*E. Beaumierana* valde affinis *E. officinarum*, sed distincta videtur spinis stipularibus non deflexis, glandulis involucri intense rubris (see. Jackson) non flavidis. Forsan sub *E. officinarum* auctorum species plures adhuc confusæ.

- “Subvar. *fasciata*.—Rami omnes vel plerique fasciati in cristas crassas contortas expansi spuiis stipularibus abortivis brevissimis vel evanidis.
 “Hinc inde cum planta normali in ditione Ksima inter Agadir et Oued Noun (rabbinus Mardochée, 1874).”

EUPHORBIA ECHINUS, (*Hook. f. et Coss. in Bull. Soc. Bot. Fr.* xxi. 164).

- “Planta sæpius 7–30 centim. alta. Caulis primarius superne ramis conformis in longitudinem exerescens, infra apicem ramos numerosos subcorymbosos ipsi subæquilongos vel breviores emittens. Rami sæpius 6goni, sinibus inter costas latiusculis parum profundis. Spinæ stipulares elongatæ validæ patentēs hand deflexæ, inferne in scutellum oblongo-lineare confluentes, scutellis linealiter in costa decurrentibus *Puncta gemmipara* scutello contigua. Cymæ pedunculatæ, pedunculo ramis cymæ subæquilongo, 4–5- rarius 3-floræ, involuero calyciformi medio subsessili masculo, cito deciduo, lateralibus pedunculatis hermaphroditis. Involuerum calyciforme campanulato-cyathiforme, glandulis carnosius minusculis transverse oblongo-subrhombeis, statu sicco vinoso-purpureis, lobis membranaceis suborbiculatis longe exsertis. Capsula parvula, subgloboso-tricocca, coccis lævibus a latere vix compressis dorso carinatis. Semen irregulariter rugulosum.”

Mar. merid.—In regno Maroccoano australiore hand procul a litore : ad Tazeroualt rabbinus Mardochée invenit, et inde specimina Horto Parisiensi a cl. Beaumier missa; in ditione Ba-Ahmran rabbinus Mardochée Augusto 1873 floriferam et fructiferam legit.

- “*E. Echinus* ab *E. Beaumierana* differt caule sæpius hexagono, spinis stipularibus longioribus et præsertim modo crescendi.

“Var. *brevispina*. Spinæ stipulares breves.”

Vir oculatus Jackson, etiamsi rei botanices inexpertus, differentias inter *E. resiniferam* et *E. Beaumieranam* olim observavit et indicavit. Opus Jacksonianum tamen scriptoribus botanicis vix cognitum.

ANDRACHNE (*L. Gen.* 973).

A. MAROCCANA (*Ball in Journ. Bot.* 1875, 205).

Ex collo radicis lignosæ ramosissima, rami tenues, sæpius indivisi, striati, 6–15-pollicares, e basi foliacei; folia anguste lanceolata, breviter petiolata, nitida, viridia, apice cartilagineo mucronata; flores omnes solitarii, masculi minimi, versus apicem ramorum conferti, breviter pedicellati, calycis segmentis membranaceis vix nervosis, late ovatis, subacutis, petalis lineari-cuneatis apice bilobis calyce dimidio brevioribus, disci glandulis (squamis N. ab E.) usque ad basin bipartitis, petalis dimidio brevioribus, filamentis monadelphis; flores fœminei segmentis calycinis foliaceis lanceolatis, acutis, dorso carinatis, carina in pedicello superne incrassato foliis demum æquilongo decurrente, petalis minimis oblongo-cuneatis, apice dentatis, glandulis disci

usque ad basin bipartitis petalis subduplo longioribus; semina fuseo-atra, superficiei granulata aspera.

Mar. oecid.—In monte Djebel Hadid haud procul Mogador!

Mar. merid.—In rupibus supra Seksaoua! et supra Milhaïn!

Primo intuitu ab *A. telephioide* differt foliis multo angustioribus, viridibus, nec glaucescentibus, petalis in flore maseulo apice bilobis, nec lanceolatis integris, calyce et pedicello fl. foeminei fere duplo longioribus, petalis apice dentatis, nec nullis vel minutissimis integris, glandulis disci ter majoribus.

In N. ab E. 'Genera Fl. German.' glandulæ disci petalis alternæ ostenduntur, sed re vera sepalis alternæ et petalis oppositæ sunt.

MERCURIALIS (*L. Gen.* 1125).

M. ANNUA (*L., DC. Pr.* xv. pars 2, 797).

Mar. sept.—Tanger! Tetuan! Inter Tetuan et Ceuta!

Mar. oecid.—Mogador (*Lowe Cat.*).

Mar. merid.—Amsmiz! Seksaoua!

—— —, var. *AMBIGUA* (*Duby, Bot. Gall.* i. 417), = *L. ambigua*, *L.*

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. merid.—In ruderatis urbis Marocco! Mzouda!

—— —, var. *SERRATIFOLIA*, foliis profunde et acute serratis.

Mar. sept.—In monte Beni Hosmar prope Tetuan!

Ar. Geog. Regio omnis Mediterranea. In Europa media occurrit passim in ruderatis et cultis, sed verisimiliter olim introducta.

RICINUS (*L. Gen.* 503, *ed. Schreb.* 1464).

R. COMMUNIS (*L., Müll. Arg. in DC. Pr.* xv. pars 2, 1817).

Mar. sept.—Tanger!

Mar. oecid.—Mogador (*Lowe Cat.*).

Cl. Müller Argoënsis *Ricinos* omnes ad unam speciem reduxit, et quoad species sic diet. Mediterraneas libenter assentio. Planta verisimiliter ex Africa tropica orta, olim culta in regione Mediterranea (non indigena), valde variabilis est. Formas e cultura enatas vel subspontaneas botanici perperam sicut species distinctas enumerarunt.

URTICACEÆ (*Endl. Prod. Norf.* 37).

URTICA (*L. Gen.* 1054).

U. URENS (*L., Wedd. in DC. Pr.* xvi. pars 1, 40).

Mar. oecid.—Mogador (*Lowe Cat.*).

Ar. Geog. In temperatis orbis veteris latissime diffusa, rarior in tropicis et subtropicis. In Americam introducta, ibi hodie spontanea.

U. PILULIFERA (*L., Wedd. in DC. Pr.* xvi. pars 1, 48).

Mar. sept.—Certe visa pr. Tanger, sed spec. non conservata!

Mar. merid.—In ruderatis urbis Marocco visa !

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia). In Europa media satis frequens sed verisimiliter introducta. Indicatur in Asia tropica et in insula Sanctæ Helenæ.

URTICA DIOICA (L., *Wedd. in DC. Pr.* xvi. pars 1, 50), var. PROCERA (*Wedd. l. c.* 52), = U. gracilis (*Ait. Hort. Kew.* iii. 341), see Weddell l. c. ; sed planta Aitoniana ex America subaretica juxta sinum Hudson orta est.

Mar. merid.—In irrigatis urbis Marocco !

Ar. Geog. Species in temperatis imprimis frigidioribus Europæ et Asiæ vulgarissima, in calidioribus regionis Mediterraneæ infrequens, nec in Macaronesia indicatur. In America fere tota diffusa, dicitur introducta sed occurrunt varietates plurimæ etiam in regione montana Americæ centralis et australis. Var. *procera* in America boreali et in Algeria obvia est.

U. MEMBRANACEA (*Poir. Dict.* iv. 638 ; *Wedd. in DC. Pr.* xvi. pars 1, 56).

Mar. sept.—Tanger ! (*Salzm. Exsicc.*) !

Mar. occid.—Juxta Mazagan visa ! In insula Mogador !

Ar. Geog. Regio Mediterranea imprimis occidentalis (inclusa Macaronesia), nec deest in insulis Azoricis. Ex Asia Mediterranea spec. nulla vidi. In Gallia occidentali (sec. Weddell l. c., deest tamen in Gren. et Godr.) verisimiliter introducta.

PARIETARIA (L. *Gen.* 1152).

P. OFFICINALIS (L., *Wedd. in DC. Pr.* xvi. pars 1, 235 [42]).

Mar. sept.—Tanger ! Cap. Spartel !

Mar. occid.—Mogador (*Lowe spec. in Herb. Kew.*) !

Specimina nostra sub var. *diffusa* (Wedd. l. c.), = *P. diffusa* (M. et K. Deutschl. Fl. i. 827) enumeranda—caules tamen sæpe suberecti.

Ar. Geog. Regio omnis Mediterranea. Europa omnis exclusa arctica.

P. MAURITANICA (*Durieu in Duchartre, Rev. Bot.* ii. 427 ; *Wedd. in DC. Pr.* xvi. pars 1, 235 [45]), var. DIFFUSA (*Wedd. l. c.*).

Mar. sept.—In monte Beni Hosmar prope Tetuan !

Mar. merid.—Prope Amsmiz !

Ar. Geog. Africa borealis. Peninsula Iberica. In monte Clarence Africæ æquatorialis.

FORSKOHLEA (L. *Amœn. Acad.* vii. 72).

F. TENACISSIMA (L., *Wedd. in DC. Pr.* xvi. 235 [55]).

Mar. merid.—In collibus saxosis ab urbi Marocco septentrionem versus legit cl. G. Maw, et spec. communicavit ! Prope urbem Marocco (*R. et Fr. spec. in Herb. Kew.*) !

Specimina nostra inter typum et *F. Cossonianam* (Webb, *Otia* *Hisp.* 49, tab. 44), a el. Weddell optimo consilio ad varietatem reductam, fere media.

Ar. Geog. Species imprimis in regione subtropica desertorum ex India occidentali usque Algeriam interiorem diffusa; insuper in Marocco meridionali obvia. Var. *Cossoniana* unico loco in Hispania austro-occidentali hucusque detecta.

ULMACEÆ (*Mirb. Elém.* 905).

ULMUS (*L. Gen.* 316).

U. CAMPESTRIS (*Sm. Eng. Fl.* ii. 20; *Planch. in DC. Pr.* xvii. 156)?

Mar. sept.—Frustulum foliaceum prope El Araïsch lectum attulerunt ell. Rein et Fritsch verisimiliter huc referendum. An sponte creseit?

Ar. Geog. Species valde ludibunda, imprimis per Europam et Asiam temperatam late diffusa. In regione tota Mediterranea hinc inde occurrit, interdum fruticeta in collibus ineultis efformans, interdum arborea, plane introducta et culta.

CELTIS (*L. Gen.* 1143).

C. AUSTRALIS (*L., Planch. in DC. Pr.* xvii. 169).

Mar. sept.—Ad radices montis Beni Hosmar pr. Tetuan visa!

Mar. merid.—Sheshaoua! Grigory (*R. et Fr. spec. in Herb. Kew.*)! an eodem loco qui nobis Gurguri scribitur?

Ar. Geog. Regio fere omnis Mediterranea inclusa Madera sed exclusis insulis Canariensibus. Nascitur etiam in calidioribus Galliae mediæ et Hungariæ australis.

CUPULIFERÆ (*Rich. Anal. Fr.* 33).

QUERCUS (*L. Gen.* 1070).

Q. HUMILIS (*Lam. Dict.* i. 719; *A. DC. in Prod.* xvi. pars 2, 16).

Mar. sept.—In monte Djebel Kebir pr. Tanger! (*Webb, ex A. DC. l. c.*). (*Salzm. Exsicc.* sub “*Q. lusitanica*”)!

Ar. Geog. Peninsula Iberica. Africa Tingitana.

Q. LUSITANICA (*Lam. Dict.* i. 719; *Webb, It. Hisp.* 11; *A. DC. in Prod.* xvi. pars 2, 17).

Mar. sept.—In agro Tingitano vulgatissima (*Webb, l. c.*)! Djebel Kebir!

Ar. Geog. Peninsula Iberica. Africa Tingitana. Var. in Asia Mediterranea late diffusa.

Q. Mirbeckii (*Durieu in Duchartre, Rev. Bot.* ii. 426), = *Q. lusitanica* subspecies *bætica* (*A. DC. l. c.* 19), = *Q. lusitanica*, var. *bætica* et var. *Salzmanniana* (*Webb, It. Hisp.* 12).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! In monte Djebel Kebir!

Ar. Geog. Peninsula Iberica. Africa borealis. Var. in Oriente.

Q. infectoria (*Oliv. Voy.* i. 252, tab. 14 et 15), planta orientalis cujus

lusus vel varietates sunt species Kotschyanae *Q. polycarpus*, *Q. brachycarpa*, *Q. Pfæffingeri*, et *Q. Cypriacæ* non *Q. Boissieri* (Reut. in Boiss. Diagn. Pl. Or. ser. 1, xii. 119), est mere varietas *Q. lusitanicæ* a planta occidentali ægre distinguenda. *Q. Mirbeckii* sicut subspeciem enumeravi, forsitan tamen pro varietate melius habenda.

QUERCUS ILEX (L., A. DC. in Prod. xvi. pars 2, 38).

Mar. merid.—Frequens in regione inferiore Atlantis Majoris, plerumque inter 1000 m. et 1500 m. s. m. Tasseremout! Ourika! Reraya! Amsmiz! pluries alibi visa!

— —, var. BALLOTA (A. DC. l. c.), = *Q. Ballota* (Desf. Fl. Atl. ii. 350; Webb, It. Hisp. l. c.).

Mar. sept.—In monte Djebel Dersa prope Tetuan (Webb, l. c.).

Mar. merid.—In monte Djebel Tezah a 2400 m. usque 2700 m. s. m.!

Arbores sparsæ sat magnæ reliquæ ut videtur, sylvarum clivos Atlantis olim late vestientium, hodie incuria incolarum dirutarum.

Ar. Geog. *Q. Ilex* per regionem Mediterraneam (exclusa Macaronesia) late diffusa, occurrit etiam in Gallia occidentali. Var. *Ballota* peninsula Ibericam et Africam borealem incolit.

Q. COCCIFERA (L., A. DC. in Prod. xvi. pars 2, 52).

Mar. sept.—In monte Djebel Kebir! Spec. ex herb. Schousboe benev. comm. cel. Cosson!

— —, var. IMBRICATA (A. DC. loc. cit.), = *Q. pseudococcifera* (Webb, It. Hisp. 15, non Desf. Fl. Atl. ii. 349, sec. A. DC.).

Mar. sept.—In monte Djebel Kebir prope Tanger (Webb, l. c.).

Hinc verisimiliter spec. prope Tetuan in monte Djebel Dersa lectum (sec. A. DC. l. c.) a beato Webb ad *Q. pseudococciferam*, Labill., non Desf., relatum. Hæc novissima, = *Q. calliprinos* (Webb, It. Hisp. 15; A. DC. in Prod. xvi. pars 2, 54), est species regionis Mediterraneæ orientalis propria et haud facile in ditioe nostra invenienda.

Ar. Geog. *Q. coccifera* regionem fere omnem Mediterraneam incolit, frequens in occidentalibus fit rarior orientem versus. Deest cum congeneribus (immo cum *Cupuliferis* omnibus) in Macaronesia. In provinciis Maroccanis meridionalibus hucusque non visa.

Quercus regionis Mediterraneæ omnes quam maxime ludibundæ a cel. A. DeCandolle in 'Prodromo' eximie tractatæ, adhuc studio diligente et diuturna observatione in vivo egent.

Vidimus prope Tanger et Tetuan *Castaneam vulgarem*, Lam., certe cultam. Spontaneam vel etiam subsponsaneam in Imperio Maroccoano nullibi vidimus.

SALICINÆ (Rich. Élém. ed. 4, 560).

SALIX (L. Gen. 1098).

S. ALBA (L., Anders. in DC. Pr. xvi. pars 2, 211).

Mar. sept.—Tanger (Webb Herb.)!

Ar. Geog. In Europa et Asia temperata late diffusa et per totam aream regionis Mediterraneam (exclusa Macaronesia) dispersa. Sec. cl. Andersson l. c. ex Asia versus occidentem migrata.

SALIX CANARIENSIS (*Chr. Smith, sec. Webb, Phyt. Canar.* iii. 270, tab. 215; *Anders. in DC. Pr.* xvi. pars 2, 216)?

Mar. merid.—In convalle Ourika circ. 1100 m., sine flore et fructu et ideo prorsus incerta!

Spec. nostra ex ramis sterilibus desumpta Canariensibus omnino conformia sunt.

Ar. Geog. Species hucusque Macaronesiæ propria.

S. PEDICELLATA (*Desf. Fl. Atl.* ii. 362; *Anders. in DC. Pr.* xvi. pars 2, 216).

Mar. sept.—Legimus in monte Djebel Kebir! ex eodem loco habui spec. ex herb. Webb!

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis. Calabria. Sicilia. Corsica.

S. PURPUREA (*L., Anders. in DC. Pr.* xvi. pars 2, 306), var. *HELIX* (*Anders. l. c.*), = *S. Helix* (*L., et auct. plurim.*).

Mar. merid.—In regione inferiore Atlantis Majoris. Ourika! Aït Mesan!

Ar. Geog. In temperatis Europæ et Asiæ late diffusa. In regione Mediterranea rarior, sed in tota area excepta Macaronesia dispersa.

POPULUS (*L. Gen.* 1123).

P. ALBA (*L., Wesmael in DC. Pr.* xvi. pars 2, 324).

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—Prope Milhain! alibi pluries visa; arbor medioeris altitudinis 10–12 m., typo Europæo conformis.

——, var. *INTEGRIFOLIA*, *nob.* Foliis ovato-orbicularibus, junioribus dentatis, demum subintegerrimis leviter angulato-sinuatis.

Mar. merid.—Crescit copiose in convalle Aït Mesan usque 1400 m. s. m. fructifera lecta! Hujus forma foliis minimis late ovatis acutiusculis, sine flore et fructu in convalle Ourika lecta! Arbor parva 5–7 m., ulteriore observatione egens.

Ar. Geog. Planta in temperatis præsertim calidioribus Europæ, Asiæ et Africæ borealis late diffusa. Deest in Macaronesia.

P. NIGRA (*L., Wesmael in DC. Pr.* xvi. pars 2, 327).

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Aït Mesan! Prope Ourika!

Ar. Geog. Europa et Asia temperata. Var. in America boreali. Præter nostra spec. Africana non vidi.

Specimina nostra flore et fructu carentia haud omnino certa, sed vix ad aliam speciem referenda.

CERATOPHYLLEÆ (*Gray, Arr. Br. Pl.* ii. 534).CERATOPHYLLUM (*L. Gen.* 1055).

C. DEMERSUM (*L., DC. Pr.* iii. 73).

Mar. merid.—In urbe Marocco vidimus subexsiccatum in fasciculo ex prov. Demenet allato et frustula servavimus!

Ar. Geog. Per Europam fere omnem et regionem Mediterraneam late diffusa. In Macaronesia nondum detecta.

GNETACEÆ (*Blume, Nov. Fam. Expos.* 23).EPHEDRA (*L. Gen.* 1136).

E. ALTISSIMA (*Desf. Fl. Atl.* ii. 371, tab. 253; *Parl. in DC. Pr.* xvi. pars 2, 356), est verisimiliter *E. fragilis* (*Lowe Cat.*, non *Desf.*).

Mar. occid.—In dunetis et fruticetis prov. Haha prope Mogador! Mogador (*Lowe Cat.*, sub *E. fragili*)?

Mar. merid.—Satis frequens in regione inferiore. Vidimus in prov. Haha! Shedma! Legimus juxta Sheshaoua! Tasseremont! In convalle Aït Mesan usque 1500 m.!

— —, var. NANA? Planta humilis subbipedalis ramis intricatis.

Mar. merid.—In saxosis aridis supra Milhaïn!

Spec. nostra flore et fructu carentia nimis incerta, sed ex modo crescendo *E. altissimæ* proxime affinia.

Planta in regione inferiore alte et late scandens ramis 4–5 metra longis.

Ar. Geog. Africa borealis ex Alexandria usque Mogador. Hispania. Ins. Canarienses.

E. PROCERA (*Fisch. et Mey. Ind. Sem.* 1844, 45; *Parl. in DC. Pr.* xvi. pars 2, 358), = *E. græca* (*C. A. M. Ephedr.* 93, tab. 5; *Coss. Index Pl. Mar.* 17).

Mar. merid.—Legi sine fl. et fr. in convalle Aït Mesan circ. 1400 m.!

Spec. ex montibus Djebel Ouensa et Djebel Afougueur ab indigeno Ibrahim lecta benevole comm. cel. Cosson!

Ar. Geog. Regio Mediterranea orientalis. Marocco. Ex Algeria non vidi.

CONIFERÆ (*Juss. Gen.* 411).PINUS (*L. Gen.* 1077).

P. HALEPENSIS (*Mill. Dict.* No. 8, tab. 216; *Parl. in DC. Pr.* xvi. pars 2, 383).

Mar. merid.—In Atlante Majore, ut videtur, rara. Legimus in convalle Anismiz a 1300 m. usque 1700 m.!

Arbores sparsæ mediocri magnitudine.

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

PINUS CEDRUS (L., *Parl. in DC. Pr.* xvi. pars 2, 407), var. **ATLANTICA** (*Parl. l. c.*), = *P. atlantica* (Endl. *Conif.* 137), = *Cedrus atlantica* (Manetti, *Cat. Hort. Madoet. Suppl.* 8, ex *Parl. l. c.*).

Mar. sept.—Ex montibus ditionis Riff ab urbe Tetuan meridiem versus indigena quidam attulit et beato Webb dedit. Frustulum ex herb. Webb in herb. nostro!

Mar. centr.—? Verisimiliter in montibus ab urbe Fez ad austro-orientem, a Gerhard Rohlfis viatore indefesso sed rei botanices inexperto, pro *Larice* indicata.

Ar. Geog. Species in montibus Asiæ Mediterraneæ, et in montibus Himalaya. Var. *atlantica* in Atlante Minore.

CALLITRIS (*Vent. Decad.*)

C. QUADRIALVIS (*Vent. Decad.* p. 10; *Parl. in DC. Pr.* xvi. pars 2, 452), = *Thuja articulata* (Vahl, *Symb.* ii. 96; *Desf. Fl. Atl.* ii. 352, tab. 252).

Mar. sept.—Prope Tetuan (1851)! (*Webb Herb.*)!

Mar. occid.—Prope Mogador, præsertim versus montem Djebel Hadid! (*Lowe Cat.*).

Mar. merid.—In regione inferiore Atlantis Majoris satis frequens, sed non ubique. Ourika! Seksaoua! et alibi visa.

Ar. Geog. Africa borealis.

Arbor medioeris 5–10 m. alta, indigenis *Arar* dicto, lignum pretiosum, et resinam *sandraccam* præbens, incuria indigenarum sæpe cruta vel igne consumpta, adeo ut rarius truncos ultra 30 centim. diametro invenies.

JUNIPERUS (*L. Gen.* 1134).

J. OXYCEDRUS (L., *Parl. in DC. Pr.* xvi. pars 2, 477).

Mar. merid.—Satis frequens, præsertim ad radices Atlantis Majoris. Amsmiz! Tasseremout! In convalle Aït Mesan usque 1800 m.!

Ar. Geog. Regio fere omnis Mediterranea (inclusa Madera, exclusis tamen ins. Canariensibus).

J. macrocarpa (*Sibth. et Sm. Prod. Fl. Gr.* ii. 263; *Parl. in DC. Pr.* xvi. pars 2, 476).

Mar. sept.—Vidi pr. Tanger! et spec. ex herb. Schousboe benev. comm. Cosson!

Mar. merid.—Legi in convalle Ourika formam huic et *P. Oxycedro* fere intermediam!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

Stirps *J. Oxycedro* nimis proxima—pro subspecie enumerata, sed forsanelius pro varietate habenda.

J. PHOENICEA (L., *Parl. in DC. Pr.* xvi. pars 2, 486).

Mar. sept.—Inter Tetuan et Ceuta!

Mar. occid.—Rabat (*spec. in Herb. Webb*)! Mogador! Agadir (*Coll. Indig.* 1871)! In monte Djebel Hadid haud procul Mogador (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—In regione inferiore Atlantis Majoris. Ourika! In convalle Aït Mesan! supra Amsmiz circ. 1300 m. s. m.!

In Atlante interdum fit arbor sat magna truncum usque 60 centim. diametro attingens. In planta juniore folia omnia sæpius libera.

— —, var. GALBULIS MOLLIBUS, = *J. lycia*, *L.*

Mar. merid.—Tasseremout! In convalle Aït Mesan usque 1500 m.!

Sec. Parlatores l. c. galbuli immaturi in *J. phænicea* duri, maturi semper molles. Ego vidi in forma vulgari galbulos etiam omnino maturos duros, et formam galbulis mollibus sicut varietatem enumeravi.

Ar. Geog. Species per regionem omnem Mediterraneam (inclusa Macaronesia) late diffusa. Sæpius occurrit in zona litorali ubi fruticem 4–8-pedalem efficit. In Atlante et in montibus ins. Canariensium fit arbor mediocris 30-pedalis.

JUNIPERUS THURIFERA (*L., Parl. in DC. Pr.* xvi. pars 2, 487).

Mar. merid.—In convalle Aït Mesan et in ascensu jugi Tagherot a 2400 m. ad 2900 m.! Arborea solitaria truncum diametro usque 70 centim. attingentes zonam latam in clivo montium altissimarum effingunt, reliquæ ut videtur sylvarum hominum malitia et cœli intemperie devastatarum.

Ar. Geog. Species in peninsula Iberica et in Africa boreali-occidentali sat rara.

ORCHIDEÆ (*Juss. Gen.* 64).

ORCHIS (*L. Gen.* 1009).

O. PAPILIONACEA (*L., Rehb. Icon.* xiii. tab. 40).

Mar. sept.—Inter Fondak et Tetuan!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

O. LONGICRURIS (*Link, in Schrad. Journ.* ii. 323; *Rehb. Icon.* xiii. tab. 23), = *O. undulatifolia* (*Biv. Bern. Cent.* ii. 44, tab. 6), = *O. tephrosanthos* (*Desf. Fl. Atl.* ii. 319, non *Vill.*), = *O. militaris* (*Poir., non L.*).

Mar. sept.—In monte Beni Hosmar prope Tetuan! (*Webb Herb.*)!

Ar. Geog. Regio Mediterranea calidior (exclusa Macaronesia).

O. TRIDENTATA (*Scop. Fl. Carn.* ed. 2, ii. 190), var. *ACUMINATA* (*Rehb. fil. in Icon.* xiii. tab. 18), = *O. acuminata* (*Desf. Fl. Atl.* ii. 318), = *O. lactea* (*Poir.*), = *O. globosa* (*Brot. non L.*).

Mar. sept.—In monte Beni Hosmar prope Tetuan!

Ar. Geog. Species in Europa media et australi. Var. *acuminata* in regione Mediterranea imprimis occidentali (exclusa Macaronesia).

ORCHIS MASCULA (L., *Rchb. Icon.* xiii. tab. 38).

Mar. sept.—In monte Beni Hosmar pr. Tetuan! (*Webb spec.*)!

Folia observavit eel. Webb maculata; in sp. nostris immaculata sunt.

Ar. Geog. Europa omnis temperata. Regio Mediterranea fere omnis (exclusa Maearonesia).

O. LATIFOLIA (L., *Rchb. Icon.* xiii. tab. 50 et 51).

Mar. merid.—In convalle Aït Mesan supra Arround (*R. et Fr. spec. in Herb. Kew.*)!

———, var. DURANDII, nob., = O. Durandii (*Boiss. et Reut. Pug.* 111).

Mar. sept.—In humidis prope Tanger! (*Durand, in Herb. Pavon, ex Boiss. et Reut.*). (*Webb Herb.*)! Juxta lacum salsum (*G. Maw*)!

Mar. merid.—In convalle Ourika circ. 1000 m. s. m. legimus specimina quadripedalia et ultra!

Ar. Geog. O. *latifolia* per Europam omnem (exclusa arctica) diffusa, in Africa boreali et insula Madera occurrit. Var. *Durandii* statura maxima et bracteis longissimis insignis, in Imp. Maroecano et in Hispania (monente Boiss. et Reuter) obvia est.

Flores in spec. nostris Atlanticis quam in Tingitanis minores. Formam hanc sicut varietatem O. *incarnatæ*, L., enumerarunt Willkomm et Lange in 'Fl. Hisp.' i. 170; Duce cl. H. G. Reichenbach *Orchidearum* sagaeissimo scutatore O. *latifoliae* conjunxi.

ACERAS (*R. Br. in Hort. Kew.* ed. 3, v. 191).

A. ANTHROPOPHORA (L., sub *Ophryde*; *R. Br. l. c.*; *Rchb. Icon.* xiii. tab. 5).

Mar. sept.—In monte Beni Hosmar prope Tetuan!

Mar. merid.—Ad radices Atlantis Majoris. Spec. unicum prope Tasscremout legit G. Maw!

Ar. Geog. Regio fere omnis Mediterranea (exclusa Maearonesia). In Europa occidentali hinc inde infrequens.

A. INTACTA (*Link, in Schrad. Journ.* 1799, sub *Orchide*; *Rchb. Icon.* xiii. 148), = A. *densiflora* (*Brot. Fl. Lus.* i. 22, sub *Satyrio*; *Boiss. Voy. Esp.* 595), = A. *secundiflora* (*Bertol. Pl. Rar. Ital.* ii. 42, sub *Orchide*; *Lindl. Bot. Reg.* tab. 1525), = *Orchis atlantica* (*Willd. Sp. Pl.* iv. 42), = *Tinea cylindrica* (*Biv. Bern.*; *Todaro, Orch. Sic.* 8).

Mar. sept.—In monte Beni Hosmar prope Tetuan specimen unicum legit J. D. H.!

Ar. Geog. In regione Mediterranea imprimis occidentali. Deest tamen in Maearonesia.

A. PYRAMIDALIS (L., sub *Orchide*; *Rich.*, sub *Anacamptide*; *Rchb. fil. in Rchb. Icon.* xiii. tab. 9).

Mar. occid.—In monte Djebel Hadid haud procul Mogador (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—Ad radices et in regione inferiore Atlantis Majoris. In convalle Aït Mesan! Seksaoua! Milhaïn! Mtouga (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. In Europa temperata late diffusa, in calidioribus regionis Mediterraneæ infrequens. Deest in Macaronesia.

PLATANThERA (*Rich. Mus. Par.* iv. 20).

P. DIPHYLLA (*Link, in Schrad. Journ.* 1799, sub *Satyrio*; *Rchb. fil. in Rchb. Icon.* xiii. tab. 84), = *Habenaria cordata*, *R. Rr.*, = *Peristylus cordatus*, *Lindl.*, = *Orchis cordata* (*Willd. Sp. Pl.* iv. 27).

Mar. sept.—In umbrosis collinis prope 'Tanger! (*Salzm. Exsicc.*)! In monte Djebel Kebir! (*Webb Herb.*)! Cap. Spartel!

Ar. Geog. Africa boreali-occidentalis. Macaronesia. Peninsula Iberica australis. Fertur etiam in Sardinia.

OPHRYS (*L. Gen.* 1011).

O. LUTEA (*Cavan. Icon.* ii. 46, tab. 160; *Rchb. Icon.* xiii. 94).

Mar. sept.—In umbrosis collinis inter Tanger et Tetuan! In monte Beni Hosmar prope Tetuan! (*Webb Herb.*)!

Ar. Geog. Per regionem Mediterraneam (exclusa Macaronesia) late dispersa.

O. SPECULUM (*Link, in Schrad. Journ.* 1799; *Rchb. Icon.* xiii. 96), = *O. Scolopax* (*Brot. non Cavan.*).

Mar. sept.—In collinis inter Tanger et Tetuan!

Ar. Geog. Distributio fere eadem ac *O. lutea*, cui nimis proxima, et forsàn cum *O. fusca* in unam speciem aggreganda.

O. BOMBYLIFLORA (*Link, in Schrad. Journ.* 1799; *Coss. Pl. Crit.* 64; *Rchb. Icon.* tab. 104), = *O. tabanifera* (*Willd. Sp. Pl.* iv. 68), = *O. picta* (*Salzm. Exsicc.*).

Mar. sept.—In monte Djebel Kebir! Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Regio Mediterranea calidior (inclusa Macaronesia). Deest in Gallia et in Italia superiore.

O. APIFERA (*Huds. Fl. Angl.* 340; *Rchb. Icon.* xiii. 105).

Mar. sept.—Prope Tanger!

Mar. merid.—Ad radices Atlantis Majoris. In convalle Aït Mesan circ. 1300 m.! Ourika (*J. D. H.*)! Reraya (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Europa media et australis. Africa borealis. Ex Asia Mediterranea non vidi.

O. SCOLOPAX (*Cavan. Icon.* ii. 46, tab. 161; *Rchb. Icon.* xiii. tab. 106).

Mar. sept.—Inter Tanger et Tetuan! Prope Tetuan!

Mar. merid.—In ditone Reraya circ. 1100 m.!

Ar. Geog. Regio fere omnis Mediterranea (exclusa Macaronesia).

OPHRYS TENTHREDINIFERA (*Willd. Sp. Pl.* iv. 67; *Rehb. Icon.* xiii. 111).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Regio Mediterranea imprimis calidior (exclusa Macaronesia).

In Oriente fit rarior.

CEPHALANTHERA (*Rich. Orchid. Eur.* 21).

C. XIPHIOPHYLLUM (*L. fil. Suppl.*, sub *Serapiade*; *Rehb. Icon.* xiii. tab. 118), = C. ensifolia (*Swartz*, sub *Epipactide*; *Rich. l. c.*).

Mar. sept.—In monte Beni Hosmar prope Tetuan spec. unicum legit J. D. II.!

Ar. Geog. In Europa omni (exclusa aretica) late diffusa. In regione Mediterranea fere omni (exclusa Macaronesia) hinc inde occurrit.

SERAPIAS (*L. Gen.* 1012).

S. LINGUA (*L.*, *Rehb. Icon.* xiii. tab. 87).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Inter Tetuan et Ceuta!

Ar. Geog. Per regionem fere omnem Mediterraneam (exclusa Macaronesia) late dispersa.

S. CORDIGERA (*L.*, *Rehb. Icon.* xiii. 88).

Mar. sept. Tanger (*spec. ex Herb. Schousboe comm. Cosson*)! Inter Tetuan et Ceuta!

Ar. Geog. Per regionem Mediterraneam (exclusa Macaronesia) late diffusa.

Orchideæ in Imperio Maroecano nobis cognito paucae et raræ, in prov. australioribus perpaucae. Specimina nostra omnia sub oculis habuit clar. II. G. Reichenbach et quæ nobis incerta fuerunt determinavit.

IRIDEÆ (*Juss. Gen.* 57).

GLADIOLUS (*L. Gen.* 57).

G. SEGETUM (*Gawl. in Bot. Mag.* tab. 719; *Klatt in Linnæa*, xxxii. 701), = G. communis (*Desf. Fl. Atl.* i. 35; *Fl. Gr.* tab. 37; *Schousb. Gew. Marok.* 14, non *L.*), = G. byzantinus (*Baker MSS.*, an *Mill.*?).

Mar. sept.—Tanger (*Webb spec.*)! (*Schsb. l. c.*). Cap. Spatel! Prope Tetuan!

Ar. Geog. Species per regionem Mediterraneam (inclusa Macaronesia) late diffusa.

G. ILLYRICUS (*Koch in Sturm Deutschl. Fl.* 83, et *Syn. Fl. Germ.* 699), = "G. byzantinus?" (*Coss. Index Pl. Maroc.* 28, non *Mill.*).

Mar. occid.—In monte Djebel Hadid haud procul Mogador (*R. et Fr. spec. in Herb. Kew.*)! Ex eodem monte spec. ab indigena Ibrahim lectum comm. cel. Cosson!

Ar. Geog. Europa Mediterranea a Dalmatia usque fretum Herculis. Ditio Mogadorensis.

Proxime affinis *G. communi*, L., per Europam mediam et australem inter segetes obvio, et sicut subspecies enumerandus.

XYPHION (*Parl. Nuov. Gen. e Spec. Monocot.* 43).

X. FILIFOLIUM (*Boiss. Voy. Esp.* 602, tab. 170, sub *Iride*; *Klatt in Linnæa*, xxxii. 571; *Baker in Journ. Bot.* ix. 14).

Mar. sept.—“In fissuris rupium montis Beni Hosmar prope Tetuan” (*spec. ex Herb. Webb*)! Inter paludes salsas et El Araisch coll. cl. G. Maw et spec. benevole communicavit!

— — —, var. **INTERMEDIUM**, *Baker, MSS.*, = *X. tingitanum* (*H. fil. in Bot. Mag.* tab. 5981; non *Boiss. et Reut.*, sub *Iride*).

Mar. sept.—Nascitur juxta paludes salsas ab urbe Tanger austro-occidentem versus, unde spec. nobis dedit Eques Drummond Hay legatus Britannicus!

Flores in hac forma pulcherrimi eis *X. tingitani* similes sed majores; foliis tamen filiformibus *X. filifolio* accedit.

Ar. Geog. *X. filifolium* in Africa boreali-occidentali et in peninsula Iberica australiore obvium est.

X. tingitanum (*Boiss. et Reut. Pug.* 113, sub *Iride*; *Baker in Journ. Bot.* ix. 13), = *Iris Xyphium* (*Schousb. Gew. Marok.* 15, nec aliorum)?

Mar. sept.—“In arvis argillosis circa Tingidem” (*Boiss. et Reut. l. c.*). “In fruticetis regionis Tingitanæ” (*Schousb. l. c.*). Inter Tanger et Tetuan nubios legit cl. Maw et in horto educavit!

A præcedente differt foliis latioribus, late nec arete canaliculatis; ut mihi videtur pro subspecie enumerandum.

Iris Xyphinum apud Schousboe, sicut et apud ipsum Linnæum, est forsitan nomen collectivum plurimas species et variatates complectens. Ccl. Cosson mihi dedit ex herbario Schousboe specimina *Iris juncea*, Poir., inter frutices montis Djebel Kebir lecta, cui “folia canaliculato-subulata” potius quam *I. tingitanæ*, Boiss. et Reut., tribuenda. Cæterum *I. juncea* (*Xyphion*, Parl. l. c.) est a *X. tingitano* probe distincta floribus luteis dimidio minoribus et foliis rectis pluries brevioribus.

X. SISYRINCHIUM (*L.*, sub *Iride*; *Baker in Journ., Bot.* ix. 42), = *Gynandriris Sisyrinchium* (*Parl. Nuov. Gen. e Spec. Monocot.* 52; *Klatt, Linnæa*, xxxii. 577).

Mar. sept.—Valde frequens tam in maritinis quam in locis siccis a mari remotis. Tanger! (*Webb Herb.*)! In monte Djebel Kebir! Cap. Spartel! Inter Tanger et Tetuan!

Mar. occid.—In monte Djebel Hadid hand procul Mogador!

Mar. merid.—In saxosis aridis prope Aïn Oumast! et alibi pluries visum!

Ar. Geog. Regio omnis Mediterranea calidior (exclusa Macaronesia). Deest in Gallia et in Italia superiore.

XYPHION FÆTIDISSIMUM (L., sub *Iride*; *Parl. Nuov. Gen. e Spec. Monocot.* 45).

Mar. sept.—“In rupibus altioribus montis Beni Hosmar prope Tetuan” (*spec. ex Herb. Webb*)!

Ar. Geog. Species occidentalis et australis. In regione sublitorea Atlantica ab Anglia usque insulas Canarienses. In regione Mediterranea præsertim occidentali diffusa, in Oriente nonnisi prope Byzantium et in Tauria detecta.

X. PSEUDACORUS (L., *Schousb. Gew. Marok.* 16, sub *Iride*; *Parl. Nuov. Gen. e Spec. Monocot.* 45).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Schsb. l. c.*). Visum inter Tanger et Tetuan!

Ar. Geog. Europa omnis (exclusa arctica). In Oriente et in Africa boreali rarius occurrit. In Macaronesia deest.

IRIS (L. *Gen.* 59).

I. FLORENTINA (L., *Desf. Fl. Atl.* i. 36; *Schousb. Gew. Marok.* 15).

Mar. sept.—Vidimus, nisi fallor, prope Tetuan, forsam cultam! Sec. *Schousboe*, l. c., colitur ornamenti gratia ad tumulos. Imp. Marocani dubia civis.

Ar. Geog. Planta ex Europa austro-occidentali et Asia Mediterranea orta, diu culta et potius subspontanea quam indigena in regione Mediterranea occidentali. Indicatur in insulis Canariensibus; an indigena?

I. GERMANICA (L., *Schousb. Gew. Marok.* 15; *Koch, Syn. Fl. Germ.* 807).

Mar. sept.—Vidimus cultam et subspontaneam juxta Tanger et Tetuan, præsertim ad tumulos Maurorum!

Mar. occid.—Vidimus cultam prope Mogador!

Mar. merid.—In convalle Aït Mesan juxta pagum Arround ad muros et aggeres agrorum!

Planta apud Arround lecta, forsam indigena, habet flores longiuscule pedicellatos.

Ar. Geog. Verisimiliter in regione Mediterranea fere omni indigena; in Europa media certe introducta. Indicatur in P. B. Webb, *Phyt. Canar.* iii. 314, sine loco natali.

ROMULEA (*Maratti, Pl. Romul. et Saturn.* 13).

Nomen genericum *Romulea* a Maratti l. c. anno 1772 impositum, a plerisque auctoribus prætervisum, quam *Trichonema* (Ker in *Ann. of Bot.* 1805, 222) antiquius et jure prioritatis gaudens.

Species Europæe hujus generis valde ludibundæ, ab auctoribus inordinate multiplicatæ, extricatu difficiles sunt; forsam omnes ad typos *Ixia Bulbocodium*, L., et *I. Columnæ*, R. et Sch., revocandæ.

ROMULEA BULBOCODIUM (*L. et Schousb. Gew. Marok.* 13, sub *Ixia*; *Seb. et Maur. Fl. Rom. Pr.* 17; *Parl. Fl. It.* iii. 242), = *Trichonema Bulbocodium* (*Ker in Ann. of Bot.* i. 222; *Koch, Syn. Fl. Germ., et plurim. auct.*).

Mar. sept.—Specimina deflorata mense Aprili prope Tanger passim vidimus et legimus! Formam grandifloram prope Tanger legit Salzmann (*Exsicc. in Herb. Gay*)! Florentem legimus in monte Beni Hosmar prope Tetnan circ. 1000 m. s. m.!

Mar. occid.—Casa Blanca! Mazagan!

Spec. plurima deflorata et subexsiccata ineerta sunt. Schousboe sub *Ixia Bulbocodium* sua tres varietates (scil. “major,” “media,” et “minor”) distinxit. Harum omnium loci natales, ut videtur, in ditione Tingitana; sub priore forsitan intellexit *R. grandiscapam* nostram.

Ar. Geog. Regio omnis Mediterranea in zona litorali et sublitorali. Deest in Macaronesia, ubi ejus locum tenet planta proxime affinis mox enumeranda.

R. grandiscapa (*Webb, Phyt. Canar.* iii. 312; *Klatt in Linnæa*, xxxii. 671, sub *Trichonema*).

Mar. sept.—Legimus defloratam in litore sinus Tingitani urbi opposito in arenosis. Hujus formam scapo brevior (T. *umbellata*, Klatt in ‘*Linnæa*’ l. c.) legerunt Plummer et Blackmore. Specimen vidi pulchrum sexflorum (*Blackmore in Herb. Kew.*)!

Structura valde proxima *R. Bulbocodium*, et hujus varietati *Clusiana* (Lange, *Exsicc. Hisp.*), sed scapi elongati, demum bipedales et ultra, et semina exacte globosa lævissima lucentia (in *T. Bulbocodio* subopaca et oculo armato minute granulata). Pro subspecie enumeranda videtur.

Ar. Geog. Planta Canariensis et Tingitana, verisimiliter alibi invenienda.

R. umbellata (Klatt, sub *Trichonema*) ex spec. Schousboeano prope Tingim lecto descripta, nulla nota stabili dignoscenda est.

R. COLUMNÆ (*R. et Sch. Syst. Veg. Mant.* 30, sub *Ixia*; *Seb. et Maur. Fl. Rom. Pr.* 18; *Parl. Fl. It.* iii. 253).

Mar. sept.—Tanger (*Schousboe*, see. Klatt in *Linnæa*, xxxii. 659).

Ar. Geog. Regio Mediterranea imprimis occidentalis. Europa occidentalis ab Anglia usque Lusitaniam. In Oriente vix aut ne vix detecta.

R. RAMIFLORA (*Ten. Fl. Neap. Prod. Add.*; *Bert. Fl. It.* i. 223).

Mar. sept.—Prope Tanger (*Plummer, spec. in Herb. Kew.*)!

Ar. Geog. Hucusque in Italia et in agro Tingitano tantum detectus.

CROCUS (*L. Gen.* 55).

C. SEROTINUS (*Salisb. Parad.* tab. 3), var. **SALZMANNI** (*J. Gay in Féruss. Bull.* xv. 220; *Il. fl. in Bot. Mag.* xcviii. tab. 6000), = *C.*

Salzmännianus (*Herb. in Bot. Reg.* 1847, tab. 4. fig. 4), = *C. tingitanus* (*Salzm. Exsicc.*; *Herb. in Not. ad Bot. Mag.* tab. 3868).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Peninsula Iberica. Var. nostra in agro Tingitano.

In Anglia culta floret mense Octobri. Quid sit *Crocus vernus* (Schonsboe, *Gew. Marok.* 12), prope Tingim indicata, dijudicare nequeo.

AMARYLLIDÆ (*R. Br. Prod.* 296).

LEUCOJUM (*L. Gen.* 402).

L. AUTUMNALE (*L., Desf. Fl. Atl.* i. 281; *Schonsb. Gew. Marok.* 139), = *Acis autumnalis* (*Herb. Amaryll.* 332; *Kunth, Enum.* v. 475).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Spec. ex herb. Schonsboe in monte Djebel Kebir lectum comm. cel. Cosson!

Ar. Geog. Regio Mediterranea imprimis occidentalis ex Cephalaria et Sicilia usque Lusitaniam. Deest in Gallia australi in Italia continentali et in Macaronesia.

L. TRICHOPHYLLUM (*Schonsb. Gew. Marok.* 140; *Brot. Fl. Lus.* i. 552; *Willk. et Lge. Fl. Hisp.* i. 148).

Mar. sept.—In campis arenosis Tingitanis loco dicto Scaarf el Kaab (*Schsb. l. c.*). Spec. ex herb. Schsb. benevole comm. cel. Cosson!

Mar. oecid.—Prope Mogador (*Carstensen in Herb. Kew.*)!

Ar. Geog. Species sat rara, in peninsula Iberica et in Imperio Maroccano hinc inde obvia.

L. TINGITANUM, *Baker*, n. sp.

“Folia linearia, carnosio-herbacea, synanthia, pedalia, superne lin. 1, inferne lin. 2 lata; scapus pro grege robustus 1–1½-pedalis; umbella 6–7-flora, spathæ valvis pallidis lanecolatis subpollicaribus, ad basin fissis, pedicellis filiformibus, demum 12–21 lin. longis; perianthii limbus albidus, 7–8 lin. longus, segmentis lanecolatis subconformibus, distincte cuspidatis, nervis pluribus (in sicco brunceis) segregatis percurtis; stamina 3 lin. longa, antheris lanecolatis filamentis æquilongis. Bulbus deest.”

Mar. sept.—Tanger (*Trove Blackmore, spec. in Herb. Kew.*)!

Species nova, cum *L. trichophyllo* in *Herb. Kew.* commixta, ab oculatissimo J. G. Baker recognita et descripta. Monente el. auctore, “a *L. trichophyllo* recedit habitu multo robustiore, foliis latioribus pedilibus, umbellis multifloris, genitalibus majoribus, cæterisque notis indicatis.”

LAPIEDRA (*Lag. Gen. et Sp.* 14).

L. GRACILIS, *Baker*, n. sp.

“Bulbus globosus, diam. 3–4 lin. latus, tunicis brunneis membranaceis supra collum productis; folia solitaria, post anthesin producta, fili-

formia, subteretia, glabra; scapus filiformis, uniflorus, 2-5-pollicaris; spatha monophylla, lanecolata, membranacea, 6-8 lin. longa; pedicellus spatha brevior, florifer erectus; ovarium ovoideum, clavatum, $1\frac{1}{2}$ lin. longum; perianthii albidii erecti tubus brevissimus, campanulatus; limbi segmenta subconformia, lanceolata, 5-6 lin. longa, nervis 4-5 obscuris medium versus percursa; stamina ad faucem tubi inserta, filamentis rectis filiformibus 4 lin. longis, antheris lanceolato-deltaideis semilineam longis, basi profunde sagittatis; stylus filiformis, curvatus, limbo æquilongus; stigma capitatum."

Mar. sept.—? Inter specimina Marocæana *Leucoji trichophylli* ex herb. Schousboe a cl. Lange communicata hanc speciem novam generis hucusque monotypici detexit et descripsit cl. J. G. Baker. Locus natalis adhuc incertus, sed verisimiliter in ditione Tingitana a Schousboe lecta fuit.

Primo aspectu *Leucojo trichophyllo* simillima, hæc stirps insigniter differt staminibus in perianthii tubo brevissimo insertis, nec epigynis, filamentis quam antheræ multoties longioribus, nec subæquilongis, antheris profunde sagittatis, nec basifixis. A *Lapiedra Martinezii* Hispanica aspectu longius distat, habitu multo graciliore, et foliis filiformibus, nec ovato-oblongis.

PANCRATIUM (*L. Gen.* 404).

P. MARITIMUM (*L., Kunth, Enum.* v. 658; *Schousb. Gew. Marok.* 145).

Mar. sept.—In arenosis pr. Tanger (*Schsb. l. c.*)! Folia visa in arenosis maritimis inter Tetuan et Ceuta!

Mar. occid.—Folia visa prope Mazagan!

Ar. Geog. Habitat in litoribus arenosis totius regionis Mediterraneæ (exclusa Macaronesia) et in Gallia occidentali.

TAPEINANTHUS (*Herb. Amaryll.* 190).

T. HUMILIS (*Cavan. Icon.* iii. tab. 207, fig. 2, sub *Pancratio*; *Herb. l. c.*; *Kunth, Enum.* v. 703) = *Amaryllis exigua* (*Schousb. Gew. Marok.* 146), = *Sternbergia exigua* (*Gawl. in Schult. Syst.* vii. 795; *Kunth, Enum.* v. 702), = *Oporanthus exiguus* (*Herb. Amaryll.* 180), = *Carregnoa humilis* (*Boiss. Voy.* 605).

Mar. sept.—In collibus et campis regionis Tingitanæ infrequens (*Schsb. l. c.*).

Ar. Geog. Nonnisi in Hispania Bætica et in agro Tingitano hucusque detectus.

Cavanilles primus hanc speciem raram cognovit et delineavit. Herbert anno 1837 genus novum, *Tapeinanthus*, proposuit, cum *Tapinantho*, Blume, non confundendum. Genus Blumeianum ab auctoribus recentioribus ad sectionem *Loranthi* reductum hodie negligendum. Botanici egregii Kunth et Herbert plantam Tingitanam verisimiliter

nounisi ex descriptione cognoverunt, dum eam in duobus generibus diversis enumerarunt. Confer J. Gay in Bull. Soc. Bot. Fr. vi. 88.

CORBULARIA (*Haw. Monogr.* 1).

C. BULBOCODIUM (*L.*, sub *Narcisso*; *Haw. Narc. Revis.* 123; *Herb. Amaryll.* 297, tab. 38. fig. 13; *Kunth, Enum.* v. 705).

Mar. sept.—Tanger (*fide Kunth, l. c.*).

Ar. Geog. Gallia austro-occidentalis. Hispania. Lusitania. Ager Tingitanus.

C. OBESA (*Salisb. Prodr.* 222, sub *Narcisso*; *Haw. Narc. Revis.* 122; *Herb. Amaryll.* 298; *Kunth, Enum.* v. 706).

Mar. sept.—Tanger (*fide Kunth, l. c.*).

Ar. Geog. Peninsula Iberica australis. Ditio Tingitana.

NARCISSUS (*L. Gen.* 403).

N. TAZETTA (*L.*, *Desf. Fl. Atl.* i. 282), = *Hermione Tazetta* (*Haw. Narc. Revis.* 142; *Kunth, Enum.* v. 740; *Schousb. Gew. Marok.* 141).

Mar. sept.—Tanger “in campis subhumidis” (*Schsb. l. c.*).

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

N. PAPYRACEUS (*Gawl. in Bot. Mag.* tab. 947; *Parl. Fl. It.* iii. 125), = *N. niveus* (*Loisel. Narc.* 27; *Gren. et Godr. Fl. Fr.* iii. 260; *an Willd. Enum. Suppl.* 16?), = *Hermione papyracea*, var. *nivea* (*Herb. Amaryll.* 408; *Kunth, Enum.* v. 746), = *Hermione unicolor* β , *tingitana* (*M. J. Ræm. Amar.* 231).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. In zona sublitorali regionis Mediterraneæ occidentalis a Sicilia usque Lusitaniam et Tingim. Ex Algeria non vidi.

N. SEROTINUS (*L. Sp. Pl.* 417; *Parl. Fl. It.* iii. 157; *J. Gay in Bull. Soc. Fr.* vi. 12), = *N. serotinus*, var. *uniflora* (*Desf. Fl. Atl.* i. 283, sec. *Parlatore*), = *Hermione serotina* (*Haw. Monogr.* 13; *Kunth. Enum.* v. 749).

Mar. occid.—Prope Saffi (*fide Leared in ‘Morocco and the Moors’*). Nisi fallor, etiam a Broussonnet in regione litorali Atlantico olim lectus.

Ar. Geog. Regio Mediterranea fere omnis calidior (exclusa Macaronesia). Dcest in Gallia australi et in Italia superiore et media.

N. elegans (*Haw. Monogr.* 13, sub *Hermione*; *Spach, ex Kunth, Enum.* v. 749; *Parl. Fl. It.* iii. 159; *J. Gay in Bull. Soc. Bot. Fr.* vi. 12), var. INTERMEDIA, *J. Gay, l. c.* = *N. serotinus* (*Salzm. Exsicc.*; *Schousb. Gew. Marok.* 141, saltem quoad pl. in herb. *Schousb.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! “In collibus et campis argillosis regionis Tingitanæ valde frequens” (*Schsb. l. c.*). Specimen ex herb. *Schousboc*, sub *N. serotino*, benev. concessit cel. Cosson!

Mar. occid.—Prope Mogador (*Carstenseu, spec. in Herb. Kew.*)!

Ar. Geog. In zona sublitorali Italiæ mediæ et inferioris, et in Sicilia. Africa borealis. Hispaniæ dubia civis.

Ex descriptione brevi, nulla nota diagnostica illustrata, in Willk. et Lge. Fl. Hisp. i. 155, utrum sub *N. serotino* auctores formam hanc cum *N. serotino* genuino semel misceverunt, vel hanc floræ Hispanicæ alienam esse judicarunt non satis liquet. Monentibus auctoribus citatis *N. serotinus* (Desf. Fl. Atl. i. 283, exclusa var. *uniflora*), et *N. oxyptetalus* (Boiss. Voy. Esp. 606) inter synonyma *N. elegantis* memoranda sunt. *Hermione elegans* (Kunth, Enum. v. 748) est nomen collectivum, *Aureliam Broussonnetii*, J. Gay, cum *N. elegante* complectens. Primo intuitu *N. elegans* a *N. serotino* differt perigonii laciniis acutioribus, corona brevissima integerrima, nec ut in specie Linnæana tripartita, ex segmentis emarginatis subsexfida. Oculatissimus J. Gay tamen coronam subintegerrimam in *N. serotino* vidit, et quoad formam perigonii planta Tingitana, monente viro optimo semper deslendo, *N. serotino* propius quam *N. eleganti* accedit. Differentiæ foliorum leves; *N. serotinus* sæpius uniflorus, *N. elegans* 2–7 florus; sed hæ notæ vix constantes. Hic mihi videtur ad subspeeiem reducendus.

NARCISSUS VIRIDIFLORUS (*Schousb. Gew. Marok.* 142, tab. 2; *Gay in Bull. Soc. Bot. Fr.* vi. 12), = *Hermione viridiflora* (*Haw. Narc. Revis.* 147; *Kunth, Enum.* v. 750), = *Chloraster fissus* (*Haw. Monogr.* 7, No. 1).

Mar. sept.—In ditione Tingitana rarissimus. “Copiose legi in loco elevato saxoso humido juxta Tingidem quando itur ab arce ad montem Shibil Kibir mox dextroversum flectendo cursum inter hortos mare versus” (*Schsb. l. c.*).

Ar. Geog. In utroque latere freti Herculis. Juxta Gibraltar et Algeiras. Floret Oct. Novemb. Planta in herbariis fere omnibus desiderata.

Foliis teretibus fistulosis, floribus viridibus a *N. serotino* satis differt.

AURELIA (*J. Gay in Bull. Soc. Bot. Fr.* vi. 87).

Genus ex forma perigonii infundibuliformi, laciniis tubo multo brevioribus in campanæ formam imbricatim conniventibus, insigne.

A. BROUSSONNETII (*Lag. Gen. et Sp.* 13, sub *Narcisso*; *J. Gay, l. c.*), = *Narcissus obliteratus* (*Willd. in Reliq., in Schult. Syst. Veg.* vii. 981), = *Hermione obliterata* (*Haw. Monogr.* 13), = *H. elegans* (*Kunth, Enum.* v. 749, *pro parte*).

Mar. occid.—Mogador (*Brouss. Herb., fide J. Gay, l. c.*). Prope Saffi (*ex bulbis a cl. Leared lectis, in Hort. Kew. cultis*).

Species parum nota Imp. Maroceani propria.

Agave Americana, L., planta in regione Mediterranea calidiore late dif-

fusa, in Imp. Maroccanum olim introducta, ab incolis præsertim prope Tingim ad sepes construendas colitur, interdum fit spontanea.

DIOSCOREÆ (*R. Br. Prod.* 294).

TAMUS (*L. Gen.* 1119).

T. COMMUNIS (*L., Kunth, Enum.* v. 453).

Mar. sept.—Prope Tanger! In monte Beni Hosmar! et alibi prope Tetuan!

Mar. merid.—In ditione Mesfioua ad radices Atlantis Majoris (*J. D. H.*)!

Ar. Geog. Europa media. Regio fere omnis Mediterranea. Asia temperata. In insulis Canariis indicata, sed, monente Webb, plantarum insul. est *T. edulis*, Lowe.

ALISMACEÆ (*Juss. Dict. Sc. Nat.* vii. 474).

ALISMA (*L. Gen.* 460).

A. PLANTAGO (*L., Kunth, Enum.* iii. 148; *Schousb. Gew. Marok.* 170).

Mar. sept.—Tanger (*Schousb. l. c.*).

Ar. Geog. Planta amphigea in temperatis tam frigidioribus quam calidioribus utriusque orbis, et in Australia, diffusa. In Macaronesia desideratur.

A. RANUNCULOIDES (*L., Kunth, Enum.* iii. 149).

Mar. sept.—Tanger! Inter Tanger et Tetuan!

Ar. Geog. Europa fere omnis (excl. arctica). Africa borealis. Deest in Oriente.

DAMASONIUM (*Juss. Gen.* 46).

D. STELLATUM (*Rich. in Pers. Syn.* i. 400; *Kunth, Enum.* iii. 155).

Mar. occid.—“Ex horto Imperatoris prope Mogador” (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Europa occidentalis. Africa borealis. Indicatur etiam in Russia media et in Persia juxta Mare Caspicum.

JUNCAGINEÆ (*Rich. Mém. Mus. Par.* ii. 365).

TRIGLOCHIN (*L. Gen.* 463).

T. BARRELIERI (*Loisel. Fl. Gull.* i. 725; *Kunth, Enum.* iii. 143),=

T. palustre, var. β , *L.*, = T. palustre (*Schousb. Gew. Marok.* 170; *Brot. Fl. Lus.* i. 600).

Mar. sept.—Tanger, in maritimis et locis humidis! (*Schsb. l. c.*).

Ar. Geog. Regio Mediterranea imprimis occidentalis. Desideratur in Asia Mediterranea et in Macaronesia.

POTAMEÆ (*Juss. Dict. Sc. Nat.* 93).POTAMOGETON (*L. Gen.* 174).

- P. NATANS* (*L., Kunth, Enum.* iii. 127; *Schousb. Gew. Marok.* 59).
 Mar. occid.—In paludosis salsis prope Mogador! (*Schsb. l.c.*) (*Lowe Cat.*).
Ar. Geog. Species per regiones temperatas hemisphæræ borealis late diffusa, nec deest forma *P. fluitans* in ins. Canariensibus.
- P. PECTINATUS* (*L., Kunth, Enum.* iii. 137), = *P. marinus* (*Desf. Fl. Atl.* i. 150; *Schousb. Gew. Marok.* 59, non *L.*).
 Mar. occid.—In aquis prope Mogador (*Schsb. l. c.*).
Ar. Geog. Europa media. Regio Mediterranea (exclusa Macaronesia).
P. marinus, *L.*, geminus habitat in frigidioribus Europæ mediæ et septentrionalis, nec in regione Mediterranea huiusmodi detectus est.

ZOSTERA (*L. Gen.* 1032).

- Z. MARINA* (*L., Kunth, Enum.* iii. 116).
 Mar. sept.—Vidimus in sinu Tingitano! legi florentem in altero litore freti Hereulis prope Algeciras.
Ar. Geog. Ad litora Atlantica Europæ, Americæ borealis, et Africæ borealis. Nascitur etiam in mari Mediterraneo, et in Baltico, et in Ponto Euxino.

PALMÆ (*Juss. Gen.* 37).CHAMÆROPS (*L. Gen.* 319).

- C. HUMILIS* (*L., Kunth, Enum.* iii. 248).
 Per territorium Maroccanum omne nobis notum valde frequens, colles incultos interdum obtegens.
 Mar. sept.—Circa Tanger! (*Salzm. Exsicc.*)! Tetuan! Vidimus in monte Beni Hosmar usque 375 m. s. m.!
 Mar. occid.—Casa Blanca! Safli! Mogador! (*Lowe Cat.*).
 Mar. merid.—In regione inferiore et in convallibus calidioribus Atlantis Majoris usque 1300 m. s. m.! Shedma! Seksaona! et alibi visa.
Ar. Geog. Species per regionem Mediterraneam, præsertim calidiorem, late diffusa. Deest in Macaronesia.

PHENIX (*L. Gen.* 1224).

- P. DACTYLIFERA* (*L., Kunth, Enum.* iii. 255).
 Mar. sept.—Cultam pluries vidimus!
 Mar. occid.—Juxta Mazagan! Safli! et Mogador! certe culta.
 Mar. merid.—In provinciis Cisatlanticis vix indigena. Juxta urbem Marocco sylvam magnam efficit, sed nullibi spontanea nascitur. In regionibus tamen a catena Atlantis Majoris meridiem versus *Phœnix* verisimiliter civis indigena est.

Ar. Geog. Patria *Phœnicis* in subtropicis Africæ borealis a Mari Rubro usque Oceanum et insulas Canarienses. Inde versus Mare Mediterraneum et Asiam occidentalem olim opere hominum diffusa est. Confer Webb, *Phyt. Canar.* iii. 189.

TYPHACEÆ (*Juss. Gen.* 25).

TYPHA (*L. Gen.* 1040).

T. ANGUSTIFOLIA (*L., Kunth, Enum.* iii. 91).

Mar. sept.—Tanger (*Webb spec.*)! Folia visa!

Ar. Geog. Species cosmopolita in temperatis et calidioribus hemisphæræ borealis late diffusa. Occurrit etiam in America meridionali et in Australia; an indigena? In ins. Canariensibus obvia est stirps huic valde proxima, *T. macranthelia*, Webb, sicut subspecies vel varietas enumeranda.

AROIDEÆ (*Juss. Gen.* 23).

ARISARUM (*Targ. Tozz. in Ann. Mus. Fir.* ii. 67).

A. VULGARE (*Targ. Tozz. l. c.; Kunth, Enum.* iii. 15; *H. fil. in Bot. Mag.* tab. 6023), = *Arum arisarum*, *L.*

Mar. sept.—Ubique circa Tanger et Tetuan et in regione collina intermedia! (*Salzm. Exsicc.*)!

Mar. oecid.—Folia vidimus frequentia circa Casa Blanca! Mazagan! Saffi! Pr. Mogador minus frequens!

Mar. merid.—Occurrit, sed infrequens, in regione inferiore a mari remota! Rhizomata diu sub aqua contusa et lavata edulia fiunt, et tempore inediae nutrimentum ineolis præstant.

Ar. Geog. Regio fere omnis Mediterranea (inclusa Macaronesia).

Planta Canariensis *A. subexsertum* (Webb, *Phyt. Can.* iii. 301) occurrit etiam in Maroeco, sed formis intermediis in typum transit, et vix pro varietate enumeranda videtur.

ARUM (*L. Gen.* 1028).

A. ITALICUM (*Mill. Dict.* No. 2; *Kunth, Enum.* iii. 24).

Mar. oecid.—Casa Blanca!

Mar. merid.—Ad radices Atlantis prope Milhain (folia tantum visa)!

Ar. Geog. Regio Mediterranea imprimis occidentalis. In Anglia australi et Gallia occidentali obvinum, in Oriente (præter Græciam) desideratur. Forma proxime affinis (*A. canariense*) in Canariis occurrit.

A. TENUIFOLIUM (*L., Bert. Fl. It.* x. 249), = *Biarum tenuifolium* (*Schott, Meletem.* 17; *Kunth. Enum.* iii. 22).

Mar. sept.—? Legimus sine flore et fructu in monte Beni Hosmar prope Tetuan!

Ar. Geog. Lusitania. Hispania Granatensis. Sicilia. Italia inferior. Dalmatia. Græcia.

Ob. spec. nimis incompleta incertum, attamen folia adamussium conformia.

LEMNACEÆ (*Duby, Bot. Gall. i. 533*).

LEMNA (*L. Gen. 1038*).

L. GIBBA (*L., Rhcb. Icon. vii. tab. 16*), = *Telmatosphace gibba* (*Schleid. in Linnæa, viii. 391; Kunth, Enum. iii. 6*).

Mar. merid.—Reraya (sine flore et fructu)!

Ar. Geog. In aquis stagnantibus hemisphæræ borealis temperatæ late diffusa. Adest in Macaronesia; sed ex Algeria spec. non vidi.

LILIACEÆ (*DC. Théor. Elém. 249*).

GAGEA (*Salisb. in Ann. Bot. ii. 555*).

G. FOLIOSA (*R. et Sch. Syst. vii. 1703; Kunth, Enum. iv. 236*), = *G. polymorpha* (*Boiss. Voy. 611*).

Mar. merid.—Specimen unicum legi in regione subalpina montis Djebel Tezah circ. 2700 m.!

Ar. Geog. Habitat in Hispania, Corsica, Sicilia, Græcia, et in Imp. Maroccano.

HYACINTHUS (*L. Gen. 170*).

H. ROMANUS (*L., Desf. Fl. Atl. i. 308; Schousb. Gew. Marok. 160*), = *Bellevalia romana* (*Rhcb. Fl. Exc. i. 105; Kunth, Enum. iv. 367*).

Mar. sept.—Tanger in agris (*Schsb. l. c.*).

Ar. Geog. In regione Mediterranea centrali et occidentali (exclusa Macaronesia) late diffusus.

MUSCARI (*Mill. Dict. v. 197*).

M. COMOSUM (*L., sub Hyacintho; Mill. Dict. No. 2; Desf. Fl. Atl. i. 309*), = *Bellevalia comosa* (*Kunth, Enum. iv. 306*).

Mar. sept.—Tanger (*J. D. II.*)!

Mar. merid.—Sat frequens in cultis reg. inferioris. Prope Seksaoua! Pluries alibi visum in convallibus Atlantis Majoris!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). In Europa media hinc inde introductum.

M. RACEMOSUM (*L., sub Hyacintho; Mill. Dict. No. 3; Schousb. Gew. Marok. 161*), = *Botryanthus odorus* (*Kunth, Enum. iv. 131*).

Mar. sept.—Tanger (*Schsb. l. c.*).

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia). Europa media, imprimis occidentalis.

SCILLA (*L. Gen.* 419).

S. AUTUMNALIS (*L., Kunth, Enum.* iv. 315; *Schousb. Gew. Marok.* 155).

Mar. sept.—Tanger “in vinetis et locis arenosis” (*Schsb. l. c.*).

Ar. Geog. Regio fere omnis Mediterranea (exclusa Macaronesia). Europa occidentalis.

S. OBTUSIFOLIA (*Poir. Voy.* ii. 149; *Kunth. Enum.* iv. 315), var. *FALLAX* (*Baker in Journ. Linn. Soc.* xiii. 235), = *S. fallax* (*Steinheil in Ann. Sc. Nat.* ser. 2, i. 103).

Mar. sept.—Tanger (*Salzm. spec. in Herb. Kew.*)!

Ar. Geog. Species in Algeria, Sardinia, et Corsica obvia. Var. *fallax* nonnisi in agro Tingitano.

S. HEMISPHERICA (*Boiss. Voy. Esp.* 613), var. *GLABRA*, *Boiss., l. c.*, = *S. peruviana* (*L., Brot. Fl. Lus.* i. 526; *Schousb. Gew. Marok.* 153), = *S. peruviana*, var. *glabra* (*Kunth, Enum.* iv. 319; *Baker in Journ. Linn. Soc.* xiii. 240), = *S. Clusii* (*Parl. Fl. It.* ii. 462), = *Calosquilla Clusii* (*Jord. et Fourr. Icon.* tab. 245).

Mar. sept.—Tanger versus montem Djebel Kebir! (*Schsb. l. c.*) (*Webb Herb.*)!

Ar. Geog. *S. hemisphærica* in regione Mediterranea occidentali a Sicilia et Melita usque Lusitaniam, et ex agro Tinctano usque Tingim, diffusa est. Varietates plurimas sicut species distinctas descripserunt Jordan et Fourreau in opere citato.

Nomen triviale Linnæum, ex errore Clusii natum, omnino falsum et hodie prætermittendum.

S. VERNA (*Huds. Fl. Angl.* 142; *Kunth, Enum.* iv. 319), var. *MAJOR* (*Boiss. Voy. Esp.* 613; *Kunth, l. c.* 320), = *S. Ramburei* (*Boiss. Elench.* 61; *Willk. et Lge. Fl. Hisp.* i. 214), = *S. italica* (*L. Herb., non Sp. Pl.* 442, fide *Baker in Journ. Linn. Soc.* xiii. 242). Huc fere certe referenda est *S. italica* (*Schousb. Gew. Marok.* 153).

Mar. sept.—Tanger (*Brouss. spec. in herb. Hooker olim Gouan*)! (*Schsb. l. c.*)? Bulbos, forsán huc referendos, invenimus prope Tetuan!

Ar. Geog. Species occidentalis ex Anglia, Hibernia, et usque Hispaniam, præsertim in zona sublitorali diffusa. Var. *Boissieriana* in peninsula Iberica et in ditioe Tingitana obvia est.

S. MONOPHYLLA (*Link, in Schrad. Journ.* ii. 319; *Willk. et Lge. Fl. Hisp.* i. 214), = *S. tingitana* (*Salzm. Exsicc., non Schsb. Gew. Marok.* ?), = *S. pumila* (*Brot. Fl. Lus.* i. 527; *Bot. Mag.* tab. 3023; *Kunth, Enum.* iv. 320).

Mar. sept.—Tanger (*Webb spec.*)! (*Salzm. Exsicc.*)! In monte Djebel Kebir! Spec. ex herb. Schousboe in eodem monte lectum benevole comm. cel. Cosson sub “*S. tingitana*, *Schsb. Herb.*”

Ar. Geog. Peninsula Iberica. Ager Tingitanus.

Quid sit *Scilla tingitana* (Schousb. Gew. Marok. 155) mihi non satis liquet. In scheda herb. Webb ad spec. *S. monophyllæ* fertur "*S. pumila*, Brot.", "*S. tingitana*, Salzm., non Schsb., sec. Boiss." Revera b. Schousboe vix speciem novam folio unico insignem descripserit verbis "foliis lanceolatis planis apice involutis," nec eam *S. mauritanicæ* suæ simillimam dixerit. Certum est b. Schousboe in herb. suo *S. monophyllam* sub nomine "*S. tingitana*" conservasse. Specimen *S. tingitanæ*, Schousb. in herb. Willdenow, verisimiliter ab ipso Schousboe receptum, ex Kunth, Enum. iv. 322, est omnino *S. monophylla*, Lk. Ex descriptione *S. tingitana*, Schousb., a *S. lusitunica*, L., vix differt. Hujus specimina nulla vidi, et deest in herbariis fere omnibus.

SCILLA LINGULATA (Poir. Voy. ii. 151; Desf. Fl. Atl. i. 208, tab. 85; Kunth, Enum. iv. 324), = *S. bipartita* (Salzm. Exsicc.).

Mar. sept.—Tanger (Salzm. Exsicc.)! (Blackmore, spec. in Herb. Kew.)! Ar. Geog. Africa boreali-occidentalis.

S. MAURITANICA (Schousb. Gew. Marok. 154; Kunth, Enum. iv. 324), = *S. vincentina* (Hffgg. et Link in N. Schrift. Ges. Naturf. iv. 17; Kunth, Enum. iv. 323; Baker in Journ. Linn. Soc. xiii. 244).

Mar. sept.—Tanger (Salzm. Exsicc.)! In monte Djebel Kebir (Schsb. l. c.).

Ar. Geog. Lusitania. Ager Tingitanus.

Nomen triviale Schousboeanum antiquius et satis aptum prætuli.

S. HISPANICA (Mill. Dict. No. 8; Baker in Journ. Linn. Soc. xiii. 255), = *S. campanulata* (Ait. Hort. Kew. i. 444; Kunth, Enum. iv. 326; Schousb. Gew. Marok. 153), = *Endymion campanulatus*, Willk.

Mar. sept.—"Passim in dumetis regionis 'Tingitanæ'" (Schsb. l. c.). A nullis recentiorum visa, an error?

Mar. merid.—Legi in regione superiore Atlantis Majoris circ. 2700 m. in monte Djebel Tezali!

Ar. Geog. Peninsula Iberica. Marocco.

S. HÆMORRHOIDALIS (Webb, Phyt. Canar. ii. 336, tab. 230).

Mar. occid.—? Mogador (Lowe Cat., cum signo interrogationis).

Deest in herbario Lowe hodie Kewensi, et forsitan errore aliquo indicatur. Species Canariensis, hucusque non alibi visa.

URGINEA (Steinheil in Ann. Sc. Nat. ser. 2, i. 321).

U. MARITIMA (L., sub *Scilla*; Baker in Journ. Linn. Soc. xiii. 221), = *U. Scilla* (Steinheil, l. c.; Kunth, Enum. iv. 332), = *Scilla maritima* (Schousb. Gew. Marok. 151, et plur. auct.).

Mar. sept.—Vidimus bulbos maximos in collibus Tingitanis sat frequentes! "Ubique in campis et collibus" (Schsb. l. c.).

Schousboe l. c. certe de regione Tingitana loquitur; in provinciis meridionalibus facile invenienda, certe non frequens.

Ar. Geog. In regione Mediterranea tota præsertim calidiore late diffusa, nec semper in maritimis.

URGINEA ANTHERICOIDES (*Poir. Voy.* ii. 149; *Kunth, Enum.* iv. 333; *Baker in Journ. Linn. Soc.* xiii. 220), = *Seilla serotina* (*Schousb. Gew. Marok.* 151; non *Gawl. in Bot. Mag.*). Verisimiliter huc etiam ducenda est *U. hesperia* (*Webb, Phyt. Canar.* iii. 339).

Mar. occid.—“Vulgaris in campis arenosis prope Saffi”! (*Schsb. l. c.*).

Ar. Geog. Africa boreali-occidentalis. Verisimiliter etiam in Canariis occurrit. Plantam Webbiam non vidi.

ORNITHOGALUM (*L. Gen.* 418).

In hoc genere a cl. J. G. Baker optime tractato, ordinem specierum in *Journ. Linn. Soc.* xiii. pp. 257 et seq., secutus sum.

O. COMOSUM (*L., Kunth, Enum.* iv. 359; *Baker, l. c.* 262), var ATLANTICUM, *Baker MSS.*

Mar. merid.—Spec. incompletum in *Herb. Kew.* asservatum in convalle Aït Mesan supra Arround legerunt cl. Rein et Fritsch! Ejusdem var. spec. in monte Djebel Afougueur ab indigena Ibrahim lectum benev. comm. Cosson!

Ar. Geog. Species in Italia, Istria, Dalmatia, Macedonia et etiam prope Vindobonam obvia. Var. nostra nonnisi in Atlante Majori hucusque detecta.

Braetæ elongatæ, basi vaginantes, flores juniores superantes.

O. TENUIFOLIUM (*Guss. Prod. Fl. Sic.* i. 413; *Baker, l. c.* 265), = O. Gussonii (*Ten. Fl. Nap.* iii. 337, tab. 226; *Kunth, Enum.* iv. 364).

Mar. merid.—Specimen unicum legi in subalpinis montis Djebel Tezah circ. 2700 m.!

Ar. Geog. Europa australis a Gallia Mediterranea ad Græciam. Atlas Major. Varietates in insula Cypro et in Ægypto.

O. ORTHOPHYLLUM (*Ten. Syll.* 594; *Kunth, Enum.* iv. 361; *Baker, l. c.* 265), = O. algeriense (*Jord. et Fourr. Icon.* tab. 121). Huc etiam sicut varietas adjungendum O. baticum (*Boiss. Elench.* 84).

Mar. sept.—In monte Djebel Kebir pr. Tanger! In saxosis prope Tetuan! et in monte Beni Hosmar!

Mar. merid.—In clivo montis Djebel Tezah circ. 1700 m. (*J. D. H.*)!

Ar. Geog. Italia media et inferior. Hispania. Africa boreali-occidentalis. Forma proxime affinis in Transylvania.

O. UMBELLATUM (*L., Kunth, Enum.* iv. 362; *Baker, l. c.* 256, et forsan *Schousb. Gew. Marok.* 150),

Mar. sept.—? In agris humidis regionis Tingitanæ (*Schsb.*).

Mar. occid.—Casa Blanca!

Synonymon Schousbœanum forsân ad *O. orthophyllum* referendum?

Ar. Geog. Europa fere omnis, exclusa arctica. Africa borealis. Oriens.
In Macaronesia deest.

ORNITHOGALUM ARABICUM (*L. ex parte; Kunth, Enum. iv. 353; Baker, l. c. 270*).

Mar. sept.—Tanger (*Webb Herb.*)! Tctuan (*Webb Herb.*)!

Mar. occid.—Rabat (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Per regionem Mediterraneam omnem, præsertim calidiorem, dispersum, sed infrequens.

O. PYRENAICUM (*L. Sp. Pl. 440, non herb.; Kunth, Enum. iv. 355; Baker, l. c. 275*), var. *FLAVESCENS* (*Baker, l. c.*), = *O. flavescens* (*Lam. Ill. tab. 242*), = *O. sulfureum* (*R. et Sch. Syst. vii. 518; Kunth, Enum. iv. 356*), = *O. pyrenaicum* (*L. herb.*).

Mar. merid.—Supra Amsiniz (*J. D. H.*)!

Ar. Geog. Species per Europam mediam et australem diffusa. *O. flavescens* in Italia, Gallia, Hispania et in Atlante Majore. In Asia Minore detexit etiam Balansa.

Specimina Africana præter nostra non vidi.

Cel. Koch *O. flavescens* pro specie distincta habuit; pace el. Baker mihi videtur pro subspecie enumerandum.

O. SESSILIFLORUM (*Desf. Fl. Atl. i. 295; Kunth, Enum. iv. 371; Baker, l. c. 276*).

Mar. merid.—Legi in saxosis supra Seksaoua!

Ar. Geog. Africa boreali-occidentalis.

O. NARBONENSE (*L., Kunth, Enum. iv. 355; Baker, l. c. 277*), = *O. stachyoides* (*Ait. Hort. Kew. i. 441; Kunth, Enum. iv. 356*).

Mar. sept.—Tanger! Tctuan (*Webb spec., sub O. pyrenaico*)!

Mar. occid.—Casa Blanca!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

O. UNIFOLIUM (*Gawl. in Bot. Mag. tab. 935; Kunth, Enum. iv. 359; Baker, l. c. 282*), = *O. nanum* (*Brot. Fl. Lus. i. 529; non Sibth. et Sm.*), = *Scilla unifolia* (*L., Salzmann. Exsicc.*).

Mar. sept.—Tanger (*Salzmann. Exsicc.*)! (*Webb Herb.*)! In monte Djebel Kebir copiose provenit!

Ar. Geog. Peninsula Iberica. Ditio Tingitana.

DIPCADİ (*Medik. in Act. Palat. vi. 431*).

D. SEROTINUM (*L., sub Hyacintho; Medik. l. c.*), = *Lachenalia serotina* (*Willd. Sp. Pl. ii. 175; Schousb. Gew. Marok. 161*), = *Uropetalum serotinum* (*Gawl. in Bot. Reg. fol. 156; Boiss. Voy. Esp. 617; Kunth, Enum. iv. 378, et auct. recent. fere omnium*), var. *FULVUM*, = *Hyacinthus fulvus* (*Cavan. Anal. Cienc. Nat. iii. 47*), = *Dipeadi fulvum* (*Webb, Phyt. Canar. iii. 340*).

Mar. sept.—Tanger (*Webb Herb.*)! In monte Djebel Kebir! (*Schsb. l. c.*). Prope Tetuan visum!

Mar. occid.—? Mogador (*Brouss. fide Kunth*, forsā enī spec. Canariensi commutatum).

Mar. merid.—In regione inferiore et ad radices Atlantis pinus visum. Sheshaoua! In saxosis supra Seksaoua!

Ar. Geog. Regio Mediterranea occidentalis a Liguria usque Lusitaniam et insulas Canarienses.

Invito animo nomen genericum barbarum, a summis scientiæ magistris prætermissum, ab optimo Webb infauste resuscitatum, conservare debui.

D. fulvum, Webb, est nil nisi var. robustior *D. serotini*, florum colore saturatiore diversa.

ALLIUM (*L. Gen.* 409).

A. AMPELOPRASUM (*L., Kunth, Enum.* iv. 383), = *A. multiflorum* (*Desf. Fl. Atl.* i. 288), = *A. mogadorese* (*Willd. Herb.; R. et Sch. Syst.* vii. 1004).

Mar. occid.—Mazagan! Saffi! Mogador (*Brouss. ex Willd. Herb.*).

Ar. Geog. Regio Mediterranea occidentalis (inclusa Macaronesia). Anglia. Sarnia.

A. ROTUNDUM (*L., Kunth, Enum.* iv. 385).

Mar. merid.—Prov. Iaha (*J. D. H.*)! Prov. Shedma!

Ar. Geog. Europa centralis et anstralis. Oriens. Hispania. Marocco. Deest in Macaronesia. Ex Algeria non vidi.

A. SPHÆROCEPHALUM (*L., Kunth, Enum.* iv. 387; *Schonsb. Gew. Marok.* 148).

Mar. sept.—“Passim in arenosis regionis Tingitanæ” (*Schsb. l. c.*).

Ar. Geog. Europa media. Regio Mediterranea (inclusa Macaronesia).

A. FLAVUM (*L., Kunth, Enum.* iv. 402), var. TAURICUM (*Kunth, l. c.*).

Mar. merid.—Specimen unicum legi in saxosis supra Seksaoua!

Ar. Geog. Species late dispersa, nullibi frequens, per regionem Mediterraneam ex Armenia usque Hispaniam centralem diffusa. Spec. Africana, præter nostrum, non vidi.

Specimen nostrum incompletum et ideo subdubium.

A. PALLENS (*L., Kunth, Enum.* v. 405; *Schonsb. Gew. Marok.* 148).

Mar. sept.—Tanger (*Webb Herb.*)! (*Schsb. l. c.*).

Mar. occid.—In provincia Iaha prope Mogador! In monte Djebel Iiadid (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—In aridis juxta Aïn Ommast!

Ar. Geog. Regio fere omnis Mediterranea (exclusa Macaronesia). Pannonia.

ALLIUM PANICULATUM (*L., Sibth. et Sm. Fl. Gr.* iv. 16; *an Kunth, Enum.* iv. 407?), = *A. monspessulanum* (*Willd. Enum. Suppl.* 16; *Kunth, Enum.* iv. 404, *non Gouan*), = *A. intermedium* (*DC. Fl. Fr.* v. 318; *Kunth, Enum.* iv. 406), = *A. dentiferum* (*Webb, Phyt. Canar.* iii. 345).

Mar. sept.—Tanger! (*Webb spec.*, sine nomine specifico)!

Mar. occid.—Mazagan (specimen nanum floribus rubris)!

Mar. merid.—Mzouda!

Ar. Geog. Regio fere omnis Mediterranea (inclusa Macaronesia). Gallia occidentalis. Indicatur etiam in Rossia australi. Spec. ex Algeria non vidi.

— — —, var. **TENUIFLORUM**, *Baker, MSS.*, = *A. tenuiflorum* (*Ten. Fl. Nap.* i. 165; *Parl. Fl. It.* ii. 548; *Kunth, Enum.* iv. 409).

Mar. merid.—Legimus in monte Djebel Tezah circ. 1800 m.! Juxta Seksaoua circ. 1000 m.!

Ar. Geog. In Italia media et inferiore, et in Sicilia, late diffusa. In Atlante Majore. Verisimiliter alibi invenienda.

A. TRIQUETRUM (*L., Kunth, Enum.* iv. 436; *Schousb. Gew. Marok.* 149).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Schsb. l. c.*). In monte Djebel Kebir!

Ar. Geog. Regio Mediterranea imprimis occidentalis (exclusa Macaronesia). Græcia excepta deest in Oriente.

A. pendulinum (*Ten. Fl. Nap.* i. 168, tab. 31; *Kunth, Enum.* iv. 437).

Mar. sept.—In monte Djebel Kebir prope Tanger!

Ar. Geog. Italia. Corsica. Sicilia. Marocco.

Forsan præcedenti nimis proximum. Pro subspecie enumeravi.

A. ROSEUM (*L., Kunth, Enum.* iv. 438).

Mar. sept.—Tanger (*Webb Herb.*)! In regione collina inter Tanger et Tetuan! In saxosis pr. Tetuan! Ad radices montis Beni Hosmar!

Mar. occid.—In apricis prope Casa Blanca!

Ar. Geog. Regio Mediterranea fere omnis (inclusa Macaronesia).

A. SUBHIRSUTUM (*L., Kunth, Enum.* iv. 440; *Schousb. Gew. Marok.* 147).

Mar. sept.—Tanger (*Schsb. l. c.*). Tetuan (*Webb Herb.*)! Inter Tetuan et Ceuta!

— — —, var. **SUBVILLOSUM**, = *A. subvillosum* (*Salzm. in R. et Sch. Syst.* vii. 1104; *Willk. et Lge. Fl. Hisp.* i. 212).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Species per regionem Mediterraneam (exclusa Macaronesia) late diffusa. Varietas in peninsula Iberica et in agro Tingitano.

ALLIUM CHAMÆMOLY (L., *Kunth, Enum.* iv. 447; *Schousb. Gew. Marok.* 150).

Mar. sept.—Tanger in campis argillossis (*Schsb. l. c.*). (*Salzm. Exsicc.*)!

Ar. Geog. Regio Mediterranea imprimis occidentalis (exclusa Macaronesia).

A. NIGRUM (L., *Kunth, Enum.* iv. 447; *Schousb. Gew. Marok.* 148) = A. magicum (*Salzm. Exsicc., Willd., et verisimiliter L. Sp. Pl.* 424).

Mar. sept.—Tanger! (*Webb Herb.*)! (*Salzm. Exsicc.*)! (*Schsb. l. c.*).

Ar. Geog. Regio fere omnis Mediterranea (exclusa Macaronesia).

ASPHODELUS (*L. Gen.* 421).

A. CERASIFERUS (*J. Gay in Ann. Sc. Nat.* ser. 4. vii. § 2, 15; *Willk. et Lge. Fl. Hisp.* i. 204), = A. ramosus (L., *Kunth, Enum.* iv. 555, pro parte; *Salzm. Exsicc.*), = A. albus (*Boiss. Voy. Esp.* 618).

Mar. sept.—Legimus in monte Djebel Kebir prope Tanger! Tanger (*Salzm. Exsicc. in Herb. Kew.*)!

Ar. Geog. Regio Mediterranea occidentalis a Gallia australi usque Tingim et Lusitaniam. Ex Algeria non vidi.

Linnæus tres species hujus sectionis semel confusit, quæ nobis, duce eel. J. Gay, probe distinctæ videntur, et a patribus, saltem pro parte, haud ignoratæ. Inter synonyma plurima a beato Gay citata notandum est "Asphodelus major flore albo ramosus et non ramosus" (*J. Bauhin, Hist.* ii. 265; edit. Chabræus, Ebroduni, 1651), ubi legitur "flores stellati *Ormithogali* majores foliis albis stria rubra notatis constantes apicibus luteis, quibus decedentibus capitula succedunt cerasis paria semen continentia copiosum triquetrum fuscum."

A. MICROCARPUS (*Viv. Fl. Cors. Spec.* 5; excl. syn. *Clusii*; *Boiss. Voy. Esp.* 618; *J. Gay, l. c.*), = A. ramosus (L., *Kunth, Enum.* iv. 555, pro parte; *Schousb. Gew. Marok.* 156), = A. æstivus (*Brot. Fl. Lus.* i. 525; *Kunth, l. c.* 557), nomen antiquius sed ineptum, flores enim vernaes sunt et mense Majo planta semper fructifera.

Mar. sept.—Tanger "in campis et fruticetis frequens" (*Schsb. l. c.*). Hanc speciem vidimus prope Tanger, sed spec. non servavi!

Mar. occid.—Mogador (*Lowe, sub A. ramoso, spec. incompletum in Herb. Kew.*)!

Mar. merid.—Ex convalle Aït Mesan (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio fere omnis Mediterranea (inclusa Macaronesia).

A. FISTULOSUS (L., *Kunth, Enum.* iv. 557).

Mar. sept.—In monte Djebel Kebir prope Tanger! In collibus inter Tanger et Tetuan!

Ar. Geog. Regio Mediterranea imprimis calidior (inclusa Macaronesia).

— —, var. *TENUIFOLIUS*, = *A. tenuifolius* (*Cavan. in Anal. Cienc. Nat.* iii. 46, tab. 27; *Kunth, Enum.* iv. 558), = *A. fistulosus* (*Schousb. Gew. Marok.* 156; *Lowe Cat.* 35).

Mar. occid.—Valde frequens in arenosis prope Mogador! (*Schsb. l. c.*). (*Lowe Cat.*).

Mar. merid.—Prov. Mtouga (*R. et Fr. spec. in Herb. Kew.*)!

Primo intuitu planta Mogadorensis a typo solenni *A. fistulosi* valde diversa est, sed intersunt formæ plurimæ intermediæ, ex gr. planta ex prov. Mtouga supra citata; immo specimina *A. fistulosi* Tingitana a typo recedunt et *A. tenuifolio* nimis vicina sunt.

ANTHERICUM (*L. Gen.* 422).

A. LILIAGO (*L., Koch, Syn. Fl. Germ.* 616), = *Phalangium Liliago* (*Schreb. Spicil.* 36; *Kunth, Enum.* iv. 594), var. *BÆTICUM*, = *A. bæticum* (*Boiss. Voy. Esp.* 619, tab. 172).

Mar. sept.—In monte Djebel Kebir prope Tanger! (*Webb spec.*)! Frequens in saxosis circa Tetuan! Ad radices montis Beni Hosmar!

Mar. merid.—In convalle Aït Mesan legit G. Maw!

Ar. Geog. Species in montanis Europæ mediæ et australis. Var. *bæticum* in peninsula Iberica et in Africa boreali-occidentali.

Culta ex radicibus Tingitanis planta nostra ad typum propius revertit, et fit *A. Liliago*, var. *australe* (*Willk. et Lge. Fl. Hisp.* i. 202).

SIMETHIS (*Kunth, Enum.* iv. 618).

S. PLANIFOLIA (*L., sub Antherico; Gren. et Godr. Fl. Fr.* iii. 222), = *S. bicolor* (*Desf. Fl. Atl.* i. 304, tab. 90, sub *Antherico; Kunth, l. c.*), = *Anthericum planifolium* (*Schousb. Gew. Marok.* 157; *Salzm. Exsicc.*).

Mar. sept.—Circa Tanger, et copiose in monte Djebel Kebir! (*Salzm. Exsicc.*)! (*Schsb. l. c.*). Tetuan (*Webb Herb.*)!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia). Europa occidentalis ex Hibernia usque Lusitaniam.

APHYLLANTHES (*L. Gen.* 408).

A. MONSPELIENSIS (*L., Kunth, Enum.* iv. 647).

Mar. sept.—In clivo montis Beni Hosmar prope Tetuan!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

MELANTHACEÆ (*R. Br. Prod.* 272).COLCHICUM (*L. Gen.* 457).

C. BIVONÆ (*Guss. Cat. H. Boccad.* 72; *Kunth, Enum.* iv. 139).

Mar. merid.—? Fructiferum in monte Djebel Tezah circ. ad 2400 m. legit J. D. II.!

Folia hysterranthia elongata, anguste linearia, sesquipedalia. Planta cujus flores nondum vidi, nimis incerta et facile ad sp. novam referenda.

Ar. Geog. Regio Mediterranea calidior. Deest in Macaronesia et in Gallia australi.

C. AUTUMNALE (*L., Kunth, Enum.* iv. 140), = *C. montanum* (*Schousb. Gew. Marok.* 170).

Mar. sept.—Tanger (*Salzin. spec. in herb. Gay, hodie Kewensi*)! (*Schsb. l. c.*).

Ar. Geog. Europa temperata fere omnis. Regio Mediterranea.

C. ARENARIUM (*W. K. Pl. rar. Hung.* ii. 195, tab. 179), var.?

Mar. merid.—In convalle Aït Mesan supra Arround, et in jugo Tagherot a 2200 m. usque 2900 m.!

Ar. Geog. *Colchicum arenarium*, species imprimis Europæ austro-orientalis, et in Hispania obvia, forsân per regionem omnem Mediterraneam dispersa est. Planta proxime affinis in Italia præsertim meridionali. Diagnoses in hoc genere difficillimæ.

Specimina nostra incerta faice *C. alpiuo* similiora, sed folia terna. Species ulterius inquirenda.

ERYTHROSTICTUS (*Schlecht. in Linnæa*, 1826, p. 90).

E. PUNCTATUS (*Cavan. Anal. Cienc. Nat.* iii. 49, tab. 26, sub *Melanthio*; *Schlecht. l. c.*; *Kunth, Enum.* iv. 154), quocum conjungendus *E. gramineus* (*Cavan. l. c.* p. 50, sub *Melanthio*; *Schlecht. l. c.*; *Kunth, l. c.* 155).

Mar. occid.—Legimus copiose in arenosis prope Mogador, ubi hanc stirpem singularem primus legit Broussonnet! Conf. *Cavan. l. c.*

Planta satis variabilis. Simul crescunt exemplaria foliis latis et angustis ex quibus species duas stabilivit cel. Cavanilles. Forma intermedia in Hispania unico loco prope Almeria a cl. Lange detecta fuit.

Ar. Geog. Marocco. Hispania austro-orientalis.

SMILACEÆ (*R. Br. Prod.* 292).CONVALLARIA (*L. Gen.* 425).

C. MULTIFLORA (*L., Koch, Syn. Fl. Germ.* 705), = *Polygonatum multiflorum* (*All. Fl. Ped.* i. 131; *Kunth, Enum.* v. 138).

Mar. sept.—In monte Beni Hosmar prope Tetuan anno 1869, a cl. G. Maw detecta. Ex radice in horto culta specimina mihi benevole misit!

Ar. Geog. Species in temperatis præsertim frigidioribus hemisphæræ borealis late diffusa. In montibus Asiæ mediæ obvia, fit rarior in calidioribus. In Hispania nonnisi in provinciis septentrionalibus. Præter spec. Mawcana ex Africa nulla vidi.

SMILAX (*L. Gen.* 1120).

S. ASPERA (*L., Kunth, Enum.* v. 214).

Mar. sept.—Tanger!

Mar. occid.—Mogador (*Lowe Cat.*). In monte Djebel Hadid haud procul Mogador! (*R. et Fr. spec. in Herb. Kew.*)! Agadir (*Coll. Indig.* 1871)!

Ar. Geog. Regio fere omnis Mediterranea (exclusa Macaronesia).

S. MAURITANICA (*Poir. Voy.* ii. 263; *Kunth, Enum.* v. 216).

Mar. sept. Tanger! magis frequens quam *S. aspera*.

Mar. merid.—In convalle Aït Mesan circ. 1500 m. s. m. (*J. D. H.*)!

Ar. Geog. Regio Mediterranea imprimis occidentalis. In Macaronesia adest.

ASPARAGINEÆ (*Kunth in Act. Acad. Berol.* 1812, 25).

ASPARAGUS (*L. Gen.* 458).

A. SCABER (*Brign. Fasc. Pl. Forojul.* 22; *Kunth, Enum.* v. 62).

Mar. sept.—Tanger (*Goudot, fide Kunth, l. c.*).

Ar. Geog. Per regionem Mediterraneam (exclusa Macaronesia) hinc inde obuius. An idem ac *A. altissimus*, Munby?

A. ACUTIFOLIUS (*L., Kunth, Enum.* v. 65; *Schousb. Gew. Marok* 160).

Mar. sept.—Tanger (*Schousb. l. c.*). In maritimis prope Tanger! Prope Tetuan!

Mar. occid.—? In insula Mogador! specimen omnino incertum.

Mar. merid.—In convalle Aït Mesan circ. 1400 m.!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

A. APIHYLLUS (*L., Kunth, Enum.* v. 66; *Schousb. Gew. Marok.* 160).

Mar. sept.—Sat frequens prope Tanger! (*Schsb. l. c.*). Ad radices montis Beni Hosmar prope Tetuan! Inter Tetuan et Centa!

Ar. Geog. Regio Mediterranea præsertim calidior (inclusa Macaronesia).

A. stipularis (*Forsk. Fl. Æg.-Arab.* 72), = *A. horridus* (*L. fil. Suppl.* 203; *Kunth, Enum.* v. 67).

Mar. sept.—Tetuan (*Webb Herb.*)!

Mar. merid.—Videtur satis frequens in regione inferiore. Prov. Haha! Sheshaoua! Tasseremout (*G. Maw*)! In planitie prope urbem Marocco (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio omnis Mediterranea calidior. Deest in Gallia et in Italia continentali.

Proximus et valde affinis *A. aphyllus*; characteres autem videntur satis constantes, et subspeciem sensu nostra sistit.

ASPARAGUS SCOPARIUS (*Lowe in Trans. Camb. Phil. Soc.* iv. 11; *Kunth, Enum.* v. 74).

Mar. merid.—Legimus pluries in regione inferiore. Prov. Shedma! Inter Ourika et Reraya! In convalle Aït Mesan circ. 1200 m.!

Specimina nostra Maderensibus simillima, sed ob defectum florum et fructus incerta.

Ar. Geog. *A. scoparius* est Macaronesiæ et prov. meridionalium Imp. Maroccani propria species.

A. FENICULACEUS (*Lowe in Journ. Linn. Soc.* v. 44).

Mar. occid.—Mogador (*Lowe, spec. in Herb. Kew.*)! Agadir (*Coll. Indig.* 1871)!

Species, ut videtur, nonnisi Maroccana, parum cognita, flores et fructus hucusque non visi.

Hic est verisimiliter *A. declinatus* (*Schousb. Gew. Marok.* 159) juxta Saffi indicatus. Nequit esse *A. declinatus* (*L., Kunth*, l. c. 59); sed ex diagnosi manca dijudicare non audeo.

ASPARAGOPSIS (*Kunth in Act. Acad. Berol.* 1842, 35).

A. ALBA (*Kunth, Enum.* v. 84), = *Asparagus albus* (*L., Schousb. Gew. Marok.* 159).

Mar. sept.—Tanger (*Schsb. l. c.*).

Mar. occid.—Saffi (*Schsb. l. c.*).

— —, var. *PASTORIANUS*, = *A. Pastorianus* (*Webb, Phyt. Canar.* iii. tab. 229).

Mar. occid.—Prope Mogador (*Lowe Cat.*). In insula Mogador spec. maneam sed certe huc referendum!

Ar. Geog. *A. albus* in regione Mediterranea occidentali a Calabria usque Lusitaniam et Imp. Maroccanum. Var. *Pastorianus* nascitur in Canariis et in Marocco.

Quid sit *Asparagus retrofractus* (*Schousb. Gew. Marok.* 159), prope Saffi lectus, non satis liquet. Species hujus nominis Linnæana est planta Capensis haud facile in ditione nostra invenienda.

Asparagi species Maroccanæ plurimæ ulterius inquirendæ, et studio diuturno in vivo egent.

RUSCUS (*L. Gen.* 1139).

R. HYPOPHYLLUM (*L., Kunth, Enum.* v. 275).

Mar. sept.—Vidi sine fl. et fr. prope Tanger! Tetuan (*Webb Herb.*)!

Mar. occid.—Mogador (*Lowe Cat.*).

Ar. Geog. Regio Mediterranea calidior. In Madera obvius, desideratur in insulis Canariensibus. Deest in Gallia et in Italia superiore.

JUNCÆE (*R. Br. Prod.* 257).

LUZULA (*DC. Fl. Fr.* iii. 158).

L. FORSTERI (*Sm. Eng. Bot.* tab. 1293, sub *Junco*; *Desv. Journ. de Bot.* i. 141; *Kunth, Enum.* iii. 297), = *Juncus pilosus* (*Schousb. Gew. Marok.* 166, non *L.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)! (*Schsb. l. c.*) In monte Beni Hosmar prope Tetuan!

Ar. Geog. Europa media. Regio Mediterranea fere omnis (exclusa Macaronesia).

JUNCUS (*L. Gen.* 437).

J. GLAUCUS (*Ehrh. Beitr.* vi. 83; *Kunth, Enum.* iii. 316).

Mar. sept.—Vidi in monte Djebel Kebir pr. Tanger, sed spec. non servavi!

Ar. Geog. Species per regiones temperatas et frigidas hemisphæræ borealis latissime diffusa.

J. CONGLOMERATUS (*L., Koch, Syn. Fl. Germ.* 726), = *J. communis*, var. α (*E. Mey. Junc.* 20; *Kunth, Enum.* iii. 320).

Mar. sept.—Pluries vidi prope Tanger et Tetuan! Specimen ex herb. Schousboe benevole comm. Cosson!

— —, var. EFFUSUS, = *J. effusus* (*L., Koch, Syn. Fl. Germ.* 726), = *J. communis*, var. β (*E. Mey. Junc.* 20; *Kunth, l. c.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Nihil dubito *J. conglomeratum* et *J. effusum* nexu genetico colligari; notæ tamen differentiales subconstantes videntur, et forsân *J. effusus* melius pro subspecie enumerandus erit.

Ar. Geog. Tam *J. conglomeratus* quam *J. effusus* sunt plantæ cosmopolitæ per orbem terrarum latissime diffusæ. Prædiligunt regiones temperatas, sed zonam subtropicam non fugiunt. *J. effusus* in Macaronesia indicatur.

J. MARITIMUS (*Lam. Dict.* iii. 264; *Kunth, Enum.* iii. 322; *Schousb. Gew. Marok.* 163).

Mar. sept.—Legi mense Junio fructiferum pr. Tanger! (*Schsb. l. c. et spec.*)! (*Salzm. Exsicc.*)! Tetuan (1851)!

Ar. Geog. Species cosmopolita, per orbem totum in maritimis regionum temperatarum diffusa.

J. ACUTUS (*L., ex parte*; *Kunth, Enum.* iii. 323; *Schousb. Gew. Marok.* 162).

Mar. sept.—Tanger (*Salzm. spec. in Herb. Kew.*) Inter Tetuan et Ceuta!

Mar. occid.—Mazagan! Mogador (*Lowe Cat.*). Aïn el Hadjar haud procul Mogador in regione sublitorali!

Ar. Geog. In arenosis maritimi Europæ et Africæ borealis ex Britannia usque Macaronesiam, et regionis omnis Mediterraneæ.

JUNCUS MULTIFLORUS (*Desf. Fl. Atl. i. 313, tab. 91; Kunth, Enum. v. 323*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! In lacustribus salsis pr. Tingim (*Webb spec.*)! Cap. Spartel (*spec. ex Herb. Schousb.*)!

Mar. occid.—Aïn el Hadjar haud procul Mogador!

Ar. Geog. Species in maritimis regionis Mediterraneæ (exclusa Macaronesia) late diffusa, hucusque in Hispania non detecta.

J. LAMPROCARPUS (*Ehrh. Calam. No. 126; Kunth, Enum. iii. 325*).

Mar. sept.—(*Salzm. Exsicc.*)!

Mar. occid.—Prope Mogador (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Per Europam et Asiam temperatam late diffusus. In Africa boreali infrequens. Desideratur in Macaronesia.

J. STRIATUS (*Schousb. in E. Mey. Junc. 27; Kunth, Enum. iii. 330*), = *J. Fontanesii* (*J. Gay in Laharpe, Monog. Junc. 42; Kunth, l. c. 329*), = *J. sylvaticus* (*Brot. Fl. Lus. i. 517, et Salzm. Exsicc.; non aliorum*).

Mar. sept.—Prope Tanger (*Webb spec.*)! (*Salzm. Exsicc.*)! (*Schousb. spec. ex Herb.*)! (*Goudot, fide Kunth, l. c.*). Specimen in monte Beni Hosmar immaturum forsitan huc referendum!

Hujus forma minor, *J. echinuloides* (*Webb, It. Hisp., non Brot.*), in Marocco nondum detecta fuit.

Ar. Geog. Occurrit hinc inde in regione Mediterranea occidentali, a Sicilia et ditone Tunetana usque fretum Herculis.

J. SUPINUS (*Mærch, Enum. Pl. Hass. 296, tab. 5; Koch, Syn. Fl. Germ. 730*), = *J. bulbosus* (*L. Sp. Pl. ed. i. 327, no. ed. ii.*), = *J. uliginosus* (*Roth, Mag. f. d. Bot. ii. 16; Kunth, Enum. iii. 334*).

Mar. sept.—Mense Junio prope Tanger visus, sed spec. non servavi! Specimen prope Cap. Spartel lectum ex herb. Schousboe benev. misit cel. Cosson!

Ar. Geog. Europa fere omnis ex Lapponia usque Calabriam. In regione Mediterranea infrequens. Deest in Macaronesia.

J. HETEROPHYLLUS (*Léon Duf. in Ann. Sc. Nat. v. 88; Parl. Fl. It. iii. 342*).

Mar. sept.—Legi in uliginosis reg. litoralis inter Tetuan et Ceuta!

Duce el. Kunth, qui hanc speciem satis distinctam inter synonyma *J. uliginosi* enumeravit, specimina supra citata ad varietatem *J. supini* in herbario retuli, donec oculatissimus Cosson errorem indicavit.

Ar. Geog. Gallia præsertim occidentalis. Corsica. Italia. Sicilia. Africa Tingitana. In Hispania nondum detectus, nec spec. ex Algeria vidi. Verisimiliter per regionem Mediterraneam occidentalem dispersus.

JUNCUS PYGMÆUS (*Thuil. Fl. Paris.* i. 178; *Kunth, Enum.* iii. 341).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Europa media et occidentalis. Regio Mediterranea occidentalis (exclusa Macaronesia). Fertur in Græciæ insula Milo, sed valde dubius.

J. fasciculatus (*Schousb. in E. Mey. Junc.* 28; *Kunth, Enum.* iii. 330 non *Bert. Fl. It.*, nec *Schrank, Bayer. Fl.*), = *J. acuminatus* (*Salzm. Exsicc.*)!

Mar. sept.—Tanger (*Salzm. Exsicc.*)! In monte Djebel Kebir (*spec. ex Herb. Schousboe*)! (*Webb Herb.*)!

Mihi videtur *J. pygmaeo* nimis proximus et hujus proles hexandra. Pro subspecie enumeravi.

Ar. Geog. Stirps, ut videtur, ditionis Tingitanæ propria.

J. CAPITATUS (*Weigel, Obs.* 28; *Kunth, Enum.* iii. 346; *Schousb. Gew. Marok.* 164).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! In monte Djebel Kebir (*Schsb. l. c.*). Juxta Cap. Spartel! Inter Tanger et Tetuan! Prope Tetuan!

Mar. occid.—Casa Blanca!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). In Europa occidentali etiam obuius.

J. TENAGEIA (*Ehrh. Phyt.* No. 63; *Kunth, Enum.* iii. 352).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! In monte Djebel Kebir (*spec. ex Herb. Schousboe comm. Cosson*)!

Ar. Geog. Europa media. Asia media. Regio Mediterranea occidentalis (exclusa Macaronesia).

In regione sublitorali Hispaniæ australis non indicatur; legi tamen in collibus prope San Roque!

J. BUFONIUS (*L., Kunth, Enum.* iii. 353; *Schousb. Gew. Marok.* 166).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)! In monte Djebel Kebir! (*Schsb. l. c.*). Juxta Cap. Spartel!

Mar. occid.—Casa Blanca!

Mar. merid.—Satis frequens ad radices Atlantis Majoris. Mesfioua! Sektana! Ourika! et in convallibus calidioribus—Aït Mesan! Amsmiz!

— — —, var. *FASCICULATUS* (*Koch, Syn. Fl. Germ.* 732), = *J. fasciculatus* (*Bert. Fl. It.* iv. 190, non *Schousb.*).

Mar. sept.—Spec. in humidis prope Tanger lectum ex herb. *Schousboe comm. cel. Cosson*!

Mar. occident.—Prope Casa Blanca! Prope Mogador! (*Lowe Cat.*).

Mar. merid.—Juxta Amsmiz!

Ar. Geog. Species cosmopolita per regiones temperatas orbis fere totius dispersa.

Juncus foliosus (*Desf. Fl. Atl.* i. 315, tab. 92).

Mar. sept.—Tanger (*Webb spec.*)! (*Salzm. Exsicc.*)! (*Spec. ex Herb. Schousboe*)!

J. bufonio valde proximus et sicut subspecies enumerandus, differt tamen foliis latoribus, anthela magis decomposita, et, fide el. Lange, seminibus (oculo fortiter armato) etiam transverse striolatis.

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis.

CYPERACEÆ (*DC. Fl. Fr.* iii. 99).

CYPERUS (*L. Gen.* 66).

C. MUNDTII (*N. ab Esen. in Linnæa*, ix. 283, sub *Pycreo*; *Kunth, Enum.* ii. 17; *Böckeler in Linnæa*, xxxv. 448), = *C. turfusus* (*Salzm. Exsicc.*), = *C. Eragrostis* (*Vahl, Enum.* ii. 322; *Kunth, Enum.* l. c. 8), var. *sec. Kunth, l. c.*

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Spec. ex Herb. Schousboe, comm. Cosson*)!

Ar. Geog. Species quoad distributionem anomala. Habitat in Africa australi ad prom. Bonæ Spei, in Hispania australi, et ad ripas rivulorum prope Tanger. Species proxime affinis *C. Eragrostis*, Vahl, etiam in India orientali et in insula Mauritio obvia est.

C. LÆVIGATUS (*L. Cod.* 61; *Böckeler, l. c.* 487), = *C. mucronatus* (*Rottb. Gram.* 10, tab. 8. fig. 4), = *C. mucronatus*, var. *albidus* (*Kunth, Enum.* ii. 17), = *C. cosyrensis* (*Tineo in Guss. Fl. Sic.* ii. 779).

Mar. occident.—In arenosis prope Mogador!

Ar. Geog. Species ampligea in tropicis et subtropicis utriusque orbis diffusa. In regione Mediterranea nonnisi in Ægypto, in insula Pantellaria, et prope Mogador hucusque detecta.

C. distachyos (*All. Auct. ad Fl. Ped.* 18; *Parl. Fl. It.* ii. 19), = *C. mucronatus*, var. γ (*Kunth, l. c.* 20), = *C. juneiformis* (*Cavan. Ic.* 3).

Mar. sept.—Prope Cap. Spartel (*spec. ex Herb. Schousboe*)!

Mar. occident.—Casa Blanca! In arenosis haud procul Mogador!

Ar. Geog. Stirps cosmopolita in regione Mediterranea (inclusa Macaronesia) late diffusa, in India orientali et in Australia certe indigena. Loci natales plurimi incerti ob synonyma confusa.

C. distachyos est *C. lævigato* proxime affinis, sed notis plurimis differt. In priore spiculæ elongatæ fere semper binæ, squamæ angustiores atro-

fusæ, achenium ovoidco-ellipticum apice mucronatum; in *C. lævigato* spicæ numerosæ dimidio breviores ovatæ, squamæ præsertim basin versus multo latiores albidæ, achenium demum obovatum muticum.

Pro subspecie enumeravi stirpem Allionii, sed forsân pro specie omnino diversa vindicanda est.

CYPERUS ÆGYPTIACUS (*Gloxin, Obs.* 20, tab. 3; *Kunth, Enum.* ii. 48), = *C. schœnoides* (*Griseb. Spic. Fl. Rum.* ii. 421), = *Schœnus mucronatus* (*L., Rchb. Icon.* viii. fig. 680; *Schousb. Gew. Marok.* 16; *Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Schsb. l. c.*). Inter Tetuan et Ceuta!

Mar. occid.—Rarius occurrit in arenosis prope Mogador!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

C. FUSCUS (*L., Kunth, Enum.* ii. 37).

Mar. sept.—In arenosis subhumidis videtur haud infrequens. Vidi prope Tetuan! Specimen prope Tanger lectum eomm. eel. Cosson!

Ar. Geog. Species per Europam et Asiam temperatam, et regionem omnem Mediterraneam (exclusa Macaronesia) late diffusa.

C. ROTUNDUS (*L., Kunth, Enum.* iii. 58), = *C. hexastachyos* (*Rottb. Gram.* 28, tab. 14), = *C. tetrastachyos* (*Desf. Fl. Atl.* i. 44, tab. 8), = *C. olivaris* (*Targ. Tozz., Bertol. Fl. It.* i. 267), = *C. esculentus* (*Gouan, Fl. Monsp.* 388; *Schousb. Gew. Marok.* 18; an *L. Cod.* 380?).

Mar. sept.—Tanger (*Schsb. l. c.*). Specimen ex herb. Schousb. comm. Cosson!

Mar. occid.—Mogador? (*Lowe Cat.*).

Mar. merid.—In ditione Mesfoua haud procul urbi Marocco!

Ar. Geog. Species per regiones tropicales orbis totius late diffusa. In regione Mediterranea (inclusa Macaronesia) satis frequens.

C. LONGUS (*L., Kunth, Enum.* ii. 60), var. *BADIUS*, = *C. badius* (*Desf. Fl. Atl.* i. 45, tab. 7; *Salzm. Exsicc., et plurim. auct.*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! Tetuan (1851)!

Mar. occid.—Mogador!

Mar. merid.—In regione inferiore haud infrequens. Sheshaoua! In irrigatis urbis Marocco! Ourika!

C. badius videtur forma meridionalis *C. longi*, foliis et radiis brevioribus, spicis aggregatis.

Ar. Geog. Species per Europam mediam et regionem Mediterraneam diffusa. Var. *badius* præsertim in regione Mediterranea calidiore (inclusa Macaronesia), et in India orientali obvia.

SCHÆNUS (*L. Gen.* 65).

S. NIGRICANS (*L., Koch, Syn. Fl. Germ.* 640), = *Chætospora nigricans* (*Kunth, Enum.* ii. 323).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! In rupibus juxta Cap. Spartel! (*Webb Herb.*)!

Ar. Geog. Europa media. Regio fere omnis Mediterranea (exclusa Macaronesia).

HELEOCHARIS (*R. Br. Prod.* 80).

II. PALUSTRIS (*L., sub Scirpo; Kunth, Enum.* ii. 147).

Mar. sept.—Tanger! Inter Tanger et Tetuan!

Mar. occid.—Mazagan!

Ar. Geog. Species amphigea in temperatis hemisphæræ borealis, in Africa australi, et in insulis Oceani obvia.

II. UNIGLUMIS (*Link, Jahrb. d. Gewk.* fasc. iii. 77, sub *Scirpo; Schult. in Syst. Veg. Mant.* ii. 88; *Koch, Syn. Fl. Germ.* ed. 2, 852), = *II. palustris* (*Kunth, l. c.* pro parte).

Mar. sept.—Legi prope Cap. Spartel! In monte Beni Hosmar pr. Tetuan!

Ar. Geog. Europa media; in regione Mediterranea nonnisi in Italia septentrionali et in Hispaniæ prov. Navarra indicata; verisimiliter sæpe cum *II. palustri* commutata.

Ab *H. palustri* differt statura minore, culmis tenuioribus, spicis minoribus spadiceis, et squama infima culmum omnino amplectente, nec semiamplexicauli. Mihi videtur species satis distincta.

SCIRPUS (*L. Gen.* 67).

S. LACUSTRIS (*L., Kunth, Enum.* ii. 164).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Europa fere omnis. Asia media et Mediterranea. Africa borealis. Fertur etiam in America boreali, in Australia et in insulis Sandvicensibus.

S. LITORALIS (*Schrad. Fl. Germ.* i. 142, tab. 5; *Kunth, Enum.* ii. 166), = *S. Tabernæmontani* (*Salzm. Exsicc., non Gmel.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Europa australi. Oriens. In Hispania et Algeria nondum detectus.

S. MARITIMUS (*L., Kunth, Enum.* ii. 167).

Mar. sept.—Tanger (*J. D. H.*)! Prope Cap. Spartel meridiem versus! Tetuan 18 51)!

Mar. occid.—Casa Blanca!

Ar. Geog. Species cosmopolita per orbem omnem late diffusus, sæpius, sed non semper, in maritimis crescens.

SCIRPUS PUBESCENS (*Poir. Voy.* ii. 254; *Desf. Fl. Atl.* i. 52, tab. 10), = *Fuirena pubescens* (*Kunth, Enum.* ii. 183), = *Scirpus ciliaris* (*Schousb. Gew. Marok.* 19, non *L.*).

Mar. sept.—“Rarissime in locis humentibus regionis Tingitanæ” (*Schsb. l. c.*). Spec. in monte Djebel Kebir lectum ex herb. Schousboe comm. cel. Cosson!

Ar. Geog. Africa boreali-occidentalis. Lusitania. Corsica, et in insulis Balearicis.

S. SAVII (*Seb. et Maur. Fl. Rom. Prod.* 22; *Bert. Fl. It.* i. 238), = *S. filiformis* (*Savi, Fl. Pis.* i. 46, non *Lam.*), = *S. numidianus* (*Vahl, Enum.* ii. 254, et *Herb.*, fide *Kunth*), = *Isolepis Saviana* (*Schult. in Syst. Veg. Maurit.* ii. 63; *Kunth, Enum.* ii. 193), = *S. setaceus* (*Schousb. Gew. Marok.* 19).

Mar. sept.—Frequens in ditioe Tingitana. Legi in maritimis juxta urbem Tanger! et prope Cap. Spartel! Tetuan (*Webb Herb.*)! In elivo montis Beni Hosmar!

Mar. merid.—In ditioe Reraya et in convalle Aït Mesan a 1000 m. ad 1200 m.!

Ar. Geog. Species per regiones temperatas totius orbis diffusa. In Europa occidentali ab Hibernia, ubi frequentissima, usque fretum Herculis, et inde in Marocco et in Macaronesiam se protendit. Per regionem omnem Mediterraneam dispersa, sed non ubique obvia. In Asia orientali, Japonia, et Australasia diffusa, et verisimiliter etiam in America obvia.

Species diu vexata et nominibus plurimis in herbariis dissimulata. Ab hac vix, aut ne vix, differt *Isolepis pygmæa* (*Kunth, Enum.* ii. 191). Catervam synonymorum citare nolui, dum plurima verificare nequeo. Nomen *S. numidianus*, Vahl, prioritatem gaudet; sed nomen triviale ab auctoribus recentioribus fere omnibus acceptum mutare nolui. Cl. Böekeler in ‘*Linnaea*,’ xxxvi. 500, *S. Savii* cum *S. setaceo*, L., semel junxit. Ego, consentientibus auctoribus gravissimis, has pro speciebus distinctis, fructus structura facile dignoscendis, teneo, et formas intermediarias nunquam vidi.

Specimina Tingitana *S. gracilis* (*Salzm. Exsicc.*), sec. Rehb. Icon. viii. 34, ad *Heleocharis carniolicam*, Koch, referenda sunt. Specimina Salzmanniana non vidi; sed monente oculatissimo Cosson meram formam *S. Savii* sistunt.

S. HOLOSCHÆNUS (*L., Desf. Fl. Atl.* i. 49; *Schousb. Gew. Marok.* 18), = *Isolepis holoschænus* (*R. et Sch. Syst.* ii. 115; *Kunth, Enum.* ii. 200).

Mar. sept.—Tanger (*Schsb. l. c.*), (*Salzm. Exsicc.*)! Tetuan (1851)!

Mar. occid.—Casa Blanca! Mazagan! Prope Mogador (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—Prope Tasseremout!

— — —, var. AUSTRALIS (*Koch, Syn. Fl. Germ.* 856), = *S. australis*, *L., sec. Vahl.*

Mar. sept.—Tanger (*spec. ex Herb. Schousboe comm. Cosson*)!

Ar. Geog. In regione omni Mediterranea (inclusa Macaronesia) late diffusus. In Europa media et occidentali rarius occurrit.

SCIRPUS MICHELIANUS (*L., Desf. Fl. Atl.* i. 51), = *Isolepis Micheliana* (*R. et Sch. Syst. Veg.* ii. 114; *Kunth, Enum.* ii. 203).

Mar. occid.—Ad ripas fluminis Sebou (*Brouss., fide Desfontaines, l. c.*).

Ar. Geog. Species in Europa media et australi infrequens. In Asia temperata necnon in India orientali obvia, et (fide Kunth) in Senegambia.

CAREX (*L. Gen.* 1046).

C. DIVISA (*Huds. Fl. Angl.* 405; *Kunth, Enum.* ii. 372).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Cap. Spartel!

Mar. merid.—In prov. Shedma!

Ar. Geog. Europa media. Regio omnis Mediterranea (inclusa Macaronesia).

C. VULPINA (*L., Kunth, Enum.* ii. 383).

Mar. sept.—Prope Tanger legi fructiferam mense Junio!

Ar. Geog. Regio Mediterranea fere omnis (inclusa Macaronesia). Europa media. Asia media.

C. DIVULSA (*Good. in Linn. Trans.* ii. 160; *Kunth, Enum.* ii. 385).

Mar. sept.—Tanger! ubi sat frequens (*Salzm. Exsicc.*)! Tetuan!

Mar. merid.—In collibus saxosis prope urbem Marocco (*G. Maw*)!

Ar. Geog. Species per hemisphæram borealem temperatam fere omnem diffusa, nec deest in Macaronesia.

C. LINKII (*Schkuhr, Caric.* ii. 39; *Kunth, Enum.* ii. 478), = *C. gynomane* (*Bertol. Pl. rar. Lig.* dec. 2, 43); et forsàn = *C. distachya* (*Desf. Fl. Atl.* ii. 336).

Mar. sept.—Mauritania Tingitana (*Kunth, l. c.*). Prope Tanger in monte Djebel Kebir! In monte Beni Hosmar pr. Tetuan!

Ar. Geog. Regio Mediterranea imprimis occidentalis (exclusa Macaronesia). In Oriente nonnisi prope Byzantium indicata.

C. BASILARIS (*Jordan, Obs. Fragn.* iii. 246, tab. 12; *Walp. Ann. Bot. Syst.* iii. 703).

Mar. sept.—Legi specimen unicum juxta Cap. Spartel!

Legi etiam in collibus supra Algeeciras!

Ar. Geog. Species hucusque in Gallia australi et in Algeria tantum indicata. Adest in Hispania Bætica et in ditione Tingitana; verisimiliter alibi invenienda.

C. HALLERIANA (*Asso, Syn.* 133, tab. 9; *Willk. et Lge. Fl. Hisp.* i.

125), = *C. gynobasis* (*Vill. Dauph.* ii. 206; *Boiss. Voy. Esp.* 630), = *C. alpestris* (*All. Fl. Ped.* ii. 270; *Kunth, Enum.* ii. 479).

Mar. sept.—In monte Beni Hosmar prope Tetuan!

Mar. merid.—In Atlante Majore; legi in convalle Amsmiz circ. 1500 m.!

Ar. Geog. Europa media et australis. Ad litora maris Caspii (fide Kunth). Spec. nulla ex Asia Minore nec ex Algeria vidi.

CAREX AMBIGUA (*Link in Schrad. Journ.* 1799, 308; *Kunth, Enum.* ii. 479).

Mar. sept.—Legi prope Cap. Spartel!

Mar. merid.—In convalle Aït Mesan, spec. nimis immatura!

Species paucis cognita, sed probe distincta utriculis glabris elliptico-ovoideis tricostatis, nec, ut in *C. Halleriana*, obovato-oblongis trigonis plurinerviis.

Spec. ex Lusitania Transtagana in Herb. Kew. nostris conformia sunt, nisi præsentia in spec. nostris fl. fœminicorum 2–4 ad basin spicæ superioris masculæ.

Spec. ex Atlante Majore nimis immatura majora, culmis fere pedem longis, sed Tingitanis similia. Planta ulterius inquirenda.

Ar. Geog. Lusitania. Marocco.

C. DISTANS (*L., Kunth, Enum.* ii. 449).

Mar. sept.—? In clivo montis Beni Hosmar prope Tetuan legi specimen nimis immaturum!

Spec. nostrum incertum, sed vix ad aliam speciem referendum. Teste cl. Willkomm *C. distans* nascitur in isthmo Gibraltarico.

Ar. Geog. Frequens occurrit in Europa media et australi; necnon in Caucaso, et in America boreali.

C. FISSIROSTRIS (*Ball in Journ. Bot.* 1875, 206).

Dense cæspitosa; culmi numerosi, humiles (subspithamæi), foliacei, læves, cum foliis elevato-striati; folia angusta sæpius undulato-recurva, scabriuscula; spica mascula solitaria, squamis dorso obscure trinerviis, pallidis vel inferioribus tricoloribus; spicæ fœminæ 3–4 oblongæ, pedicellatæ, squamis dorso viridibus apice mucronatis, vel inferioribus in quavis spica membranaceis obtusis; utriculi (vix satis maturi) virides, glabri, nervis duobus extramarginalibus prominentibus et aliis subobsoletis instructi, in rostrum breve, apice in dentes duos acutos divergentes fissum, acuminati; bractæ spicarum inferiorum foliaceæ, elongatæ, spicam masculam superantes.

Mar. merid.—Legimus in regione media Atlantis Majoris, in rupestribus siccis montis Djebel Tezah circ. 2000 m. ! et verisimiliter eandem plantam in convalle Aït Mesan supra Arround, ultra 2000 m. s. m. !

Species in grege *C. distantis* collocanda. Ab hac differt imprimis statura multo minore, foliis angustioribus, spicis minoribus, squamis spicæ fœmin. inferioribus obtusis, utriculo minus inflato, nervis minus pro-

minentibus, rostro glabro apice insigniter fisso, et bracteis pluribus longioribus.

CAREX GLAUCA (*Scop. Fl. Carn.* ii. 233; *Kunth, Enum.* ii. 459), = *C. recurva* (*Huds. Fl. Angl.* 413), var. *SERRULATA, nob.*, = *C. serrulata* (*Biv. Sic. Pl. Manip.* iv. 9; *Parl. Fl. It.* ii. 180).

Mar. sept.—In rupibus prope Cap. Spartel! In monte Beni Hosmar prope Tetuan!

—— —, var. *ATTENUATA, nob.*, vel forsā sp. nova?

Mar. sept.—In dumetis montis Beni Hosmar prope Tetuan!

Planta nimis immatura, in grege *C. glaucae* certe collocanda, culmis 2-3 pedalibus, foliis angustissimis subpedalibus, spicis longis attenuatis insignis. Planta ulterius inquirenda.

Ar. Geog. *C. glauca* in Europa fere omni obvia, in America boreali et in Africa boreali rarius occurrit. Var. *serrulata* in regione Mediterranea haud infrequens.

C. serrulata mihi videtur forma meridionalis *C. glaucae*. Species genuina in Europa calidiore rarius occurrit.

GRAMINEÆ (*Juss. Gen.* 28).

LEERSIA (*Soland. ap Swartz. Fl. Ind. Occ.* i. 132).

L. HEXANDRA (*Swartz, Prod.* 21; *Kunth, Enum.* i. 6), = *L. mauritanica* (*Salzm. Exsicc.*; *Steud. Syn. Glum.* i. 2).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Spec. ex herb. Schousboe comm. ecl. Cosson!

Ar. Geog. Nonnisi in insula Jamaica et in agro Tingitano ab auctoribus indicata. Ex spec. in Herb. Kew. non satis exploratis videtur species late diffusa.

LYGEUM (*L. Gen.* 31).

L. SPARTUM (*L., Kunth, Enum.* i. 19).

Mar. merid.—In ditone Reraya circ. 1000 m. semel lectum.

Ar. Geog. In regione Mediterranea calidiore, præsertim centrali, satis frequens. In Ægypto et in Marocco rarius occurrit. Deest in Gallia et in Macaronesia.

CRYPsis (*Ait. Hort. Kew.* i. 48).

C. SCHÆNOIDES (*L., sub Phleo; Lam. Ill.* i. 166, tab. 42; *Kunth, Enum.* i. 22).

Mar. sept.—(*Salzm. Exsicc.*)! (*Webb Herb.*)!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia). Asia media.

C. ACULEATA (*Ait. Hort. Kew.* i. 48; *Kunth, Enum.* ii. 22; *Schousb. Gew. Marok.* 11).

Mar. sept.—Tanger in arenosis maritimis (*Schsb. l. c.*).

Ar. Geog. Regio Mediterranea (exclusa Macaronesia). Europa media. Sibiria.

ALOPECURUS (*L. Gen.* 78).

A. PRATENSIS (*L., Kunth, Enum. i. 24*), var., = *A. caudicans* (*Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Per hemisphæram borealem temperatam late diffusus.

PHALARIS (*L. Gen.* 74).

P. CANARIENSIS (*L., Kunth, Enum. i. 31*).

Mar. sept.—Tanger (*Salzm., spec. in Herb. Kew.*)! (*Webb Herb.*)!

Ar. Geog. Ex insulis Canariensibus orta see. auctores; in regione Mediterranea hinc inde subspontanea. In Europa media advena.

P. BRACHYSTACHIYS (*Link in Schrad. Journ. i. pars 3, 134; Kunth, Enum. i. 32*), = *P. quadrivalvis* (*Lag. G. et Sp. 3*), = *P. nitida* (*Presl, Cyp. et Gram. Sic. 26*).

Mar. sept.—Tanger! Tetuan (1851)!

Ar. Geog. Peninsula Iberica. Italia. Marocco. Insulæ Canarienses.

Nisi fallor, hæc species per regionem omnem Mediterraneam late diffusa est. Spec. in herb. nostro in Macedonia a Friwaldsky lectum verisimiliter huic referendum est.

De notis diagnosticis quibus hæc a *P. canariensi* distinguitur, confer Parlatores (*Fl. It. i. 69*). Ego tamen in spec. nostris video flosculos steriles magnitudine variabiles, et diagnosi mihi incerta manet. Forsan hæc est forma spontanea unde orta est *P. canariensis*, quæ, extra insulas Canarienses, semper culta, vel ex cultura evasa.

P. MINOR (*Retz. Obs. iii. 8; Kunth, Enum. i. 32*), = *P. aquatica* (*Ait. Hort. Kew. i. 56, non L.*), = *P. bulbosa* (*Desf. Fl. Atl. i. 35; Schousb. Gew. Marok. i. 20*).

Mar. sept.—Tanger (*Schsb. l. c.*). Tetuan (1851)!

Mar. occid.—Casa Blanca! Prope Mogador!

Mar. merid.—Sat frequens in regione inferiore. Legi prope Aïn Oumast! In irrigatis urbis Marocco! In glareosis Oued en Fisk! et alibi visa!

Ar. Geog. Regio Mediterranea omnis (inclusa Macaronesia). Spontanea etiam, ut fertur, ad prom. B. Spei.

Species valde variabilis: in pinguibus enata, culmi erecti bipedales et ultra, folia latiora, spicæ sat magnæ; in aridis fit multo minor, spicis abbreviatis, culmis plurimis basi subnodosis geniculatis.

P. NODOSA (*L., Kunth, Enum. i. 32*), = *P. aquatica* (*L. Sp. Pl. 97, et Herb., fide Parlatores*), = *P. tuberosa* (*L. Mant. 557*).

Mar. sept.—Tetuan ad ostium fluminis (1851)!

Mar. occid.—Casa Blanca!

Mar. merid.—Frequens in regione inferiore et ad radices Atlantis Majoris. Sheshaoua! Mesfioua! Sektana! In convalle Aït Mesan usque 1400 m.!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

Phalaris cærulescens (*Desf. Fl. Atl.* i. 56; *Kunth, Enum.* i. 32), = *P. bulbosa* (*Cavan. Icon.* i. tab. 64, non *L.*, nec *Desf.*), = *P. tuberosa* (*Salzm. Exsicc.*, non *L.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb spec.*)!

Ar. Geog. Regio Mediterranea occidentalis (inclusa Macaronesia).

Notæ diagnosticæ quibus hæc a *P. nodosa* distinguitur mihi videntur instabiles, et re vera hæc species perennes a sp. annuis ejusdem gregis (*P. minor*, Retz., et *P. truncata*, Guss.) structura partium floralium ægre distinguuntur. *P. cærulescens* a *P. nodosa* differt imprimis glumarum ala dorsali dilatata eroso-denticulata, nec angusta integerrima; quoad hanc notam *P. minor* intermedia est secundum eel. Parlatore, qui Gramineas Italicas accurate exposuit. *P. minor* et *P. nodosa* gaudent flosculo sterili minimo solitario, dum in *P. truncato* adsunt flosculi neutri duo, et in *P. cærulescente* omnino desunt. Contra beatus Kunth (*Enum.* i. Supplem. 23) flosculos neutros duos in *P. cærulescente* vidit, et ego vidi rudimenta bina in spec. *P. nodosæ* ex Mesfioua. Interea *P. cærulescens* post *P. nodosam* sicut subspecies enumeranda videtur.

P. PARADOXA (*L.*, *Kunth, Enum.* i. 33).

Mar. sept.—Tanger! ex regione collina inter Tanger et Tetuan! Tetuan (*Webb Herb.*)! Ad ostium fluminis Tetuanensis (1851)!

Mar. occid.—Casa Blanca!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

HOLCUS (*L. Gen.* 1146).

H. LANATUS (*L.*, *Kunth, Enum.* i. 34).

Mar. sept.—Tanger (*Webb spec.*)! In monte Djebel Kebir!

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Ourika (forma panicula densa, aristis rectis)!

— —, var. **TUBEROSUS**, = *H. tuberosus* (*Salzm. Exsicc.*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Legi copiose in altero latere freti Hereulis prope S. Roque!

Ab hæc varietate vix differt *H. Reuteri* (Boiss., in Boiss. et Rent. Pug. 119).

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). Europa media. America borealis. Var. in dit Tingitana et in Hispania centrali.

H. SETIGLUMIS (*Boiss. et Reut. Diagn. Pl. Nov. Hisp.* No. 5; *Willk. et Lge. Fl. Hisp.* i. 74), = *H. annuus* (*Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Hispania centralis. Ditio Tingitana.

ANTHOXANTHUM (*L. Gen.* 42).

A. odoratum (*L., Kunth, Enum.* i. 38), var. γ , glumis pubescentibus (*Kunth, l. c.*).

Mar. sept.—Tanger (*Webb Herb.*)! In monte Beni Hosmar pr. Tetuan (ubi hanc var. jam legerat b. Webb)!

Ar. Geog. Species per Europam mediam et Sibiriam late diffusa. In Europa australi et in Africa boreali rarius occurrit. In America boreali forsitan introducta.

A. ovatum (*Lag. Gen. et Spec.* 2; *Kunth, Enum.* i. 38), = *A. lasianthum* (*Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salzm. Exsicc. in Herb. Kew.*, sub *A. lasiantho* et *A. ovato*)!

Mar. occid.—Prope Casa Blanca! Crescit copiose prope S. Roque!

—— —, var. panicula laxiore pauciflora, facie *A. gracili*, Biv., accedens, vix ab *A. Puelii* (Lee. et Lam. Cat. Pl. Auv. 385) diversa.

Mar. sept.—Legi prope Tetuan!

Mar. occid.—Post imbres mensis Maji iterum florens legi juxta Aïn el Hadjar haud procul Mogador!

Ar. Geog. Hispania, ditio Tingitana. *A. Puelii* nascitur in peninsula Iberica et in Gallia media.

A. ovatum indicatur in Sicilia et in Græcia, verisimiliter eum affini, sed satis diverso, *A. gracili*, Biv., commutatum.

Anthoxanthum ovatum ab *A. odorato* statim dignoscitur radice annua, spica abbreviata oblongo-ovata; sed hoc et arete affine *A. Puelii*, mihi, videntur formæ australes sp. vulgaris Europæ. Radix fibrosa *A. odoratæ* in calidioribus Europæ sæpius æstate exsiccata est; inde proles radice annua donata temperiem meridionalem facilius tolerat. Cæterum radix in *A. ovato* interdum perennans. Haud sine dubio pro subspecie enumeravi.

PANICUM (*L. Gen.* 76).

P. repens (*L., Kunth, Enum.* 103; *Schousb. Gew. Marok.* 21), = *P. arenarium* (*Brot. Phyt. Lus.* tab. 6).

Mar. sept.—Tanger (Sehse. l. c., et spec. ex dono Cosson)! (*Salzm. Exsicc.*)!

Ar. Geog. Species ampligea, in regione omni Mediterranea (exclusa Macaronesia), in Africa australi, et in America subtropica diffusa.

P. Crus-Galli (*L., Desf. Fl. Atl.* i. 58), = *Oplismenus Crus Galli* (*Kunth, Enum.* i. 143).

Mar. sept.—Tanger; vidi mense Junio mox floribundum!

Ar. Geog. Species cosmopolita in temperatis (præsertim calidioribus) orbis totius diffusa.

SETARIA (*Pal. Beauv. Agrost.* 51, tab. 13).

S. VERTICILLATA (*L.*, sub *Panico*; *Pal. Beauv. l. c.*; *Kunth, Enum.* i. 152).

Mar. sept.—Vidi prope Tanger!

Mar. occid.—Mazagan!

Ar. Geog. Species amphigea in regione omni Mediterranea, in Africa australi, in India orientali, et in America boreali diffusa. In Europa media hinc inde occurrit, cum seminibus introducta.

DIGITARIA (*Scop. Fl. Carn.* i. 42).

D. SANGUINALIS (*L.*, sub *Panico*; *Scop. Fl. Carn.* i. 52), = *Panicum sanguinale* (*Schousb. Gew. Marok.* 21; *Kunth, Enum.* i. 82).

Mar. sept.—In agris et hortis frequens (*Schsb. l. c.*).

Nisi fallor, vidi prope Tanger, sed spec. non servavi!

Ar. Geog. Species cosmopolita in temperatis et regionibus calidioribus orbis fere totius diffusa.

PENNISETUM (*Pal. Beauv. Agrost.* 59, tab. 13).

P. CILIARE (*L.*, sub *Cenchro*; *Link, fide Steud.*), = *P. cenchroides* (*Rich. in Pers. Syn.* i. 72; *Kunth, Enum.* i. 162), = *P. distylum* (*Guss. Syn. Fl. Sic.* i. 115), = *Cenchrus ciliaris* (*Desf. Fl. Atl.* ii. 387; *Schousb. Gew. Marok.* 20), = *C. rufescens* (*Desf. Fl. Atl.* ii. 388).

Mar. occid.—In arenosis prope Mogador!

Ar. Geog. Species late dispersa, in Africa boreali et australi, in ins. Canariensibus et in Arabia obvia. In Sicilia rarius occurrit. Fertur etiam in America tropica, sed vix genuinum.

PIPTATHERUM (*Pal. Beauv. Agrost.* 18).

P. CÆRULESCENS (*L.*, sub *Milio*; *P. B. l. c.*; *Kunth, Enum.* i. 176), = *Milium cærulescens* (*Schousb. Gew. Marok.* 22).

Mar. occid.—In monte Djebel Hadid haud procul Mogador! Agadir (*Schsb. l. c.*).

Mar. merid.—Prov. Ithala (*Schsb. l. c.*). In regione inferiore Atlantis Majoris haud infrequens. Tasseremout! Supra Amsmiz! Sektana! In convalle Aït Mesan a 1400 m. ! usque 2100 m. !

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

In spec. nostris rami paniculæ interdum linea scabra notati, et pedicelli undique scabri, aristæ, glumis æquilongæ, vel eas paullo superantes. Pedicelli sæpe incrassati, apice turbinato, ut fit in *P. paradoxo*, var. *Balansæ* (*J. Gay in Herb.*).

P. MILIACEUM (*L.*, sub *Agrostide*; *Coss. Not. Crit.* 129), = *P. multiflorum* (*Pal. Beauv. Agrost.* 18; *Kunth, Enum.* i. 177), = *Milium multiflorum*, *Cavan.*, = *Agrostis miliacea* (*Schousb. Gew. Marok.* 24).

“Undique in maceritiis” (*Schsb. l. c.*).

Mar. sept.—Tanger (*Webb Herb.*)! Vidi prope Tetuan!

Mar. occid.—Mogador (*Lowe Cat.*). Aïn el Hadjar haud procul Mogador!

Mar. merid.—Prov. Shedma! Sheshaoua! In umbrosis urbis Marocco!
Ourika! Sektana (*J. D. H.*)!

Rarius ludit glumis purpurascensibus.

Ar. Geog. Regio omnis Mediterranea. Fertur etiam in Sibiria, sed forsitan erronee; deest in Ledebour, Fl. Rossica.

MACROCHLOA (*Kunth, Gram. i. 59*).

M. TENACISSIMA (*L.*, sub *Stipa*; *Kunth, Enum. i. 179*).

Mar. sept.—Tetuan! (*Webb Herb.*)!

Mar. occid.—Prope Mogador visa! In monte Djebel Hadid (*J. D. H.*)!

Mar. merid.—Ad radices Atlantis Majoris haud infrequens. Reraya!
Sektana! Seksaoua! Ex prov. Haha spec. attulerunt Rein et Fritsch!

Ar. Geog. Peninsula Iberica. Africa boreali-occidentalis. Fertur in Attica, sed spec. nulla vidi—an genuina?

In provinciis occidentalibus Imp. Maroccani *M. tenacissima* magna copia ab incolis legitur et in fascias colligata in Angliam ad chartam fabricandam transvehitur.

STIPA (*L. Gen. 90*).

S. ARISTELLA (*L. Syst. Nat. ed. 12, iii. 229*; *Kunth, Enum. i. 180*),
= *Agrostis bromoides* (*L. Mant. 30*; *Schousb. Gew. Marok. 24*), = *Aristella bromoides* (*Bert. Fl. It. i. 690*).

Mar. sept.—Tanger (*Schsb. l. c.*).

Ar. Geog. Europa Mediterranea. Asia Minor. In Algeria rara.

S. TORTILIS (*Desf. Fl. Atl. i. 99, tab. 31*; *Kunth, Enum. i. 180*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Tetuan (1851)!

Mar. occid.—Casa Blanca! Mogador (*Lowe Cat.*). Agadir (*Coll. Indig. 1871*)!

Mar. merid.—In saxosis et aridis regionis inferioris ubique occurrit.
Prov. Shedma pluries lecta! Tasseremout! Reraya! Sektana!

— —, var. *PUBESCENS*. Foliis utrinque basin versus, cum vagina folii supremi, velutino-pubescentibus.

Mar. merid.—Legimus prope Sektana!

Ar. Geog. Regio omnis Mediterranea præsertim calidior (inclusa Macaronesia). Nascitur etiam ad prom. B. Spei.

S. GIGANTEA (*Lag. Gen. et Spec. 3*; *Kunth, Enum. i. 180*).

Mar. merid.—? Specimen defloratum et fructu jam elapso, ob. formam glumarum fere certe huc referendum, ex convalle Ait Mesan attulerunt ell. Rein et Fritsch!

Ar. Geog. Hispania. Italia meridionalis. Sicilia. Africa borealis.

STIPA PARVIFLORA (*Desf. Fl. Atl.* i. 98, tab. 29; *Kunth, Enum.* i. 181).
Mar. merid.—Ad radices Atlantis Majoris. Legimus in ditione Reraya
cire. 1200 m. ! In convalle Aït Mesan usque 1400 m. ! Taradut (*R.
et Fr. spec. in Herb. Kew.*) !

Ar. Geog. Africa borealis. Creta. Hispania.

S. NITENS (*Ball in Journ. Bot.* 1875, 206).

Perennis, dense cæspitosa, foliis rigidis erectis filiformibus, culmo æqui-
longis, arcte convolutis, extus lævibus, facie interna striatis asperulis,
vagina brevi laud inflata, ligula brevissima sub lente fimbriata; pa-
nicula ex ramis brevibus adpressis scabris subcylindraeca; glumæ ex-
teriores inæquales, hyalinæ, nitentes, inferior florem duplo superans
basi trinervosa, nervo medio ineolori ad apicem producto; stipes floris
solitarii brevis plumosus; gluma florifera (palea inferior) paleam
pluries excedens tenuiter villosa; arista pro grege brevis ($1\frac{1}{2}$ -2-polli-
caris), medium versus geniculata, infra genu plumoso-barbata, superne
nuda scabriuscula.

Mar. merid.—Legimus in regione subalpina Atlantis Majoris in convalle
Aït Mesan supra Arround cire. 2200 m. s. m. !

Ex structura partium floralium *S. tortili* valde proxima et pro subspecie
a me olim enumerata. Nimis differt tamen radice perenni habitu
dense cæspitoso, vagina folii superioris non inflata, aristis dimidio
brevioribus. Nullæ aliæ nisi forte *S. compressæ*, R. Br. ex Australia,
vel *S. chrysophyllæ*, ex America australi, comparanda videtur.

ARISTIDA (*L. Gen.* 94).

A. ADSCENSIONIS (*L. Sp. Pl.* 121; *Kunth, Enum.* 190), = *A. gigantea*
(*L. fil. Suppl.* 121), = *A. cærulescens* (*Desf. Fl. Atl.* i. 109, tab. 21;
(*Kunth, l. c.*), = *A. canariensis* (*Willd. Enum.* 99). *Chætariæ* species
(*Pal. Beauv., R. et Sch., Nees ab Esenb., Parlatore*).

Mar. occid.—Legi juxta Aïn el Hadjar haud procul Mogador !

Mar. merid.—Ad radices Atlantis Majoris haud infrequens. Ourika !
Seksaua ! Gurguri ! Aïn Tarsil ! et alibi visa !

Ar. Geog. Africa borealis. Ins. Canarienses et Capitis Viridis. Arabia
Petræa. Insula Adscensionis. Brasilia. In Europa nonnisi in His-
pania australi et in Sicilia detecta.

Statura valde variabilis, interdum planta pumila, sæpe gaudet culmis bi-
pedalibus et ultra.

A. PLUMOSA (*L., Kunth, Enum.* i. 195), = *A. lanata* (*Forsk. Fl. Arab.-
Æg.* 25).

Mar. occid.—In monte Djebel Hadid haud procul Mogador legerunt
ell. Rein et Fritsch (*spec. in Herb. Kew.*) !

Mar. merid.—Legimus in ditione Mesfioua !

Ar. Geog. Algeria interior. Marocco. Ægyptus. Asia Mediterranea.

SPOROBOLUS (*R. Br. Prod.* 170).

S. PUNGENS (*Schreb. Gram.* ii. 46, tab. 27, sub *Agrostide*; *Kunth. Enum.* i. 210)!

Mar. sept.—In maritimis pr. Tanger (spec. ex herb. Schousboe benev. comm. eel. Cosson).

— —, var. *GADITANUS*, = *S. gaditanus* (*Boiss. et Reut. Pug.* 125).

Mar. sept.—“In arenosis maritimis Tingitanis” (*Durand in Herb. Pavon, fide Boiss. et Reut. l. c.*).

Ar. Geog. *S. pungens* in maritimis regionis Mediterraneæ (exclusa Macaronesia) late diffusa est. Var. *gaditanus* nonnisi prope Gades et Tingim detecta.

Ex descriptione *S. gaditanus* differt culmis elatioribus, ligula oblonga, non ad setas reducta. Haud sine dubio pro varietate enumeravi.

AGROSTIS (*L. Gen.* 80).

A. ALBA (*L., Kunth, Enum.* i. 219), var. *COARCTATA* (*Coss. et Dur. in Expl. Scient. Alger.* 63), = *A. maritima*, var. *pseudopungens* (*Lge. Pug.* 31).

Mar. sept.—Tanger (*Salzm. spec. in Herb. Kew.*)! Specimen ex herb. Schousboe comm. Cosson!

Mar. occid.—Mogador (*Lowe Cat.*, sub *A. stoloniferæ*).

— —, var. *MARITIMA* (*Mey. Chlor. Han.* 655), = *A. maritima* (*Lam. Dict.* i. 61; *Salzm. Exsicc.*)!

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

— —, var. *FONTANESII* (*Coss. et Dur. in Expl. Scient. Alg.* 63), = *A. capillaris* (*Desf. Fl. Atl.* i. 69, non *L.*), = *A. divaricata* (*Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! “In sepibus *Agaves americanæ* prope Tanger” (*spec. ex Herb. Schousboe*)!

Ar. Geog. Species in Europa et Asia temperata, et in regione omni Mediterranea frequens. Var. *Fontanesii* in reg. Mediterranea.

Stirps quam maxime ludibunda, in species plurimas ab auctoribus dilaniata, de quibus amplius disserere hic non fas est. Gramineæ Algerienses, et imprimis *Agrostideæ* et *Avenaceæ*, in *Coss. et Dur. ‘Expl. Scient. Algérie’* eximie tractatæ, lectori assiduo commendandæ.

A. verticillata (*Vill. Dauph.* i. 74; *Schousb. Gew. Marok.* 25), = *A. stolonifera* (*Kunth, Enum.* i. 219; *Salzm. Exsicc.*; *L. Herb. sec. Parlatores, an Sp. Pl.* 93?).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb IIerb.*)! (*Schsb. l. c.*). Tetuan (1851)!

Mar. occid.—Mogador!

Mar. merid.—In irrigatis urbis Marocco! In convalle Ait Mesan circ. 1400 m.!

Ab *A. alba* distinguitur glumis obtusis, glumellis subæqualibus. Pro subspecie, ut videtur, enumeranda.

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). Ins. Capitis Viridis. Gallia occidentalis.

AGROSTIS CASTELLANA (*Boiss. et Reut. Diagn. Pl. Hisp.* 26; *Willk. et Lge. Fl. Hisp.* i. 53).

Mar. sept.—Specimina ex herb. Schousboe in dumetis prope Cap. Spartel lecta benev. comm. eel. Cosson! Legi in clivo montis Beni Hossmar prope Tetuan *Agrostidem* immaturam viviparam verisimiliter huc referendam!

Legi in collibus prope S. Roque spec. plantæ genuinæ et varietatem floribus majoribus aristis brevioribus in sequentem transeuntem.

— —, var. HISPANICA, = *A. hispanica* (*Boiss. et Reut. Pug.* 120; *Willk. et Lge. l. c.*).

Mar. sept.—Legi mense Junio in monte Djebel Kebir!

Differt ab *A. castellana*, sec. ell. auctores, flosculis majoribus basi non penicillatis et aristis brevioribus; hæ notæ tamen variabiles.

Ar. Geog. Tam species quam varietas in Hispania et in ditione Tingitana occurrunt.

A. olivetorum (*Gren. et God. Fl. Fr.* iii. 483), species Galliæ australis mihi non satis nota, videtur *A. castellanæ* valde proxima.

A. PALLIDA (*DC. Fl. Fr.* v. 251; *Kunth, Enum.* 221).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! In monte Djebel Kebir!

Mar. occid.—Casa Blanca! In prov. Haha haud procul Mogador!

Ar. Geog. Regio Mediterranea occidentalis (exclusa Macaronesia).

A. SETACEA (*Curt. Fl. Lond. fasc. vi. tab.* 12; *Kunth, Enum.* i. 222).

Mar. sept.—Specimen ex dumetis montis Djebel Kebir a Schousboe lectum comm. eel. Cosson!

Legi in altero latere freti Herculis prope San Roque! Species in Hispania Bætica hucusque non indicata.

Ar. Geog. Species occidentalis ex Anglia usque fretum Herculis dispersa.

GASTRIDIDIUM (*Pal. Beauv. Agrost.* 21, tab. 6).

G. LENDIGERUM (*L. Sp. Pl.* i. 91, sub *Milio*; *Boiss. Voy. Esp.* 647), = *G. australe* (*Pal. Beauv. l. c.*; *Kunth, Enum.* i. 230), = *Agrostis australis* (*L. Mant.* i. 30).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)!

Mar. occid.—In monte Djebel Hadid haud procul Mogador (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—Ad radices Atlantidis Majoris satis frequens. Seksaoua! Mzouda! et alibi, e. g. prope Mskala, lectum!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). Europa occidentalis. Adest etiam in America australi; an introductum?

POLYPOGON (*Desf. Fl. Atl.* i. 66).

P. MONSPELIENSIS (*L.*, sub *Alopecuro*; *Desf. l. c.*; *Kunth, Enum.* i. 232), = *Agrostis panieea* (*Willd. Sp. Pl.* i. 128; *Schousb. Gew. Marok.* 24).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)! (*Schsb. spec. ex. Herb.*)!

Mar. occid.—Mogador! (*Lowe Cat.*). Aïn el Hadjar haud procul Mogador!

Mar. merid.—Satis frequens in regione inferiore, etiam a mari remota. Misra ben Kara! Mesfiona! Sektana! Amsmiz! Ourika (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Regio omnis Mediterranea. Europa occidentalis. In America australi (ad Monte Video) forsan introductus.

P. maritimus (*Willd. in Nov. Act. Soc. Nat. Cur. Berol.* iii. 443; *Kunth, Enum.* i. 233).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. occid.—Casa Blanea!

Præcedenti valde proximus, notæ tamen diagnosticæ subeconstantes, et inde pro subspecie enumerandus.

Ar. Geog. Regio omnis Mediterranea. Africa australis. Asia media. India orientalis.

P. subspathaceus, Req., *P. maritimo* valde proxima species, et a viris oculatissimis J. Gay et Cosson semel conjunctus, differt articulo superiore pedicelli latitudinem suam pluries excedente, nec ut in *P. maritimo* et *P. monspeliensi* latitudinem suam subæquante. In planta nostra juxta Casa Blanea lecta video pedicellos intermedios articulo superiore quam latitudo sua duplo longiore. An igitur hæc nota diagnostica fallax?

LAGURUS (*L. Gen.* 104).

Genus a cl. Kunth inter *Avenaceas* perperam collocatum, eerte inter *Agrostideas* enumerandum.

L. ovatus (*L.*, *Kunth, Enum.* i. 295; *Schousb. Gew. Marok.* 34).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Schsb. l. c.*).

Mar. occid.—Casa Blanea! Mogador (*Lowe Cat.*).

Mar. merid.—Satis frequens in regione inferiore. Sektana! Seksaoua! et alibi pluries visus!

Ar. Geog. Regio omnis Mediterranea. In Europa occidentali rarius occurrit.

AMMOPHILA (*Host, Gram. Austr.* iv. 24).

A. ARENARIA (*L.*, sub *Arundine*; *Link, Enum. H. Berol.* i. 105), = *A. arundinacea* (*Host, l. c.*; *Kunth, Enum.* i. 245).

Mar. sept.—In arenosis litoris Atlantici a prom. Spartel meridiem versus!

Ar. Geog. Frequens in maritimis Europæ septentrionalis et occidentalis. Per litora Mediterranea dispersa, sed minus frequens. Nascitur in America boreali, sed deest in Macaronesia.

ARUNDO (*L. Gen.* 93).

A. DONAX (*L., Kunth, Enum.* i. 246; *Schousb. Gew. Marok.* 34).

Per territorium omne Maroccanum nobis notum ab incolis cultum. Schousboe non vidit spontaneam. In regione inferiore et ad radices Atlantis Minoris pluries vidimus nondum florentem in locis a cultura remotis subspontaneam.

Ar. Geog. Species verisimiliter ex Africa boreali subtropica orta, et inde in temperatas calidiores orbis veteris advecta, in regione Mediterranea sæpe subspontanea.

A. PLINIANA (*Turra, Fl. It. Prod.* i. 63; *Parlat. Fl. It.* i. 218), = A. mauritanica (*Desf. Fl. Atl.* i. 106; *Salzm. Exsicc.*; *Kunth, Enum.* i. 246).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Ar. Geog. Regio Mediterranea imprimis calidior (exclusa Macaronesia).

AMPELODESMOS (*Link, Hort. Reg. Berol.* i. 136).

A. TENAX (*Link, l. c.*; *Kunth, Enum.* i. 249), = Arundo Ampelodesmos (*Cyr. Pl. rar. Neap. fasc.* ii. 30), = A. festucoides (*Desf. Fl. Atl.* i. 108).

Mar. sept.—Vidi prope Tanger nondum florentem, sed foliis facile dignoscendum. Spec. fructiferum ex herb. Schousboe comm. cel. Cosson!

Ar. Geog. Regio Mediterranea imprimis occidentalis (exclusa Macaronesia). Ex Oriente non vidi specimina.

PHRAGMITES (*Trin. Fund. Agrost.* 134).

P. COMMUNIS (*Trin. l. c.*; *Kunth, Enum.* i. 251), = Arundo Phragmites (*L. Sp. Pl.* 120).

Mar. merid.—In convalle Aït Mesan circ. ad 1400 m. legimus nondum florentem culmis et paniculis exsiccatis anni præcedentis!

Ar. Geog. Species cosmopolita in temperatis totius orbis diffusa. Ex Macaronesia non vidi.

ECHINARIA (*Desf. Fl. Atl.* ii. 385).

E. CAPITATA (*L., sub Cenchro*; *Desf. l. c.*; *Kunth, Enum.* i. 257).

Mar. merid.—In regione inferiore et media Atlantis Majoris. In con-

valle Aït Mesan eire. 1400 m., et iterum supra Arround ad 2200 m.!

In monte Djebel Tezah eire. 2000 m.!

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia).

CYNODON (*Rich. in Pers. Syn.* i. 85).

C. DACTYLON (*Rich. l. c.*; *Kunth, Enum.* i. 259), = *Panicum Dactylon* (*L., Schousb. Gew. Marok.* 21).

“In arvis totius imperii frequentissime occurrit” (*Schsb. l. c.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. oecid.—Casa Blanca! Mazagan! Mogador (*Lowe Cat.*)!

Mar. merid.—Ocurrit, sed minus frequens, in terris interioribus. Legimus prope Sheshaona! et pluries alibi vidimus. Onrika (*R. et Fr. spec. in Herb. Kew.*)!

Ar. Geog. Species cosmopolita, in temperatis et calidioribus orbis totius diffusa.

SPARTINA (*Schreb. in L. Gen. Pl.* i. 43).

S. STRICTA (*Roth, Catal. Bot.* iii. 9; *Kunth, Enum.* i. 278), = *Lin. pungens* (*Pers. Syn.* i. 72; *Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Spec. ex Herb. Schousboe*)!

Ar. Geog. Europa occidentalis. Hispania Gaditana. Litora maris Adriatici. Ditio Tingitana.

AIRA (*L. Gen.* 81).

A. CARYOPHYLLEA (*L., Kunth, Enum.* i. 289), var. α (*Coss. in Expl. Scient. Alg.* 95).

Mar. sept.—Tanger (*Salzm. Exsicc.*)!

Mar. merid.—In regione media Atlantis Majoris. In convalle Amsmiz eire. 1600 m. In monte Djebel Tezah usque 2200 m.!

— — —, var. β , MICROSTACHYA (*Coss. l. c.*).

Mar. sept.—Tanger (*Webb spec.*)! Tetuan in saxosis!

Mar. merid.—Ad radices Atlantis Majoris in ditone Reraya (*J. D. H.*)!

— — —, var. CUPANIANA (*Coss. l. c.*), = *A. Cupaniana* (*Guss. Syn. Fl. Sic.* i. 145).

Mar. oecid.—Legi in prov. Haha laud procul Mogador!

Ar. Geog. Species amphigea. In Europa et Asia media, in regione omni Mediterranea, necnon in Africa australi et in America meridionali obvia. Varietates enumeratæ in regione Mediterranea occurrunt.

A. CAPILLARIS (*Host, Gram. Austr.* iv. 20), = *A. elegans* (*Gaud. Agrost. Helv.* i. 130; *Kunth, Enum.* i. 289).

Mar. sept.—Tanger (*Salzm. spec. in Herb. Kew.*)!

Ar. Geog. In Europa australi fere omni. Ex Algeria non vidi. Deest, ut videtur, in Asia Mediterranea.

AIROPSIS (*Pal. Beauv. Agrost.* 71, tab. 14).

- A. MINUTA (*Læfl. It. Hisp.*, sub *Aira*; *Desv. in R. et Sch. Syst.* ii. 578),
 = *Catabrosa minnta* (*Triu. Fuud. Agrost.* 136; *Kunth, Enum.* i. 370),
 = *Moliniera minuta* (*Parlat. Fl. It.* 237).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Ar. Geog. Peninsula Iberica. Ditione Tingitana. Italia. Græcia.
 Persia.

TRISETUM (*Pers. Syn.* i. 97).

- T. PANICEUM (*Lam. Ill.* 202, sub *Avena*; *Pers. l. c.*), = T. neglectum
 (*R. et Sch. Syst.* ii. 660; *Kunth, Enum.* 296).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Spec. ex Herb. Schousboe*)!

Mar. occid.—Casa Blanca! Mazagan! In provincia Haha prope
 Mogador!

Mar. merid.—In regione inferiore vulgatissimum, sed vix supra 900 m. s. m.,
 nec in convallibus interioribus se protendit. Shedma! Aïn Oumast in
 saxosis aridissimis! Sheshaoua! Misra ben Kara! Mesfioua! Mzouda!
 Mskala!

Ludit panicula contracta facie fere *Kæleria phleoidis*.

Ar. Geog. In regione Mediterranea occidentali (inclusa Macaronesia)
 late diffusa.

- T. PUMILUM (*Desf. Fl. Atl.* i. 103, sub *Avena*; *Kunth, Enum.* i.
 297).

Mar. occid.—Legi in arenosis maritimis prope Mogador mense Aprili
 jam defloratum!

Ar. Geog. Species, sec. cel. Cosson, late diffusa per Africam borealem
 usque Arabiam et Persiam. Occurrit etiam in Hispania austro-
 orientali, in insulis Canariensibus et ad prom. B. Spei.

- T. FLAVESCENS (*L.*, sub *Avena*; *Pal. Beauv. Agrost.* 88, tab. 18;
Kunth, Enum. i. 298).

Mar. merid.—In regione inferiore Atlantis Majoris. In convalle Aït
 Mesan! in monte supra Seksaoua!

Planta nostra vaginis villosa-pubescentibus fere media est inter typum
 Linnæanum et *T. velutinum* (*Boiss. Voy.* 653, tab. 174).

Ar. Geog. Species in temperatis frigidioribus hemisphæræ borealis late
 diffusa; in calidioribus fit rarior.

GAUDINIA (*Pal. Beauv. Agrost.* 97, tab. 19).

- G. FRAGILIS (*L.*, sub *Avena*; *Pal. Beauv. l. c.*; et auctt. plurim.), =
Avena fragilis (*Kunth, Enum.* i. 303; *Schousb. Gew. Marok.* 34).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! (*Schsb. l. c.*).

Mar. occid.—Casa Blanca! Mogador!

Mar. merid.—Ourika! Sektana!

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

Avena (*L. Gen.* 91).

A. SATIVA (*L., Kunth, Enum.* i. 301).

Mar. occid.—In arenosis multis prope Mogador!

Mar. merid.—Shedma! Misra ben Kara!

E cultura evasa hodie subspontanea.

Ar. Geog. De patria *A. sativæ* hic disserere non oportet.

A. STERILIS (*L., Kunth, Enum.* i. 303; *Schousb. Gew. Marok.* 33).

“In agris et aggeribus hand raro” (*Schsb. l. c.*).

Mar. occid.—Casa Blanca!

Mar. merid.—Reraya! Sektana!

Ar. Geog. In regione Mediterranea (inclusa Macaronesia) verisimiliter spontanea. In Europam mediam cum seminibus introducta.

A. BARBATA (*Brot. Fl. Lus.* i. 108; *Willk. et Lge. Fl. Hisp.* i. 68), =

A. hirsuta (*Roth, Catal. Bot.* iii. 19; *Kunth, Enum.* i. 302), = *A. fatua* (*Salzm. Exsicc., non L.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Inter Tetuan et Centa!

Mar. occid.—In monte Djebel Hadid hand procul Mogador!

Mar. merid.—Satis frequens in regione inferiore et ad radices Atlantis Majoris. Prov. Shedma! Aïn Oumast! Mesfioua! Tasseremout! Sektana (*R. et Fr. spec. in Herb. Kew.*)!

In aridis prov. Shedma ludit multiculmis, culmis brevioribus, glumella inferiore insigniter nervosa (*A. deusta*, nob. olim. in Herb.).

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). Etiam in Gallia occidentali obvia.

— —, var.? Panicula ramosa diffusa, glumella inferiore longe bisetosa.

Mar. occid.—Casa Blanca!

Fere media inter *A. barbatam* et *A. longiglumem* (*Dur. in Duch. Rev. Bot.* i. 359). Hæc differt glumis longioribus, sed mihi videtur nimis proxima *A. barbata*.

A. PRATENSIS (*L., Kunth, Enum.* i. 300).

Mar. sept.—Prope Tetuan!

Ar. Geog. Europa media; in regione Mediterranea hinc inde rarius occurrit. Deest in Macaronesia.

A subspecie sequente differt imprimis rachis pilosa.

A. bromoides (*Gouan, Hort. Monsp.* 52; *Willk. et Lge. Fl. Hisp.* i. 69)

Var. *α*, GENUINA, *Willk. l. c.*, = *A. australis* (*Parl. Fl. It.* i. 285).

Mar. merid.—In ditione Reraya! Occurrit lusus foliis rigidis subconvolutis, vaginis cum culmo scabris in rupibus supra Seksaoua!

Var. *β*, MICROSTACHYA, *Willk.*

Mar. merid.—Mesfioua! Milhaïn! Seksaoua!

A. bromoides mihi videtur proles australis *O. pratensis*, et sicut subspecies enumeranda.

Ar. Geog. Regio Mediterranea (exclusa Macaronesia). Ex Oriente non vidi. *Var. microstachya* non nisi in Hispania indicatur.

ARRHENATHERUM (*Pal. Beauv. Agrost.* 55, tab. 11).

A. ELATIUS (*M. K. Deutschl. Fl.* i. 546; *Parl. Fl. It.* i. 293), = *A. avenaceum* (*Pal. Beauv. l. c.*; *Kunth, Enum.* i. 307), = *Avena elatior* (*L., Desf. Fl. Atl.* i. 100).

Mar. merid.—In convalle Amsmiz!

Ar. Geog. Europa media. Regio omnis Mediterranea (inclusa Macaronesia).

POA (*L. Gen.* 83).

P. ANNUA (*L., Kunth, Enum.* i. 349).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! Cap. Spartel!

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—In regione inferiore, et in Atlante Majore usque reg. subalpinam. In ruderatis urbis Marocco! Ourika! Sektana! In convalle Amsmiz usque 1500 m. s. m.! In convalle Aït Mesan usque in ascensum jugi Tagherot circ. 2700 m.!

Ar. Geog. Species cosmopolita per orbem fere totum diffusa.

P. BULBOSA (*L., Kunth, Enum.* i. 352).

Mar. sept.—Inter Tanger et Tetuan! Tetuan (*Webb Herb.*)! In monte Beni Hosmar!

Mar. merid.—Ad radices Atlantis Majoris, unde in regionem alpinam ascendit. Ourika! In convalle Aït Mesan et in ascensu jugi Tagherot usque 2800 m.! In monte Djebel Tezah usque 2900 m.!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). Europa et Asia media.

P. TRIVIALIS (*L., Kunth, Enum.* i. 352).

Mar. sept.—Tanger!

Mar. merid.—Ourika! In convalle Aït Mesan circ. 1400 m.! et iterum circ. 2100 m.!

Ar. Geog. In temperatis hemisphæræ omnis borealis. In Macaronesia non deest.

P. PRATENSIS (*L., Kunth, Enum.* i. 352).

Mar. merid.—In regione inferiore et media Atlantis Majoris. In convalle Aït Mesan usque 2100 m.! In convalle Amsmiz 1400–1600 m.! In monte Djebel Tezah usque 2400 m.!

Ar. Geog. Europa et Asia temperata. In montosis regionis Mediterraneæ calidioris infrequens. Deest in Macaronesia. Ex Hispania meridionali non vidi, nec ex Algeria.

POA COMPRESSA (*L.*, *Kunth, Enum.* i. 355; *Schousb. Gew. Marok.* 26).
Mar. sept.—“In pratis siccis regionis Tingitanæ raro occurrit” (*Schsb. l. c.*).

Ar. Geog. Species in temperatis hemisphæræ borealis late diffusa.
Deest in Macaronesia.

GLYCERIA (*R. Br. Prod.* 179).

G. FLUITANS (*R. Br. l. c.*; *Kunth, Enum.* i. 367),=Festuca fluitans,
L.,=Poa fluitans (*Scop. Fl. Carn.* 106).

Mar. sept.—Tanger (specimen typo Linnæano conforme)!

Ar. Geog. Species fere cosmopolita, in Europa omni, in regione Mediterranea, in utraque America et in Australia diffusa. Ex Macaronesia non vidi.

— —, var. PLICATA, *Griseb.*,=G. plicata (*Fries, Mant.* iii. 176;
Koch, Syn. Fl. Germ. ed. 2, 932).

Mar. sept.—Præcedenti magis frequens. Tanger (*J. D. II.*)!

Mar. merid.—Leginus prope Onrika!

Ar. Geog. Europa fere omnis. Africa borealis.

Hæc et *G. spicata* (*Guss. Syn. Fl. Sic.* ii. 784) notis diagnosticis ad speciem delimitandum idoneis non carent; characteres tamen variabiles sunt, et ambo, monente *cel. Cosson*, pro varietatibus enumerandæ.

G. DISTANS (*L.*, sub *Poa*; *Wahlenb. Fl. Upsal.* 36; *Rchb. Icon. Fl. Germ.* fig. 1609),=Festuca distans (*Kunth, Enum.* i. 393),=Atropis distans (*Griseb. in Led. Fl. Ross.* iv. 388; *Coss. in Expl. Scient. Alg.* 139),=Puccinellia distans (*Parlat. Fl. It.* i. 367), var. FESTUCÆFORMIS (*Coss. l. c.*),=G. festucæformis (*Heynh. in Rchb. Fl. Excurs.* i. 45).

Mar. sept.—Tanger (spec. mancum)! (Specimen ex herb. Schousboe benev. comm. *Cosson*)!

— —, var. TENUIFOLIA (*Coss. l. c.*),=G. tenuifolia (*Boiss. et Reut. Pug.* 127, non *Presl*),=G. leptophylla (*Steud. Syn. Pl. Glum.* i. 286).

Mar. sept.—Prope Tanger (*Webb spec.*)!

Ar. Geog. Species variabilis per Europam et Asiam mediam et regionem Mediterraneam diffusa. Var. *tenuifolia* in Hispania et in Africa boreali.

BRIZA (*L. Gen.* 84).

B. MAXIMA (*L.*, *Kunth, Enum.* i. 371; *Schousb. Gew. Marok.* 27).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Schsb. l. c.*). In monte Djebel Kebir! Tetuan (1851)! (*Webb Herb.*)!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). In dit. Maroccana infrequens. Obvia etiam in Africa australi, et in India orientali.

BRIZA MINOR (L., *Kunth, Enum.* 372), = *B. virens* (L., *Kunth, l. c.*; *Schousb. Gew. Marok.* 26).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb spec.*)! (*Schsb. l. c.*). In monte Djebel Kebir!

Mar. merid.—In ditione Mesfiona!

Ar. Geog. Species amphigea, per regionem Mediterraneam omnem diffusa. In Africa australi, America meridionali, et in insula Mauritii obvia.

MELICA (L. *Gen.* 82).

M. MINUTA (L., *Kunth, Enum.* i. 376), = *M. pyramidalis* (*Poir. Voy.* ii. 95; *Schousb. Gew. Marok.* 25; *Coss. in Expl. Scient. Alg.* 136).

Mar. sept.—Tanger! (“In fruticetis rarissime occurrit,” *Schsb. l. c.*) (*Salzm. Exsicc.*)! In rupibus prope Tetuan! In monte Beni Hosmar!

— —, var. LATIFOLIA (*Coss. l. c.* 137), = *M. major* (*Parlat. Fl. It.* i. 306; *Willk. et Lge. Fl. Hisp.* i. 84; *an Sibth. et Sm. Fl. Gr. Prod.* i. 51?), = *M. pyramidalis* (*Bert. Amæn. It.* 329, non *Poir.*), = *M. arrecta* (*Kmze, Chlor. Hisp.* No. 621 a).

Mar. sept.—Tanger (*Webb spec.*)! In monte Djebel Kebir! In regione collina inter Tanger et Tetuan! Prope Tetuan (1851)!

Ar. Geog. Tam species quam varietas per regionem Mediterraneam (exclusa Macaronesia) late diffusæ sunt.

Stirps variabilis, ab auctoribus diu vexata, et in species plurimas ægre extricandas dilaniata. Genus omne a cel. Cosson optime illustratum.

M. CILIATA (L., *Kunth, Enum.* i. 375; *Coss. in Expl. Scient. Alg.* 133).

Mar. merid.—In regione inferiore videtur potius frequens. Legimus in prov. Shedma! prope Sektana! et in rupibus supra Seksaoua!

Spec. supra Seksaoua lecta proxime accedunt *M. nebrodensi* (*Parlat. Fl. Palerm.* 1, 120). Hanc tamen, duce cel. Cosson, habeo pro forma rupestri *M. ciliatæ*.

— —, var. MAJOR, thyrso lobato, = *M. Magnolii* (*Gren. et Godr. Fl. Fr.* iii. 550).

Mar. sept.—Legi prope Tanger mense Junio! Prope Tetuan (1851)!

Mar. occid.—In prov. Haha haud procul Mogador!

Mar. merid.—In convalle Aït Mesau cire. 1400 m.!

Ar. Geog. Species variabilis per regionem fere omnem Mediterraneam (inclusa Macaronesia) late diffusa. Occurrit etiam in Europa media et in Sibiria. Var. *major* in reg. Mediterranea hinc inde occurrit.

M. CUPANI (*Guss. Pr. Fl. Sic. Suppl.* 17; *Coss. in Expl. Scient. Alg.* 134).

Mar. merid.—In rupibus prope Seksaoua! et Milhaïn!

— —, var. MINOR, foliis brevioribus rigidis subdistichis, = *M. humilis* (*Boiss. Voy. Esp.* 662, tab. 177).

Mar. merid.—In regione inferiori et media Atlantis Majoris. In convalle Aït Mesau a 1600 m. usque 2300 m.! In convalle Amsmiz usque in elivo montis Djebel Tezali (1500–1800 m.)!

Ar. Geog. Species, ut videtur, per regionem Mediterraneam calidiorem ex Imp. Maroccano per Africam borealem, Hispaniam, Siciliam, Asiam Minorem, Syriam, et Persiam usque in montes Himalaya diffusa. Vidi specimen ex Aghanistan in Herb. Kew., nec satis differt *M. Jacquemonti*, Decaisne. Var. nostra in Hispania et in Atlante.

M. Cupani a *M. ciliata* præcipue distinguitur glumella inferiore flosculi inferioris in quavis spicula dorso undique dense pilosa, nec tantum margine ciliata. Nota equidem non gravis, sed mihi videtur bene distincta species.

KÆLERIA (*Pers. Syn.* i. 97).

K. VILLOSA (*Pers. l. c.*; *Kuuth, Enum.* i. 382, et plurim. auct.), = *K. pubescens* (*Pal. Beauv.*; *Coss. Expl. Scient. Alger.*), = *Alopecurus litoricus* (*All. Fl. Ped.* ii. 235), = *Aira pubescens* (*Vahl, Symb.* iii. 9), var. LONGEARISTATA (*Coss. l. c.*), = *K. hispida* (*Salzm. Exsicc.*, non DC.), = *K. Webbiana* (*Parlat. in Giorn. Bot. Ital.* ii. 376), = *K. Salzmanni* (*Boiss. in Boiss. et Reut. Pug.* 123).

Mar. sept.—Tanger (*Webb spec.*)! Ex monte Djebel Kebir (spec. ex herb. Schousboe commun. cel. Cosson)!

Ar. Geog. Species per regionem Mediterraneam imprimis centram dispersa. Deest in Macaronesia. Varietatem nonnisi ex agro Tingitana vidi.

Cel. Cosson nomen triviale *pubesceas* antiquiori *villosa* prætulit; nomen tamen Allionii prioritatis jure gaudet, sed nolui mutare nomen ab auctoribus præcipuis acceptum.

K. PHLEOIDES (*Vill. Dauph.* ii. 95, tab. 2, sub *Festuca*; *Pers. Syn.* i. 97; *Kuuth, Enum.* i. 383), = *K. aristata* (*Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! Inter Tanger et Tetuan!

Mar. occid.—Prope Mogador!

Mar. merid.—Frequens in regione inferiore. Prov. Shedma! Misra ben Kara! In ruderatis urbis Marocco! Mesfoua! Ourika!

— —, var. SUBMUTICA, nob. Glumella inferiore puberula submutica, vel breviter aristulata, glumella flosculi superioris dorso recurva. Refert *K. berytheam*, Boiss. et Blanche.

Mar. merid.—Legi prope Amsmiz!

Ar. Geog. Species in regione Mediterranea (inclusa Macaronesia) late diffusa.

SCHISMUS (*Pal. Beauv. Agrost.* 74).

S. CALYCINUS (*L.*, sub *Festuca*; *Coss. in Expl. Scient. Alg.* 138), =
S. marginatus (*Pal. Beauv. l. c.*; *Kunth, Enum.* i. 385).

Mar. occid.—Prope Mogador (*Lowe Cat.*).

Mar. merid.—Prov. Shedma! In planitie prope urbem Marocco! Prov.
Mtouga!

Ar. Geog. Species per regionem Mediterraneam dispersa. Deest in
Italia et in Macaronesia.

DACTYLIS (*L. Gen.* 86).

D. GLOMERATA (*L.*, *Kunth, Enum.* i. 386).

Mar. occid.—Casa Blanca (spec. typo Europæo conforme)!

— —, var. AUSTRALIS (*Willk. et Lge. Fl. Hisp.* i. 88), = D. his-
panica (*Roth, Catal. Bot.* i. 8; *Kunth, l. c.*).

Mar. sept.—Tanger (*Webb spec.*)!

Mar. occid.—Casa Blanca (*J. D. H.*)!

Mar. merid.—Satis frequens in regione inferiore et in Atlante Majore.
Prov. Shedma! Amsmiz! Ourika (*J. D. H.*)! Seksaoua! In
convalle Aït Mesan usque 2400 m. s. m.!

— —, var. JUNCINELLA (*Boiss. Voy. Esp.* 665), = D. juncinella,
Bory.

Mar. merid.—In regione media Atlantis Majoris. In convalle Aït
Mesan circ. 1500 m.! In monte Djebel Tezah circ. 2000 m.!

Spica in spec. nostris valde diminuta; nisi vidissem spec. Hispanica in-
termedia pro sp. diversa habuerim.

Ar. Geog. Species per hemisphæram borealem late diffusa. Var. *australis*
in calidioribus frequens; var. *juncinella* in montanis Hispaniæ et Africæ
borealis.

CYNOSURUS (*L. Gen.* 87).

C. ECHINATUS (*L.*, *Kunth, Enum.* i. 388; *Schousb. Gew. Marok.* 27).

Mar. sept.—Tanger (*Webb Herb.*)! (*Schsb. l. c.*).

Ar. Geog. Species in regione Mediterranea (inclusa Macaronesia) late
diffusa, etiamsi in territorio nostro rara videtur. Occurrit in calidiori-
bus Europæ mediæ, et indicatur ad promont. B. Spei, et ad Monte
Video, sed verisimiliter ex Europa allata.

C. ELEGANS (*Desf. Fl. Atl.* i. 82, tab. 17; *Kunth, Enum.* i. 389), = C.
polybracteatus (*Willk. et Lge. Fl. Hisp.* i. 90, non *Poir.*).

Mar. merid.—In regione inferiore et media Atlantis Majoris. In ditione
Reraya ad 1100 m.! In convalle Amsmiz et in clivo montis Djebel
Tegah usque 2200 m.!

Ar. Geog. Regio fere omnis Mediterranea (inclusa Macaronesia).

Monente ecl. Cosson C. *polybracteatus*, *Poir. Voy.* ii. 97, est species a
C. *elegante* probe diversa paucis cognita.

CYNOSURUS AUREUS (*L., Desf. Fl. Atl. i. 83; Willk. et Lge. Fl. Hisp. i. 90; Schousb. Gew. Marok. 27*), = *Lamarekia aurea* (*Mœnch, Method. 201; Kunth, Enum. i. 389*).

“In saxosis per totum imperium” (*Schsb. l. c.*).

Mar. sept.—Tetuan!

Mar. occid.—Aïn el Hadjar haud procul Mogador (*J. D. H.*)!

Mar. merid.—In regione inferiore satis frequens. Aïn Oumast! Mskala!

Seksaona! Amsniz!

Ar. Geog. Regio Mediterranea (inclusa Macaronesia).

FESTUCA (*L. Gen. 88*).

F. MEMPHITICA (*Spreng. Hort. Hal. Add. i. 20, sub Dactylide; Coss. Not. Crit. 183*), = *F. divaricata* (*Clem., fide Willk.; Kunth, Enum. i. 392, pro parte, non Desf.*), = *Scleropoa memphitica* (*Parlat. Fl. It. i. 471*), = *Cutandia scleropoides* (*Willk. in Bot. Zeit. 1860, 130*), genus pessimum.

Mar. occid.—Legimus in arenosis prope Mogador!

Ar. Geog. Species per regionem desertorum ex Arabia usque Algeriam Saharensis diffusa, inde per Bithyniam et Transeucasiam dispersa. Habitat insuper in Hispania australi ad Cabo de Gata, et in plaga Mogadorensi.

F. MARITIMA (*DC. Fl. Fr. iii. 47; Kunth, Enum. i. 392*), = *Triticum maritimum* (*L., et auct. plurim.*).

Mar. sept.—Tanger! (spec. ex herb. Schoushoe comm. eel Cosson)!

Ar. Geog. Regio Mediterranea imprimis occidentalis (exclusa Macaronesia). Ex Oriente non vidi.

F. RIGIDA (*L., sub Poa; Kunth, Enum. i. 392*).

Mar. sept.—Tanger! Inter Tetuan et Ceuta!

Mar. occid.—Casa Blanca!

Mar. merid.—Valde frequens in planitie et in regione inferiore Atlantis Majoris. Aïn Oumast! Misra ben Kara! Mesfioua! Reraya! In convalle Aït Mesan usque 1400 m.!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). Europa occidentalis.

F. hemipoa (*Del. in Spreng. Syst. Veg. iv. pars 2, 36; Kunth, Enum. i. 411*), = *Scleropoa hemipoa* (*Parlat. Fl. It. i. 472*), = *F. rigida*, var. *hemipoa* (*Coss. in Expl. Scient. Alg. 182*).

Mar. occid.—Legimus in arenosis prope Mogador.

Pace eel. Cosson hæc stirps, præcedenti valde proxima, differt tamen notis satis stabilibus, et mihi videtur sicut subspecies enumeranda.

Ar. Geog. Africa boreali-occidentalis. Gallia anstralis; Italia.

F. TUBERCULOSA (*Moris in Atti terz. Riun. Sc. Ital. 489, sub Catapodio; Coss. in Expl. Scient. Alg. 189, tab. 41*), = *Castellia tuberculata* (*Tineo, Pl. rar. Sic. fasc. ii. 18; Parlat. Fl. It. i. 480*).

Mar. merid.—Legi in planitie Maroecana prope Sheshaoua!

Ar. Geog. Species in Africa boreali-occidentali, in Siciliæ insula Linosa, et in Sardinia, ubique rara.

FESTUCA UNILATERALIS (*Schrad. Cat. Gött.* 1814; *Kunth, Enum.* i. i. 395), var. ARISTATA, = *F. tenuiflora* (*Schrad. Fl. Germ.* i. 345; *Kunth, l. c.*), = *Nardurus unilateralis*, var. *aristatus* (*Boiss. Voy. Esp.* 667), = *Triticum Nardus* (*DC. Fl. Fr.* iii. 87).

Mar. merid.—Legi in convalle Ait Mesan circ. 1300 m. s. m. lusus spiculis glabris et pilosis semel crescentes!

Ar. Geog. Regio Mediterranea (exclusa Maearonesia).

F. BROMOIDES (*L., Coss. in Expl. Scient. Alg.* 172; *Schousb. Gew. Marok.* 27), = *F. uniglumis* (*Soland. in Ait. Hort. Kew.* i. 108; *Kunth, Enum.* i. 196), = *Vulpia membranaea* (*Link, Hort. Berol.* i. 147).

Mar. sept.—Tanger!

Ar. Geog. Europa occidentalis. Regio Mediterranea fere omnis (exclusa Maearonesia).

F. ALOPECURUS (*Schousb. Gew. Marok.* 28; *Kunth, Enum.* i. 397).

Mar. sept.—Tanger! (*Schsb. l. c., spec. comm. Cosson*)! (*Webb spec.*)! (*Salzm. Exsicc.*)! Ad ostium fluminis Tetuanesis!

Mar. occid.—Casa Blanca! Mazagan! Aïn el Hadjar hand procul Mogador!

Ar. Geog. Marocco. Hispania. Italia occidentalis. Fertur etiam in Græcia, sed spec. non vidi.

Præcedenti structura proxime affinis; sed glumæ semper majores pluries latiores. Species valde ludibunda, cujus varietates plurimæ in Willk. et Lge. *Fl. Hisp.* i. 92, enumerantur.

Inter specimina plurima juxta Mazagan lecta video nonnulla glumellis minute seabris, nec eiliatis; et juxta Aïn el Hadjar legi specimen glumellis undique villosis, aristis brevissimis.

F. MYURUS (*L., Kunth, Enum.* i. 396), = *Vulpia Myurus* (*Gmel. Fl. Bad.* i. 8).

Mar. sept.—Juxta Tetuan!

—— —, var. SCIUROIDES, = *F. seiuroides* (*Roth, Catal. Bot.* ii. 11), = *F. bromoides* (*Kunth, Enum.* i. 396, et *auct. plurim., non L.*).

Mar. sept.—Tanger (*Webb Herb.*)!

Mar. occid.—Mogador (*Lowe Cat.*).

—— —, var. BROTERI, = *Vulpia Broteri* (*Boiss. et Reut. Pug.* 128), = *Vulpia seiuroides*, var. *longearistata* (*Willk. et Lge. Fl. Hisp.* i. 91).

Mar. sept.—Inter Tetuan et Centa!

Ar. Geog. Species valde ludibunda; forma eum typica eum var. *seiuroide* in Europa media et in regione omni Mediterranea. Fertur etiam in America boreali et ad prom. B. Spei. Var. *Broteri* per regionem Mediterraneam diffusa.

Festuca ciliata (*Danth. in DC. Fl. Fr.* iii. 55; *Kunth, Enum.* i. 396),
= *F. Myurns*, var. γ (*Coss. in Expl. Scient. Alg.* 175).

Mar. merid.—Prope Seksaoua!

Ar. Geog. Stirps verisimiliter regionis Mediterraneæ propria. Fertur in Europa media sed spec. genuina non vidi.

Planta *F. Myurus* valde proxima; sed videtur satis constans, nec vidi formas intermedias. Pro subspecie enumeravi.

F. GENICULATA (*L.*, sub *Bromo*; *Willd. Enum.* i. 118), = *F. stipoides* (*Desf., pro parte*; *Salzm. Exsicc.*).

Occurrit frequentissime in omni territorio Maroccano nobis noto.

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (spec. ex Herb. Schousboe)! In Monte Djebel Kebir!

Mar. occid.—Legi prope Casa Blanca lusum aristis longioribus (*Vulpia attenuata*, *Parlat. Fl. Palerm.* i. 195)!

Mar. merid.—Fere ubique in regione inferiori, usque in regionem mediam Atlantis Majoris ascendens. Mesfioua! Ourika! Reraya! Seksaoua! In convalle Amsmiz usque 2000 m. s. m.! In monte Djebel Tezah eire. 1900 m.!

— —, var. *PUMILA*, *nob.* Planta pumila multieulmis, subæspitosa; panicula contracta ramis abbreviatis; gluma inferior superiore tertia parte brevior; glumella inferior latior abrupte in aristam brevem contracta, superior bicarinata, earinis viridibus seabris vel breviter pilosulis. Confer var. *confertam*, *Coss. Pl. Crit.* 48. Nostra tamen notis indicatis a typo longius reedit.

Mar. occid.—Legimus prope Mogador! et in prov. Iiana finitima! et juxta Aïn el Hadjar.

Mar. merid.—Satis frequens in aridis regionis inferioris. Legimus prope Aïn Oumast! In convalle Amsmiz! Juxta Aïn Tarsil!

Specimen ex convalle Amsmiz panicula subsecunda ramis divaricatis rigidiusculis refert *F. delicutulam*, *Lagasca*. Planta ad Aïn Tarsil lecta panicula valde attemmata, glumis paullo longioribus, propior videtur *F. Myurus*, sed mihi potius ad hanc varietatem *F. geniculata* referenda videtur.

Ar. Geog. *F. geniculata* per regionem Mediterraneam occidentalem late diffusa, deest in Oriente et in Macaronesia.

F. DURIUSCULA (*L., Kunth, Enum.* i. 399), var. *INDIGESTA* (*Boiss. Voy. Esp.* 671), = *F. indigesta* (*Boiss. El.* 64; *Wilk. et Lge. Fl. Hisp.* i. 94).

Mar. merid.—In regione superiore Atlantis Majoris. In convalle Aït Mesan et in jugo Tagherot a 2300 m. usque 3200 m.! In monte Djebel Tezah eire. 3000 m.!

Ar. Geog. Species per Europam omnem et in montibus regionis Mediterraneæ diffusa. Var. *indigesta* in summis montibus Hispaniæ, et in Pyrenæis obvia est.

FESTUCA ARUNDINACEA (*Schreb. Spec. Fl. Lips.* 57; *Koch, Syn. Fl. Germ.* ed. 2, 943), = *F. elatior* (*L. Sp. Pl., non Fl. Succ.; Kunth, Enum.* i. 404, *pro parte*).

Mar. merid.—Satis frequens in humidis reg. inferioris. Mesfioua! Ourika! Sheshaoua! In convalle Aït Mesan circ. 1300 m.! Visa in convalle Amsmiz!

— —, var. β , *Coss. in Expl. Scient. Alg.* 170, = *F. interrupta* (*Desf. Fl. Atl.* i. 89).

Mar. occid.—Casa Blanca!

Mar. merid.—Amsmiz! Mzouda!

Ar. Geog. Europa media. Sibiria. Regio Mediterranea occidentalis (exclusa Macaronesia). Ex Oriente non vidi. Var. in calidioribus hinc inde occurrit.

F. CÆRULESCENS (*Desf. Fl. Atl.* i. 87; *Kunth, Enum.* i. 406), = *Kœleria tunicata* (*Presl, Cyp. et Gram. Sic.* 34).

Mar. sept.—In monte Djebel Kebir prope Tanger! In monte Beni Hosmar prope Tetuan! Prope Ceuta!

Mar. merid.—Legi in monte supra Seksaoua circ. 1400 m.!

Ar. Geog. Africa boreali-occidentalis. Hispania. Sicilia. Legi in collibus prope Algeciras!

F. DURANDII (*Clauson in Billot Annot.* 163).

Mar. sept.—In monte Djebel Kebir prope Tanger (sec. Cosson, *Expl. Scient. Alg. Add.* 299), ad varietatem *F. spadiceæ*, *L.*, referenda?

Ar. Geog. Species dubia in Africa boreali-occidentali indicata.

BROMUS (*L. Gen.* 89).

B. MOLLIS (*L., Kunth, Enum.* 413).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)!

Mar. occid.—Casa Blanca!

Mar. merid.—Misra ben Kara!

— —, var. *LEIOSTACHYS*, *M. K.*? Planta glabrior, panicula depauperata, spicis paucifloris.

Mar. merid.—In convalle Aït Mesan 1800–2100 m. (*J. D. H.*)!

— —, var. *RAMOSUS*, *nob.* Paniculae ramis verticillatis insigniter inæqualibus.

Mar. merid.—Supra pagum Amsmiz, ad circ. 1200 m. s. m.!

Ar. Geog. Species in Europa fere omni, et in regione Mediterranea (inclusa Macaronesia) late diffusa. In America meridionali, forsitan introducta?

B. ALOPECURUS (*Poir. Voy.* i. 100, *non Vahl*), = *B. alopecuroides* (*Poir. Dict. Suppl.* i. 703; *Kunth, Enum.* i. 415), = *B. contortus* (*Desf. Fl. Atl.* i. 95, tab. 25), = *B. seoparius* (*Guss., non L.*).

Mar. occid.—Casa Blanca!

Ar. Geog. Africa boreali-occidentalis. Sicilia. Fertur in Hispania (*Kunth, l. c.*), sed fide Willk. et Lgc. *Fl. Hisp.*, species hucusque non detecta.

BROMUS MACROSTACHYS (*Desf. Fl. Atl.* i. 96, tab. 10; *Kunth, Enum.* i. 415), = *B. lanceolatus* (*Roth, Catal. Bot.* i. 18; *Kunth, l. c.*), = *B. divaricatus* (*Rohde in Loisel. Not.*; *Kunth, l. c.*; *Salzm. Exsicc.*).
 Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb spec.*)! Tetuan (1851)! (*Webb spec.*)!

Mar. ooeid.—Casa Blanca!

Mar. merid.—Ourika (*R. et Fr. spec. in Herb. Kew.*)!

— —, var. *LANUGINOSUS* (*Parlat. Fl. It.* i. 398, sub *Serrafalco*), = *B. lanuginosus* (*Poir. Dict. Suppl.* i. 703).

Mar. merid.—Occurrit frequentissime in regione calidiore, necnon in reg. inferiore Atlantis Majoris. Prov. Shedma! Prope urbem Marocco! Reraya! Seksaoua! Amsmiz! Mzouda! In convalle Aït Mesan usque 1400 m.!

Ar. Geog. Species per regionem fere omnem Mediterraneam (inclusa Macaronesia) dispersa. Varietas in Sicilia et in Africa boreali obvia est.

B. TECTORUM (*L., Kunth, Enum.* i. 419).

Mar. sept.—Prope Tanger (*J. D. H.*)!

Mar. merid.—A radicibus Atlantis Majoris usque in regionem subalpinam. Reraya! In convalle Aït Mesan a 1400 m. ad 2500 m. s. m.!

Ar. Geog. Europa fere omnis (exceptis Lapponia et Britannia). Asia media. Regio omnis Mediterranea (inclusa Macaronesia).

B. MATRITENSIS (*L., Kunth, Enum.* i. 419; *Schousb. Gew. Marok.* 31).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! In elivo montis Beni Hosmar prope Tetuan!

Mar. occid.—Casa Blanca! Mogador! (*Lowe Cat.*).

Mar. merid.—Frequens in regione inferiore. In ruderatis urbis Marocco! Reraya! Seksaoua! Amsmiz! Mesfioua! Vidimus etiam in convallibus Atlantis Majoris usque 1400 m. s. m.!

Ar. Geog. Regio omnis Mediterranea. In calidioribus Europæ mediæ.

B. MAXIMUS (*Desf. Fl. Atl.* i. 95, tab. 26; *Kunth, Enum.* i. 419), = *B. rigidus* (*Roth in Roem. et Ust. Mag. Bot. fasc. x.* 21; *Kunth, l. c.*).

Mar. sept.—Tanger (*Webb spec.*)!

Mar. occid.—Casa Blanca! Mogador!

Mar. merid.—Amsmiz!

Ar. Geog. Regio Mediterranea (inclusa Macaronesia). Occurrit in Gallia occidentali.

B. RUBENS (*L., Kunth, Enum.* i. 420).

Mar. sept.—Tetuan (1851)!

Mar. ooeid.—Casa Blanca! In monte Djebel Hadid haud procul Mogador (*R. et Fr. spec. in Herb. Kew.*)!

Mar. merid.—In regione inferiore satis frequens. Aïn Oumast ! Mesfoua ! Ourika ! Seksaoua ! Milhaïn !

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

LOLIUM (*L. Gen.* 95).

L. PERENNE (*L., Kunth, Enum.* i. 436 ; *Schousb. Gew. Marok.* 35).

Mar. sept.—Tanger (*Schsb. l. c.*).

Mar. occid.—Mogador (*Lowe Cat.*).

Mar. merid.—In convalle Aït Mesan circ. 1300 m. legi formam pumilam spicis depauperatis !

— —, var. *ITALICUM*, = *L. italicum* (*Al. Braun in Flora*), = *L. Boucheanum* (*Kunth, l. c.*) ; verisimiliter huc etiam var. *aristatum*, *Schousb. l. c.*

Mar. sept.—Tanger (*Webb Herb.*) ! (*Schsb. l. c.*) ?

Mar. occid.—Casa Blanca !

Mar. merid.—Ourika !

— —, var. *RIGIDUM*, = *L. rigidum* (*Gaud.*), = *L. strictum* (*Presl*).

Mar. occid.—Mogador !

Mar. merid.—Sheshaoua ! Misra ben Kara !

Ar. Geog. Species per regiones temperatas hemisphæræ borealis late diffusa. Varietates nostræ in calidioribus occurrunt.

TRITICUM (*L. Gen.* 99).

De *Tritici* speciebus in territorio Maroccano cultis parum discere potui. *Triticum æstivum* (*L., Schousb. Gew. Marok.* 38) sæpissime colitur. Ad radices Atlantis (juxta Ourika et alibi) vidimus in agris speciem diversam verisimiliter ad *T. turgidum*, *L.*, referendam.

T. VILLOSUM (*L., sub Secale ; Pal. Beauv. Agrost.* 103 ; *Kunth, Enum.* i. 440).

Mar. merid.—In glarcosis Oued en Fisk prope Gurguri !

Ar. Geog. Regio Mediterranea fere omnis (exclusa Macaronesia).

T. HORDEACEUM (*Coss. et Dur. MSS.*).

Mar. merid.—Ad radices Atlantis Majoris. Sektana ! Seksaoua ! Visum etiam in convalle Aït Mesan !

Species Algeriensis mihi non bene cognita, monente cl. Coss. in Indice Pl. Marocc., jampridem a cl. Balansa in monte Djebel Sidi Fars prope Keira lecta.

T. JUNCUM (*L., Kunth, Enum.* i. 444 ; *Schousb. Gew. Marok.* 39).

Mar. sept.—Tanger ! (*Salzm. Exsicc.*) ! (*Schsb. l. c.*).

Mar. occid.—Mazagan ! Saffi !

Ar. Geog. Species per hemisphæram borealem temperatam, præsertim in arenosis maritimis, late diffusa. Decet in Macaronesia.

TRITICUM ELONGATUM (*Host, Gram. Austr. ii. 18*), = *T. rigidum* (*Schrad. Sem. Hort. Gott. 1803, 23; Kunth, Enum. i. 445*).

Mar. sept.—Tanger! (*Webb spec.*)! (*Salzm. Exsicc.*)!

Ar. Geog. Europa media. Occurrit etiam hinc inde in regione Mediterranea. Deest in Macaronesia.

BRACHYPODIUM (*Pal. Beauv. Agrost. 100, tab. 19*).

B. PINNATUM (*L. Sp. Pl. 115, sub Bromo; P. B. l. c. 101, et auct. recent.*), = *Triticum pinnatum* (*Mœnch, En. Pl. Hass. 102; Kunth, Enum. i. 445*).

Mar. sept.—Tanger (*Webb Herb.*)! (*Schsb. l. c.*).

—— —, var. AUSTRALE (*Gren. et Godr. Fl. Fr. iii. 610*), = *Triticum phœnicoides* (*DC. Fl. Gall. vi. 284, et auct. plurim.*), = *Bromus ramosus* (*Schousb. Gew. Marok. 31?, non L., nec Vahl*).

Mar. occid.—Casa Blanca! Mazagan (*J. D. H.*)!

Mar. merid.—Reraya (*J. D. H.*)!

Ar. Geog. Europa media. Regio Mediterranea (exclusa Macaronesia).

Var. *australe* in calidioribus magis frequens.

B. RAMOSUM (*L. Mant. i. 34, sub Bromo; R. et Sch. Syst. ii. 737*), = *Bromus pinnatus*, var. β (*L. Sp. Pl. 115*), = *Festuca cæspitosa* (*Desf. Fl. Atl. i. 91, tab. 24*), = *Triticum cæspitosum* (*DC. Cat. Monsp. 153; Kunth, Enum. i. 446*), = *Festuca phœnicoides* (*All. Fl. Ped. ii. 253, vix L.*), = *Brachypodium phœnicoides* (*Coss. in Expl. Scient. Alg. 191*), = *B. Plukenetii* (*Link, Hort. R. Berol. i. 40*).

Mar. merid.—Legi in saxosis supra Seksaoua ad radices Atlantis Majoris! nec alibi vidimus.

Ar. Geog. Regio fere omnis Mediterranea (exclusa Macaronesia).

Bromus ramosus (*Schousb. Gew. Marok. 31*), “in arenosis maritinis Tingitanis abunde creescens,” nequit huc referri; species genuina enim nemini in agro Tingitano post Schousbœum obvia fuit. Quid sit *Bromus longifolius* (*Schousb. l. c.*), *B. ramoso* suo affinis, et nisi habitu et foliis longioribus non distinguendus, erucere non potui. Verisimiliter inter varietates *Brachypodii pinnati* collocandus est.

B. DISTACHYUM (*L., sub Bromo; R. et Sch. Syst. ii. 741*), = *Triticum ciliatum* (*DC. Fl. Fr. iii. 85; Salzm. Exsicc.; Kunth, Enum. i. 447*).

In omni territorio nobis cognito valde frequens.

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! Tetuan (1851)!

Mar. occid.—Casa Blanca! Mogador!

Mar. merid.—Passim in regione calidiore et in regione inferiore Atlantis Majoris. Prov. Shedna! Mesfioua! Reraya! In convalle Aït Mesan usque 1500 m.! In couvalle Amsmiz! Seksaoua! Sektana (*R. et Fr. spec. in Herb. Kew.*)!

Variat insigniter statura pygmæa vel ultrapedali, culmis lævibus vel scabriuseulis.

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

SECALE (*Pal. Beauv. Agrost.* 105, tab. 20).

S. MONTANUM (*Guss. Ind. Sem. H. Boccadif.* 1825; *Parlat. Fl. It.* i. 517; *Kunth, Enum.* i. 449).

Mar. merid.—In convalle Aït Mesan circ. 1600 m.!

Ar. Geog. Africa boreali-occidentalis. Sicilia. Hispania.

ELYMUS (*L. Gen.* 96).

Genus forsán cum *Hordeo* conjungendum.

E. CAPUT MEDUSÆ (*L., Kunth, Enum.* i. 452).

Specimen ex Imperio Maroccano conservatur in herb. Webb.

Mar. merid.—Prope Amsmiz! Prope Mzouda (*J. D. H.*)!

— —, var. CRINITUS, = E. erinitus (*Schreb. Gram.* 15, tab. 24; *Kunth, Enum.* i. 452). Differt glumis dimidio brevioribus, suberectis, nec patentissimis.

Mar. merid.—Specimen unicum legi in convalle Aït Mesan circ. 1400 m.!

Spec. nostrum gaudet glumellis pubescentibus ut fiunt in spec. prope Batna lectis (*Balansa, Pl. d'Algérie*, 1853, No. 733).

Ar. Geog. Species in Africa boreali-occidentali et in peninsula Iberica nascitur. Var. *crinitus* per Africam borealem, Orientem et Europam austro-orientalem hinc inde occurrit.

HORDEUM (*L. Gen.* 98).

H. BULBOSUM (*L., Kunth, Enum.* i. 455), = H. strictum (*Desf. Fl. Atl.* i. 113; *Schousb. Gew. Marok.* 36; *Salzm. Exsicc.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)! (*Schsb. l. c.*).

Ar. Geog. Regio omnis Mediterranea (exclusa Macaronesia) præsertim occidentalis.

In Hispania Bætica non indicatur; legi tamen copiose crescens prope S. Roque!

H. MURINUM (*L., Kunth, Enum.* i. 456).

Valde frequens in territorio omni nobis noto.

Mar. sept.—Tanger!

Mar. oecid.—Casa Blanea! Mogador!

Mar. merid.—Shedma! In ruderatis urbis Marocco! Mzouda! In convalle Aït Mesan usque 1400 m.! Vidimus in convalle Amsmiz!

Ar. Geog. Europa omnis temperata. Regio omnis Mediterranea. In America extratropica tam boreali quam meridionali, sed forsán ex Europa allata.

H. MARITIMUM (*With. Bot. Arrang.* 172; *Kunth, Enum.* i. 456).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)!

Mar. occid.—Mazagan !

Ar. Geog. Species amphigea in maritimis hemisphæræ borealis et Americæ australis late diffusa.

ÆGILOPS (*L. Gen.* 1150).

Æ. OVATA (*L., Kunth, Enum.* i. 458).

Frequentissime occurrit in territorio omni nobis cognito.

Mar. sept.—Tanger ! (*Salzm. Exsicc.*) ! (*Webb Herb.*) !

Mar. occid.—Casa Blanca !

Mar. merid.—Prov. Shedma ! Reraya ! In convalle Aït Mesan usque 1400 m. s. m. ! Seksaoua ! In convalle Amsmiz visa ! Ludit glumis glabrescentibus, pubescentibus, vel villosis.

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

Æ. TRIUNCIALIS (*L., Kunth, Enum.* i. 458).

Mar. sept.—Tanger ! (*Salzm. Exsicc.*) !

Mar. merid.—Seksaoua ! Mesfiousa !

Spec. ex Mesfiousa ludit glumis glabris costis scabriusculis.

Ar. Geog. Regio Mediterranea imprimis occidentalis (exclusa Macaronesia).

Æ. VENTRICOSA (*Tausch in Flora*, 1837 ; *Boiss. Voy. Esp.* 632), = *Æ. squarrosa* (*Cavan. Icon.* tab. 90. fig. 2 ; *Salzm. Exsicc.*, non *L.*).

Mar. sept.—Tanger (*Salzm. Exsicc.*) ! (*Webb Herb.*) !

Mar. merid.—In convalle Aït Mesan circ. 1200 m. s. m. !

Ar. Geog. Italia. Hispania. Africa boreali-occidentalis.

LEPTURUS (*R. Br. Prod.* 207).

L. FILIFORMIS (*Trin. Fund. Agrost.* 123 ; *Kunth, Enum.* i. 462), = *Rottbœllia crecta* (*Savi, Fl. Pis.* iv. 230), = *R. subulata* (*Salzm. Exsicc.*, non *Savi*).

Mar. sept.—Tanger (*Salzm. Exsicc.*) ! (*Webb spec.*) ! (*Spec. ex Herb. Schousboe*) !

Ar. Geog. In maritimis Europæ mediæ et regionis omnis Mediterraneæ (exclusa Macaronesia) dispersa.

MONERMA (*Pal. Beauv. Agrost.* 117).

M. CYLINDRICA (*Willd. Sp. Pl.* i. 464, sub *Rottbœllia* ; *Coss. et Dur. in Expl. Scient. Alg.* 209), = *M. subulata* (*Pal. Beauv. l. c.*), = *Lepturus cylindricus* (*Trin. Fund. Agrost.* 123), = *Rottbœllia subulata* (*Savi, Due Cent.* 35 ; *DC. Fl. Fr.* vi. 281), = *Lepturus subulatus* (*Kunth, Enum.* i. 462).

Mar. sept.—In monte Djebel Kebir prope Tanger (*spec. ex Herb. Schousboe benev. comm. Cosson*) !

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia). In Africa australi etiam obvia.

IMPERATA (*Cyr. Pl. rar. Neap. fasc. ii. 26*).

I. CYLINDRICA (*L.*, sub *Laguro*; *Pal. Beauv. Agrost. 7*), = I. arundinacea (*Cyr. l. c.*; *Kunth, Enum. i. 477*).

Mar. merid.—Ad ripas fluminis prope Misra ben Kara!

Ar. Geog. Species amphigea in temperatis calidioribus et in reg. tropicalibus orbis totius diffusa.

ANTHISTIRIA (*L. Gen. 1359*).

A. PALEACEA (*Vahl, Symb. ii. 24*, sub *Stipa*), = A. glauca (*Desf. Fl. Atl. ii. 380*, tab. 254; *Kunth, Enum. i. 482*), = *Stipa paleacea* (*Schousb. Gew. Marok. 33*).

“In collibus et locis aridis raro” (*Schsb. l. c.*), loco natali non indicato.

Ar. Geog. Species in Africa boreali continentali, necnon in Africa tropica occidentali, obvia.

SORGHUM (*Pers. Syn. i. 101*).

S. HALEPENSE (*L.*, sub *Holco*; *Pers. l. c.*), = *Andropogon halepensis* (*Sibth. et Sm. Prod. Fl. Gr. i. 47*; *Kunth, Enum. i. 502*).

Mar. merid.—Mzouda! Prope Frouga (*G. Maw*)!

Ar. Geog. Regio Mediterranea (exclusa Macaronesia). In America et Australia introducta.

ANDROPOGON (*L. Gen. 1145*).

A. HIRTUS (*L.*, *Kunth, Enum. i. 492*).

Mar. sept.—Prope Cap. Spartel!

Mar. occid.—Mogador (*Lowe Cat.*). In monte Djebel Hadid haud procul Mogador!

Mar. merid.—Ourika (*G. Maw*)!

Ar. Geog. Regio omnis Mediterranea (inclusa Macaronesia).

— —, var. LONGEARISTATUS, *Willk.*, = A. pubescens (*Vis. in Flora, 1829, 1*).

Mar. merid.—In rupestribus prope urbem Marocco (*G. Maw*)! In convalle Aït Mesan usque 1800 m. s. m.! Milhaïn (*J. D. H.*)!

Ar. Geog. Occurrit cum specie hinc inde in regione Mediterranea.

A. LANIGER (*Desf. Fl. Atl. ii. 379*; *Kunth, Enum. i. 493*).

Mar. merid.—In rupibus apricis prope urbem Marocco (*G. Maw*)! Prope Gurguri in via qua itur versus Amsmiz!

Ar. Geog. Species imprimis regionis desertorum, ab Indiæ orientalis ditione Pantjab usque prov. meridionales Imp. Maroccani.

A. ANNULATUS (*Forsk. Fl. Æg.-arab. 173*; *Kunth, Enum. i. 498*).

Mar. merid.—Prope Tasseremout detexit oculatissimus G. Maw et specimen mihi benevole dedit!

Ar. Geog. Per regiones subtropicales Asiæ occidentalis et Africæ borealis diffusus. Occurrit etiam in Africa australi.

POLLINIA (*Spreng. Pug.* ii. 12).

P. DISTACHYA (*Spreng. l. c.*; *Bert. Fl. Ital.* i. 463), = *Andropogon distachyon* (*L., Kunth, Enum.* i. 491; *Salzm. Exsicc.*).

Mar. sept.—Tanger! (*Salzm. Exsicc.*)! (*Webb Herb.*)! Tetuan!

Ar. Geog. Regio Mediterranea fere omnis (inclusa Macaronesia).

ENUMERATIO FILICUM, EQUISETACEARUM et LYCOPODIACEARUM in territorio Maroccano nobis cognitarum.

DAVALLIA CANARIENSIS, *Sm., H. et Bak. Syn.* 47.

Mar. sept.—In monte Djebel Kebir! Prope Cap. Spartel!

CISTOPTERIS FRAGILIS, *Bernh., H. et Bak. Syn.* 103.

Mar. merid.—In jugo Tagherot, 2700 m.! In monte Djebel Tezali circ. 2800 m.!

ADIANTUM CAPILLUS VENERIS, *L., H. et Bak. Syn.* 123.

Mar. sept.—In monte Beni Hosmar prope Tetuan!

Mar. occid.—Prope Mogador!

Mar. merid.—In hortis urbis Marocco!

CHEILANTHES FRAGRANS, *Webb, H. et Bak. Syn.* 134, = *C. odora*, *Sw.*

Mar. merid.—Mskala! Reraya! In couvalle Aït Mesan! Seksaoua!

PTERIS ARGUTA, *Ait., H. et Bak. Syn.* 160.

Mar. sept.—Prope Tanger (*Blackmore spec.*)!

P. AQUILINA, *L., H. et Bak. Syn.* 162.

Mar. sept.—Fronde emarcidæ in monte Beni Hosmar prope Tetuan!

Mar. merid.—Vidimus in couvalle Aït Mesan ultra 2000 m. s. m.!

ASPLENIUM HEMIONITIS, *L., H. et Bak. Syn.* 194, = *A. palmatum*, *Lam.*

Mar. sept.—In monte Djebel Kebir prope Tanger! (*Salzm. Exsicc.*)!

De *A. Hemionitis*, *L.*, genuino conf. *Hook.* in *Bot. Mag.* lxxxii. tab. 4911. Planta Linnæana, a plurimis auctoribus cum *Scolopendrio Hemionitis* (*Hemionitis vera*, *Clus.*) confusa, citatur in Hispania in Cavan. prælect., sed nemini recentiorum obvia fuit. Habitat in rupibus graniticis prope Cintra, ubi cum olim legi; nec alibi (quoad sciam) in Europa hodie inveniunda.

A. VIRIDE, *Huds., H. et Bak. Syn.* 195.

Mar. merid.—Specimina duo nana deteximus in regione superiore Atlantis Majoris in jugo Tagherot circ. 2800 m.!

ASPLENium TRICHOMANES, *L., H. et Bak. Syn.* 196.

Mar. sept.—In monte Beni Hosmar prope Tetuan !

Mar. ooeid.—In monte Djebel Hadid haud procul Mogador !

Mar. merid.—In convalle Aït Mesan a 1500 m. usque 2200 m. s. m. !

A. PETRARCHÆ, *DC., H. et Bak. Syn.* 197.

Mar. merid.—Vidi specimen unicum nanum in fissuris rupium supra Milhaïn ; frondes avulsæ elheu perierunt, una excepta hodie in Herb. Kew. conservata !

A. MARINUM, *L., H. et Bak. Syn.* 207.

Mar. sept.—Vidimus in sic dicto antro Hereulis a Cap. Spartel meridiem versus, loco inaccessibili ereseens !

A. ADIANTUM NIGRUM, *L., H. et Bak. Syn.* 214.

Mar. merid.—In regione superiore Atlantis Majoris. In jugo Tagherot circ. 2800 m. s. m. !

—— —, var. VIRGILII, = A. Virgilii, *Bory et Chaub.*

Mar. sept.—In monte Djebel Kebir prope Tanger satis frequens !

A. LANCEOLATUM, *Huds., H. et Bak. Syn.* 217.

Mar. sept.—Prope Cap. Spartel ! spec. Anglicis conforme.

—— —, var. OBOVATUM, = A. obovatum, *Viv.*

Mar. sept.—Prope Tanger ! (*Webb Herb.*) !

A. FILIX-FÆMINA, *Bernh., H. et Bak. Syn.* 227.

Mar. sept.—In collibus inter Tanger et Tetuan !

A. CETERACH, *L., H. et Bak. Syn.* 245, = Ceterach officinarum (*Willd. et auctt. plurim.*).

Mar. sept.—Vidimus prope Tanger ! Legi in rupibus juxta Tetuan !

Mar. ooeid.—In monte Djebel Hadid haud procul Mogador !

Mar. merid.—Satis frequens ad radices Atlantis Majoris et usque in regionem superiorem. In convalle Aït Mesan et in ascensu jugi Tagherot 1400–2700 m. ! Amsmiz ! Seksaoua ! Milhaïn ! In monte Djebel Tezah (*J. D. H.*) !

Connubium generis *Ceterach* cum *Asplenio* naturæ abhorrens ; paece tantorum virorum vix posteris gratum erit.

SCOLOPENDRIUM HEMIONITIS, *Clem. in Anal. Cienc. Nat.* v. (1802) ; *H. et Bak. Syn.*, = S. sagittatum, *DC.* Var. frondibus acute hastatis lobis triangulari-aeuminatis.

Mar. sept.—In fissuris rupium montis Beni Hosmar !

POLYPODIUM VULGARE, *L., H. et Bak. Syn.* 334.

Mar. sept.—In monte Beni Hosmar prope Tetuan !

NOTHOCHLÆNA VELLEA (*Ait. Hort. Kew.* ed. 1, iii. 457, sub *Acrosticho* ;

Desv. in Journ. Bot. iii. 92), = *N. lanuginosa* (*Desf. Fl. Atl.* ii., sub *Acrosticho*; *Desv. in Dict. Suppl.* iv. 110; *H. et Bak. Syn.* 370).

Mar. merid.—In rupestribus aridis haud infrequens. In rupibus prope urbem Marocco (*G. Maw*)! In convalle Aït Mesan 1500–1700 m.! prope Milhaïn! Aïn Tarsil!

GYMNOGRAMME LEPTOPHYLLA, *Desv., H. et Bak. Syn.* 383.

Mar. sept.—Tanger (*J. D. H.*)!

OSMUNDA REGALIS, *L., H. et Bak. Syn.* 427.

Mar. sept.—Legi in rupibus maritimis (mirum!) prope Tanger! In umbrosis versus Cap. Spartel!

OPHIOGLOSSUM LUSITANICUM, *L., H. et Bak. Syn.* 445.

Mar. sept.—Tanger (*Salzm. Exsicc.*)! (*Webb Herb.*)!

Filices Maroccanas paucas, et fere omnes admodum raras, secundum ordinem operis citati—Hooker et Baker, 'Synopsis Filicum,' edit. 2—enumeravi.

EQUISETUM TELMATEJA, *Ehrh.*

Mar. sept.—Tanger (*J. D. H.*)!

E. RAMOSISSIMUM, Desf.

Mar. sept.—Vidimus prope Tanger!

Mar. merid.—In convalle Aït Mesan!

Cum spec. typo conformibus legimus formam seu varietatem eulmis majoribus paucistriatis, vaginarum dentibus magis numerosis.

ISOETES HYSTRIX (*Dur. in Expl. Scient. Alg.* tab. 36. fig. 1).

Mar. sept.—Legimus in paseuis subhumidis Tingitanis ad radices montis Djebel Kebir meridiem versus!

SELAGINELLA RUPESTRIS, *Spreng.*

Mar. merid.—In ditione Reraya circ. 1100 m. s. m.! In convalle Aït Mesan usque 1800 m.!

S. DENTICULATA, Spreng.

Mar. sept.—Tanger (*Webb Herb.*)! Inter Tanger et Tetuan!

MUSCI MAROCCANI.

A cl. W. Mitten determinati et enumerati.

DICRANELLA VARIA, *Hedw.* Beni Hosmar prope Tetuan!

GRIMMIA APOCARPA, *L.* Beni Hosmar prope Tetuan!

G. PULVINATA, L. In jugo Tagherot, alt. 3000 m.!

G. ORBICULARIS, B. & S. Aït Mesan circ. 2500 m.!

G. COMMUTATA, Hüben. Aït Mesan circ. 2500 m.!

TRICHOSTOMUM TOPIACEUM, *Brid.* Tanger et Tetuan!

T. CRISPULUM, Bruch. Tanger et Tetuan! In monte Beni Hosmar!

- TRICHOSTOMUM MUTABILE, *Bruch.* Tanger et Tetuan !
 T. BARBULA, *Schw.* Tanger et Tetuan !
 TORTULA MURALIS, *L.* Tanger et Tetuan !
 T. INERMIS, *Bruch.* Aït Mesan 2000–3000 m. !
 T. RURALIS, *L.* Aït Mesan circ. 2000 m. !
 ENCALYPTA VULGARIS, *Hedw.* In jugo Tagherot, 3000 m. !
 ORTHOTRICHUM CUPULATUM, *Hoffm.* Aït Mesan 2000–3000 m. !
 ENTOSTHODON OBTUSUS (*Dicks.*) = *E. erieetorum*, *De Notaris.* Tanger et Tetuan !
 FUNARIA HYGROMETRICA, *L.* Tanger et Tetuan ! Aït Mesan 1400–1500 m. !
 BARTRAMIA (PHILONOTIS) RIGIDA, *Brid.* Tanger et Tetuan !
 B. FONTANA, *L.* Djebel Tezah, alt. 1700–3000 m. !
 BRYUM ALPINUM, *L.* Aït Mesan 2000–2600 m. !
 B. PALLESCENS, *Schw.* Aït Mesan 2000–2600 m. !
 B. CAPILLARE, *L.* Tanger et Tetuan !
 B. TURBINATUM, *Hedw.* In jugo Tagherot 2700–3000 m. !
 MNIMUM UNDULATUM, *Hedw.* Beni Hosmar prope Tetuan !
 LEUCODON SCIUROIDES (*L.*). Beni Hosmar prope Tetuan !
 PTEROGONIUM GRACILE (*L.*). Beni Hosmar prope Tetuan !
 HOMALOTHECIUM SERICEUM (*L.*). Beni Hosmar prope Tetuan !
 LEPTODON SMITHII (*Dicks.*). Beni Hosmar prope Tetuan !
 THAMNIUM ALOPECURUM (*L.*). Beni Hosmar prope Tetuan !
 HYPNUM (RHYNCHOSTEGIUM) TENELLUM, *Dicks.* Tanger et Tetuan !
 H. (EURHYNCHIUM) CIRCINATUM, *Brid.* Beni Hosmar prope Tetuan !
 H. (SCLERPODIUM) ILLECEBRUM, *Schw.* Tanger et Tetuan !
 H. (BRACHYTHECIUM) VELUTINUM, *Dill.* Tagherot 2500–3200 m. !
 H. (BRACHYTHECIUM) RUTABULUM, *L.* Aït Mesan 2000–3000 m. !
 In monte Djebel Tezah !
 AMBLYSTEGIUM COMMUTATUM (*Hedw.*). Beni Hosmar prope Tetuan !
 Aït Mesan 2000–2500 m. !
 FISSIDENS INCURVUS, *W. & M.* Beni Hosmar prope Tetuan !
 F. TAXIFOLIUS, *L.* Tanger et Tetuan !

HEPATICÆ.

- CEPHALOZIA TURNERI (*Hook.*). Beni Hosmar prope Tetuan !
 C. BICUSPIDATA (*L.*). Beni Hosmar prope Tetuan !
 CALYPOGEIA TRICHOMANIS (*L.*). Beni Hosmar prope Tetuan !
 FOSSOMBRONIA ANGULOSA, *Raddi.* Tanger et Tetuan !

ASTERELLA HEMISPHERICA, *Beauv.* Aït Mesan 1400–2000 m.!

ANTHOCEROS PUNCTATUS. Beni Hosmar prope Tetuan!

POSTSCRIPTUM.

Clarus A. Leared, M.D., qui plagas meridionales Imperii Maroeani jampridem peragravit et illustravit, nuper mihi frustula nonnulla plantarum in itinere a Tingi ad urbes Mequinez et Fez anno 1877 suscepto benevole communicavit. Harum adjeci enumerationem brevem, et ob specimina nimis incompleta, pluribus locis incertam.

CLEMATIS FLAMMULA, *L.* Prope Aleazar.

*NIGELLA HISPANICA, *Coss.* Inter Tanger et Mequinez.

*DELPHINIUM, an sp. nov.? Flores in spica longa conferti; pedunculi tennes demum elongati, braeteis duabus linearibus longiusculis instructi. A *D. peregrino*, *L.*, certe diversum. Inter Tanger et Mequinez.

FUMARIA CAPREOLATA, *L.*, var. Inter Tanger et Mequinez.

CAPPARIS SPINOSA, *L.* Inter Fez et Mequinez.

*RESEDA COLLINA, *J. Gay?* Inter Tanger et Mequinez.

CISTUS POPULIFOLIUS, *L.* Inter Tanger et Mequinez.

SILENE OBTUSIFOLIA, *W.* Inter Tanger et Mequinez.

S. LÆTA (*Ait.*). Inter Tanger et Mequinez.

HYPERICUM PERFOLIATUM, *L.* Inter Tanger et Mequinez.

LINUM CORYMBIFERUM, *Desf.*, var.? Inter Tanger et Mequinez.

ZIZYPHUS LOTUS (*L.*). Inter Tanger et Mequinez.

CYTISUS LINIFOLIUS (*L.*). Inter Tanger et Mequinez.

ONONIS VISCOSA, *L.* Inter Tanger et Mequinez.

TRIFOLIUM PROCUMBENS, *L.*, var. CAMPESTRE. Inter Tanger et Mequinez.

LOTUS HISPIDUS, *Desf.* Inter Tanger et Mequinez.

PSORALEA BITUMINOSA, *L.* Inter Tanger et Mequinez.

LYTHRUM FLEXUOSUM, *Lag.* Inter Tanger et Mequinez.

COTYLEDON HISPANICA, *L.*, var. SALZMANNI. Inter Tanger et Mequinez.

ECBALIUM ELATERIUM (*L.*). Prope Mequinez.

CENTRANTHUS CALCITRAPA (*L.*). Inter Tanger et Mequinez.

ODONTOSPERMUM AQUATICUM (*L.*). Inter Tanger et Mequinez.

ANACYCLUS RADIATUS, *Lois.* Inter Tanger et Mequinez.

ANTHEMIS MIXTA, *L.* Inter Tanger et Mequinez.

*ECHINOPS RITRO, *L.* Prope Muley Edris Serone.

GALACTITES TOMENTOSA, *L.* Inter Tanger et Mequinez.

CENTAUREA PULLATA, *L.* Inter Tanger et Mequinez.

SCOLYMUS HISPANICUS, *L.* Inter Tanger et Mequinez.

TOLPIS BARBATA, *Gertn.* Inter Tanger et Mequinez.

CAMPANULA LÆFLINGII, *Brot.*, var. DECUMBENS? Inter Tanger et Mequinez.

TRACHELIUM CÆRULEUM, *L.* Inter Tanger et Mequinez.

T. ANGUSTIFOLIUM, *Schousb.* Inter Tanger et Mequinez.

STATICE THOUINI, *Viv.* Inter Tanger et Mequinez.

ANAGALLIS ARVENSIS, *L.*, var. LATIFOLIA. Inter Tanger et Mequinez.

A. LINIFOLIA, *L.* Inter Tanger et Mequinez.

*CONVOLVULUS, sp. nov. Egrege *C. undulati*, *Cav.*, sed certe diversus. Inter Tanger et Mequinez.

ECHIUM ANGUSTIFOLIUM, *Lam.* Inter Tanger et Mequinez.

ANCHUSA GRANATENSIS, *Boiss.*? Inter Tanger et Mequinez.

VERBASCUM SINUATUM, *L.* Inter Tanger et Mequinez.

ANTIRRHINUM ORONTIUM, *L.*, var. GRANDIFLORUM. Inter Tanger et Mequinez.

ACANTHUS MOLLIS, *L.* Muley Edris Serone.

MENTHA PULEGIUM, *L.* Inter Tanger et Mequinez.

ORIGANUM —? Inter Tanger et Mequinez.

*MELISSA OFFICINALIS, *L.*, var. Inter Tanger et Mequinez.

STACHYS HIRTA, *L.* Inter Tanger et Mequinez.

CHENOPODIUM AMBROSIoidES, *L.* Prope Mequinez.

DAPHNE GNIDIUM, *L.* Inter Tanger et Mequinez.

OPHRYS APIFERA, *Huds.* Inter Tanger et Mequinez.

XYPHION SISYRHYNCHIUM, *L.* Inter Tanger et Mequinez.

ASPHODELUS MICROCARPUS, *Salzm.* Inter Tanger et Mequinez.

ORNITHOGALUM ARABICUM, *L.* Inter Tanger et Tetuan.

ASPLENIUM HEMIONITIS, *L.* Inter Tanger et Mequinez.

GYMNOGRAMME LEPTOPHYLLA, *Desv.* Inter Tanger et Mequinez.

Inter species 53 enumeratas, sex, quorum nomina asterisco notata, in Imperio Maroccoano nondum indicatæ sunt.

INDEX ICONUM.

PLATE IX.

Fig. 1. *Nasturtium atlanticum*, Ball, magnit. nat. 2. Flos magnit. auct.
3. Andrœcium cum pistillo, magnit. auct. 4. Siliqua, magnit. auct.

PLATE X.

Fig. 1. *Arabis decumbens*, Ball, magnit. nat. 2. Flos, magnit. auct. 3. Pistillum, magnit. auct. 4. Racemus fructifer, magnit. nat. 5. Siliqua, magnit. auct.

PLATE XI.

Fig. 1. *Arabis conringioides*, Ball, magnit. nat. 2. Flos, magnit. auct. 3. Andrœcium cum pistillo, magnit. auct. 4. Racemus fructifer, magnit. nat. 5. Siliqua, magnit. auct.

PLATE XII.

Fig. 1. *Brassica elata*, Ball. Pars caulis inferior cum foliis 2 inferioribus, et pars superior florifera. Exemplar pumilum, magn. nat. 2. Flos, magnit. dupl. auct. 3. Andrœcium cum pistillo, magnit. auct. 4. Silicula immatura, magnit. auct. 5. Silicula matura, magnit. dupl. auct.

PLATE XIII.

Fig. 1. *Brassica rerayensis*, Ball. Exemplar pumilum florifer, magnit. nat. 2. Folia caulina duo ex exemplari majore desumpta, magnit. nat. 3. Racemus fructu nondum maturo, magnit. nat. 4. Flos, magnit. auct. 5. Andrœcium cum pistillo, magnit. auct. 6. Pistillum magnit. nat. 7. Siliqua nondum omnino matura, magnit. bis auct.

PLATE XIV.

Fig. 1. *Reseda attenuata*, Ball. Caulis pars inferior et superior fructifera, magnit. nat. 2. Spica florifera exemplaris maerioris, magnit. nat. 3. Flos, magnit. auct. 4. Idem petalis demptis, magnit. auct. 5. Pistillum, magnit. auct. 6. Petalum inferius, magnit. auct. 7. Petalum medium, et superius, magnit. auct.

PLATE XV.

Fig. 1. *Lotononis maroccana*, Ball. Frustulum, magnit. nat. 2. Flos, magnit. auct. 3. Vexillum, magnit. auct. 4. Ala, magnit. auct. 5. Carina magnit. auct. 6. Andrœcium cum pistillo, magnit. auct. 7. Pistillum, magnit. auct. 8. Legumen immaturum, magnit. nat.

PLATE XVI.

Fig. 1. *Ononis polyphylla*, Ball, magnit. nat. 2. Calyx cum legumine nondum maturo, magnit. auct. 3. Calyx florifer cum pedunculo aristato, magnit. auct. 4. Vexillum, magnit. auct. 5. Ala, magnit. auct. 6. Carina, magnit. auct. 7. Andrœcium, magnit. auct. 8. Pistillum, magnit. auct. 9. Legumen magnit. auct.

PLATE XVII.

Fig. 1. *Ononis atlantica*, Ball. Ramus florifer et fructifer, magnit. nat. 2. Folium caulinum, magnit. auct. 3. Flos cum pedunculo spinescens, magnit. auct. 4. Vexillum, magnit. auct. 5. Carina, magnit. auct. 6. Andrœcium, magnit. nat. 7. Ala, magnit. auct.

PLATE XVIII.

Fig. 1. *Trifolium atlanticum*, Ball, magnit. nat. 2. Flos, magnit. auct. 3. Vexillum, magnit. auct. 4. Ala, magnit. auct. 5. Carina, magnit. auct. 6. Andræcium, magnit. auct. 7. Pistillum, magnit. auct.

PLATE XIX.

Fig. 1. *Rubus debilis*, Ball, magnit. nat. 2. Flos, petalis delapsis, magn. nat.

PLATE XX.

Fig. 1. *Sedum modestum*, Ball. Exemplar plantæ macrioris, magnit. nat. 2. Flos hexamerus, magnit. auct. 3. Calyx fructifer, magnit. auct. 4. Exemplar pumilum (forma frequentior), magnit. nat. 5. Folium, magnit. auct. 6. Petalum, magnit. auct.

PLATE XXI.

Fig. 1. *Galium acuminatum*, Ball. Forma contracta, magnit. nat. 2. Idem; forma laxa, ramus florifer, magnit. nat. 3. Flos, magnit. auct. 4. Parum accurata, deleatur. 5. Fructus immaturus, magnit. auct. 6. Folium ex verticillo inferiore, magnit. auct.

PLATE XXII.

Fig. 1. *Galium Bourgæanum*, var. *maroccanum*, Ball, magnit. nat. 2. Flos, magnit. auct. 3. Parum accurata, deleatur. 4. Fructus, magnit. auct. 5. Folium ex verticillis superioribus, magnit. auct. 6. Folium ex verticillis mediis, magnit. auct.

PLATE XXIII.

Fig. 1. *Gnaphalium helichrysoides*, Ball, magnit. nat. 2. Flosculus fœmineus, magnit. auct. 3. Stylus fl. fœminci, magnit. auct. 4. Flosculus hermaphroditus, magnit. auct. 5. Antheræ ejusdem, magnit. auct. 6. Stylus ejusdem, magnit. auct.

PLATE XXIV.

Fig. 1. *Anacyclus depressus*, Ball, magnit. nat. 2. Flosculus fœmineus, magnit. auct. 3. Stylus ejusdem, magnit. auct. 4. Flosculus hermaphroditus exterior, magnit. auct. 5. Antheræ ejusdem, magnit. auct. 6. Stylus ejusdem, magnit. auct.

PLATE XXV.

Fig. 1. *Anthemis tenuisecta*, Ball, magnit. nat. 2. Flosculus fœmineus, magnit. auct. 3. Stylus ejusdem, magnit. auct. 4. Flosculus hermaphroditus, magnit. auct. 5. Anthera ejusdem, magn. auct. 6. Stylus ejusdem, magn. auct.

PLATE XXVI.




Fig. 1. *Chrysanthemum atlanticum*, Ball. Forma scapis foliaceis, magnit. nat. 2. Idem. Forma scapis nudis, magnit. nat. 3. Flosculus fœmineus, magnit. auct. 4. Stylus ejusdem, magnit. auct. 5. Flosculus hermaphroditus, magnit. auct. 6. Antheræ ejusdem, magnit. auct. 7. Stylus ejusdem, magnit. auct.


PLATE XXVII.

Fig. 1. *Thymus maroccanus*, Ball. Ramus florifer, magnit. nat. 2. Flos, magnit. auct. 3. Flos expansus, magnit. auct. 4. Stamen, magnit. auct. 5. Calyx cum pistillo corolla delapsa, magnit. auct. 6. Ovarium, magnit. auct.


PLATE XXVIII.

Fig. 1. *Salvia maurorum*, Ball. Caulis pars superior florifera, cum ramo foliaceo, magnit. nat. 2. Flos, magnit. auct. 3. Stamina, magnit. auct. 4. Calyx cum pistillo corolla delapsa, magnit. auct. 5. Ovarium, magnit. auct.













W.H. Fitch lith

BRASSICA FRUTICULOSA Ba.

Fitch del.










L. TONONIS MAROCCANA Ball


















GALIUM EGYPTIACUM
ZIZIPHORA CLINOPODIODES



GNAPHALIUM HELICHRYSOIDES, Ball.

W.H. Fitch lith

etch imp



W H Fitch hth

ANACYCLUS DEPRESSUS, Ball.

Fitch imp







W.H. Fitch del.

THYMUS MAROCCANUS, B. & H.

Fitch imp.



INDEX.

	Page		Page
Abrotanella rhynehoearpa, <i>Balf.</i>		African tropical Rubiaceæ, W. P.	
<i>fil. sp. n.</i>	9, 11, 16	Hiern on	248
—, variation of leaves of	24	Agaricus campanulatus, 38, 41;	
Acacia Farnesiana	442	campestris, 38; edodes, 50; gue-	
— gummifera	442	piniiformis, <i>B.</i> , (footnote) 50;	
Acanthus mollis	606, 740	lachnocephalus, <i>B.</i> , 48; melleus,	
Acaste, subgen. characterized	166	var. minor, 38; micromegas, <i>B.</i> ,	
Acer monspessulanum	392	51; sapineus, 38, 41; scabrius-	
Aceras anthropophora, 672; in-		culus, <i>B.</i> , 48; semisupinus, <i>B.</i>	
tacta, 672; pyramidalis, 672.		& <i>Br.</i> , 49; subfunereus, <i>B.</i> , (foot-	
Achillea ligustica, 505; var. foliosa,		note) 50; vaginatus, 38.	
<i>Ball</i> , 505; <i>Santolina</i> , 505.		Agrimonia odorata	444
Achyranthes argentea	643	Agrostis alba, var. coarctata, 713,	
Acidantha, characters of	78, 159	var. maritima, 713, var. <i>Fonta-</i>	
— æquinoctialis, 160; bicolor,		nesii, 713; castellana, 714, var.	
160; brachystachys, 160; brevi-		hispanica, 714; pallida, 714; se-	
collis, 160; flexuosa, 160; gra-		tacca, 714; verticillata, 713.	
minifolia, 160; <i>Huttoni</i> , 160;		Aira capillaris, 717; caryophyllea,	
platypetala, 160; tubulosa, 160;		717, var. β . microstachya, 717,	
unicolor, 160.		var. <i>Cupaniana</i> , 717.	
Aerostichum aureum, 34, 37; lati-		Airopsis minuta	718
folium, 34, 37, 205; lineare, 205;		Aizoon canariense, 460; hispani-	
mucosum, 37; platyrhynchos,		cum, 460.	
34; quercifolium, 34; scolopen-		Ajuga Iva	634
drifolium, 37; sorbifolium, 205;		Albertæ, tribe of Rubiaceæ	268
spathulatum, 37; spicatum, 34;		Algæ, freshwater, of Cape of Good	
squamosum, 37, var. leucocolepis,		Hope, Reinsch on	232
<i>Baker</i> , 205.		— of Rodriguez	6
Adenocarpus anagryifolius	397	Alisma plantago	682
— telonensis	396	— ranunculoides	682
Adiantum æthiopicum, 35; <i>Capil-</i>		Allium ampeloprasum, 690; <i>Cha-</i>	
lus-veneris, 27, 735; caudatum,		mæmoly, 692; flavum, var.	
27; flabellatum, 27; hispidu-		tauricum, 690; nigrum, 692;	
lum, 199; <i>Kaulfussii</i> , 34; lmu-		pallens, 690; paniculatum, 691,	
latum, 27, 199; macrophyllum,		var. tenuiflorum, 691; penduli-	
35; <i>Steerei</i> , <i>Harrington</i> , <i>n. sp.</i> ,		num, 691; roseum 691; rotun-	
34.		dum, 690; sphærocephalum,	
Adina microcephala	263	690; subhirsutum, 691, var.	
Admiralty Islands, Fungi of	48	subvillosum, 691; triquetrum,	
—, Lichens of	226	691.	
Adonis æstivalis, 304; autumnalis,		Alœ lomatophylloides, <i>Balf. fl.</i>	
303; microearpa, 304, var.		<i>sp. n.</i>	11, 22
dentata, 304.		Alopecurus pratensis	707
Ægilops ovata, 733; triuncialis,		Alophia, characters of	76, 125
733; ventricosa, 733.		— Drummondiana, 126; genieu-	
Æruea congesta, <i>Balf. fl. sp. n.</i> 11, 20		lata, 126; linearis, 126.	

- | | Page | | Page |
|--|---------|--|---------|
| <i>Alsophila ferox</i> , 34; <i>latebrosa</i> , 25. | | <i>Anomatheca cruenta</i> , 154; <i>grandiflora</i> , 154; <i>junceae</i> , 154. | |
| <i>Althæa hirsuta</i> , var. <i>grandiflora</i> , <i>Ball</i> | 375 | <i>Anthemis fuscata</i> , 506; <i>heterophylla</i> , 507; <i>mixta</i> , 506, 739; <i>tenuisecta</i> (Pl. XXV.), 506; <i>tubereulata</i> , 506. | |
| <i>Alyssum alpestre</i> , 320, var. <i>macrosepalum</i> , <i>Ball</i> , 320; <i>calycinum</i> , 320; <i>campestre</i> , 321; <i>libyeum</i> , 321; <i>maritimum</i> , 321, var. <i>lepidoides</i> , <i>Ball</i> , 321; <i>montanum</i> , var. <i>atlanticum</i> , <i>Ball</i> , <i>an Bois.</i> ?, 320; <i>spinosum</i> , 321. | | <i>Anthericum Liliago</i> , var. <i>bætieum</i> | 693 |
| <i>Amaralia bigoniæflora</i> | 267 | <i>Anthistiria palcacea</i> | 734 |
| <i>Amarantus albus</i> , 643; <i>Blitum</i> , 643, var. <i>græcizans</i> , 643; <i>cholorostachys</i> , 643; <i>deflexus</i> , 643. | | <i>Anthoeros minutus</i> , <i>Mitt. n. sp.</i> (Pl. IV.), 195; <i>punctatus</i> , 739. | |
| <i>Amblystegium commutatum</i> | 738 | <i>Antholyza</i> , characters of | 79, 178 |
| <i>Ammi majus</i> , 467; var. <i>tenuis</i> , <i>Ball</i> , 468; <i>Visnaga</i> , 468. | | — <i>abyssiniea</i> , 179; <i>æthiopica</i> , 179, var. <i>bicolor</i> , 179, var. <i>ringens</i> , 179; <i>caffra</i> , 179; <i>Cunonia</i> , 179; <i>fucata</i> , 180; <i>huillensis</i> , 179; <i>intermedia</i> , 180; <i>nervosa</i> , 179; <i>paniculata</i> , 180; <i>quadrangularis</i> , 179; <i>saccata</i> , 180. | |
| <i>Ammophila arenaria</i> | 715 | <i>Anthospermeæ</i> , tribe of <i>Rubiaceæ</i> | 274 |
| <i>Ampelodesmos tenax</i> | 716 | <i>Anthospermum asperuloides</i> , 274; <i>muricatum</i> , 274; <i>pachyrrhizum</i> , 274. | |
| <i>Amphiroa fragilissima</i> , var. <i>linearis</i> | 7 | <i>Anthoxanthum odoratum</i> , 709; <i>ovatum</i> , 709, and var., 709. | |
| <i>Anaeyclus clavatus</i> , 504; <i>depressus</i> , <i>Ball</i> (Pl. XXIV.), 503; <i>maroccanus</i> , 504; <i>radiatus</i> , 504, 739, var. <i>ochroleucus</i> , <i>Ball</i> , 504; <i>Valentinus</i> , 505. | | <i>Anthyllis Gerardi</i> , 421; <i>hamosa</i> , 422; <i>Tejedensis</i> , 422; <i>tetraphylla</i> , 422; <i>vulneraria</i> , 421, var. <i>Dillenii</i> , 421. | |
| <i>Anagallis arvensis</i> , 562, 740, var. <i>latifolia</i> , 562, 740; <i>crassifolia</i> , 563; <i>linifolia</i> , 562, 740, var. <i>collina</i> , <i>Ball</i> , 562, var. <i>microphylla</i> , <i>Ball</i> , 562; <i>tenella</i> , 563. | | <i>Antiaris toxicaria</i> , detachment of branches of | 181 |
| <i>Anagyris foetida</i> | 394 | <i>Antirrhæa frangulacea</i> | 9 |
| — — —, var. <i>unnamed</i> | 394 | <i>Antirrhinum intricatum</i> , 596; <i>majus</i> , 595; <i>Orontium</i> , 595, var. <i>indicum</i> , 595, var. <i>grandiflorum</i> , 595, 740; <i>tortuosum</i> , var. <i>calyce pubescente</i> , 596. | |
| <i>Anarrhinum fruticosum</i> | 595 | <i>Antrophyum callæfolium</i> , 33; <i>immersum</i> , 33; <i>lineatum</i> , 37; <i>plantagineum</i> , 33; <i>semicostatum</i> , 33. | |
| — <i>pedatum</i> | 595 | <i>Aphloia mauritiana</i> , <i>heterophylla</i> of | 24 |
| <i>Anchusa atlantica</i> , 571; <i>granatensis</i> , 740; <i>italica</i> , 571; <i>macrophylla</i> , 571; <i>undulata</i> , 571. | | <i>Aphyllanthes monspeliensis</i> | 693 |
| <i>Aneylanthos Bainesii</i> , 270; <i>cine-rascens</i> , 270; <i>cistifolius</i> , 270; <i>fulgidus</i> , 270; <i>rubiginosa</i> , 270. | | <i>Api</i> (New Hebrides), <i>Fungi</i> from | 40 |
| <i>Andraehne maroccana</i> | 663 | <i>Apium graveolens</i> | 467 |
| <i>Andropogon annulatus</i> , 734; <i>hirtus</i> , 734, var. <i>longearistatus</i> , 734; <i>laniger</i> , 734. | | — <i>nodiflorum</i> | 467 |
| <i>Androsolen</i> , sect. of <i>Sisyrinchium</i> , defined | 116 | <i>Apogon</i> , subgen. characterized | 137 |
| <i>Andryala integrifolia</i> , 541, var. <i>arenaria</i> , 541; <i>laxiflora</i> , 541; <i>mogadorensis</i> , 540; <i>ragusina</i> , 539, var. <i>lyrata</i> , 540. | | <i>Aquilegia vulgaris</i> | 309 |
| <i>Anemia phyllitidis</i> | 37 | <i>Arabis albidæ</i> , 317; <i>auriculata</i> , 318; <i>conringioides</i> , <i>Ball</i> (Pl. XI.), 319; <i>decumbens</i> , <i>Ball</i> (Pl. X.), 318; <i>crubeseens</i> , 318; <i>pubesceens</i> , 318; <i>verna</i> , 317. | |
| — <i>tomentosa</i> | 37 | <i>Arbutus Unedo</i> | 557 |
| <i>Anemone palmata</i> | 303 | <i>Arcea aurea</i> | 10 |
| <i>Aneura multifida</i> | 194 | <i>Arenaria emarginata</i> , 365; <i>fasciculata</i> , 365; <i>procumbens</i> , 365; <i>pungens</i> , 364, var. <i>glabresceens</i> , <i>Ball</i> , 364; <i>serpylli folia</i> , 364; <i>setacea</i> , 366; <i>spatulata</i> , 364; | |
| <i>Angiopteris erecta</i> | 34 | | |
| <i>Angræcum Gilpinæ</i> , <i>Reich. fl. & S. Moore, n. sp.</i> | 206 | | |
| <i>Anomatheca</i> , characters of | 78, 153 | | |
| — <i>angolensis</i> | 154 | | |

- Page
- tenuifolia, 365, var. glandulosa, 365; trinervia, 365; verna, 367.
- Argania Sideroxylon 563
- Argostemma punilum 264
- Argyrolobium Linnaeanum, 396, var. fallax, 396; microphyllum, 396; stipulaceum, 396.
- Arisarum vulgare 684
- Aristea, characters of 75, 110
- anceps, 111; angolensis, 111; capitata, 111; compressa, 111; cyanea, 111; dichotoma, 111; Eckloni, 112; juncifolia, 111; madagascariensis, 111; melaleuca, 112; pusilla, 111; racemosa, 111; schizolæna, *Harv. MS.*, 111; spiralis, 111; Wredowia, 112.
- Aristeæ, tribe of Iridaceæ, characters of 75
- Aristida adscensionis 712
- plumosa 712
- Aristolochia bætica, 656; longa, 656; Pistolochia, 656.
- Armeria gaditana, var. tingitana, *Ball*, 560; mauritanica, 560; plantaginica, 561.
- Arrhenatherum elatius 720
- Artemisia absinthium 513
- herba-alba 513
- Arthonia polymorphoides, *Nyl.*, *sp. n.* 216, 226
- rubella inferiuscula 217
- Arthrocnemum glaucum 646
- Aru, Fungi from 41
- Arum italicum, 614; tenuifolium, 684
- Arundo Donax, 716; Pliniana, 716.
- Asparagopsis alba 696
- —, var. pastorianus 696
- Asparagus acutifolius, 695; aphyllus, 695; fœniculaceus, 696; madagascariensis, *Baker, n. sp.*, 206; scaber, 695; scoparius, 696; stipularis, 695.
- Asperococcus sinuosus 6
- Asperula aristata, 488, var. scabra, 488; hirsuta, 489; species unnamed, 489.
- Asphodelus cerasiferus, 692; fistulosus, 692, var. tenuifolius, 693; microcarpus, 692, 740.
- Aspidium aculeatum, 29; amabile, 29; aristatum, 29; falcatum, 29; varium, 29.
- Asplenium aculeatum, 36; adiantum nigrum, 736, var. Virgilii, 736; affine, 29, var. pecten, *Baker*, 199; auritum, 35; bantamense, 29; brevipes (Diplazium), *Baker, n. sp.*, 201; Ceterach, 736; cicutarium, 36; cultratum, 29; extensum, 33; filix fœmina, 36, 736; fragile, 35; fragrans, *Sw.*, var. fœniculaceum, *H. B. K.*, 35; furcatum, 35; Gilpinæ, *Baker, n. sp.*, 200; Hemionitis, 735, 740; herpetopteris, *Baker, n. sp.*, 200; japonicum, *Thunb.*, var. coreanum, *Baker*, 29; lætum, 199; lanceolatum, 736, var. obovatum, 736; lanceum, 29; latifolium, 29; lunulatum, 28, 35, 199; marinum, 736; monanthemum, 35; nidus, 28; nigripes, 200; oligophyllum, 199; Petrarchæ, 736; resectum, 29; rhizophyllum, 55; Steerci, *Harrington, n. sp.*, 28; tenerum, 28; trichomanes, 35, 736; triphyllum, 36; viride, 735; viviparum, 200; Wightianum, 28.
- Asterella hemisphærica 739
- Asterolinum Linum stellatum 561
- Astragalus atlanticus, 433; bæticus, 432; chlorocyaneus, 434; epiglottis, 431; glaux, 433, var. rostrata, *Ball*, 433; hamosus, 431; incurvus, 434; lanigerus, var. glabrescens, *Ball*, 432; lusitanicus, 432; ochroleucus, 434; pentaglottis, 431; Reinii, *Ball, n. sp.*, 432; sesamus, 431; Solandri, 431.
- Atavism or reversion of plants 207
- Atractylis cæspitosa, 519; cancellata, 519; flava, 519; humilis, 519; macrophylla, 520.
- Atriplex Halimus, 645; laciniata, 645; parviflora, 645; patula, 645.
- Aulacocalyx jasminiflora 268
- Aurelia Bronssonnetii 681
- Avena barbata, 719, and var. 719; bromoides, var. genuina, 719, var. microstachya, 719; pratensis, 719; sativa, 719; sterilis, 719.
- Averrhoa Bilimbi 231
- Azores (Furnas), Lichens from 217
- Babiana, characters of 78, 164
- Bainesii, 165; cuneifolia, 165; disticha, 165; Dregei, 165; fimbriata, 166; flabellifolia, *Harv. MS.*, 165; hypogæa, 165; mucronata, 165; plicata, 165; pygmæa, 165; ringens, 166; sambucina, 165; secunda, 166; spa-

- | | Page | | Page |
|--|---------|---|------|
| thacea, 165; Sprengelii, 165; | | Boerhavia maroccana | 638 |
| stricta, 166, var. angustifolia, 166, | | — plumbaginea | 638 |
| var. obtusifolia, 166, var. purpu- | | Boletus ananæeeps | 38 |
| rea, 166, var. rubrocyanca, 166, | | — subtomentosus | 38 |
| var. sulphurea, 166, var. villosa, | | Borago officinalis | 571 |
| 166; Thunbergii, 165; tubata, | | Botrychium, rhizome of | 5 |
| 165; tubiflora, 165. | | Botryococcus giganteus, <i>Reinsch</i> , | |
| Babiana et Acaste, characters of . . | 78 | <i>sp. n.</i> | 239 |
| — et Eubabiana, characters of . . | 78 | Bouccrosia maroccana | 566 |
| Bæomyces chilensis | 223 | Boulger, G. S., on distribution of | |
| Bahia, Lichens from | 219 | perfoliate penny-cress (<i>Thlaspi</i> | |
| Baker, J. G., on coll. of Ferns made | | <i>perfoliatum</i>) in Britain | 183 |
| by Miss H. Gilpin in Madagascar | 197 | Brachypodium distachyum, 731; | |
| —, Systema Iridacearum | 61 | pinnatum, 731, var. australe, | |
| Balfour, I. B., Algæ collected by, | | 731; ramosum, 731. | |
| at Rodriguez | 6 | Branches, on the disarticulation of | 180 |
| —, Phænogamic vegetation of | | Brassica adpressa, 327, var. lasio- | |
| Rodriguez, with descriptions of | | earpa, <i>Ball</i> , 327; alba, 327; | |
| new plants | 7 | arvensis, 327; elata, <i>Ball</i> | |
| Ball, J., Spicilegium Floræ Maroc- | | (Pl. XII.), 326; fruticulosa, 325; | |
| canæ | 281 | geniculata, 327; hispida, 327; | |
| Ballota hirsuta, 628; nigra, 629. | | nervosa, 328; nigra, 326; rera- | |
| Bartramia (Philonotus) fontana . | 738 | ensis, <i>Ball</i> (Pl. XIII.), 326; | |
| — rigida | 738 | spec. nov.? <i>Ball</i> , 328; torulosa, | |
| Bartsia aspera, 602; purpurea, 602; | | 325; varia, 325. | |
| Trixago, 601; viscosa, 601. | | Briza maxima, 721; minor, 722. | |
| Batrachospermum moniliforme, | | Bromus alopecurus, 728; macro- | |
| var. æquinoctiale | 7 | stachys, 729, var. lanuginosus, | |
| Beatonia, subgen. characterized . | 136 | 729; matritensis, 729; maximus, | |
| Beilia, subgen., characters of . . | 158 | 729; mollis, 728, var. leiosta- | |
| Belemcanda, characters of . . | 76, 113 | chys, 728, var. ramosus, <i>Ball</i> , | |
| — chinensis | 113 | 728; rubens, 729; tectorum, 729. | |
| Bellis annua, 494, var. minuta, | | Bryonia acuta, 459; dioica, 459. | |
| 494; cærulescens, 495; sylves- | | Bryum alpinum, 738; pallescens, | |
| tris, var. pappulosa, 494. | | 738; capillare, 738; turbinatum, | |
| Belonophora coffeoides | 268 | 738. | |
| Benzonia corymbosa | 275 | Buchanan, J., on root-stock of <i>Ma-</i> | |
| Berberis cretica | 311 | <i>rattia fraxinea</i> | 2 |
| Berkeley, Rev. M. J., on Fungi | | Buffonia tenuifolia | 367 |
| of 'Challenger' Expedition . . | 38 | Bupleurum acutifolium, 466; ca- | |
| Bermuda, Lichens from | 214 | nescens, 466; dumosum, 465; | |
| Bertiera æthiopica, 266; africana, | | foliosum, 466; fruticescens, 465; | |
| 266; bracteolata, 266; brevifo- | | fruticosum, 467; gibraltarium, | |
| lia, 266; laxa, 266; macrocarpa, | | 467; glaucum, 464; lateriflorum, | |
| 266; montana, 266; subsessilis, | | 467; oblongifolium, <i>Ball</i> , 466; | |
| 266. | | Odontites, 464; oligactis, 465; | |
| Beta vulgaris, var. maritima . . | 644 | paniculatum, 464; procumbens, | |
| Betonica officinalis, var. algeriensis | 624 | 464; protractum, 464; spino- | |
| Biflora testiculata | 475 | sum, 464; tenuissimum, var. co- | |
| Biscutella apula, 332, var. macro- | | lumnæ, 464. | |
| carpa, 332; lyrata, 332. | | Cakile maritima | 336 |
| Biserrula Pelecinus | 434 | Calamintha Acinos, 614; alpina, | |
| Blechnum occidentale, 35; orien- | | var. parviflora, <i>Ball</i> , 614; at- | |
| tale, 28; serrulatum, 35. | | lantica, 614; gravecolens, 614; | |
| Blimbing, movements of leaves of . | 231 | menthæfolia, 613. | |
| Bobartia, characters of . . | 76, 114 | Calendula arvensis, 516, var. bi- | |
| — aphylla, 114; filiformis, 114; | | color, 516, var. echinata, <i>Ball</i> , | |
| gladiata, 114; spathacea, 114. | | | |

	Page		Page
516, var. mierantha, <i>Ball</i> , 516 ;		<i>Capparis spinosa</i> , 739, var. <i>Ægyptia</i> ,	
maroecana, 517 ; officinalis, 516 ;		337, var. canescens, 337.	
stellata, 516 ; suffruticosa, 517,		<i>Capsella Bursa-Pastoris</i>	330
var. tomentosa, 517.		<i>Cardamine hirsuta</i>	319
<i>Callipetis cueullaria</i>	483	— — —, var. <i>sylvatica</i>	319
<i>Callitriche vernalis</i>	455	<i>Cardiostigma</i> , characters of	74, 102
<i>Callitris quadrivalvis</i>	670	— — — <i>longispatha</i>	102
<i>Calluna vulgaris</i>	556	<i>Carduncellus lueens</i>	532
<i>Calocasia indica</i> , how used by		— — — <i>pinnatus</i>	532
<i>Samoans</i>	57	<i>Carduus Ballii</i> , 520 ; <i>macrocephalus</i> ,	
<i>Calophyllum inophyllum</i>	56	520 ; <i>myriacanthus</i> , 521 ;	
<i>Calycotome villosa</i>	398	<i>pycnoccephalus</i> , 521, var. <i>Sardois</i> ,	
— — —, var. <i>intermedia</i> , <i>Ball</i>	398	521, var. <i>tenuiflorus</i> , <i>Ball</i> , 521.	
<i>Calydorea</i> , characters of	74, 110	<i>Carex ambigua</i> , 705 ; <i>basilaris</i> ,	
— — — <i>campestris</i> , 101 ; <i>fureata</i> , 102 ;		704 ; <i>distans</i> , 705 ; <i>divisa</i> , 704 ;	
<i>Gardneri</i> , 101 ; <i>luteola</i> , 102 ;		<i>divulsa</i> , 704 ; <i>fissirostris</i> , 705 ;	
<i>nuda</i> , 102 ; <i>punctata</i> , 102 ; <i>speciosa</i> ,		<i>glauca</i> , var. <i>attenuata</i> , <i>Ball</i> , 706,	
101 ; <i>texana</i> , 101.		var. <i>serrulata</i> , <i>Ball</i> , 706 ; <i>Halleriana</i> ,	
<i>Calypogeia bidentula</i>	191	704 ; <i>Linkii</i> , 704 ; <i>vulpina</i> , 704.	
— — — <i>Trichomanis</i>	738	<i>Carissa xylopicron</i> , heterophylly	
<i>Calystegia sepium</i>	578	of	23, 24
— — — <i>Soldanella</i>	577	<i>Carlina involuerata</i> , 519 ; <i>lanata</i> ,	
<i>Camiguin</i> (Philippines), Fungi from	45	519 ; <i>racemosa</i> , 519.	
<i>Campanula diohotoma</i> , 553, var.		<i>Caroxylon articulatum</i>	647
<i>parviflora</i> , 553 ; <i>Erinus</i> , 554 ;		<i>Carrihtera Vellæ</i>	330
<i>Lœflingii</i> , 554, var. <i>decumbens</i> ?,		<i>Carthamus cœruleus</i>	532
740 ; <i>maroccana</i> , 554 ; <i>mollis</i> ,		— — — <i>lanatus</i>	532
553 ; <i>Rapunculus</i> , 554, var.		<i>Carum ammoides</i> , 470 ; <i>dichotomum</i> ,	
<i>calyec strigosus</i> , 554.		469 ; <i>mauritanicum</i> , 469 ;	
Cane named "Whangec," note on	1	<i>Petroselinum</i> , 469 ; <i>Ridolfia</i> ,	
<i>Cantharellus cibarius</i> , 38 ; <i>concinus</i> ,		469 ; <i>tennifolium</i> , 470.	
<i>B.</i> , 38 ; <i>flabellatus</i> , <i>B.</i>		<i>Carumbium acuminatum</i> , 57 ; <i>pedicellatum</i> ,	
(fnote), 50.		57, 58 ; <i>populifolium</i> ,	
<i>Canthium acutiflorum</i> , 269 ; <i>Afzelianum</i> ,		57 ; species of, 57.	
269 ; <i>anomoearpum</i> ,		<i>Castilloa</i> , detachment of branches	
269 ; <i>Barteri</i> , 269 ; <i>bibraetatum</i> ,		of	180
269 ; <i>caudatiflorum</i> ,		— — — <i>elastica</i> , leafy shoots of (Pl.	
269 ; <i>coassum</i> , 269 ; <i>congensc</i> ,		III.)	182
269 ; <i>Cornelia</i> , 269 ; <i>discolor</i> ,		<i>Casuarina equisetifolia</i>	57, 59
269 ; <i>fetidum</i> , 269 ; <i>glabriflorum</i> ,		<i>Catanauche cœrulea</i> , 533, var.	
269 ; <i>glaucum</i> , 269 ; <i>gracile</i> , 269 ;		<i>tenuis</i> , 533 ; <i>cæspitosum</i> , 533 ;	
<i>Heudelotii</i> , 269 ; <i>hispidum</i> , 269 ;		<i>lutea</i> , 534 ; <i>montana</i> , 534.	
<i>horizontalis</i> , 269 ; <i>lividum</i> , 269 ;		<i>Caucalis cœrulescens</i> , 478 ; <i>daucoides</i> ,	
<i>lanceiflorum</i> , 269 ; <i>lividum</i> , 269 ;		477 ; <i>heterophylla</i> , 479 ;	
<i>Mannii</i> , 269 ; ? <i>multiflorum</i> ,		<i>infesta</i> , 478, var. <i>neglecta</i> , 478 ;	
269 ; <i>neglectum</i> , 269 ; <i>nervosum</i> ,		<i>latifolia</i> , 477 ; <i>leptophylla</i> , 478,	
269 ; <i>nitens</i> , 269 ; <i>oligocarpum</i> ,		var. <i>heteroearpa</i> , <i>Ball</i> , 478 ;	
269 ; <i>polycarpum</i> , 269 ; <i>rubens</i> ,		<i>nodosa</i> , 478, var. <i>heteroearpa</i> , <i>Ball</i> ,	
269 ; <i>Schimperianum</i> , 269 ; <i>setiflorum</i> ,		478.	
269 ; <i>setosum</i> , 269 ;		<i>Caulerpa Boryana</i> , 7 ; <i>crucifolia</i> , 7 ;	
<i>subcordatum</i> , 269 ; <i>Thonningii</i> ,		<i>mamillosa</i> , 7 ; <i>plumaris</i> , 7.	
269 ; <i>vangueroides</i> , 269 ; <i>Vatkeanum</i> ,		<i>Celastrus senegalensis</i>	391
269 ; <i>venosum</i> , 269 ;		<i>Celsia Barnadesii</i> , 585 ; <i>betonicæ-</i>	
<i>zanzibaricum</i> , 269.		<i>folia</i> , 585 ; <i>crotia</i> , var. <i>Cavanillesii</i> ,	
Cape Hepaticæ	187	584 ; <i>maroecana</i> , 586 ; <i>ramosissima</i> , 586.	
Cape of Good Hope, freshwater		<i>Celtis australis</i>	666
Algae of	232	<i>Centaurea aspera</i> , 531 ; <i>caleitrapa</i> ,	
Cape Verde, Lichens from	211		
<i>Capnophyllum dichotomum</i>	474		

- | | Page | | Page |
|--|----------|---|----------|
| 529; <i>Chamaerhaponticum</i> , <i>Ball</i> , | | <i>Chlora imperfoliata</i> , 567; perfo- | |
| 526; <i>Clementi</i> , 528; <i>Cosso-</i> | | liata, 567, var. <i>mascariensis</i> , 567. | |
| niana, 527; <i>diluta</i> , 531; <i>crio-</i> | | <i>Chlorochytrium</i> , spec. ? | 247 |
| phora, 529; <i>fragilis</i> , 531, var. | | <i>Chlorococcum africanum</i> , <i>Reinsch</i> , | |
| <i>integrifolia</i> , <i>Ball</i> , 531; <i>incana</i> , | | <i>sp. n.</i> | 239 |
| 528, var. <i>Hookeriana</i> , <i>Ball</i> , | | <i>Chnoospora implexa</i> | 6 |
| 529; <i>maroccana</i> , 539; <i>meliten-</i> | | <i>Chrococlonium procumbens</i> | 244 |
| <i>sis</i> , 529; <i>polyacantha</i> , 539; <i>pul-</i> | | <i>Chorisis</i> , fission, or bifurcation in | |
| <i>lata</i> , 527, 740, var. <i>minor</i> , 528; | | in plants | 207, 209 |
| <i>salmantica</i> , 531, var. <i>Clusii</i> , 531, | | <i>Chroococcus cohaerens</i> | 234 |
| var. <i>leptolonehus</i> , 532; <i>seridis</i> , | | — <i>turgidus</i> | 233 |
| var. <i>auriculata</i> , 539; <i>sulphurea</i> , | | <i>Chroolepus Montis Tabulae</i> , | |
| 539; <i>tagana</i> , 526. | | <i>Reinsch, n. sp.</i> | 245 |
| <i>Centranthus angustifolius</i> | 490 | <i>Chrysanthemum atlanticum</i> , <i>Ball</i> | |
| — <i>calcitrapa</i> | 490, 739 | (Pl. XXVI.), 510; <i>carinatum</i> , | |
| — var. <i>orbiculata</i> | 490 | 508; <i>catananche</i> , 511; <i>corona-</i> | |
| <i>Cephaelis clausa</i> , 273; <i>congensis</i> , | | <i>rium</i> , 508; <i>Gayanum</i> , 509, var. | |
| 273; <i>cornuta</i> , 273; <i>fuscescens</i> , | | <i>depressum</i> , <i>Ball</i> , 510; <i>glabrum</i> , | |
| 273; <i>latifolia</i> , 273; <i>Mannii</i> , 273; | | 509; <i>macrotum</i> , 509; <i>Marsii</i> , | |
| <i>peduncularis</i> , 273; <i>rubescens</i> , | | var. <i>Hosmariense</i> , <i>Ball</i> , 510; | |
| 273; <i>spathacea</i> , 273; <i>suaveo-</i> | | <i>myconis</i> , 508; <i>scgetum</i> , 508; | |
| <i>lens</i> , 273. | | <i>Webbianum</i> , 509. | |
| <i>Cephalanthera xiphophyllum</i> | 674 | <i>Chthonoblastus salinus</i> , <i>Kütz.</i> , var. | 234 |
| <i>Cephalanthum</i> , sect. of <i>Sisyrinchium</i> | | <i>Chylocladia rigens</i> | 7 |
| defined | 119 | <i>Cicendia pusilla</i> | 568 |
| <i>Cephalozia bicuspidata</i> , 738; <i>te-</i> | | <i>Cichorium Intybus</i> | 534 |
| <i>nuissima</i> , 187; <i>Turneri</i> , 738. | | — <i>pumilum</i> | 534 |
| <i>Cerastium arvense</i> , 363; <i>brachy-</i> | | <i>Cinchonae</i> , tribe of <i>Rubiaceae</i> | 263 |
| <i>petalum</i> , 363; <i>glaucum</i> , 362; | | <i>Cipura</i> , characters of | 76, 125 |
| <i>glomeratum</i> , 362; <i>pumilum</i> | 362 | — <i>cubensis</i> , 125; <i>paludosa</i> , 125. | |
| <i>Ceratocnemum</i> , genus characterized | 334 | <i>Cistopteris fragilis</i> | 735 |
| — <i>rapistroides</i> | 335 | <i>Cistus albidus</i> , 342; <i>crispus</i> , 342; | |
| <i>Ceratonia siliqua</i> | 442 | <i>ladaniferus</i> , 343; <i>monspeliensis</i> , | |
| <i>Ceratophyllum demersum</i> | 669 | 343; <i>polymorphus</i> , 342, var. | |
| <i>Cerithe major</i> | 577 | <i>corsicus</i> , 342, var. <i>creticus</i> , 342; | |
| —, var. <i>gymnandra</i> | 577 | <i>populifolius</i> , 343, 739; <i>salviae-</i> | |
| 'Challenger' Expedition, Fungi of | 38 | <i>folius</i> , 343, var. <i>macrocalyx</i> , | |
| —, <i>Lichens</i> of | 211 | 343. | |
| <i>Chamaerops humilis</i> | 683 | <i>Citrullus Colocynthis</i> | 459 |
| <i>Chantransia caeruleascens</i> | 7 | <i>Cladanthus arabicus</i> | 507 |
| <i>Characium coronatum</i> , <i>Reinsch</i> , | | <i>Cladia aggregata</i> | 223 |
| <i>sp. n.</i> | 247 | <i>Cladonia acuminata hebescens</i> , 214; | |
| <i>Cheilanthes farinosa</i> , 27; <i>fragrans</i> , | | <i>adspersa</i> , 227; <i>cornuta</i> , 220; | |
| 735; <i>madagascariensis</i> (<i>Adiant-</i> | | <i>fibula</i> , 214; <i>fimbriata</i> , 217, f. <i>cos-</i> | |
| <i>opsis</i>), <i>Baker, n. sp.</i> , 198; <i>Mat-</i> | | <i>tata</i> , <i>Flk.</i> , 222; <i>mitrula</i> , 214; | |
| <i>thewsii</i> , 35; <i>microphylla</i> , 35; | | <i>sobolifera</i> , 217. | |
| <i>myriophylla</i> , 35; <i>scariosa</i> , 35; | | <i>Cladophora pannosa</i> , 7; <i>Roettleri</i> , | |
| <i>tenuifolia</i> , 27. | | 7; <i>setiformis</i> , <i>Kütz.</i> , var. <i>pachy-</i> | |
| <i>Cheiranthus semperflorens</i> | 316 | <i>derma</i> , 244. | |
| <i>Chenolea canariensis</i> | 646 | <i>Clavaria aurea</i> , 40; <i>botrytis</i> , 40; | |
| <i>Chenopodium album</i> , 644; <i>Ambro-</i> | | <i>flava</i> , 40. | |
| <i>sioides</i> , 644, 740; <i>murale</i> , 644; | | <i>Cleanthe</i> , series of <i>Aristea</i> , cha- | |
| <i>olidum</i> , 644; <i>opulifolium</i> , 644. | | acters of | 112 |
| <i>Chilosecyphus expansus</i> | 188 | <i>Clematis balcarica</i> , 303; <i>cirrhusa</i> , | |
| <i>Chiodecton rubrocinctum</i> | 220 | 303; <i>flammula</i> , 303, 739. | |
| <i>Chlamylostylus</i> , characters of, 75, 107 | | <i>Cleonia lusitanica</i> | 620 |
| — <i>bractcolatus</i> , 107; <i>Medusa</i> , | | <i>Clerodendron</i> , 9; <i>heterophyllum</i> | 19 |
| 107; <i>multiflorus</i> , 107; <i>tenuis</i> , | | — <i>laciniatum</i> , <i>Balf. fil. sp. n.</i> 11, 19 | |
| 107; <i>triflorus</i> , 107. | | — —, <i>heterophylly</i> of | 24 |

	Page
* <i>Cnicus Casabonæ</i> , 523; <i>chrysacanthus</i> , 522; <i>celinatus</i> , 522; <i>giganteus</i> , 521; <i>ornatus</i> , 522; <i>syriacus</i> , 522.	
<i>Cocos nucifera</i>	10
<i>Codium tomentosum</i>	7
— <i>adhaerens</i>	7
<i>Cœlastrum scabrum</i> , <i>Reinsch</i> , <i>sp. n.</i>	238
— <i>verrucosum</i> , <i>Reinsch</i>	238
<i>Cœnogonium confervoides</i>	222
<i>Coffea Afzelii</i> , 271; <i>arabica</i> , 9, 271; <i>brevipes</i> , 271; <i>hypoglaucæ</i> , 271; <i>jasminoides</i> , 271; <i>liberica</i> , 271; <i>melanocarpa</i> , 271; ? <i>microcarpa</i> , 271; <i>racemosa</i> , 271; <i>rupestris</i> , 271; <i>stenophylla</i> , 271; <i>subcordata</i> , 271; <i>Zanguebariæ</i> , 271.	
<i>Colchicum arnarium</i> , 694; <i>autumnale</i> , 694; <i>Bivonæ</i> , 694.	
<i>Collema byrsinum</i> (<i>Dichodium</i>), 224, 226	
<i>Collemopsis pelia</i>	211
<i>Colutea arborescens</i>	430
<i>Conferva Ansonii</i> , 7; <i>bombycina</i> , var. <i>inequalis</i> , <i>Ktze.</i> , 7; <i>lævis</i> , 244; <i>moluccana</i> , 7.	
<i>Conium maculatum</i>	463
Contributions to the Botany of H.M.S. 'Challenger,' Fungi, 38; Lichens, 211.	
<i>Convallaria multiflora</i>	694
<i>Convolvulus althæoides</i> , 579; <i>arvensis</i> , 578; <i>Cantabrica</i> , 578; <i>sabatius</i> , 578, var. <i>atlanticus</i> , <i>Ball</i> , 578; <i>siculus</i> , 579; <i>sp. n.</i> ? 740; <i>tricolor</i> , 578; <i>undulatus</i> , 578.	
<i>Coprinus nychthemercus</i>	40
<i>Cora pavonia</i>	53
<i>Corbularia Bulbocodium</i>	680
— <i>obesa</i>	680
<i>Coriandrum sativum</i>	475
<i>Coriaria myrtifolia</i>	394
<i>Coris Monspelienensis</i>	561
<i>Coronella glauca</i> , 426; <i>juncæ</i> , 427; <i>minima</i> , 427; <i>pentaphylla</i> , 427; <i>ramosissima</i> , 427; <i>repanda</i> , 428; <i>scorpioides</i> , 428; <i>viminalis</i> , 427.	
<i>Corrigiola telephiifolia</i>	639
<i>Corydalis heterocarpa</i>	314
<i>Corynanthe paniculata</i>	264
<i>Cosmariium abnorme</i> , 241; <i>asperulum</i> , <i>Reinsch</i> , <i>n. sp.</i> , 212; <i>hammeri</i> B, <i>intermedium</i> , <i>Reinsch</i> , 240; <i>parvulum</i> , 240; <i>plicatum</i> , <i>Reinsch</i> , B, <i>minus</i> , 241; <i>pulcherrimum</i> , 241; <i>punctulatum</i> , 241.	
<i>Cotyledon Cossoniana</i> , 451; <i>hispanica</i> , 450, var. <i>Salzmanni</i> ,	

	Page
<i>Ball</i> , 451, 739, <i>mucizona</i> , 450; <i>umbilicus</i> , 449, var. <i>horizontalis</i> , 450.	
<i>Crambe hispanica</i>	334
<i>Cratægus oxyacantha</i>	447
—, var. <i>foliis obovato-cuneatis</i>	447
<i>Craterispermum brachynematum</i> , 270; <i>cerinanthum</i> , 270; <i>laurinum</i> , 270; <i>montanum</i> , 270; <i>Schweinfurthii</i> , 270.	
<i>Cremaspora africana</i>	268
— ? <i>Thomsoni</i>	268
<i>Crepis bulbosa</i> , 538; <i>Hookeriana</i> , 538; <i>tingitana</i> , 537; <i>taraxacifolia</i> , 537, var. <i>intybacca</i> , 538, var. <i>stellata</i> , <i>Ball</i> , 538; <i>virens</i> , 538.	
<i>Crithmum maritimum</i>	472
Croceæ, tribe of Irids, characters of	73
<i>Crocisma</i> , sect. of <i>Tritonia</i> , defined	163
<i>Crocus</i> , characters of	73, 79
— <i>acrius</i> , 83, var. <i>cyprius</i> , 83, var. <i>pulchricolor</i> , 83; <i>alatavicus</i> , 82; <i>aleppicus</i> , 83; <i>banaticus</i> , 81; <i>biflorus</i> , 80, var. <i>Adami</i> , 81, var. <i>argenteus</i> , 81, var. <i>nubigenus</i> , 81, var. <i>pusillus</i> , 81, var. <i>Weldeni</i> , 81; <i>Boryii</i> , 84, var. <i>cretensis</i> , 84, var. <i>orphanidis</i> , 84; <i>byzantinus</i> , 85; <i>Cambessedesii</i> , 83; <i>carpetanus</i> , 82; <i>Cartwrightianus</i> , 82; <i>caspicus</i> , 84; <i>chrysanthus</i> , 80, var. <i>fusco-lineatus</i> , 80, var. <i>fusco-tinctus</i> , 80; <i>Clusii</i> , 84; <i>Crewei</i> , 80; <i>damascenus</i> , 84; <i>etruscus</i> , 82; <i>Fleischeri</i> , 84; <i>gargaricus</i> , 80; <i>hadriaticus</i> , 83; <i>hyemalis</i> , 84, var. <i>Gaillardotii</i> , 84; <i>Imperati</i> , 83; <i>Kotschyianus</i> , 83; <i>laziens</i> , 83; <i>longiflorus</i> , 83, var. <i>melitensis</i> , 83; <i>mæsiacus</i> , 80, var. <i>aureus</i> , 80, var. <i>lacteus</i> , 80, var. <i>stellaris</i> , 80, var. <i>sulphureus</i> , 80; <i>medius</i> , 85; <i>minimus</i> , 81; <i>nudiflorus</i> , 85, var. <i>asturicus</i> , 85; <i>ochroleucus</i> , 84; <i>Olivieri</i> , 83; <i>Pallasii</i> , 82; <i>parviflorus</i> , 82; <i>peloponnesiacus</i> , 83; <i>pulehellus</i> , 85; <i>reticulatus</i> , 82, var. <i>albicans</i> , 82, var. <i>Dalmaticus</i> , 82; <i>sativus</i> , 82; <i>Scharojani</i> , 83; <i>Schimperi</i> , 84; <i>scrobinus</i> , 83, var. <i>Salzmanni</i> , 84, 677; <i>Sieberi</i> , 81, var. <i>Atticus</i> , <i>Orphanides MS.</i> , 82, var. <i>exiguus</i> , 82, var. <i>veluchensis</i> , 82; <i>speciosus</i> , 85; <i>suaveolens</i> , 81; <i>susianus</i> , 80, var. <i>ancyrensis</i> , 80,	

- var. *immaculatus*, 80; *suterianus*, 83; *syriacus*, 84; *Thomasii*, 82; *vallicola*, 84; *vernus*, 81, var. *albiflorus*, 81, var. *nervifolius*, 81, var. *nivigenus*, 81, var. *obovatus*, 81, var. *Tommasiannus*, 81; *versicolor*, 81, var. *crestensis*, 81; *visianicus*, 82; *vitellinus*, 84.
- Crombie, Rev. J. M., the Lichens of the 'Challenger' Expedition (with a revision of those enumerated by Dr. J. Stirton in Linn. Journ. Bot. xiv. pp. 366-375), 211.
- Crossopteryx Kotschyana* . . . 264
- Crucianella augustifolia* . . . 489
- *maritima* . . . 489
- Crupina vulgaris* . . . 525
- Crypsis aculeata* . . . 706
- *schenoides* . . . 706
- Cuscuta planiflora* . . . 580
- Cuviera acutiflora*, 270; *longiflora*, 270; *subuliflora*, 270; *trilocularis*, 270.
- Cyathea spinulosa* . . . 25
- Cynara Cardunculus*, 524; *humilis*, 524; *Hystrix*, 524.
- Cynodon Dactylon* . . . 717
- Cynoglossum cheirifolium*, 570, var. *heterocarpum*, 570; *clandestinum*, 570; *Dioseoridis*, var. *nebrodense*, 569; *pietum*, 570.
- Cynosurus aureus*, 725; *echinatus*, 724; *elegans*, 724.
- Cypella*, characters of . . . 77, 128
- *gracilis*, 129; *Herberti*, 128, var. *Polia ramosa*, 129.
- Cyperus ægyptiacus*, 701; *distachyos*, 700; *fuscus*, 701; *lævigatus*, 700; *longus*, var. *badius*, 701; *Mundtii*, 700; *rotundus*, 701.
- Cystopteris fragilis* . . . 34
- Cytinus Hypocistis* . . . 656
- Cytisus albidus*, 403; *arbores*, 402; *bætieus*, 402, var. *macranthus*, *Ball*, 402; *Balansæ*, var. *atlanticus*, *Ball*, 403; *candicans*, 403; *cinnatus*, *Ball*, 404; *Fontanesii*, 405; *Hosmariensis*, 404; *linifolius*, 405, 739; *tridentatus*, var. *lasiantha*, 405; *triflorus*, 403.
- Dactylis glomerata*, 724, var. *australis*, 724, var. *juncinella*, *B.*, 724.
- Dædalia glabrescens*, *B.*, 39, *indurata*, *B.*, 43; *unicolor*, 51.
- Damasonium stellatum* . . . 682
- Dammara robusta* . . . 180
- Danais corymbosa*, *Balf. fil. sp. n.*, 11, 13; *fragrans*, 14.
- Daphne Gnidium*, 653, 740; *Laureola*, 653, var. ? *Hosmariensis*, *Ball*, 653.
- Datura Stramonium* . . . 582
- Daucus aureus*, 477; *Carota*, 476; *crinitus*, 477; *Gingidium*, 476, var. *hispidus*, *Ball*, 477; *maximus*, 476; *muricatus*, 475; *parviflorus*, 476; *pumilus*, 477; *setulosus*, 475.
- Davallia affinis*, 26; *canariensis*, 735; *contigua*, 26; *elegans*, 27; *Hookeriana*, 27; *hymenophylloides*, 26; *inæqualis*, 34; *paralela*, 26; *pedata*, 26; *pentaphylla*, 27; (*Microlepis*) *philippensis*, *Harrington, n. sp.*, 27; *pulehella*, 26; *repens*, 26; *strigosa*, 27; *tenuifolia*, 27; *vestita*, 26.
- Delphinium Ajacis*, 310; *halteratum*, 310, var. *cardiopetalum*, 310, var. *macropetalum*, 310; *sp. nov.*? *Ball*, 310; *pentagynum*, 311.
- Desmodium gyrans*, movements of, 231
- Deverra juncea*, *Ball* . . . 468
- *scoparia* . . . 468
- Dialysis of organs of plants . . . 207
- Dianthus attenuatus*, 354; *gaditanus*, 355; *siculus*, 354; *virginicus*, 354.
- Dichone*, sect. of *Tritonia*, defined . 163
- Dickie, G., on Algæ collected by Dr. I. B. Balfour at Rodriguez . 6
- Dicksonia Barometz*, 25; *flabellifolia*, 197; *Henriettæ*, 197; *hypolepidoides*, *Baker, n. sp.*, 197.
- Dicranella varia* . . . 737
- Dietyandra arborescens* . . . 266
- *involucrata* . . . 266
- Dietyosperma alba*, var. *aurea* . . 10
- Dietyosphæria favulosa* . . . 7
- Dietyota dichotoma* . . . 6
- Didymochlæna lunulata* . . . 29
- Dicrama*, characters of . . . 74, 99
- *pendula* . . . 99
- *pulcherrima* . . . 99
- Dietes*, series of *Iris*, characters of . 147
- Digenea simplex* . . . 6
- Digitalis laciniata* . . . 599
- *lutea* . . . 599
- Digitaria sanguinalis* . . . 710
- Diodia arenosa*, 274; *brevisetata*, 274; *Kirkii*, 274; *maritima*, 274; *rubricosa*, 274; *vaginalis*, 274.
- Diospyros diversifolia* . . . 11

- | | Page | | Page |
|---|----------|--|-----------|
| <i>Diotis maritima</i> | 506 | <i>Enteromorpha compressa</i> , 7, var. | |
| <i>Dipcadi serotinum</i> , var. <i>fulvum</i> | 689 | <i>trichodes</i> , <i>Kütz.</i> 246; <i>intestina-</i> | |
| <i>Diplarrhena</i> , characters of | 77, 148 | <i>lis</i> , 7; <i>ramulosa</i> , 7. | |
| — <i>latifolia</i> , 149; <i>Moræa</i> , 149, | | <i>Enterospermum littorale</i> | 266 |
| var. <i>alpina</i> , 149. | | <i>Entosthodon obtusus</i> | 738 |
| <i>Diplostaxis catholica</i> , 329; <i>crucoides</i> , 329; <i>Harra</i> , 329; <i>siifolia</i> , 329; <i>tenuifolia</i> , 329; <i>tenuisiliqua</i> , 329, var. <i>rupestris</i> , 330; <i>virgata</i> , 329. | | <i>Ephedra altissima</i> , 669, var. <i>nana</i> ?, 669; <i>procera</i> , 669. | |
| <i>Dipsacus sylvestris</i> | 492 | <i>Epilobium hirsutum</i> , 458; <i>parviflorum</i> , 458; <i>tetragonum</i> , 458, var. ? <i>tingitanum</i> , 459. | |
| <i>Dirichletia glabra</i> , 264; <i>glaucescens</i> , 264; <i>pubescens</i> , 264. | | <i>Equisetum ramosissimum</i> | 737 |
| <i>Dirina repanda</i> , var. | 213 | — <i>Telmateja</i> | 737 |
| Disarticulation of branches, literature on | 180 | <i>Erica arborea</i> , 556; <i>australis</i> , 556; <i>ciliaris</i> , 556; <i>multiflora</i> , 556; <i>scoparia</i> , 557; <i>umbellata</i> , 556. | |
| <i>Dombeya ferruginea</i> , heterophylly of, 24; variation of leaves of, 24. | | <i>Erigeron linifolium</i> | 495 |
| <i>Doryenium rectum</i> , 422, var. <i>pauciflorum</i> , <i>Ball</i> , 422. | | <i>Erodium atlanticum</i> , 384; <i>Botrys</i> , 387; <i>chium</i> , 387; <i>ciconium</i> , 387; <i>cicutarium</i> , 385, var. <i>bipinnatum</i> , 385, var. γ , 385, var. <i>maculatum</i> , 385; <i>glaucophyllum</i> , 388; <i>guttatum</i> , 388; <i>Jacquinianum</i> , 386; <i>laciniatum</i> , 387; <i>litoreum</i> , 387; <i>malacoides</i> , 387; <i>moschatum</i> , 386, var. <i>dissectum</i> , 387; <i>petraeum</i> , 384; <i>romanum</i> , 386. | |
| <i>Draba hispanica</i> | 322 | <i>Erophila vulgaris</i> | 322 |
| <i>Drosophyllum lusitanicum</i> | 455 | <i>Eruca sativa</i> | 330 |
| <i>Drymoglossum carnosum</i> , 33, var. <i>obovatum</i> , <i>Harrington</i> , 33, var. <i>subcordatum</i> , <i>Hook. & Baker</i> , 33; <i>piloselloides</i> , 33. | | <i>Eryngium aquifolium</i> , 462; <i>Bourgati</i> , var. <i>atlanticum</i> , 461; <i>campestre</i> , 461; <i>dichotomum</i> , 461; <i>dilatatum</i> , 461; <i>ilicifolium</i> , 462; <i>maritimum</i> , 461; <i>tenue</i> , 462; <i>tricuspidatum</i> , 462; <i>triquetrum</i> , 461; <i>variifolium</i> , 461. | 324 |
| Eaton's, Rev. A. E., freshwater Algæ of Cape, <i>Reinsch</i> on | 232 | <i>Erysimum australe</i> | |
| —, <i>Hepaticæ</i> collected by | 187 | <i>Erythraea Centaurium</i> , 568, var. <i>suffruticosa</i> , 568; <i>maritima</i> , 568; <i>ramosissima</i> , 567; <i>spicata</i> , 568. | |
| <i>Ebenus pinnata</i> | 436 | <i>Erythrodictus punctatus</i> | 694 |
| <i>Ecbalium Elaterium</i> | 459, 739 | <i>Euantholyza</i> , subgen. characterized | 179 |
| <i>Echinaria capitata</i> | 716 | <i>Euaristia</i> , series of <i>Aristia</i> , characters of | 111 |
| <i>Echinops Ritro</i> , 739; <i>spinosus</i> , 518; <i>strigosus</i> , 518. | | <i>Euastrum subincisum</i> , <i>Reinsch</i> , <i>sp.</i> <i>n.</i> | 242 |
| <i>Echinospermum barbatum</i> | 570 | <i>Eubabiana</i> , subgen. characterized | 165 |
| <i>Echium angustifolium</i> , 575, 740; <i>flavum</i> , 577; <i>longifolium</i> , 576, var. <i>maroccanum</i> , <i>Ball</i> , 576; <i>maritimum</i> , 577; <i>modestum</i> , 576; <i>plantagineum</i> , 575, var. <i>grandiflorum</i> , 576; <i>pomponium</i> , 577; <i>sericeum</i> , var. <i>diffusum</i> , 575; <i>tuberculatum</i> , 574. | | <i>Euclilamydostylus</i> , subgen. characterized | 107 |
| <i>Elæodendron orientale</i> , heterophylly of | 24 | <i>Eugenia Balfouri</i> | 9, 11, 13 |
| <i>Elæoselinum exinvolucratum</i> , 480; <i>fœtidum</i> , 481; <i>humile</i> , 479; <i>meoides</i> , 479. | | — <i>cotinifolia</i> , variation of leaves of | 24 |
| <i>Eleutherine</i> , characters of | 74, 100 | <i>Eugladiolus</i> , subgen. characterized | 171 |
| — <i>anomala</i> , 101; <i>plicata</i> , 100. | | <i>Euixea</i> , subgen. characterized | 90 |
| <i>Elymus Caput Medusæ</i> | 732 | <i>Eumontbretia</i> , subgen. characterized | 168 |
| — —, var. <i>crinitus</i> | 732 | <i>Eumoræa</i> , subgen. characterized | 129 |
| <i>Emex spinosa</i> | 648 | <i>Eunemastylus</i> , subgen. characterized | 103 |
| <i>Empogona Kirkii</i> | 267 | | |
| <i>Enation</i> in plants | 207 | | |
| <i>Encalypta vulgaris</i> | 738 | | |
| <i>Endococcus thalamita</i> , <i>Nyl., sp. n.</i> | 217 | | |

- | | Page | | Page |
|---|---------|--|----------|
| <i>Euphorbia akenocarpa</i> , 657; <i>Beau-</i> | | nacea, 728; <i>bromoides</i> , 726; | |
| <i>micrana</i> , 662; <i>Chamæsyce</i> , 656, | | <i>cærulescens</i> , 728; <i>ciliata</i> , 727; | |
| var. <i>canescens</i> , 657; <i>Characias</i> , | | <i>Durandii</i> , 728; <i>dnriuscula</i> , var. | |
| 661; <i>daphnoides</i> , <i>Balf. fil. sp.</i> | | <i>indigesta</i> , 727; <i>geniculata</i> , 727, | |
| <i>n.</i> , 21; <i>Echinus</i> , 663; <i>exigua</i> , | | var. <i>pumila</i> , 727; <i>hemipoa</i> , 725; | |
| 658; <i>falcata</i> , 658, var. <i>rubra</i> , | | <i>maritima</i> , 725; <i>memphitica</i> , 725; | |
| 658; <i>helioscopia</i> , 657; <i>inconspi-</i> | | <i>Myurus</i> , 726, var. <i>Broteri</i> , 726, | |
| <i>cua</i> , 658; <i>medicaginea</i> , 659, var. | | var. <i>sciuroides</i> , 726; <i>rigida</i> , 725; | |
| <i>oblongifolia</i> , 659; <i>megalatlantica</i> , | | <i>tuberculosa</i> , 725; <i>unilateralis</i> , | |
| 659; of <i>Mascarenes</i> , <i>n. sp.</i> , 9; | | var. <i>aristata</i> , 726. | |
| <i>Paralias</i> , 661; <i>Peplis</i> , 656; <i>pe-</i> | | <i>Filago gallica</i> , 497; <i>germanica</i> , | |
| <i>ploides</i> , 659; <i>Peplus</i> , 659; <i>pinca</i> , | | 496, var. γ , <i>lanuginosa</i> , 496; <i>he-</i> | |
| 659; <i>pterococca</i> , 657; <i>pubes-</i> | | <i>terantha</i> , 497; <i>spathulata</i> , 496. | |
| <i>cens</i> , 657; <i>resinifera</i> , 661; <i>rima-</i> | | <i>Fimbriaria marginata</i> | 194 |
| <i>rum</i> , 660; <i>rupicola</i> , 657; <i>terra-</i> | | <i>Fissidens incurvus</i> | 738 |
| <i>cina</i> , 660. | | — <i>taxifolius</i> | 738 |
| <i>Euriridæ</i> , tribe of <i>Irids</i> , characters | | <i>Flora of Marocco</i> , J. Ball on . . . | 281 |
| of | 77 | —, Introductory observations on | 281 |
| <i>Eurydice</i> , subgenus characterized . | 92 | —, percentage of <i>Nat. orders</i> , | |
| <i>Eusolenomelus</i> , subgen. character- | | table of | 298 |
| ized | 121 | —, <i>Præmonenda</i> | 302 |
| <i>Eutigridia</i> , subgen. characterized . | 136 | —, tabular view of species &c., | |
| <i>Eutritonia</i> , section of <i>Tritonia</i> , de- | | and <i>nat. orders</i> &c. | 295 |
| fined | 162 | <i>Fœniculum vulgare</i> | 471 |
| <i>Euxiphion</i> , subgen. characterized . | 122 | — —, var. <i>piperitum</i> | 472 |
| <i>Evansia</i> , subgen. characterized . . | 143 | <i>Fœtidia mauritiana</i> , variation of | |
| <i>Evax Heldreichii</i> | 496 | leaves of | 24 |
| — <i>pygmæa</i> | 495 | <i>Forskohlea tenacissima</i> | 665 |
| <i>Excæcaria agallochia</i> | 57, 58 | <i>Fossombronia angulosa</i> , 738; <i>cris-</i> | |
| <i>Fadogia agrestis</i> , 270; <i>ancylantha</i> , | | <i>pa</i> , 193; <i>leucoxantha</i> , 193; <i>tum-</i> | |
| 270; <i>Cienkowskii</i> , 270; <i>fuchsi-</i> | | <i>mida</i> , <i>Mitt.</i> , <i>n. sp.</i> (<i>Pl. IV.</i>), | |
| <i>oides</i> , 270; <i>glaberrima</i> , 270; <i>lac-</i> | | 193. | |
| <i>tiflora</i> , 270; <i>leucophlœa</i> , 269; | | <i>Frankenia hirsuta</i> , 353, var. <i>ve-</i> | |
| <i>stenophylla</i> , 270; <i>Zeyheri</i> , 269. | | <i>lutina</i> , <i>Ball</i> , 353, var. <i>revoluta</i> , | |
| <i>Fagonia cretica</i> | 381 | 353; <i>pulverulenta</i> , 352. | |
| <i>Favolus brasiliensis</i> , 43; <i>saccha-</i> | | <i>Fraxinus dimorpha</i> | 564 |
| <i>rinus</i> , <i>B.</i> , 43; <i>scaber</i> , 48, var. | | — <i>oxyphylla</i> | 564 |
| <i>platyporus</i> , 48. | | <i>Freesia</i> , characters of | 78, 163 |
| <i>Fedia cornucopiæ</i> | 491 | — <i>Leichtlinii</i> , 164; <i>refracta</i> , 164. | |
| — —, var. = <i>F. graciliflora</i> . | 491 | <i>Frullania brunnea</i> , 193; <i>capensis</i> , | |
| <i>Feretia apodanthera</i> | 267 | 192; <i>trinervis</i> , 193. | |
| — ? <i>canthioides</i> | 267 | <i>Fumana calycina</i> , 349; <i>glutinosa</i> , | |
| <i>Fernando de Noronha</i> , Lichens | | 348, var. <i>viridis</i> , 348; <i>lævipes</i> , | |
| from | 219 | 348. | |
| <i>Fernclia buxifolia</i> | 9, 23 | <i>Fumaria africana</i> , 314; <i>agraria</i> , | |
| — —, heterophylly of | 23 | 315, var. <i>atlantica</i> , <i>Ball</i> , 316, | |
| <i>Ferns</i> , collection of, made by Miss | | var. <i>clata</i> , <i>Ball</i> , 315; <i>capreolata</i> , | |
| Helen Gilpin in Madagascar . . | 197 | var. <i>condensata</i> , <i>Ball</i> , 315, var. | |
| —, tropical, collected by Prof. | | 739; <i>macrosepala</i> , 315; <i>media</i> , | |
| Steere, M. W. Harrington on . . | 25 | 315, var. <i>prox.</i> <i>F. sepium</i> , 315; | |
| <i>Ferraria</i> , characters of | 75, 105 | <i>officinalis</i> , 314; <i>parviflora</i> , 314; | |
| — <i>atrata</i> , 106; <i>divaricata</i> , 106; | | <i>tenuisecta</i> , 316; <i>Vaillantii</i> , 314. | |
| <i>Ferrariola</i> , 106, var. <i>angustifolia</i> , | | <i>Fumaria hygrometrica</i> | 738 |
| 106; <i>obtusifolia</i> , 106; <i>uncinata</i> , | | <i>Fungi</i> , of 'Challenger' Expedition | 38 |
| 106; <i>undulata</i> , 106. | | <i>Gagea foliosa</i> | 685 |
| <i>Ferula communis</i> | 474 | <i>Gaillonia calycoptera</i> | 274 |
| — <i>tingitana</i> | 474 | <i>Galactites tomentosa</i> | 525, 740 |
| <i>Festuca Alopæcurus</i> , 727; <i>arundi-</i> | | <i>Galaxea</i> , characters of | 75, 103 |

	Page
<i>Galaxia graminea</i> , 104; <i>ovata</i> , 103, var. <i>grandiflora</i> , 104, var. <i>muconularis</i> , 104, var. <i>obtusa</i> , 103, var. <i>versicolor</i> , 104.	
<i>Galaxiæ</i> , tribe of Irids, defined . . .	75
<i>Galieæ</i> , tribe of Rubiaceæ . . .	274
<i>Galiniera coffeoides</i>	267
<i>Galium acuminatum</i> , <i>Ball</i> (Pl. XXI.), 485; <i>Aparine</i> , 275, 488; <i>aureum</i> , 485; <i>Bourgæanum</i> , var. <i>maroccanum</i> (Pl. XXII.), 486; <i>concatenatum</i> , 486; <i>corrudæfolium</i> , 484; <i>Decaisnei</i> , 275; <i>divaricatum</i> , 487; <i>ellipticum</i> , 484; <i>erectum</i> , 484; <i>glomeratum</i> , 487, var. <i>campestre</i> , <i>Ball</i> , 487; <i>Mollugo</i> , 275; <i>murale</i> , 488; <i>nolitangere</i> , 486; <i>Parisiense</i> , var. <i>lasiocarpum</i> , 487; <i>Poirertianum</i> , <i>Ball</i> , 484; <i>rotundifolium</i> , 275; <i>saccharatum</i> , 487; <i>setaceum</i> , 487; <i>simense</i> , 275; <i>spurium</i> , var. <i>Vaillantii</i> , 488; <i>sylvestre</i> , 485; <i>tricorne</i> , 488; <i>tunetanum</i> , 486.	
<i>Gardenia Abbeokuta</i> , 267; <i>Annæ</i> , 367; <i>assimilis</i> , 267; <i>fernandensis</i> , 267; <i>Jovis-tonantis</i> , 267; <i>konguensis</i> , 267; <i>Manganjæ</i> , 267; <i>nitida</i> , 267; <i>resiniflua</i> , 267; <i>Thunbergia</i> , 267; <i>urcelliformis</i> , 267; <i>Vogelii</i> , 267; ? <i>zanguebarica</i> , 267.	
<i>Gastridium lendigerum</i>	714
<i>Gaudinia fragilis</i>	718
Gawler's 'Genera Iridearum' . . .	61
<i>Geaster dubius</i> , <i>B.</i> , 40; <i>minimus</i> , 40; <i>saccatus</i> , 40.	
<i>Geissorhiza</i> , characters of . . .	74, 93
— <i>alpina</i> , 94; <i>Bojeri</i> , 94; <i>ciliaris</i> , 94; <i>erecta</i> , 93; <i>excisa</i> , 95; <i>filifolia</i> , 94; <i>filiosa</i> , 95; <i>furva</i> , 94; <i>geminata</i> , 95; <i>grandis</i> , 95; <i>hirta</i> , 94; <i>humilis</i> , 93; <i>imbricata</i> , 94; <i>inflexa</i> , 94; <i>juncæa</i> , 93; <i>latifolia</i> , 94; <i>lutea</i> , 95; <i>minima</i> , 95; <i>obtusata</i> , 94; <i>ornithogaloides</i> , 94; <i>purpureo-lutea</i> , 93; <i>quadrangula</i> , 95; <i>Rochensis</i> , 94; <i>secunda</i> , 93, var. <i>ramosa</i> , 94, var. <i>setifolia</i> , 94; <i>setacea</i> , 95; <i>Wrightii</i> , 94.	
<i>Gelasine</i> , characters of	75, 106
— <i>azurea</i> , 107; <i>grandiflora</i> , 107; <i>purpureolucana</i> , 107.	
<i>Gelidium rigidum</i>	7
<i>Genista clavata</i> , 400, var. <i>capitata</i> ?, 400, var. <i>casuarinoides</i> , 401, var. <i>Goudotiana</i> , 401, var. <i>Webbiana</i> , 401; <i>dasycarpa</i> , 398; <i>ferox</i> , 399; <i>florida</i> , var. <i>maroccana</i> , <i>Ball</i> , 400; <i>gibraltarica</i> ,	

	Page
399; <i>juniperina</i> , 399; <i>monosperma</i> , var. <i>Bovei</i> , 398; <i>myriantha</i> , 399; <i>pseudopilosa</i> , 400; <i>triacanthos</i> , 399; <i>tricuspidata</i> , 399; <i>tridens</i> , 399.	
<i>Geophila Afzelii</i> , 273; <i>hirsuta</i> , 273; <i>involucrata</i> , 273; <i>lancistipula</i> , 273; <i>obvallata</i> , 273; <i>reniformis</i> , 273; <i>uniflora</i> , 273.	
<i>Geranium dissectum</i> , 383; <i>lucidum</i> , 383; <i>malvæflorum</i> , 382; <i>molle</i> , 382; <i>pyrenaicum</i> , 382; <i>Robertianum</i> , 383, var. β , <i>purpureum</i> , 383; <i>rotundifolium</i> , 383, var. <i>foliis palmatipartitis</i> , <i>Ball</i> , 383.	
Gilpin, Miss H., ferns collected by	197
<i>Gladiolæ</i> , divisional series of Irids, characters of	78
<i>Gladiolus</i> , characters of	79, 170
— <i>alatus</i> , 177, var. <i>namaquensis</i> , 177; <i>aleppicus</i> , 172; <i>andongensis</i> , <i>Welw. Herb.</i> , 176; <i>angolensis</i> , <i>Welw. Herb.</i> , 176; <i>angustus</i> , 173; <i>arenarius</i> , 178; <i>atropurpureus</i> , 174; <i>atroviolaceus</i> , 172; <i>aurantiacus</i> , 176; <i>benguellensis</i> , 174; <i>bicolor</i> , 178; <i>blandus</i> , 175, var. <i>albidus</i> , 175, var. <i>campanulatus</i> , 175, var. <i>corneus</i> , 175, var. <i>excelsus</i> , 175, var. <i>Gucinzi</i> , 175; <i>Brenckleyensis</i> , 177; <i>brevicaulis</i> , 174; <i>brevifolius</i> , 173, var. <i>Orobanche</i> , 174; <i>byzantinus</i> , 171; <i>cærulescens</i> , 174; <i>candidus</i> , 177; <i>cardinalis</i> , 176; <i>cochleatus</i> , 174; <i>Colvillei</i> , 177; <i>communis</i> , 171, var. <i>Notarisii</i> , 171; <i>Cooperi</i> , 176; <i>corneus</i> , 175; <i>crassifolius</i> , 175; <i>cruentus</i> , 176; <i>cuspidatus</i> , 173, var. <i>ventricosus</i> , 173; <i>debilis</i> , 174; <i>decoratus</i> , 176; <i>delicatus</i> , 177; <i>draecocephalus</i> , 176; <i>Eckloni</i> , 175; <i>floribundus</i> , 176; <i>formosus</i> , 178; <i>fragrans</i> , 177; <i>gandavensis</i> , 177; <i>gracilis</i> , 173; <i>grandis</i> , 172, var. <i>elongatus</i> , 172, var. <i>lævis</i> , 172, var. <i>suaveolens</i> , 173; <i>gregarius</i> , <i>Welw. Herb.</i> , 171; <i>halophilus</i> , 172; <i>hastatus</i> , 173; <i>Haylockianus</i> , 177; <i>Herbertianus</i> , 177; <i>hirsutus</i> , 176; <i>hyalinus</i> , 174; <i>ignescens</i> , 175; <i>illyricus</i> , 171, 674, var. <i>Reuteri</i> , 171; <i>imbricatus</i> , 171, var. <i>crispiflorus</i> , 171, var. <i>hygrophilus</i> , 171, var. <i>libanoticus</i> , 171; <i>incarnatus</i> , 177; <i>inflatus</i> , 174; <i>involutus</i> , 174; <i>Kotschyannus</i> , 172; <i>Lambda</i> , 173;	

- | | Page | | Page |
|---|---------|--|------|
| laxiflorus, 174; longicollis, 173; luridus, <i>Welw. Herb.</i> , 174; luteus, 178; Melleri, 175; Milleri, 176; Mitchamiensis, 177; montanus, 178, var. ramosus, 178; Mortonianus, 176; multiflorus, 175; Newii, 175; ochroleucus, 175; odoratus, 177; oppositiflorus, 176; orchidiflorus, 177; palustris, 171; Papilio, 175; permeabilis, 178; persicus, 172; petræus, 172; propinquus, 177; psittacinus, 176; pubescens, 173; pudibundus, 177; pulchellus, 178; purpureo-aureatus, 175; Quartinianus, 176; ramosus, 177; recurvus, 173; rigidus, 177; saltatorum, 174; Saundersii, 176; segetum, 171, 674, var. commutatus, 172, var. caucasicus, 172, var. Guepini, 172, var. inarinensis, 172, var. Ludovicæ, 172; sericeo-villosus, 175, var. Ludwigii, 175; spathaceus, 172; spicatus, 175; splendens, 176; Spofforthianus, 177; striatus, 174; tenellus, 174; tenuis, 174; Thunbergii, 173; trichonemifolius, 173; triphyllus, 172; tristis, 172, var. concolor, 172; undulatus, 176; unguiculatus, 178; villosus, 174; vittatus, 174; Welwitschii, 175; Willmoreanus, 177. | | Great-Atlas flora, remarks on . . . | 301 |
| Glaucium corniculatum | 313 | Grimmia apocarpa, 737; commutata, 737; orbicularis, 737; pulvinata, 737. | |
| — luteum | 313 | Grumilea articulata, 273; Kirkii, 273; macrocarpa, 273; micrantha, 273; psychotrioides, 273; sphærocarpa, 273; succulenta, 273; sulphurea, 273; venosa, 273. | |
| Gleichenia dichotoma | 25, 197 | Guepinia spathularia | 40 |
| — pubescens | 34 | Guettarda speciosa | 268 |
| Globularia Alypum | 607 | Guettardæ, tribe of Rubiaceæ . . . | 268 |
| Gloeocapsa conglomerata | 233 | Gymnogramme angustifrons, 37; calomelanos, var. denudata, <i>Harrington</i> , 37; elliptica, 33; lanceolata, 33; leptophylla, 205, 737, 740; macrophylla, 33; pumila, 37; tartarea, 37; tomentosa, 37; Wrightii, 33. | |
| — magma | 233 | | |
| Gleothece confluens | 233 | Halimeda opuntia, 7; tuna, 7. | |
| Glyceria distans, var. festuaceformis, 721, var. tenuifolia, 721; fluitans, 721, var. plicata, 721. | | Hamelicæ, tribe of Rubiaceæ . . . | 266 |
| Glycyrrhiza fœtida | 434 | Haplophyllum Broussonetianum . . | 389 |
| Glyphis cicatricosa, f. favulosa . . | 220 | Harrington, M. W., the tropical Ferns coll. by Prof. Steere in the years 1870-75 | 25 |
| Gnaphalium helichrysoides, <i>Ball</i> (Pl. XXIII.) | 499 | Hebea, subgen. characterized . . . | 177 |
| — luteo-album | 499 | Hedera Helix | 481 |
| Gomphocarpus fruticosus | 566 | Hedysarum capitatum, 434; flexuosum, 434; membranaceum, 435. | |
| Gonionema velutinum, <i>Stirton</i> . . . | 211 | Heinsia benguelensis, 266; densiflora, 266; jasminiflora, 266. | |
| Gracilaria corticata | 7 | Hekistocarpa minutiflora | 265 |
| Graphis Afzelii, 216, 219; assimilis, 225; dendrogramma, <i>Nyl.</i> , sp. n., 226; lobata (<i>Lecanactis</i>), 216; scripta, 216, 228, f. pulverulenta, 226; striatula, 226; tenella, 220. | | Helcocharis palustris, 702; uniglumis, 702. | |
| | | Helianthemum ægyptiacum, 346; canariense, 346; glaucum, 347, var. stœchadifolium, 347; guttatum, 345, var. inconspicuum, 345, var. macrosepalum, 345; halimifolium, 344, var. lasiocalyceum, 344, var. sepalis stellato-tomentosis, 344; lavandulæfolium, 346; Libanotis, 343; Lippii, 346; niloticum, 345, var. microcarpum, 345, var. ? pumilum, 345; papillare, 346; rubellum, 346; Tuberaria, 344; umbellatum, var. verticillatum, 344; virgatum, 347, var. strictum, 347. | |
| | | Helichrysum Stœchas | 500 |
| | | Heliotropium europæum, 569; supinum, 569; undulatum, 569. | |
| | | Helixyra, subgen. characterized . . | 132 |
| | | Helminthia aculeata | 537 |
| | | — echioides | 537 |
| | | Helminthostachys zeylanica | 34 |

- | | Page | | Page |
|---|----------------|---|----------------|
| <i>Hemicrambe fruticulosa</i> | 336 | <i>Hirneola rufa</i> , <i>B.</i> | 41, 44, 48, 50 |
| <i>Hemionitis cordata</i> | 33 | <i>Holcus lanatus</i> , 708, var. <i>tuberosus</i> , 708; <i>setiglumis</i> , 708. | |
| — <i>Zollingeri</i> | 33 | <i>Holostium umbellatum</i> | 362 |
| <i>Hemitelia Melleri</i> | 197 | <i>Holostigma</i> , a section of <i>Crocus</i> , defined | 80 |
| Henslow, the Rev. Gt., note on the causes of the numerical increase of parts of plants | 206 | <i>Homalothecium sericeum</i> | 738 |
| <i>Hepaticæ</i> , list of, coll. by the Rev. A. E. Eaton at Cape of Good Hope | 187 | <i>Homeria</i> , characters of | 75, 104 |
| <i>Heppia meiolepis</i> , <i>Cromb.</i> | 212 | — <i>aurantiaca</i> , 105; <i>collina</i> , 105, var. <i>miniata</i> , 105, var. <i>ochroleuca</i> , 105; <i>elegans</i> , 105; <i>lineata</i> , 105, var. <i>porrifolia</i> , 105; <i>maculata</i> , 105. | |
| <i>Heracleum sphondylium</i> | 475 | <i>Homoglossa</i> , subtribe of <i>Irids</i> , characters of | 78 |
| <i>Herbertia</i> , characters of | 77, 133 | <i>Homoglossum</i> , characters of | 78, 161 |
| — <i>brasiliensis</i> , 134; <i>cærulea</i> , 134; <i>Drummondiana</i> , 134; <i>pulchella</i> , 134; <i>umbellata</i> , 134; <i>unguiculata</i> , 134. | | — <i>lucidior</i> , 161; <i>Merianella</i> , 161; <i>revolutum</i> , 161, var. <i>Gawleri</i> , 161. | |
| <i>Hermodyctylus</i> , characters of | 77, 147 | <i>Hordeum bulbosum</i> , 732; <i>maritimum</i> , 732; <i>murinum</i> , 732. | |
| — <i>tuberosus</i> , 148, var. <i>bispathaceus</i> , 148, var. <i>longifolius</i> , 148. | | <i>Houttuynia</i> , subgen. characterized | 168 |
| <i>Herniaria annua</i> , 639, var. <i>virescens</i> , 639; <i>hirsuta</i> , 639; <i>polygonoides</i> , 639. | | <i>Hutchinsia petraea</i> | 333 |
| <i>Hesperantha</i> , characters of | 74, 95 | <i>Hyacinthus romanus</i> | 685 |
| — <i>angusta</i> , 96; <i>Baurii</i> , 96; <i>bulbifera</i> , 96; <i>cinnamomea</i> , 95; <i>falcata</i> , 96; <i>graminifolia</i> , 95; <i>humilis</i> , 95; <i>longituba</i> , 96; <i>Petitioniana</i> , 96; <i>pilosa</i> , 95; <i>radiata</i> , 96, var. <i>caricina</i> , 96; <i>rubella</i> , 96; <i>uniflora</i> , 96. | | <i>Hydnum nigrum</i> , 39; <i>olidum</i> , <i>B.</i> , (fnote) 51; <i>xanthum</i> , 39. | |
| <i>Hesperoxiphon</i> , characters of | 76, 127 | <i>Hydroclathrus cancellatus</i> | 6 |
| — <i>peruvianum</i> , 127; <i>pusillum</i> | 127 | <i>Hydrocoleum homœotrichum</i> | 233 |
| <i>Heteroglossa</i> , subtribe of <i>Irids</i> , characters of | 78 | <i>Hydrocotyle vulgaris</i> | 460 |
| <i>Heterophylly</i> of plants in Rodriguez | 23 | <i>Hydrostaehys Goudotiana</i> | 206 |
| <i>Hexaglottis</i> , characters of | 74, 99 | <i>Hydrotænia</i> , characters of | 77, 135 |
| — <i>longifolia</i> , 100; <i>virgata</i> , 100. | | — <i>Meleagris</i> | 135 |
| <i>Hexagona affinis</i> , 43, 50; <i>albida</i> , <i>B.</i> , 47; <i>arata</i> , <i>B.</i> , 43; <i>cladophora</i> , <i>B.</i> , 47; <i>crinigera</i> , 47; <i>flabelliformis</i> , <i>B.</i> , 47; <i>pieta</i> , 50; <i>polygramma</i> , 44; <i>rigida</i> , <i>B.</i> , 54; <i>tenuis</i> , 41. | | <i>Hymenangium Moselei</i> , <i>B. & Br.</i> | 40 |
| <i>Hexapogon</i> , subgen. characterized | 147 | <i>Hymenodictyon biafranum</i> | 263 |
| <i>Hibiscus</i> (<i>Paritium</i>) <i>liliaceus</i> | 24 | — <i>kurria</i> | 263 |
| — <i>liliiflorus</i> , <i>heterophylly</i> of | 24 | <i>Hymenophyllum dilatatum</i> , 26; <i>fraternum</i> , <i>Harrington, n. sp.</i> , 26; <i>javanicum</i> , 26; <i>Neesii</i> , 26; <i>polyanthos</i> , 26, 34; <i>thuidium</i> , <i>Harrington, n. sp.</i> , 25; <i>tunbridgense</i> , 26. | |
| — <i>tricuspis</i> , supposed insular form of <i>H. liliaceus</i> | 25 | <i>Hyophorbe amaricaulis</i> , 10; <i>indica</i> , 10; <i>Verschaffeltii</i> , 10, 11. | |
| <i>Hieracium Pilosella</i> | 539 | <i>Hyoseyanus albus</i> | 582 |
| Hiern, W. P., on the peculiarities and distribution of <i>Rubiaceæ</i> in Tropical Africa | 248 | — —, var. <i>major</i> | 582 |
| <i>Hippocrepis atlantica</i> , 429; <i>multisiliquosa</i> , 429, var. <i>major</i> , 429; <i>Salzmanni</i> , 430; <i>scabra</i> , 429. | | <i>Hyoseris radiata</i> | 534 |
| <i>Hippomarathrum cristatum</i> , 462; <i>Libanotis</i> , 463; <i>pterochlaenum</i> , 462. | | <i>Hypecoum pendulum</i> | 313 |
| <i>Hirneola polytricha</i> | 41, 50, 52, 53 | <i>Hypericum australe</i> , 373; <i>coadunatum</i> , 374; <i>perfoliatum</i> , 735; <i>perforatum</i> , 373; <i>pubescens</i> , 374; <i>tomentosum</i> , 374; <i>undulatum</i> , 373. | |
| | | <i>Hypertrophy</i> of plants | 207, 208 |
| | | <i>Hypnea hamulosa</i> , 7; <i>Valentiae</i> , 7. | |
| | | <i>Hypnum circinatum</i> (<i>Eurhynchium</i>), 738; <i>illecebrum</i> (<i>Scleropodium</i>), 738; <i>rutabulum</i> (<i>Brachythecium</i>), 738; <i>tenellum</i> (<i>Rhynchostegium</i>), 738; <i>velutimum</i> (<i>Brachythecium</i>), 738. | |

	Page		Page
Hypochaeris æthnensis, 542; glabra, 541, var. 541, var. crostris, 541, var. Salzmanniana, 541; lævigata, 542; leontodontoides, 542; radicata, var. heterocarpa, 542.		nea, 139, var. inodora, 139, var. latifolia, 139; Hartwegii, 138; Haynei, 145; hexagona, 141; humilis, 137; hybrida, 146; iberica, 142, var. Heylandiana, 142, var. ochracea, 142; japonica, 143; Korolkowi, 145; kumaonensis, 144; lacustris, 143; lævigata, 142; longicarpa, 147; longipetala, 142; lurida, 145; lutescens, 144; macrosiphon, 138; maculata, 140; Monnieri, 141; neglecta, 146; nepalensis, 143; ochroleuca, 141; pallida, 146, var. australis, 146, var. cucullata, 146, var. Junonia, 146, var. Mandraliscæ, 146, var. sicula, 146, var. Tinæi, 146; paradoxa, 142; plicata, 146; prismatica, 138; Pseudacorus, 140, var. acoriformis, 140, var. acoroides, 140, var. Bastardi, 140; pseudopumila, 143, var. panormitana, 143; pumila, 143, var. æquiloba, 143, var. attica, 143, var. cærulea, 143, var. stenoloba, 143, var. tristis, 143; Redouteana, 145; Robinsoniana, 147; rubro-marginata, 144; ruthenica, 138; Saarii, 142; sambucina, 146, var. concolor, 146; setosa, 140; sibirica, 139, var. flexuosa, 139, var. orientalis, 139, var. trigonocarpa, 140; songarica, 138; speculatrix, 143; spuria, 140, var. daenensis, 140, var. desertorum, 140, var. Dierinckii, 140, var. notha, 140, var. sogdiana, 140; squalens, 145, var. atro-purpurea, Baker, 146, var. lavandulacea, Baker, 145; suaveolens, 143; susiana, 142, var. livida, 142; Swertii, 146; tenax, 138; tenuifolia, 138; Tigridia, 144; Tolmieana, 138; tridentata, 140; tripetala, 138; unguicularis, 138; variegata, 145, var. belgica, 145; ventricosa, 138; verna, 144; versicolor, 141, var. flaccida, 141, var. virginica, 141; virescens, 144.	
Hypodematium sphærostigma	274	Irpex consors, B., 51; maximus, 44; zonatus, B., 48.	
Hypoëstes inconspicua, Balf. fil. sp. n., 11, 18; rodriguesiana, Balf. fil. sp. n., 11, 18.		Isatis tinctoria, L., var. lætevirens, Ball	334
Hypoxylon concentricum	44	Isoëtes Hystrix	737
— vernicosum	45		
Hyssopus officinalis	613		
Iberis gibraltarica	333		
— parviflora	333		
Ifloga spicata	497		
Ilcebrum verticillatum	640		
Imperata cylindrica	734		
Inaccessible Island, lichens of	222		
Inula crithmoides, 500; montana, 500; viscosa, 500.			
Iridacæ, bounds of the order	70		
—, conspectus, tribes and gen. of	73		
—, fruit and seeds of	69		
—, genera, table of	72		
—, geograph. distrib. of	70		
—, inflorescence and bracts of	64		
—, leaf of	64		
—, perianth of	66		
—, pistil of	68		
—, rootstock and root of	63		
—, stamens of	68		
—, stem of	63		
—, table of geogr. distrib. of	71		
Iridacearum Systema	61		
Iridæ, divisional series of Irids, characters of	76		
Iris, characters of	77, 136		
— acutiloba, 142; aphylla, 145; arenaria, 144; aurea, 141; bicolor, 147; biflora, 144; Bloudovii, 144; catenulata, 147; Chamæiris, 144, var. balkana, 144, var. italica, 144, var. olbiensis, 144; compressa, 147; cretensis, 138; cristata, 143; dichotoma, 142; Douglasiana, 138, var. Beecheyana, 138; ensata, 139; Guldenstædtiana, 141, var. subbarbata, 141; falcifolia, 147; flavescens, 145; flavissima, 144; florentina, 146, 676, var. albicans, 146; foetidissima, 141; fulva, 142; germanica, 146, 676; glaucescens, 144, var. elongata, Fischer, MS., 144, var. longiflora, Herb. MS., 144; goniocarpa, 145; gracilipes, 143; grami-			

- | | | | |
|---|--------|---|---------|
| | Page | | Page |
| <i>Ixia</i> , characters of | 74, 90 | <i>Klattia</i> , characters of | 75, 109 |
| — <i>anemonæflora</i> , 91; <i>aristata</i> , 91; <i>campanulata</i> , 91; <i>columellaris</i> , 91, var. <i>rhodolarynx</i> , <i>Baker</i> , 91, var. <i>rubro-cærulea</i> , <i>Baker</i> , 92, var. <i>rubro-lilacina</i> , <i>Baker</i> , 92; <i>curta</i> , 92; <i>flexuosa</i> , 90; <i>fragrans</i> , 91; <i>fureata</i> , 91; <i>hybrida</i> , 90; <i>leucantha</i> , 91; <i>lutea</i> , 91; <i>maculata</i> , 91, var. <i>fusco-citrina</i> , 91, var. <i>ochroleuca</i> , 91; <i>micandra</i> , 90; <i>monadelphæa</i> , 92; <i>patens</i> , 91; <i>polystachya</i> , 90, var. <i>flavescens</i> , 90; <i>speciosa</i> , 91; <i>viridiflora</i> , 92, var. <i>cæsia</i> , 92, var. <i>cana</i> , 92. | | <i>Klatt's papers on the Iridacæ</i> , reference to | 62 |
| <i>Ixiæ</i> , series of Irids, characters of | 73 | <i>Kobe Kiotos</i> , lichens from | 227 |
| <i>Ixora brachysiphon</i> , 270; <i>breviflora</i> , 270; <i>coccinea</i> , 275; <i>foliosa</i> , 270; <i>guineensis</i> , 270; <i>inundata</i> , 270; <i>laxiflora</i> , 270; <i>minutiflora</i> , 270; <i>modesta</i> , 270; <i>odorata</i> , 270; <i>radiata</i> , 270; <i>riparia</i> , 270; <i>Soyauxii</i> , 270; <i>Thomsoni</i> , 270. | | <i>Kœleria phleoides</i> , 723, var. <i>submutica</i> , <i>Ball</i> , 723; <i>villosa</i> , var. <i>longearistata</i> , 723. | |
| <i>Ixoreæ</i> , tribe of <i>Rubiaceæ</i> | 270 | <i>Kundmannia sicula</i> | 472 |
| <i>Jackson, J. R.</i> , note on the uses of a commercial cane termed "Whangee," a species of <i>Phyllostachys</i> | 1 | <i>Lachnocladium furecellatum</i> | 44 |
| <i>Japan</i> , fungi from | 50 | — <i>semivestitum</i> | 44 |
| <i>Jasione atlantica</i> , 553; <i>blepharodon</i> , 552; <i>cornuta</i> , 552; <i>corymbosa</i> , 552. | | <i>Lactarius subtomentosus</i> | 38 |
| <i>Jasminum fruticans</i> | 563 | <i>Lactuca saligna</i> , 546; <i>tenerrima</i> , 546; <i>viminea</i> , 546. | |
| <i>Juan Fernandez</i> , fungi from | 52 | <i>Lagurus ovatus</i> | 715 |
| —, lichens from | 223 | <i>Lanium album</i> , var. <i>parviflora</i> ? 628; <i>amplexicaule</i> , 628; <i>flexuosum</i> , 628. | |
| <i>Juncus acutus</i> , 697; <i>bufonius</i> , 699, var. <i>fasciculatus</i> , 699; <i>capitatus</i> , 699; <i>conglomeratus</i> , 697, var. <i>effusus</i> , 697; <i>fasciculatus</i> , 699; <i>foliosus</i> , 700; <i>glaucus</i> , 697; <i>heterophyllus</i> , 698; <i>lamprocarpus</i> , 698; <i>maritimus</i> , 697; <i>multiflorus</i> , 697; <i>pygmæus</i> , 699; <i>striatus</i> , 698; <i>supinus</i> , 698; <i>Tenageia</i> , 699. | | <i>Lamprothamnus zanguebaricus</i> . . | 268 |
| <i>Jungermannia colorata</i> | 180 | <i>Lansbergia</i> | 76, 126 |
| <i>Juniperus macrocarpa</i> , 670; <i>Oxycedrus</i> , 670; <i>phœnicea</i> , 670, var. <i>galbulis mollibus</i> , 671; <i>thurifera</i> , 671. | | — <i>cathartica</i> , 126; <i>juncifolia</i> , 126; <i>martinicensis</i> , 127; <i>monticola</i> , 126; <i>purgans</i> , 126; <i>spathata</i> , 127; <i>violacea</i> , 126. | |
| <i>Juno</i> , subgen. characterized | 123 | <i>Lapeyrousia</i> , characters of | 78, 154 |
| <i>Ké, Little</i> , fungi from | 35 | — <i>abyssinica</i> , 155; <i>anceps</i> , 156, var. <i>aculeata</i> , 156; <i>Bainesii</i> , 156; <i>breviflora</i> , 156; <i>corymbosa</i> , 155; <i>cyanescens</i> , 156; <i>divaricata</i> , 155; <i>erythrantha</i> , 155; <i>Fabricii</i> , 156; <i>fasciculata</i> , 155; <i>fissifolia</i> , 155; <i>fistulosa</i> , 155; <i>fragrans</i> , 156; <i>littoralis</i> , 156; <i>macrochlamys</i> , 155; <i>micrantha</i> , 156; <i>odoratissima</i> , 156; <i>purpureo-lutea</i> , 155; <i>silénoides</i> , 156. | |
| <i>Ker's 'Genera Iridearum'</i> | 61 | <i>Lapiedra gracilis</i> , <i>Baker, n. sp.</i> . | 678 |
| <i>Kerguelen's Land</i> , lichens from . . | 220 | <i>Laschia longipes</i> , <i>B.</i> | 43 |
| | | — <i>tremellosa</i> | 50 |
| | | <i>Lasianthus africanus</i> | 273 |
| | | <i>Latania Commersonii</i> , 10; <i>Lodigesii</i> , 10, 11; <i>Verschaffeltii</i> , 10. | |
| | | <i>Lathyrus annuus</i> , 441; <i>aphaca</i> , 440; <i>articulatus</i> , 441; <i>cærulescens</i> , 440; <i>cicera</i> , 440; <i>clymenum</i> , 441; <i>fissus</i> , 440; <i>latifolius</i> , 440; <i>ochrus</i> , 441; <i>sativus</i> , var. <i>amphicarpus</i> , 440; <i>sphaericus</i> , 440; <i>tingitanus</i> , 441. | |
| | | <i>Laurencia flagellifera</i> , 6; <i>obtusâ</i> , 6; <i>papillosa</i> , 6. | |
| | | <i>Laurentia Michellii</i> | 552 |
| | | <i>Laurus nobilis</i> | 655 |
| | | <i>Lavandula dentata</i> , 608, var. <i>unnamed</i> , 608; <i>multifida</i> , 608, var. <i>abrotanoides</i> , <i>Ball</i> , 608, var. <i>intermedia</i> , <i>Ball</i> , 608; <i>pedun-</i> | |

- | | Page | | Page |
|---|------|--|---------|
| culata, 608; Stœchas, 607; tenuisecta, 609. | | 192; tabularis, 192; xanthocarpa, 191. | |
| Lavatera cretica, 376, var. acutiloba, <i>Ball</i> , 377; maritima, 376; Olbia, 376, var. hispida, 376; trimestris, 376, var. malvæformis, <i>Ball</i> , 376. | | <i>Lemna gibba</i> | 685 |
| <i>Lecanora Acunhana</i> , <i>Nyl.</i> , <i>n. sp.</i> , 222; bermudensis, 215; chilatronella, 215; cyphelliformis, <i>Cromb. sp. n.</i> , 220; flavorubens, 212; gelida (<i>Placopsis</i>), <i>L.</i> , 222, ff. lateritia, <i>Nyl.</i> , et vitellina, <i>Bab.</i> , 220; glaucomodes, 215; granifera, var. leucotropa, 226; lepidoplacum, <i>Nyl. sp. n.</i> , 212; macrophthalma, 220; melanophthalma, 218; parella, 218; pyracea, 212; recubans, 219; saxicola (<i>Squamaria</i>), 218; sublegans (<i>Placodium</i>), <i>Nyl. sp. n.</i> , 218; subfusca sylvestris, 215, var. subgranulata, <i>Nyl.</i> , 222; Tencriffensis, 218; vinentina, <i>Nyl. sp. n.</i> , 212. | | <i>Lentinus abnormis</i> , <i>B.</i> , 45; axilis, 45; fasciatus, 38; Hookerianus, 42; hygrometricus, <i>B.</i> , 49; Levcklei, 42, 45; strigosus, 42; villosus, 49. | |
| <i>Lecchlora</i> , subgen. characterized . | 121 | <i>Lenzites applanata</i> | 43, 48 |
| <i>Lecidea atrobrunnea</i> , 219; canescens, 213; disjungenda, 221; endoleuca, 218; endomelæna, <i>Leight.</i> , <i>n. sp.</i> , 2; epipasta, <i>Strn. sp. n.</i> , 213; epiplacodia, 213; euporiza, <i>Strn. sp. n.</i> , 216; Fernandezii, <i>Cromb. n. sp.</i> , 223; fusciorubescens, 216; geographica, 219, 221; glaucomeria, <i>Leight. n. sp.</i> , 2; heteroloba, <i>Cromb. sp. n.</i> , 214; imponens, <i>Leight. n. sp.</i> , 2; impressula, <i>Leight. n. sp.</i> , 7; insperata, 216; leptocheloides, <i>Nyl. sp. n.</i> , 225; leucoblephara, 225; myriocarpa, 214; natalensis, 216; parellaria, <i>Leight. n. sp.</i> , 2; revertens, <i>Strn. sp. n.</i> , 216; rissolea, <i>Leight. n. sp.</i> , 2; semiusta, <i>Strn. n. sp.</i> , 216; stephanodes, <i>Strn. n. sp.</i> , 221; superjecta, <i>Nyl. sp. n.</i> , 221; taitensis, 225; tephrizans, <i>Leight. n. sp.</i> , 2; thyrsodes, <i>Strn. sp. n.</i> , 214. | | <i>Leontodon autumnalis</i> , 543; helminthioides, 544; hispidulus, 545; maroccanus, 544; Rothii, 543, var. major, 543; Salzmanni, 545; tingitanus, 544; tuberosus, 546. | |
| <i>Leersia hexandra</i> | 706 | <i>Leperoma ochroleuca</i> | 191 |
| <i>Leibleinia moniliformis</i> , <i>Reinsch, n. sp.</i> | 235 | <i>Lepidium latifolium</i> , 332; nebroidense, 331; sativum, 331. | |
| Leighton, Rev. W. A., New British Lichens (abstract) | 2 | <i>Lepidozia truncatella</i> | 191 |
| <i>Lejeunia capensis</i> , 192; Eckloniana, 192; hamatifolia, 192; minutissima, 192; rufescens, | | <i>Lepipogon obovatum</i> | 275 |
| | | <i>Leptactina densiflora</i> , 266; heinsoides, 266; Mannii, 266; senegambica, 266. | |
| | | <i>Leptodon Smithii</i> | 738 |
| | | <i>Leptogium azurcum</i> , 223; cyanizum, 227; diaphanum, 224; laceratulum, <i>Nyl. sp. n.</i> , 224; tremelloides, 219. | |
| | | <i>Leptothrix lutea</i> | 7 |
| | | <i>Lepturus filiformis</i> | 733 |
| | | <i>Leucodon sciuroides</i> | 738 |
| | | <i>Leucojum autumnale</i> , 678; tingitanum, <i>Baker, n. sp.</i> , 678; trichophyllum, 678. | |
| | | <i>Leuzea conifera</i> | 526 |
| | | <i>Leyssera capillifolia</i> | 500 |
| | | <i>Liagora viscida</i> | 7 |
| | | <i>Libertia</i> , characters of | 78, 152 |
| | | — cærulescens, 153; elegans, 153; formosa, 153, var. crassa, 153; grandiflora, 153; ixioides, 152; paniculata, 153; pulchella, 152; tricocca, 152. | |
| | | Lichens of 'Challenger' Exped., conspectus of | 228 |
| | | —, New British | 2 |
| | | <i>Limnodietyon</i> , spec. | 244 |
| | | <i>Limoniastrum monopetalum</i> | 561 |
| | | <i>Linaria amethystea</i> , 593; arvensis, var. β , simplex, 591; bipartita, 491; Broussonnetii, 593; galioides, 590, var. pseudosupina, <i>Ball</i> , 590; glabrescens, 594, var. γ , minor, 594; græca, 586; heterophylla, 588, var. tingitana, <i>Ball</i> , 589; latifolia, 587; lurida, 592; marginata, 592; maroccana, 589; Munbyana, 593; rubrifolia, 594; sagittata, 587; | |

- | | Page | | Page |
|--|--------|--|---------|
| spartea, 590; spuria, 586; Tournefortii, var. triphylla, 587; ventricosa, 587; viscosa, 589. | | pera, 361; diurna, 362; macrocarpa, 362. | |
| Lindigina renifolia, <i>Mitt., n. sp.</i> (Pl. V.) | 190 | Lycium afrum, 581; mediterraneum, 582; vulgare, 581. | |
| — scariosa (Pl. V.) | 189 | Lycopodium cernuum, 34, 205; clavatum, 206; dichotomum, 205; linifolium, 37; phlegmaria, 34. | |
| Lindsaya concinna, 27; davalloides, 27; flabellata, 27; lobata, 27; madagascariensis, <i>Baker, n. sp.</i> , 198, var. davallioides, <i>Baker</i> , 198, var. intermedia, <i>Baker</i> , 198; trapeziformis, var. falcata, 34. | | Lycopus europæus | 610 |
| Linum angustifolium, 380; bicolor, 380; corymbiferum, var. ? 379, var. Mumbyanum, 379; gallicum, 378, perenne, 380; setaceum, 379; strictum, 378; suffruticosum, 380; tenue, 379. | | Lygeum spartum | 706 |
| Listrostachys aphrodite | 11 | Lygodium dichotomum, 34; scandens, 34; venustum, 37. | |
| Lithospermum apulum, 574; arvense, 573; fruticosum, 574; in-crassatum, 573; microspermum, 574. | | Lynch, R. I., note on the Blimbing (<i>Averrhoa Bilimbi</i> , Linn.) | 231 |
| Lithothamnion mamillare | 7 | — on disarticulation of branches | 180 |
| Lobelia vagans, <i>Balf. fil. sp. n.</i> 11, 16 | | Lyngbya æruginosa, var. africana, 234; majuscula, 7. | |
| Lœflingia hispanica | 371 | Lythrum flexuosum, 457, 739; hys-sopifolium, 457, var. thymifolium, 457; salicaria, var. tomentosum, 458; Salzmanni, 457. | |
| Lolium perenne, 730, var. italicum, 730, var. rigidum, 730. | | Macrochloa tenacissima | 711 |
| Lomaria areolaris, <i>Harrington, n. sp.</i> 28 | | Macrosphyra longistyla | 267 |
| — biformis | 199 | Madagascar, ferns of | 197 |
| Lomenia borbonica . . . (ftnote) | 157 | Magydaris panacina | 472 |
| Lonchitis pubescens | 198 | — tomentosa | 472 |
| Lonicera biflora, 482; etrusca, 481; implexa, 482; Periclymenum, var. hispida, 482. | | Malamon (Philippines), fungi from | 45 |
| Lophocolea bidentata | 188 | Malanipa (Philippines), fungi from | 45 |
| — semiteres | 188 | Malcolmia lacera, 322, var. Broussonnetii, 323; litorea, 322; nana, 323. | |
| Lord Howe's Island, fungi from . | 53 | Malope malacoides, 375; trifida, 375. | |
| Lotononis maroccana, <i>Ball</i> (Pl. XV.) | 394 | Malva hispanica, 377; nicæensis, 378; parviflora, 378; rotundifolia, 377, var. intermedia, <i>Ball</i> , 377; sylvestris, 377, var. mauritiana, 377. | |
| Lotus arenarius, 422, var. Webbii, <i>Ball</i> , 423; creticus, 423; commutatus, 423; conjugatus, 425; cytisoides, 423; edulis, 422; hispidus, 424, 739; lamprocarpus, 424; maroccanus, 424; ornithopodioides, 422; parviflorus, 424; siliquosus, 425; tetragonolobus, 425; tingitanus, 425. | | Mandragora officinarum | 582 |
| Ludia sessiliflora, heterophylly of, | 23, 24 | Marasmius aleurocephalus, <i>B.</i> , 42; aruensis, <i>B.</i> , 41; direptus, <i>B.</i> , 42; fulviceps, 49; hispidulus, <i>B.</i> , 49; opacus, 38; primulinus, <i>B.</i> , 38. | |
| Lunularia cruciata | 194 | Marattia fraxinea, <i>Sm.</i> (Pl. I.), 2; on the rootstock of, 3; propagation of, 4. | |
| Lupinus angustifolius, 395; hirsutus, 395; luteus, 395; pilosus, 395. | | Marica, characters of | 77, 149 |
| Luzula Forsteri | 697 | — brachypus, 150; cærulea, 149; glauca, 149; gracilis, 150; humilis, 150; longifolia, 150; lutea, 150; Northiana, 149; Sabini, 149. | |
| Lychnis Cœli Rosa, 361. var. as- | | Marion Island, lichens from | 221 |
| | | Marocco Flora, <i>J. Ball</i> on | 281 |
| | | Marrubium Alysson, 623; echinatum, 623; vulgare, 623, var. lanatum, 623. | |
| | | Mathurina penduliflora, <i>Balf. fil.</i> . | 11 |
| | | — —, heterophylly of | 24 |

- | | Page | | Page |
|---|------|--|---------|
| <i>Matricaria Chamomilla</i> , 511; glabra, 511; fuscata, 512; maroccana, 512; pubescens, 512. | | Mitten, W., list of <i>Hepaticæ</i> coll. by the Rev. A. E. Eaton at the Cape of Good Hope Hope. . . | 187 |
| <i>Matthiola parviflora</i> , 317; <i>tricuspidata</i> , 316; <i>tristis</i> , 316. | | <i>Mnium undulatum</i> | 738 |
| <i>Medicago denticulata</i> , 415; <i>Gerardi</i> , 414; <i>Helix</i> , 412, var. <i>lævis</i> , 412, var. <i>spinulosa</i> , 412; <i>intertexta</i> , 416; <i>laciniata</i> , 416; <i>lappacea</i> , var. <i>pentacycla</i> , 415; <i>littoralis</i> , 413, var. <i>α</i> , 413, var. <i>longeaculcata</i> , 413; <i>lupulina</i> , 411; <i>maculata</i> , 416; <i>marina</i> , 413; <i>minima</i> , 416; <i>orbicularis</i> , 412; <i>sphærocarpa</i> , 414; <i>suffruticosa</i> , 412; <i>tornata</i> , 413; <i>tribuloides</i> , 413, var. <i>longeaculcata</i> , <i>Ball</i> , 413; <i>turbinata</i> , var. <i>aculcata</i> , 414, var. <i>muricata</i> , 414. | | <i>Mohria caffrorum</i> | 205 |
| <i>Melanotheca aciculifera</i> | 220 | <i>Monanthes atlantica</i> | 454 |
| <i>Melasphærule</i> , characters of . 78, 167 | | <i>Monerma cylindrica</i> | 733 |
| — <i>graminea</i> | 167 | <i>Monogramme Junghuhnii</i> | 33 |
| <i>Melica ciliata</i> , 722, var. <i>major</i> , 772; <i>Cupani</i> , 722, var. <i>minor</i> , 723; <i>minuta</i> , 722, var. <i>latifolia</i> , 722. | | <i>Montbretia</i> , characters of . 79, 167 | |
| <i>Melilotus indica</i> , 416; <i>sulcata</i> , 417, var. <i>major</i> , 417. | | — <i>capensis</i> , 168; <i>dubia</i> , 169; <i>flava</i> , 169; <i>kamisbergensis</i> , 169; <i>lacerata</i> , 168, var. <i>pectinata</i> , 168; <i>laxifolia</i> , 169; <i>lincata</i> , 169; <i>pallida</i> , 168; <i>pauciflora</i> , 168; <i>roscæ</i> , 169; <i>scurigera</i> , 169; <i>striata</i> , 168; <i>strictifolia</i> , 169; <i>viridis</i> , 169. | |
| <i>Melissa officinalis</i> , <i>L.</i> , var. | 740 | <i>Montia fontana</i> | 372 |
| <i>Meniscum triphyllum</i> | 33 | <i>Moræa</i> , characters of | 77, 129 |
| <i>Mentha Pulegium</i> , 609, 740, var. <i>villosa</i> , 610; <i>rotundifolia</i> , 609; <i>sylvestris</i> , 609; <i>Timija</i> , 609. | | — <i>andongensis</i> , 131; <i>angusta</i> , 130; <i>barbigera</i> , 130; <i>bituminosa</i> , 131; <i>Burchellii</i> , 132; <i>candelabrum</i> , 131; <i>ciliata</i> , 131; <i>crispa</i> , 131, var. <i>undulata</i> , 131; <i>diversifolia</i> , 130; <i>edulis</i> , 130, var. <i>longifolia</i> , 131, var. <i>odora</i> , 131; <i>fimbriata</i> , 130; <i>galaxioides</i> , 130; <i>gigantea</i> , 131; <i>glutinosa</i> , 131; <i>gracilis</i> , 130; <i>juncæa</i> , 130; <i>longiflora</i> , 132; <i>lurida</i> , 130; <i>minuta</i> , 130; <i>papilionacea</i> , 129; <i>plumaria</i> , 130; <i>polyanthos</i> , 130; <i>polyphylla</i> , 130; <i>polystacha</i> , 131; <i>ramosa</i> , 131; <i>setacea</i> , 130; <i>Sisyrinchium</i> , 132, var. <i>ægyptia</i> , 132, var. <i>fugax</i> , 132, var. <i>minor</i> , 132, var. <i>Samaritanii</i> , 132; <i>spathacea</i> , 131; <i>spithamæa</i> , 131; <i>textilis</i> , 130; <i>tricolor</i> , 130; <i>tristis</i> , 131; <i>viscaria</i> , 131; <i>Welwitschii</i> , 130; <i>zambesiaca</i> , 130; | |
| <i>Mercurialis annua</i> , 664, var. <i>ambigua</i> , 664, var. <i>serratifolia</i> , 664. | | <i>Morelia senegalensis</i> | 267 |
| <i>Merulius corium</i> | 53 | <i>Morinda citrifolia</i> | 271 |
| <i>Mesembryanthemum nodiflorum</i> . 460 | | — <i>longiflora</i> | 271 |
| — <i>crystallinum</i> | 460 | <i>Morindeæ</i> , tribe of <i>Rubiaceæ</i> . . | 271 |
| <i>Mesogloia virescens</i> | 6 | <i>Morphixia</i> , characters of . . 74, 96 | |
| <i>Metzgeria furcata</i> | 194 | — <i>aulica</i> , 97; <i>capillaris</i> , 97, var. <i>incarnata</i> , 97, var. <i>lancea</i> , 97; <i>Cooperi</i> , 98; <i>juncifolia</i> , 98; <i>linearis</i> , 97; <i>nervosa</i> , 98; <i>odorata</i> , 97; <i>paniculata</i> , 97, var. <i>rochensis</i> , 97, var. <i>tenuiflora</i> , 97; <i>trichorhiza</i> , 97. | |
| <i>Meum atlanticum</i> | 474 | <i>Musa troglodytarum</i> , use of spathe of | 59 |
| <i>Micranthus</i> , subgen., characters of 159 | | <i>Muscari comosum</i> | 685 |
| <i>Microcala filiformis</i> | 567 | — <i>racemosum</i> | 685 |
| <i>Microdictyon Agardhianum</i> | 7 | <i>Mussaenda Afzelii</i> , 265; <i>arcuata</i> , 265; <i>elegans</i> , 265; <i>erythrophylla</i> , | |
| <i>Micromeria microphylla</i> | 613 | | |
| — —, var. <i>villosa</i> | 613 | | |
| <i>Micropogon</i> , subgen., characterized 124 | | | |
| <i>Micropus bombycinus</i> | 496 | | |
| — <i>supinus</i> | 496 | | |
| <i>Microrhynchus nudicaulis</i> | 550 | | |
| — <i>spinosus</i> | 550 | | |
| <i>Microthamnion exiguum</i> , <i>Reinsch</i> , <i>sp. n.</i> | 245 | | |
| <i>Mimosa pudica</i> , movements of . . . | 231 | | |
| <i>Mitracarpum scabrum</i> | 274 | | |
| <i>Mitragyne africana</i> | 263 | | |
| — <i>macrophylla</i> | 263 | | |
| <i>Mitriostigma</i> ? <i>Barteri</i> | 267 | | |
| — <i>subpunctatum</i> | 267 | | |

- | | Page | | Page |
|---|---------|--|------|
| 265; grandiflora, 265; heinsioi- | | Nolletia chrysocomoides | 495 |
| des, 265; lsertiana, 265; luteola, | | Nonnea nigricans | 572 |
| 265; platyphylla, 265; polita, 265; | | — phaneranthera | 572 |
| stenocarpa, 265; tenuiflora, 265. | | Nostoc commune | 7 |
| Mussændeæ, tribe of Rubiaceæ . . . | 265 | Nothoehlaena doradilla, 37; ferru- | |
| Myosotis hispida, 573; lingulata, | | ginea, 37; hirsuta, 33; nivea, 37; | |
| var. foliolosa, 572; stricta, 573; | | sinuata, 37; Streetiæ, <i>Baker</i> , | |
| sylvatica, 572, var. alpestris, | | <i>n. sp.</i> , 204; tenera, 37; villeda, 736. | |
| 572; versicolor, 573. | | Notoceras canariense | 319 |
| Myriophyllum spicatum | 456 | Notoscyphus variifolius, <i>Mitt.</i> | |
| Myrtus communis | 456 | <i>n. sp.</i> (Pl. IV.) | 188 |
| Narcissus elegans, var. intermedia, | | Numerical increase of parts of plants | 206 |
| 680; papyraceus, 680; scrotinus, | | | |
| 680; Tazetta, 680; viridiflorus, | | Obione portulacoides | 645 |
| 681. | | Octodon filifolium | 274 |
| Nasturtium atlanticum, <i>Ball</i> | | — setosum | 274 |
| (Pl. IX.), 317; officinale, 317. | | Odontidium | 6 |
| Naucleæ, tribe of Rubiaceæ . . . | 263 | Odontospermum aquaticum, 502, | |
| Nemastylus, characters of . . . | 74, 102 | 739; imbricatum, 503; mariti- | |
| — acutus, 103; cælestinus, 103; | | num, 502, var. microphylla, | |
| purpureus, 103. | | 502; odorum, 502. | |
| Nepeta Apulei, 619; atlantica, 619; | | Odontostigma, a section of <i>Crocus</i> , | |
| multibracteata, 619. | | defined | 83 |
| Nephrodium anateinophlebium, | | Odontotrema majus, <i>Leight. n. sp.</i> | 2 |
| <i>Baker, n. sp.</i> , 202; <i>Bakeri, Har-</i> | | Ædogonium fonticola | 7 |
| <i>rington, n. sp.</i> , 29; <i>Buchanani</i> , | | Enanthe apiifolia, 473; callosa, | |
| 203; eicutarium, 30; eoncinn- | | 473; globulosa, 473. | |
| um, 36; costulare (<i>Eunephro-</i> | | Oldenlandia abyssiniea, 264; <i>Bo-</i> | |
| dium), <i>Baker, n. sp.</i> , 203; de- | | jeri, 264; calfra, 264; cæspitosa, | |
| curso-pinnatum, 29; <i>Eatoni</i> , | | 264; capensis, 265; corymbosa, | |
| <i>Baker, var. formosanum, Har-</i> | | 265; decumbens, 264; effusa, | |
| <i>rington, 29; effusum, 36;</i> | | 264; flosculosa, 264; globosa, | |
| <i>Filix-mas, 29, 36; immersum,</i> | | 264; grandiflora, 264; hedytoi- | |
| 29; latifolium, 30; longicuspe, | | des, 265; Heynei, 264; lanci- | |
| <i>Baker, n. sp.</i> , 202; <i>Luersseni</i> , | | folia, 265; lasiocarpa, 264; ma- | |
| (<i>Lastrea</i>) <i>Harrington, n. sp.</i> , | | crophylla, 265; monanthos, 264; | |
| 29; molle, 30; odoratum, 29; | | noctiflora, 264; obtusiloba, 264; | |
| patens, var. madagascariense, | | parviflora, 264; peltospermum, | |
| <i>Baker, 202; pennigerum, 203;</i> | | 264; rigida, 264; <i>Schimperi</i> , | |
| sophoroides, 30; subcrenulatum, | | 264; senegalensis, 264; <i>Sieberi</i> , | |
| <i>Baker, n. sp.</i> , 202; subpedatum, | | 9; strumosa, 264; tenuissima, | |
| <i>Harrington, n. sp.</i> , 30; tricho- | | 265; trinervia, 265; virgata, 264; | |
| phlebium, <i>Baker, n. sp.</i> , 201; | | wauensis, 265. | |
| truncatum, 30, 203. | | Olea europæa | 565 |
| Nephrolepis acuta, 30; cordifolia, | | Olpidium, spec. ? | 246 |
| 30, 201; floecigera, 30; ramosa, | | Oncoecylus, subgen. characterized . | 142 |
| 30. | | Onobrychis Crista-Galli, 436; erio- | |
| Nerium Oleander | 565 | phora, 436; sativa, 435, var. | |
| Nesogenes, 9; decumbens, <i>Balf.</i> | | pseudosupina, 435. | |
| <i>fil. sp. nov.</i> | 11, 19 | Ononis alopecuroides, 410; angus- | |
| Neuberia, subgen., characters of . | 158 | tissima, 406; antiquorum, 409; | |
| Neuropogon melaxanthus, 222, var. | | atlantica, <i>Ball</i> (Pl. XVII.), 408; | |
| ciliata, 220; <i>Taylori</i> , 220. | | cintrana, 407; <i>Cossoniana</i> , 410; | |
| Nigella arvensis, 308, var. <i>Cosso-</i> | | filicaulis, 409; laxiflora, 409; | |
| niana, <i>Ball</i> , 308; <i>Damascena</i> , | | <i>Maweana</i> , 407; minutissima, | |
| 309; <i>hispanica</i> , 309, 739. | | 411; mitissima, 410; <i>Natrix</i> , | |
| Nivenia, characters of | 75, 109 | 405, var. tomentosa, 405, var. mi- | |
| — corymbosa, 109; fruticosa, 109. | | crophylla, 405; pendula, 408, | |
| | | var. <i>Broussonnetii</i> , 409; poly- | |

- | | Page | | Page |
|--|------|--|------|
| phylla, <i>Ball</i> (Pl. XVI.), 407 ;
porrigens, 406 ; pubescens, 407 ;
reclinata, 409 ; serrata, 410, var.
minor, 410 ; sicula, 407 ; <i>Tour-</i>
<i>nefortii</i> , 410 ; viscosa, 739, var.
fruticescens, <i>Ball</i> , 406. | | mosus, 267 ; gracilis, 267 ; ma-
crophyllus, 267 ; ? platystylis,
267 ; querimbensis, 267 ; rubriflo-
rus, 267 ; <i>Smithii</i> , 267 ; speciosus,
267 ; ? sulcatus, 267 ; tubiflorus,
267 ; unilocularis, 267. | |
| <i>Onopordon illyricum</i> , 523 ; <i>macra-</i>
<i>canthum</i> , 523 ; <i>Sibthorpium</i> ,
523. | | <i>Padina Pavonia</i> 6 | |
| <i>Onychium japonicum</i> 27 | | <i>Pædericæ</i> , tribe of <i>Rubiaceæ</i> . . 274 | |
| <i>Opographa dialeuca</i> , 214 ; subvul-
gata, 226 ; undulata, <i>Strn. sp. n.</i> ,
214. | | <i>Pallenis spinosa</i> 503 | |
| <i>Ophioglossum lusitanicum</i> , 737 ;
pendulum, 34 ; rhizome of, 5. | | <i>Palmella Brebissonii</i> 237 | |
| <i>Ophrys apifera</i> , 673, 740 ; <i>bomby-</i>
<i>liflora</i> , 673 ; lutea, 673 ; <i>Scolopax</i> ,
673 ; <i>Speculum</i> , 673 ; <i>tenthredi-</i>
<i>nifera</i> , 674. | | <i>Palmogloea chilanydospora</i> , var.
minor 240 | |
| <i>Opuntia vulgaris</i> 460 | | — <i>macrococca</i> 239 | |
| Orchidaceæ, paucity of, in <i>Rodri-</i>
<i>guez</i> 10 | | <i>Pancratium maritimum</i> 679 | |
| <i>Orchis latifolia</i> , 672, var. <i>Durandii</i> ,
<i>Ball</i> , 672 ; <i>longieruris</i> , 671 ;
mascula, 672 ; <i>papilionacea</i> , 671 ;
<i>tridentata</i> , var. <i>acuminata</i> , 671. | | <i>Pandanus heterocarpus</i> , <i>Balf. fil.</i>
<i>sp. n.</i> , 11, 22 ; <i>tenuifolius</i> , <i>Balf.</i>
<i>fil. sp. n.</i> , 11, 22 ; <i>utilis</i> , 22. | |
| <i>Origanum</i> , 740 ; <i>compactum</i> , 610. | | <i>Panicum Crus-Galli</i> , 709 ; <i>repens</i> 709 | |
| <i>Ornithogalum arabicum</i> , 689, 740 ;
comosum, var. <i>atlanticum</i> , 688 ;
narbonense, 689 ; <i>orthophyllum</i> ,
688 ; <i>sessiliflorum</i> , 689 ; <i>tenui-</i>
<i>folium</i> , 688 ; <i>umbellatum</i> , 688 ;
<i>unifolium</i> , 689. | | <i>Pannaria dichroa</i> , 220 ; <i>nigro-cincta</i> ,
225 ; <i>pannosa</i> , 225 ; <i>rubiginosa</i> ,
225. | |
| <i>Ornithopus compressus</i> , 426 ; <i>ebrac-</i>
<i>teatus</i> , 426 ; <i>isthmocarpus</i> , 426. | | <i>Papaver dubium</i> , 311 ; <i>hybridum</i> ,
311 ; <i>Rhœas</i> , 312 ; <i>rupifragum</i> ,
var. <i>atlanticum</i> , <i>Ball</i> , 313 ; <i>som-</i>
<i>niferum</i> , var. <i>setigerum</i> , <i>Ball</i> ,
312 ; <i>tenue</i> , 312. | |
| <i>Orobanche barbata</i> , 606 ; <i>densi-</i>
<i>flora</i> , 604 ; <i>foetida</i> , 604, var. <i>co-</i>
<i>mosa</i> , <i>Ball</i> , 604 ; <i>Hookeriana</i> ,
<i>Ball, n. sp.</i> , 605 ; <i>pruinosa</i> , 605,
var. <i>speciosa</i> , 605 ; <i>reticulata</i> ,
604 ; <i>tetuanensis</i> , 606 ; <i>variegata</i> ,
604. | | <i>Paramatta</i> , fungi from <i>Pennant</i>
<i>Hills</i> 38 | |
| <i>Orthotrichum cupulatum</i> 738 | | <i>Parietaria mauritanica</i> , var. <i>diffusa</i> ,
666 ; <i>officinalis</i> , 665. | |
| <i>Orthrosanthus</i> , characters of . 76, 112 | | <i>Parmelia glabra</i> , 218 ; <i>latissima</i> ,
224 ; <i>perforata</i> , 212, 218, 222 ;
<i>perlata</i> , 215, 218, 222 ; <i>persul-</i>
<i>phurata</i> , 219 ; <i>revolutella</i> , <i>Nyl.</i>
<i>sp. n.</i> , 222 ; <i>subaurulenta</i> , 228 ;
<i>tinctorum</i> , 211, 215. | |
| — <i>chimboracensis</i> , 113, var. <i>gla-</i>
<i>dioloides</i> , 113 ; <i>gramineus</i> , 113 ;
<i>laxus</i> , 112 ; <i>Muelleri</i> , 112 ; <i>mul-</i>
<i>tiflorus</i> , 112 ; <i>polystachyus</i> , 113 ;
<i>spicatus</i> , 113. | | <i>Paronychia argentea</i> , 640, var. <i>mau-</i>
<i>ritanica</i> , 640, var. <i>velutina</i> , 641 ;
<i>capitata</i> , 641, var. <i>atlantica</i> , <i>Ball</i> ,
641 ; <i>cymosa</i> , 640 ; <i>echinata</i> , 640 ;
<i>macrosepala</i> , 641, var. <i>queriodes</i> ,
<i>Ball</i> , 642. | |
| <i>Osmunda javanica</i> 34 | | <i>Passerina arvensis</i> , 653 ; <i>canescens</i> ,
654 ; <i>hirsuta</i> , 654 ; <i>villosa</i> , 654 ;
<i>virgata</i> , 654, var. <i>Broussonnetii</i> ,
<i>Ball</i> , 654. | |
| — <i>regalis</i> 737 | | <i>Patersonia</i> , characters of . . 77, 150 | |
| <i>Osyris alba</i> 655 | | — <i>babianoides</i> , 152 ; <i>Drummon-</i>
<i>dii</i> , 152 ; <i>glabrata</i> , 151 ; <i>glauca</i> ,
150 ; <i>graminea</i> , 152 ; <i>inæqualis</i> ,
152 ; <i>junceæ</i> , 151, var. <i>elongata</i> ,
151 ; <i>lanata</i> , 151, var. <i>pannosa</i> ,
151 ; <i>limbata</i> , 151 ; <i>longifolia</i> , 151 ;
<i>longiscapa</i> , 151 ; <i>macrantha</i> , 151 ;
<i>Maxwellii</i> , 151 ; <i>occidentalis</i> , 151,
var. <i>angustifolia</i> , 151, var. <i>erio-</i>
<i>stephana</i> , 151, var. <i>latifolia</i> , 151 ;
<i>pygmæa</i> , 151 ; <i>rudis</i> , 151 ; <i>scri-</i> | |
| — <i>lanceolata</i> 655 | | | |
| <i>Otomeria dilatata</i> , 264 ; <i>guineen-</i>
<i>sis</i> , 264 ; <i>lanccolata</i> , 264 ; <i>ma-</i>
<i>diensis</i> , 264. | | | |
| <i>Oxalis corniculata</i> , 388 ; <i>sericea</i> ,
388. | | | |
| <i>Oxyanthus breviflorus</i> , 267 ; <i>for-</i> | | | |

- | | Page | | Page |
|--|--------|--|---------|
| cea, 151, var. latifolia, 151; umbrosa, 151; xanthina, 151. | | Phalocallis, characters of . . . | 76, 127 |
| Patersoniæ, tribe of Irids, characters of | 77 | — plumbea | 128 |
| Pauridiantha canthiifolia . . . | 265 | Phelipæa cærulea, 603; lutea, 603; Muteli, 603, var. nana, 603; Schultzi, 603; tinctoria, 603. | |
| Pavetta abyssinica, 271; Baconia, 271; bidentata, 270; brachyelyx, 270; canescens, 271; erebriifolia, 270; dolichosepala, 271; gardiniaefolia, 271; genipæfolia, 271; glaucescens, 270; gracilipes, 271; gracilis, 271; hispida, 271; Hookeriana, 271; macrosepala, 270; Mannii, 270; mollis, 271; monticola, 270; neurocarpa, 271; Oliveriana, 271; owariensis, 270; puberula, 270; rigida, 271; subcana, 271; ternifolia, 271. | | Phillyrea angustifolia, 564; latifolia, 564; media, 564. | |
| Paxillus Mülleri | 38 | Phlomis crinita | 629 |
| Pedistrum pertusum | 239 | — Herba-Venti | 629 |
| Peganum Harmala | 390 | Phoenix dactylifera | 10, 683 |
| Pellæa angulosa, 199; cordata, 35; geraniifolia, 35; marginata, 35; pectiniformis, 199; ternifolia, 35. | | Phragmitis communis | 716 |
| Peltigera canina, f. membranacea, 228; polydactyla, f. hymenina, 222. | | Phyllanthus, 9; Conami, 183; detachment of branches of, 182; dumetosus, 11; glaucescens (Pl. III.), 182; juglandifolius (Pl. III.), 182. | |
| Penium Brebissonii | 240 | Phyllostachys nigra, 1; uses of a species of, 1. | |
| — rupestre | 240 | Physcia adscensionis, 212; flavicans, 212; galactophylla, 228; Leana, 215; leucomela, 212; obsessa, 215; podocarpa, 225; podocarpoides, <i>Nyl. sp. n.</i> , 225; speciosa, 228. | |
| Pennisetum eiliare | 710 | Pieridium tingitanum, 547, var. maritimum, 547, var. maroccanum, <i>Ball</i> , 547; vulgare, 547. | |
| Penny-cress, perfoliate, distribution of, in Britain | 183 | Pieris albida, 536; pilosa, 536. | |
| Pentaloneha humilis | 266 | Pilea Balfourii, 11, 20; euneiformis, 20. | |
| Pentansia Schweinfurthii . . . | 268 | Pimpinella bubonoides, var. villosa, 471; tragium, 471. | |
| Pentas arvensis, 264; carnea, 264; occidentalis, 264; parvifolia, <i>Hiern, sp. n.</i> (Pl. VII.), 262, 264, 275; purpurea, 264; Schimperiana, 264. | | Pinguicula lusitanica | 606 |
| Peperomia, 9; elliptica, 11, 21; hirta, <i>Balf. fil. sp. n.</i> , 11, 20; reticulata, <i>Balf. fil. sp. n.</i> , 21; rodriguensiana, <i>Balf. fil. sp. n.</i> , 11, 21. | | Pinus Cedrus, var. atlantica, 670; halepensis, 669. | |
| Peplis nummulariæfolia | 456 | Piptatherum cæruleseens, 710; miliaceum, 710. | |
| Periploca lævigata | 565 | Pisonia lanceolata, 20; ovalifolia, 20; viscosa, <i>Balf. fil. sp. nov.</i> , 11, 19. | |
| Pertusaria aleianta, <i>Nyl. sp. n.</i> , 213; multipuncta, 228; papillata, 215; pustulata, 216; velata, 225. | | Pistacia atlantica | 392 |
| Peucedanum Munbyi | 475 | — Lentiscus | 393 |
| Peziza Hindsii | 44, 50 | Pisum elatius | 442 |
| Phæcasium pulehrum | 539 | Plagiochila javanica | 187 |
| Phagnalon atlanticum, 498; rupestre, 498, var. calycinum, <i>Ball</i> , 498; saxatile, 498, var. intermedium, 498. | | — sarmentosa | 187 |
| Phalaris brachystachys, 707; cærulescens, 708; canariensis, 707; minor, 707; nodosa, 707; paradoxia, 708. | | Plantago albicans, 635, var. humilis, <i>Ball</i> , 635; amplexicaulis, 636; Coronopus, 637, var. Cupani, 637, var. maroccana, <i>Ball</i> , 637; decumbens, 635; Lagopus, 636, var. lusitanica, 636; macrorhiza, 636; major, 635; manritanica, 637; Psyllium, 637; serraria, 636; stricta, 638. | |
| | | Plants, numerical increase of parts of | 206 |
| | | Platanthera diphylla | 673 |
| | | Platyterium madagascariense . . . | 205 |

- | | Page | | Page |
|--|----------|--|------|
| <i>Platygrapha stenogramma</i> , <i>Nyl.</i> | | matum, 33; percutsum, 37; per- | |
| <i>sp. n.</i> | 226 | ludens, <i>Baker.</i> , <i>n. sp.</i> , 204; per- | |
| <i>Platysterigma</i> , subgen. characterized | 107 | sicariæfolium, 37; piloselloides, | |
| <i>Pleiophylly</i> in plants | 207, 209 | <i>L.</i> , var. β , <i>H. & B.</i> , 36, var. γ , | |
| <i>Pleiotaxy</i> in plants | 207, 208 | <i>H. & B.</i> , 36; plebeium, 36; | |
| <i>Plication</i> in plants | 207, 209 | plumula, 36; pteropus, 33; | |
| <i>Plumbago europæa</i> | 561 | Schenkii (<i>Eupolypodium</i>), <i>Harr-</i> | |
| <i>Poa annua</i> , 720; <i>bulbosa</i> , 720; | | ington, <i>n. sp.</i> , 31; sphenodes, 36; | |
| <i>compressa</i> , 720; <i>pratensis</i> , 720; | | Steerei (<i>Phymatodes</i>), <i>Harring-</i> | |
| <i>trivialis</i> , 720. | | ton, <i>n. sp.</i> , 32; <i>synsorum</i> (<i>Eupolypodium</i>), <i>Baker</i> , <i>n. sp.</i> , 203; | |
| <i>Pogoniris</i> , subgen. characterized . | 143 | squamatum, 36; <i>torulosum</i> , | |
| Poison for spears and arrows, com- | | <i>Baker</i> , <i>n. sp.</i> , 204; <i>triquetrum</i> , | |
| position of | 58 | 32; <i>urophyllum</i> , 30; <i>vulgare</i> , | |
| — effects of the, and means of | | 736; <i>xantholepis</i> (<i>Goniophleb-</i> | |
| cure | 59 | bium), <i>Harrington</i> , <i>n. sp.</i> , 36. | |
| Poisoned spears and arrows . . . | 55 | <i>Polypogon monspeliensis</i> . . . | 715 |
| — application of poison to the | | — <i>maritimus</i> | 715 |
| points of | 59 | <i>Polyporus abruptus</i> , <i>B.</i> , 42; <i>adus-</i> | |
| — preparation of the points of . | 55 | tus, 51, 53; <i>affinis</i> , 42, 45, 49; | |
| — preparation of the poison for | 56 | <i>amboinensis</i> , 40; <i>aratus</i> , <i>B.</i> , 53; | |
| Poisons, vegetable, of the Samoan | | <i>armeniacus</i> , <i>B.</i> , 42; <i>aruensis</i> , | |
| Islanders | 55 | <i>B.</i> , 43; <i>australis</i> , 39, 42, 49, | |
| <i>Pollinia distachya</i> | 735 | 53, var. <i>applanata</i> , 41, 44; | |
| <i>Polycarpia nivea</i> | 371 | <i>benetostus</i> , <i>B.</i> , 52; <i>caliginosus</i> , | |
| <i>Polycarpon alsinefolium</i> , 370; <i>Bi-</i> | | <i>B.</i> , 46, 50; <i>easperatus</i> , <i>B.</i> , 47; | |
| <i>vouæ</i> , 371; <i>tetraphyllum</i> , 370. | | <i>cinnabarinus</i> , 39; <i>cochlear</i> , 42; | |
| <i>Polynemum Fontanesii</i> | 643 | <i>Dickinsii</i> , <i>B.</i> , (ftnote) 50; <i>dictyopus</i> , 41; <i>elegans</i> , 49; <i>elongatus</i> , 51; <i>endotheus</i> , <i>B.</i> , 46; <i>exoteph-</i> | |
| <i>Polyedrium muticum</i> | 238 | <i>phrus</i> , <i>B.</i> , 49; <i>flabelliformis</i> , 53; | |
| <i>Polygala Balansæ</i> , 352; <i>monspeli-</i> | | <i>floridanus</i> , <i>B.</i> , 43; <i>fedatus</i> , <i>B.</i> , | |
| <i>aca</i> , 350; <i>rosea</i> , 350; <i>rupestris</i> , | | 41; <i>Fullageri</i> , <i>B.</i> , 54; <i>gibbosus</i> , | |
| 350; <i>Webbiana</i> , 351. | | 46; <i>grammocephalus</i> , <i>B.</i> , 45; <i>hirsutus</i> , 33; <i>igniarius</i> , 47, 49; <i>incrassatus</i> , <i>B.</i> , 41; <i>laticolor</i> , <i>B.</i> , | |
| <i>Polygonum aviculare</i> , 652; <i>equisetiforme</i> , var. <i>trigynum</i> , 652; <i>litorale</i> , 652; <i>maritimum</i> , 652; <i>serrulatum</i> , var. <i>salicifolium</i> , 653. | | 46; <i>lilacino-gilvus</i> , 39, <i>lucidus</i> , | |
| <i>Polyphyly</i> , various kinds of, in | | 46, 52; <i>luteus</i> , 42; <i>meleagris</i> , | |
| plants | 207, 208 | <i>B.</i> , 42; <i>oblectans</i> , 38; <i>obliquus</i> , | |
| <i>Polypodium accedens</i> , 32; <i>adnascens</i> , 32; <i>adnatum</i> , 36; <i>angustifolium</i> , 36; <i>aoristisorum</i> (<i>Goniopteris</i>), <i>Harrington</i> , <i>n. sp.</i> , 30; <i>attenuatum</i> , 36; <i>aureum</i> , 36; <i>conjugatum</i> , 33; <i>crassifolium</i> , 37; <i>craterisorum</i> (<i>Eupolypodium</i>), <i>Harrington</i> , <i>n. sp.</i> , 31; <i>encullatum</i> , 31; <i>delicatulum</i> , 36; <i>dipteris</i> , 33; <i>Fortunei</i> , 33; <i>fragile</i> (<i>Phegopteris</i>), <i>Baker</i> , <i>n. sp.</i> , 203; <i>fraxinifolium</i> , 36; <i>Gilpinæ</i> , (<i>Eupolypodium</i>), <i>Baker</i> , <i>n. sp.</i> , 204; <i>hammatisorum</i> (<i>Phymatoides</i>), <i>Harrington</i> , <i>n. sp.</i> , 32; <i>hastatum</i> , 33; <i>heterocarpum</i> , var. <i>Zippellii</i> , 32; <i>hirtellum</i> , 31; <i>incanum</i> , 36; <i>lævigatum</i> , 36, var. <i>rigidum</i> , <i>Harrington</i> , <i>n. var.</i> , 36; <i>lanceolatum</i> , 37; <i>lingua</i> , 32; <i>loriceum</i> , 36; <i>minutum</i> , 31; <i>Moseleyi</i> , 33; <i>murorum</i> , 36; <i>pal-</i> | | | |
| | | matum, 33; <i>peroxydatus</i> , <i>B.</i> , 39; <i>Persoonii</i> , <i>Fr.</i> , 45, 50; <i>pocas</i> , <i>B.</i> , (ftnote) 51; <i>portentosus</i> , 39, 51; <i>rasipes</i> , <i>B.</i> , 49; <i>rhypidium</i> , 39; <i>rufescens</i> , 51; <i>rugosus</i> , 39; <i>russo-gramme</i> , <i>B.</i> , 45; <i>sanguineus</i> , 39, 45, 49, 53; <i>seausilis</i> , <i>B.</i> , 53; <i>seruposus</i> , 39, 43; <i>semidigitaliformis</i> , <i>B.</i> , 39; <i>senex</i> , 42, 46; <i>sterceinus</i> , 49; <i>subpellucidus</i> , <i>B.</i> , 51; <i>tabacinus</i> , 43; <i>tephronotus</i> , 39; <i>vernicipes</i> , <i>B.</i> , (ftnote) 50; <i>versicolor</i> , 51; <i>xanthopus</i> , 41, 42, 45, 49; <i>zonalis</i> , 42, 46, var. <i>semilaceatus</i> , 46. | |
| | | <i>Polysphæria lanceolata</i> , 268; <i>multiflora</i> , 268; <i>parvifolia</i> , 268; <i>Schweinfurthii</i> , 268. | |

- | | Page | | Page |
|--|------|---|------|
| <i>Populus alba</i> , <i>L.</i> , 668, var. <i>integrifolia</i> , <i>Ball</i> , 668; <i>nigra</i> , 668. | | subpunctata, 272; <i>trachystyla</i> , 273; <i>umbellata</i> , 272; <i>umbrati-eola</i> , 273; <i>virens</i> , 273; <i>Vogeliana</i> , 272; <i>zambesiana</i> , 272; <i>zangnebarica</i> , 273. | |
| <i>Portulaca oleracea</i> | 372 | <i>Psychotriacæ</i> , tribe of <i>Rubiaceæ</i> | 272 |
| <i>Potamogeton natans</i> | 683 | <i>Pteris aquilina</i> , 28, 35, 735; <i>arguta</i> , 735; <i>cretica</i> , 35, 139, var. <i>stenophylla</i> , 27, 35; <i>ensiformis</i> , 27; <i>incisa</i> , 28, 35; <i>longifolia</i> , 27; <i>pedata</i> , 35; <i>quadriaurita</i> , 128; <i>remotifolia</i> , <i>Baker</i> , <i>n. sp.</i> , 199; <i>scnipinnata</i> , 27, var. <i>dispar</i> , 27; <i>serrulata</i> , 27; <i>triplicata</i> , 199. | |
| — <i>pectinatus</i> | 683 | <i>Pterocephalus depressus</i> | 493 |
| <i>Potentilla reptans</i> | 444 | <i>Pterogonium gracile</i> | 738 |
| <i>Poterium anceps</i> , 445; <i>ancistroides</i> , 445; <i>Duriei</i> , 446; <i>multicaule</i> , 445; <i>sanguisorba</i> , 444; <i>verrucosum</i> , 444, var. <i>Magnolii</i> , 444. | | <i>Puccinia corticioides</i> , <i>B. & Br. n. sp.</i> (<i>Pl. II.</i>) | 52 |
| <i>Pouchetia africana</i> | 268 | <i>Pulicaria arabica</i> , 500; <i>longifolia</i> , 501; <i>odora</i> , 501, var. <i>macrocephala</i> , <i>Ball</i> , 501; <i>Mauritanica</i> , 501; <i>sicula</i> , 502. | |
| — <i>parviflora</i> | 268 | <i>Punica Granatum</i> | 458 |
| <i>Powell</i> , Rev. T., on the nature and mode of use of vegetable poisons employed by the Samoan Islanders | 55 | <i>Putoria calabrica</i> | 482 |
| <i>Prasium majus</i> | 629 | <i>Pyrostria trilocularis</i> , <i>Balf. fil. sp. n.</i> , 11, 14. | |
| <i>Primula vulgaris</i> , various changes in flower of | 210 | — — — — —, variation of leaves of | 24 |
| <i>Protoderma viride</i> | 245 | <i>Pyrus cydonia</i> | 447 |
| <i>Prunella vulgaris</i> | 620 | <i>Pyxine</i> — — ? | 226 |
| <i>Prunus domestica</i> | 443 | | |
| — <i>prostrata</i> | 443 | <i>Quercus coccifera</i> , 667, var. <i>imbricata</i> , 667; <i>humilis</i> , 666; <i>Ilex</i> , 667, var. <i>Ballota</i> , 667; <i>lusitana</i> , 666; <i>Mirbeckii</i> , 666. | |
| <i>Psiadia Coronopus</i> , 11; <i>rodriguesiana</i> , <i>Balf. fil. sp. n.</i> , 11, 15. | | <i>Quivisia</i> | 9 |
| <i>Psilanthus</i> ? <i>ebracteolatus</i> , 271; <i>Mannii</i> , 271; ? <i>tetramerus</i> , 271. | | — <i>laciniata</i> , <i>Balf. fil. sp. n.</i> , 11, 12 | |
| <i>Psoralea bituminosa</i> , 430, 739; <i>bracteata</i> , 430; <i>dentata</i> , 430, var. <i>calycibus villosis</i> , 430. | | — — — — —, heterophylly of | 29 |
| <i>Psychotria abrupta</i> , 272; <i>Afzelii</i> , 272; <i>anctoclada</i> , 272; <i>Ansellii</i> , 273; <i>arborea</i> , 272; <i>Benthamiana</i> , 272; <i>bidentata</i> , 272; <i>bifaria</i> , 272; <i>brachyantha</i> , 272; <i>bracteosa</i> , 272; <i>Brassii</i> , 272; <i>brunnea</i> , 272; <i>calva</i> , 272; <i>cornuta</i> , 272; <i>crispa</i> , 273; <i>cristata</i> , 272; <i>Doniana</i> , 272; <i>foliosa</i> , 272; <i>gabonica</i> , 272; <i>globosa</i> , 272; <i>humilis</i> , 272; <i>infundibularis</i> , 272; <i>insidens</i> , 272; <i>Kirkii</i> , 272; <i>Kolly</i> , 272; <i>konguensis</i> , 272; <i>lanceolata</i> , <i>Balf. fil. sp. n.</i> , 11, 15; <i>latisipula</i> , 272; <i>leptophylla</i> , 272; <i>longevaginalis</i> , 272; <i>longistylis</i> , 272; <i>lophoclada</i> , 272; <i>lucens</i> , 272; <i>Mannii</i> , 272; <i>monticola</i> , 272; <i>micronata</i> , 273; <i>nigropunctata</i> , 272; <i>nubica</i> , 273; <i>obscura</i> , 273; <i>obtusifolia</i> , 272; <i>owariensis</i> , 272; <i>pauridiantha</i> , 272; <i>pumila</i> , 272; <i>punctata</i> , 272; <i>recurva</i> , 272; <i>reptans</i> , 273; <i>Schweinfurthii</i> , 272; <i>sciadephora</i> , 272; <i>setacea</i> , 272; <i>Soyauxii</i> , 273; <i>stictophylla</i> , 273; <i>subherbacea</i> , 272; <i>subnuda</i> , 272; <i>subobliqua</i> , 272; | | | |
| | | <i>Radiola linoides</i> | 378 |
| | | <i>Radula complanata</i> | 191 |
| | | — <i>physoloba</i> | 191 |
| | | <i>Ramalina calicaris</i> , 227; <i>complanata</i> , 215; <i>cuspidata</i> , var. <i>crassa</i> , 211; <i>intermedia</i> , 211, 222; <i>minuscula</i> , 218; <i>rigida</i> , 211; <i>subfraxinea</i> , 226, 227; <i>Yemenensis</i> , 222. | |
| | | <i>Randia acuminata</i> , 266; ? <i>caudata</i> , 266; <i>dumetorum</i> , 266; <i>gambica</i> , 266; <i>genipæflora</i> , 267; <i>heterophylla</i> , <i>Balf. fil. sp. n.</i> , 11, 14—variation of leaves of, 24; <i>macrantha</i> , 267; <i>macrocarpa</i> , 267; <i>maculata</i> , 267; <i>malleifera</i> , 267; <i>Munsæ</i> , 267; <i>octomera</i> , 267; <i>pallens</i> , 266; <i>rubens</i> , 266. | |
| | | <i>Ranunculus aquatilis</i> , 304; <i>arvensis</i> , 307; <i>atlanticus</i> , 306; <i>Bau-</i> | |

- | | Page | | Page |
|--|---------|---|----------|
| dotii, 304; bulbosus, 306, var. giganteus, 307; bullatus, 304; calthæfolius, 304; chærophyllos, 305; Ficaria, 304, var. intermedius, 304; gramineus, 305; leucothrix, 305; muricatus, 307; ophioglossifolius, 308; palustris, 306; parviflorus, 308; philonotis, var. intermedius, 307; procerus, 306; spicatus, 305, var. blepharicarpos, 305; trilobus, 307. | | Rodriguez, changes in flora of . . . | 8 |
| Raphanus Raphanistrum . . . | 337 | —, new plants from . . . | 7 |
| Rapistrum rugosum . . . | 334 | Rœmeria hybrida . . . | 313 |
| Reinsch, P. F., Contributiones ad floram Algarum aquæ dulcis Promontorii Bonæ Spæ . . . | 232 | Romulca, characters of . . . | 73, 86 |
| Reseda alba, 337; attenuata, <i>Ball</i> (Pl. XIV.), 338; collina, 739; diffusa, 339; elata, 340; lanecolata, 340; lutea, var. mucronata, 339; Luteola, 341, var. australis, 341, var. crispata, 341; media, 339; neglecta, 339; Phyteuma, 338; propinqua, 337; stricta, var. subsessilis?, 340. | | — arenaria, 89; barbata, 89; bulbocodioides, 88; bulbocodiun, 86, 677, var. crocea, 87, var. flaveola, 87, var. grandiflora, 86, var. nivale, 87, var. pulchella, 86, var. pylum, 87, var. subpalustre, 87, var. syrtica, 86, var. umbellatum, 87; camerooniana, 88; chloroleuca, 89; Clusiana, 87; Columnæ, 88, 677, var. modesta, 88, var. subalbida, 88; corsica, 87; crocifolia, 87; cruciata, 89; cuprea, 89; dichotoma, 89; elongata, 87; filifolia, 88; grandiscapa, 87, 697; hirsuta, 89; latifolia, 89; ligustica, 87; Linaresii, 87; longifolia, 89; Macowani, 88; neglecta, 87; numidica, 87; pudica, 89; purpurascens, 87, var. edulis, 87; ramiflora, 87, 677; Requiennii, 88; Revelieri, 88; Rollii, 87; rosca, 88, var. Celsii, 89; similis, 89; speciosa, 89; spiralis, 90; sublutea, 88; tortuosa, 88. | |
| Rhabdostigma Kirkii . . . | 268 | Rosa canina, 446, var. subinermis, <i>Ball</i> , 446; micrantha, 446, var. atlantica, <i>Ball</i> , 446; sempervirens, 446; Seraphini, 447. | |
| Rhagadiolus arenarius, 536; Hedychnois, 535; pendulus, 535; stellatus, 535. | | Rubia cordifolia, 275; discolor, 275; peregrina, 483, var. angustifolia, 484, var. lucida, 483; tinctorum, 483. | |
| Rhamnus Alaternus, 391; lycioides, 392; oleoides, 391, var. amygdalinus, 392, var. angustifolia, 392. | | Rubiaceæ, differences in the andrœcium . . . | 255 |
| Rhizophydium, spec. ? . . . | 246 | —, differences in the calyx . . . | 256 |
| Rhus oxycantha, 393, var. albida, 393, var. unnamed, 393, var. zizyphina, 393; pentaphylla, 394. | | —, differences in the corolla . . . | 255 |
| Rhynchosnema angulare . . . | 7 | —, differences in the fruit . . . | 253 |
| Ribes grossularia, <i>L.</i> , var. atlantica. <i>Ball</i> . . . | 449 | —, differences in the habit and stature . . . | 259 |
| Ricasolia glomulifera . . . | 228 | —, differences in the inflorescence and bracts . . . | 257 |
| Riccia albomarginata, 195; limbata, 195; purpurescens, 195 | | —, differences in the leaves and stipules . . . | 258 |
| Richardia scabra . . . | 274 | —, differences in pistil . . . | 252 |
| Ricinus communis . . . | 664 | —, economic and superstitious uses . . . | 260 |
| Rigidella, characters of . . . | 77, 134 | —, flowers of . . . | 252 |
| — flammea, 135; immaculata, 135; orthantha, 135. | | —, frequency of dimorphism in . . . | 252 |
| Roeccella phycopsis . . . | 217 | —, general distribution of . . . | 251, 276 |
| — tinctoria . . . | 211 | —, new species of African . . . | 262 |
| Rochea, section of Geissorhiza, defined . . . | 93 | — of Mascarene Islands . . . | 277 |
| Rochelia stellulata . . . | 571 | — of North Africa &c. . . . | 278 |
| Rodriguez, Algæ collected at . . . | 6 | — of South Africa | 277 |
| —, aspect of the phænogamic vegetation of . . . | 7 | | |

- | | Page | | Page |
|---|------|--|----------|
| ubiaceæ of South Europe, Asia | | Sarcostemma Odontolepis, <i>Balf. fil.</i> | |
| Minor, Arabia, &c. | 279 | <i>sp. n.</i> , 11, 17; <i>vininale</i> , 18. | |
| — of Tropical America | 280 | Sargassum ilicifolium | 6 |
| — of Tropical Asia and Pacific | | <i>vulgare</i> | 6 |
| Islands | 279 | Satureja inodora | 613 |
| —, on the peculiarities and dis- | | Saxifraga globulifera, var. <i>gibraltarica</i> , | |
| tribution of, in Tropical Africa . | 248 | 417; <i>granulata</i> , 448; | |
| —, organic peculiarities of | 252 | <i>Mawcana</i> , 448; <i>tridactylites</i> , | |
| — tabular view of tribes, genera, | | 448. | |
| and species of | 263 | Scabiosa maritima, 492, var. | |
| Rubus debilis, <i>Ball</i> (Pl. XIX.) . | 443 | <i>minor</i> , 492; <i>rutæfolia</i> , 493; | |
| — fruticosus, var. <i>discolor</i> | 443 | <i>semipapposa</i> , 492; <i>stellata</i> , 493, | |
| Rumex bucephalophorus, 649; | | var. <i>nonspelensis</i> , 493. | |
| <i>conglomeratus</i> , 649; <i>crispus</i> , | | Scandix australis | 471 |
| 648; <i>intermedius</i> , 649; <i>Papilio</i> , | | — <i>Peeten-Veneris</i> | 471 |
| 651; <i>pictus</i> , 650; <i>pulcher</i> , 649; | | Scenedesmus aculeolatus, <i>Reinsch</i> , | |
| <i>scutatus</i> , <i>L.</i> , var. <i>induratus</i> , | | <i>sp. n.</i> | 238 |
| <i>Ball</i> , 650; <i>thyrsoides</i> , 649; | | — <i>quadricaudatus</i> | 238 |
| <i>tingitanus</i> , 649; <i>vesicarius</i> , <i>L.</i> , | | Schismus calycinus | 724 |
| var. <i>rhodophysa</i> , <i>Ball</i> , 651. | | Schizaea dichotoma | 205 |
| Ruscus hypophyllum | 696 | Schizophyllum commune, 38, 40, 45, 53 | |
| Ruta chalepensis, 389, var. <i>bracteosa</i> , | | Schizostigma, sect. of <i>Crocus</i> , de- | |
| 389; <i>montana</i> , 389. | | <i>finid</i> | 84 |
| Rutidea decorticata, 271; <i>ferruginea</i> , | | Schizostylis, characters of | 75, 108 |
| 271; <i>fuscescens</i> , 271; | | — <i>coccinea</i> | 108 |
| <i>glabra</i> , 271; <i>hispida</i> , 271; <i>membranacea</i> , | | — <i>paniciflora</i> | 108 |
| 271; <i>olenotricha</i> , 271; | | Schœnus nigricans | 702 |
| <i>parviflora</i> , 271; <i>rufipilis</i> , 271; | | Schweiggera, subgen., characterized | 178 |
| <i>Smithii</i> , 271. | | Scilla autumnalis, 686; <i>hæmorrhoidalis</i> , | |
| Sabicea calycina, 265; <i>capitellata</i> , | | 687; <i>hemisphærica</i> , | |
| 265; <i>cauliflora</i> , 265; <i>ferruginea</i> , | | var. <i>glabra</i> , 686; <i>hispanica</i> , 687; | |
| 265; <i>geantha</i> , 265; <i>pilosa</i> , 265; | | <i>lingulata</i> , 687; <i>mauritanica</i> , 687; | |
| <i>segregata</i> , 265; <i>venosa</i> , 265; | | <i>monophylla</i> , 686; <i>obtusifolia</i> , | |
| <i>Vogelii</i> , 265. | | var. <i>fallax</i> , 686; <i>verna</i> , var. | |
| Sagina apetala, 367; <i>ciliata</i> , 367; | | <i>major</i> , 686. | |
| <i>Linnei</i> , 367; <i>maritima</i> , 367; | | Scirpus Holoschoenus, 703, var. | |
| <i>procumbens</i> , 367. | | <i>australis</i> , 704; <i>lacustris</i> , 702; | |
| Salicornia, fruticosa | 646 | <i>litoralis</i> , 702; <i>maritimus</i> , 702; | |
| Salix alba, 657; <i>canariensis</i> , 668; | | <i>Michelianus</i> , 704; <i>pubescens</i> , | |
| <i>pedicellata</i> , 668; <i>purpurea</i> , var. | | 703; <i>Savii</i> , 703. | |
| <i>Helix</i> , 668. | | Scleranthus annuus, var. <i>verticillatus</i> | 642 |
| Salsola Kali, 648; <i>longifolia</i> , 647, | | Sclerocarya caustanea | 11, 12 |
| var. <i>verticillata</i> , 647; <i>vermiculata</i> , | | Sclerosciadium nodiflorum | 473 |
| var. <i>flavescens</i> , 647, var. | | Scolopendrium Hemionitis | 736 |
| <i>microphylla</i> , 647. | | Scolymus hispanicus | 533, 740 |
| Salvia ægyptiaca, 619; <i>argentea</i> , 616; | | — <i>maculatus</i> | 533 |
| <i>bicolor</i> , 616; <i>interrupta</i> , 615; | | Scorpiurus subvillosa, 426; <i>sulcata</i> , | |
| <i>Maurorum</i> , <i>Ball</i> (Pl. XXVIII.), | | 425; <i>vermiculata</i> , 426. | |
| 615; <i>ochroleuca</i> , 616; <i>tarasacifolia</i> , | | Scorzonera deliciosa, var. <i>tetuanensis</i> , | |
| 618; <i>Verbenaca</i> , var. <i>oblongifolia</i> , | | 551; <i>laciniata</i> , 552; <i>pygmæa</i> , | |
| 617, var. <i>præcox</i> , 617. | | 551; <i>undulata</i> , 551. | |
| Sambucus Ebulus | 481 | Scrophularia arguta, 596; <i>auriculata</i> , | |
| — <i>nigra</i> | 481 | <i>L.</i> , 597, var. <i>parviflora</i> , | |
| Samolus Valerandi | 563 | <i>Ball</i> , 597; <i>canina</i> , 598, var. | |
| Sandwich Islands, lichens from | 224 | <i>pinnatifida</i> , 598, var. <i>vel sp. nov.</i> , | |
| Saponaria vaccaria | 355 | 598; <i>Durandii</i> , 598; <i>frutescens</i> , | |
| Sarcoccephalus esculentus | 263 | 599; <i>hispida</i> , 597; | |
| — <i>Russeggeri</i> | 263 | <i>lævigata</i> , 598; <i>sambucifolia</i> , | |
| | | 597; <i>Scorodonia</i> , 597. | |

- | | Page | | Page |
|---|--------|--|---------|
| <i>Seyphoehlamys</i> , <i>Balf. fil. gen. nov.</i> | 11, 14 | <i>gibraltarica</i> , 361; <i>museipula</i> , 360; <i>noeturna</i> , 357; <i>obtusifolia</i> , 357, 739; <i>parviflora</i> , 357; <i>psammitis</i> , 358; <i>rubella</i> , 360; <i>setacea</i> , 359; <i>tridentata</i> , 356. | |
| — <i>revoluta</i> , <i>Balf. fil. sp. unica</i> | 15 | <i>Silybum Marianum</i> | 524 |
| — —, variation of leaves of | 24 | <i>Simethis planifolia</i> | 693 |
| <i>Seytonema</i> , sp.?, 211; <i>elegans</i> | 236 | <i>Siphomeris foetens</i> | 274 |
| <i>Sceale montanum</i> | 732 | <i>Sirosiphon alpinus</i> , 237; <i>Olivieri</i> , 237; <i>panniformis</i> , 236; <i>pulvinatus</i> , 236, var. <i>intermedius</i> , 236, var. 237; <i>secundatus</i> , 236. | |
| <i>Seeurigera Coronilla</i> | 425 | <i>Sisymbrium erysimoides</i> , 324; <i>Irio</i> , 323; <i>officinale</i> , 323; <i>runcinatum</i> , 324, var. <i>hirsutum</i> , <i>Ball</i> , 324; <i>thalianum</i> , 324. | |
| <i>Sedum acre</i> , 453; <i>album</i> , 453, var. <i>glanduliferum</i> , 453, var. <i>mieranthum</i> , 453; <i>altissimum</i> , 453; <i>brevifolium</i> , 452, var. <i>induratum</i> , 452; <i>caespitosum</i> , 452; <i>dasyphyllum</i> , 452, var. <i>glanduliferum</i> , 452, var. <i>oblongifolium</i> , <i>Ball</i> , 452; <i>hispidum</i> , 454; <i>modestum</i> , <i>Ball</i> (Pl. XX.), 452; <i>pruinatum</i> , 453; <i>rubens</i> , 451. | | <i>Sisyrinchia</i> , tribe of <i>Irids</i> , characters of | 76 |
| <i>Selaginella atroviridis</i> , 34; <i>cuspidata</i> , 37; <i>denticulata</i> , 737; <i>exaltata</i> , 37; <i>geniculata</i> , 37; <i>Lyalli</i> , 206; <i>molliceps</i> , 206; <i>rupestris</i> , 737; sp.? near <i>S. caulescens</i> , 34. | | <i>Sisyrinchium</i> , characters of | 76, 115 |
| <i>Sempervivum atlanticum</i> | 454 | — <i>acre</i> , 118; <i>alatum</i> , 119, var. <i>paucifolium</i> , 119; <i>avenaceum</i> , 117; <i>bermudianum</i> , 117, var. <i>anceps</i> , 117, var. <i>geniculatum</i> , 117, var. <i>gramineum</i> , 117, var. <i>indioides</i> , 117, var. <i>mucronatum</i> , 117; <i>bogotense</i> , 117; <i>californicum</i> , 118; <i>chilense</i> , 117; <i>convolutum</i> , 118; <i>cuspidatum</i> , 120; <i>fasciculatum</i> , 119; <i>filifolium</i> , 116; <i>Glaziovii</i> , 119; <i>gracile</i> , <i>Klotsch MS.</i> , 118; <i>graminifolium</i> , 118, var. <i>ascendens</i> , 118, var. <i>maculatum</i> , 118; <i>grandiflorum</i> , 116; <i>Hartwegii</i> , 119; <i>ineurvatum</i> , 119; <i>iridifolium</i> , 118; <i>Jamesoni</i> , 117; <i>junceum</i> , 116, var. <i>asperulum</i> , 116, var. <i>colehaguense</i> , 116, var. <i>gracile</i> , 116, var. <i>scirpiforme</i> , 116; <i>laterale</i> , 119; <i>Luzula</i> , 119; <i>Mandoni</i> , 118; <i>marginatum</i> , 119; <i>mieranthum</i> , 118; <i>minutiflorum</i> , 117; <i>monostachyum</i> , 120; <i>nervosum</i> , 120; <i>Nuno</i> , 116, var. <i>leueanthum</i> , 116; <i>paehyrhizum</i> , 119; <i>palmifolium</i> , 119, var. <i>congestum</i> , 120; <i>patagonicum</i> , <i>Philippi MS.</i> , 117; <i>Peareei</i> , 118; <i>Philippii</i> , 116; <i>pusillum</i> , 117; <i>restioides</i> , 119; <i>scabrum</i> , 117; <i>secundiflorum</i> , 116; <i>Sellowianum</i> , 119; <i>sessiliflorum</i> , 120; <i>setaceum</i> , 116; <i>striatum</i> , 120; <i>tenuifolium</i> , 118; <i>tinctorum</i> , 118; <i>trinerve</i> , 116; <i>unispataecum</i> , 117; <i>vaginatum</i> , 119; <i>Weirii</i> , 119. | |
| <i>Senebiera coronopus</i> | 331 | | |
| <i>Senecio coronopifolius</i> , 514; <i>erassifolius</i> , 514; <i>Doronicum</i> , var. <i>Hosmariense</i> , <i>Ball</i> , 515; <i>erraticus</i> , 515; <i>foliosus</i> , 515; <i>gallieus</i> , 514, var. <i>sonchifolius</i> , <i>Ball</i> , 514; <i>giganteus</i> , 515; <i>lividus</i> , 513, var. <i>pinguis</i> , <i>Ball</i> , 513; <i>leueanthemifolius</i> , 513, var. <i>major</i> , <i>Ball</i> , 514; <i>pteroneura</i> , 515; sp.? of <i>Mascarenes</i> , 9; <i>vulgaris</i> , 513, var. <i>teneriffæ</i> , 513. | | | |
| <i>Sepedonium chrysospermum</i> | 40 | | |
| <i>Serapias cordigera</i> | 674 | | |
| — <i>Lingua</i> | 674 | | |
| <i>Setaria verticillata</i> | 710 | | |
| <i>Sherardia arvensis</i> | 490 | | |
| <i>Sideritis Cossoniana</i> , 622; <i>grandiflora</i> , 620; <i>hirsuta</i> , 621; <i>montana</i> , 621; <i>scordioides</i> , var. <i>angustifolia</i> , 620; <i>romana</i> , 621; <i>villosa</i> , 621. | | | |
| <i>Silene apetal</i> , 359; <i>cerastoides</i> , var. <i>anomala</i> , 356; <i>colorata</i> , 358, var. <i>lasioealyx</i> , 358; <i>corrugata</i> , 359, var. <i>adusta</i> , 360; <i>decipiens</i> , 357; <i>disticha</i> , 356; <i>gallica</i> , 356, var. <i>minor</i> , 356; <i>glaucia</i> , 359; <i>hirsuta</i> , var. <i>tuberculata</i> , <i>Ball</i> , 357; <i>inaptera</i> , 360; <i>inflata</i> , 355, var. <i>rubriflora</i> , <i>Ball</i> , 356; <i>italica</i> , 360; <i>læta</i> , 361, 739; <i>mollissima</i> , 361, var. | | | |

- | | Page | | Page |
|--|---------|---|--------|
| Sisyringium, sect. of Sisyrinchium,
defined | 116 | Spicilegium Floræ Maroccanæ | 281 |
| Smilax aspera | 695 | Spirogyra insignis | 243 |
| — mauritanica | 695 | Sporobolus pungens | 713 |
| Smyrnium Olusatrum | 463 | — —, var. gaditanus | 713 |
| Solanum nigrum, 580, var. humile,
580, var. miniatum, 580, var.
suffruticosum, 580; sodomæum,
581; villosum, 580. | | Stachys arenaria, 627; arvensis,
626; circinata, 625; Durandiana,
624; germanica, 624, var. cre-
tica, 624; hirta, 626, 740;
Maweana, <i>Ball</i> , 626; mollis,
627; saxicola, 625, var. villosis-
sima, <i>Ball</i> , 626. | |
| Solenomelus, characters of | 76, 120 | Stæhelina dubia, and var. ? | 525 |
| — acaulis, 121; andinus, 121;
biflorus, 121; Chamælum, 121;
chilensis, 121, var. punctatus,
121; Lechleri, 121; nigricans | 121 | Statice Beaumieriana, 557; cchi-
oides, 559; ferulacea, 560; gum-
mifera, var. corymbulosa, 559;
lychnidifolia, 558; mucronata,
558; occidentalis, 559; olæ-
folia, 558; ornata, 559; ovali-
folia, 558; sinuata, 557, var. in-
tegrifolia, 557; spathulata, var.
β, emarginata, 559; Thouini,
557, 740. | |
| Solution of adherent organs of
plants | 207 | Staurogenia cubica, <i>Reinsch</i> , <i>n. sp.</i> | 238 |
| Somerset, Cape York, fungi from | 41 | Staurospermum capucinum | 7 |
| Sonchus acidus, 549; asper, 548;
fragilis, 549; glaucescens, 548;
maritimus, 549; oleraceus, 548;
tenerimus, 548, var. arbores-
cens, 549, var. maritimus, <i>Ball</i> ,
548, var. tuberculatus, <i>Ball</i> , 548. | | Steere, Prof., tropical ferns coll. by | 25 |
| Sorghum halepense | 734 | Stellaria arenaria, 363; media,
363, var. major, 363; uliginosa,
363. | |
| Sparaxis, characters of | 74, 98 | Stereocaulon —? 221; cymo-
sum, 220; mixtum, 222; proxi-
mum, 224, 227; ramulosum, 223. | |
| — bulbifera, 98; grandiflora, 98,
var. Liliago, 99, var. lineata, 99,
var. stellaris, 99; tricolor, 99, var.
blanda, 99, var. Griffini, 99, var.
versicolor, 99. | | Stereum contrarium, <i>B.</i> , 52; effu-
sum, <i>B.</i> , 44; elegans, 50; hir-
sutum, 40; illudens, 40; indu-
ratum, <i>B.</i> , 44; lobatum, 44;
Mosclei, <i>B.</i> , 48; ostrca, 48;
prolificans, <i>B.</i> , 41; vespillo-
neum, 44. | |
| Spartina stricta | 714 | Sticta aurata, 218; damæcornis,
217; hypopsila, 223; lineari-
loba, 223; pulmonacea, 217;
subvariabilis, 223. | |
| Spatalanthus, characters of | 75, 104 | Stictina caperata, 224; carpoloma,
224; dichotomoides, 224; in-
tricata, f. Thouarsii, 222. | |
| — speciosus | 100 | Stipa Aristella, 711; gigantea,
711; nitens, 712; parviflora,
712; tortilis, 711, var. pubes-
cens, 711. | |
| Spathirachis, sect. of Sisyrinchium,
defined | 120 | Stipularia africana, 266; elliptica,
266; gabonica, 266. | |
| Specularia falcata | 555 | Stirton, Dr. J., revision of paper of,
by the Rev. J. M. Crombie | 211 |
| Spergula arvensis | 368 | Streptanthra, characters of | 74, 92 |
| — pentandra | 368 | — cuprea, 93; elegans, 92. | |
| Spergularia diandra, 369; fim-
briata, 368, var. condensata, 368,
var. tenuis, 369, var. typica, 368;
marina, 368; rubra, 369, var.
pinguis, 369, var. sperguloides,
<i>Ball</i> , 369. | | Suæda fruticosa | 647 |
| Spermacoce compacta, 274; com-
pressa, 274; dibrachiata, 274;
filiformis, 274; globosa, 274;
hebecarpa, 274; Kotschyana,
274; leucadca, 274; phyteuma,
274; ? pilosa, 274; radiata, 274;
ramisparsa, 274; Ruelliae, 274;
senensis, 274; stachydea ?, 274;
stricta, 274; tenuissima, 274. | | — maritima | 647 |
| Spermacocæ, tribe of Rubiacæ | 274 | | |
| Spermosira macrospora, <i>Reinsch</i> ,
<i>sp. n.</i> | 235 | | |
| Sphacclaria rigida | 6 | | |
| Sphenostigma, characters of | 76, 124 | | |
| — Sellowianum | 124 | | |

- | | Page | | Page |
|---|----------|--|----------|
| Suceowia balearica | 330 | Theleophora lamellata | 44, 50 |
| Supplemental developments in
plants | 207, 209 | Thelotrema — ? | 223 |
| Symphogyna podophylla | 194 | Thelygonum Cynocrambe | 648 |
| Symphostemon, subgen., character-
ized | 121 | Thesium humile | 655 |
| Synnottia, characters of | 79, 169 | Thlaspi perfoliatum | 183, 332 |
| — bicolor, 170; galeata, 170;
variegata, 170. | | — —, var. rotundifolium, <i>Ball</i> | 333 |
| Syringodea, characters of | 73, 85 | Thorca violacea | 7 |
| — bicolor, 86; filifolia, 86;
pulehella, 85. | | Thuja dolabrata with Uromyces
growing on it (Pl. II.) | 54 |
| Systema Iridacearum, J. G. Baker
on | 61 | Thymus Broussonnetii, 612; capi-
tatus, 611; lanceolatus, 611, var.
crispus, <i>Ball</i> , 611; maroccanus,
612; Munbyanus, 612; satu-
reoides, <i>Coss.</i> , 610, var. pseudo-
mastichina, <i>Ball</i> , 611; Serpyll-
lum, var. atlanticus, <i>Ball</i> , 611. | |
| Tabernæmontana orientalis | 57, 58 | Tigrida, characters of | 77, 135 |
| Tacea pinnatifida, use of flower-
stalk of | 59 | — atrata, 136; curvata, 136;
lutea, 136; pavonia, 136, var.
conchiflora, 136; violacea, 136;
Van-Houttei, 136. | |
| Tænitis angustifolia | 37 | Tigridæ, tribe of Irids, characters
of | 77 |
| Tahiti, fungi from | 52 | Tillæa muscosa | 449 |
| — lichens from | 224 | Tinguarra sicula | 471 |
| Tamarix africana, 372; articulata,
372; gallica, 372; speciosa, 372. | | Tolpis barbata, 535, 740, var. gran-
diflora, <i>Ball</i> , 535; umbellata,
535. | |
| Tamus communis | 682 | Tongatabu, fungi from | 40 |
| Tanaetum annuum | 513 | —, lichens from | 227 |
| Tanulepis, <i>Balf. fil. gen. nov.</i> | 11, 17 | Tortula inermis, 738; muralis, 738;
ruralis, 738. | |
| — sphenophylla, <i>Balf. fil. sp.</i>
<i>unica</i> | 17 | Trachelium angustifolium | 555, 740 |
| Tapeinanthus humilis | 678 | — cæruleum | 555, 740 |
| Tapenia, characters of | 76, 114 | Tragopogon glaber | 550 |
| — magellanica | 115 | — porrifolius | 550 |
| Taraxacum officinale | 546 | Trametes bicolor, <i>B.</i> , 43; cognatus,
<i>B.</i> , 43; conchatus, <i>B.</i> , 47; de-
vexa, 39; lactineus, 50; mollis,
39, 47; occidentalis, 39, 43, 45,
47, 50; Moselei, <i>B.</i> , 40; rigida,
39. | |
| Tarennia angolensis, 266; conferta,
266; congensis, 266; grandiflora,
266; mossambiensis, 266;
nigrescens, 266; nilotica, 266;
nitidula, 266; pallidula, 266;
tetramera, 266. | | Tremella lutescens | 40 |
| Targionia capensis | 194 | Triainolepis africana | 273 |
| Taxodium distichum | 180 | — Hildebrandtii | 273 |
| Temnopteryx sericea | 265 | Tribulus terrestris | 381 |
| Teneriffe, lichens from | 218 | Trielaysia ajurensis, 268; angolen-
sis, 268; biafrana, 268; braete-
ata, 268; buxifolia, 268; coriacea,
268; congesta, 268; crassifolia,
268; gabonica, 268; jasminiflora,
268; Kirkii, 268; mierantha,
268; microphylla, 268; niam-
niamensis, 268; Nyassæ, 268;
okelensis, 268; ovalifolia, 268;
pallens, 268; reticulata, 268;
Sonderiana, 268; syrmanthera,
268. | |
| Terminalia Benzoni, variation of
leaves of | 24 | Trichoderma viride | 40 |
| Ternate (Moluccas), fungi from | 44 | | |
| Teuerium bracteatum, 630; bulla-
tum, 634; eollinum, 630; deci-
piens, 631; fruticans, 630; gra-
natense, var. atlanticum, <i>Ball</i> ,
633; Polium, 633, var. angusti-
folium, 633, var. atlanticum,
<i>Ball</i> , 633, var. gnaphalodes, 633;
Pseudo-scorodonia, 631, var. bæ-
tium, 631; resupinatum, 631;
rupestre, 632; Scorodonia, 631. | | | |
| Thalictrum glaucum | 363 | | |
| Thamnum alopecurum | 738 | | |
| Thapsia garganica | 479 | | |
| — villosa | 479 | | |

	Page		Page
<i>Trichomanes apiifolium</i> , 26; auriculatum, 26; brachypus, 34; crispum, 34; digitatum, 26; flabellatum, 197; pallidum, 26; parvulum, 26; rigidum, 26, 197; saxifragoides, 26; Smithii, 26.		<i>Ulva latissima</i>	7
<i>Trichostachys aurca</i> , 273; ciliata, 273; longifolia, 273; petiolata, 273; vaginalis, <i>Hn. sp. n.</i> (Pl. VIII.), 263, 275.		<i>Uncaria africana</i>	263
<i>Trichostomum Barbula</i> , 738; crispulum, 737; mutabile, 738; tophaceum, 737.		<i>Urceolaria scruposa</i>	223
<i>Trifolium angustifolium</i> , 417; arvense, 417, var. <i>Preslianum</i> , <i>Ball</i> , 417; atlanticum, <i>Ball</i> (Pl. XVIII.), 418; Bocconi, 417; bracteatum, 418; Cherleri, 419; fragiferum, 421; glomeratum, 419, var. <i>condensatum</i> , <i>Ball</i> , 419; humile, 419; isthmocarpum, 420; lappaceum, 417; ligusticum, 417; maritimum, 418; procumbens, var. <i>campestre</i> , 421, 739; repens, 419; resupinatum, 421; scabrum, 418; spumosum, 420; stellatum, 419; strictum, 419; subterraneum, 420; tomentosum, 421.		— <i>scruposa bryophila</i>	219
<i>Triglochin Barrelieri</i>	682	<i>Urginea anthericoides</i>	688
<i>Trigonella monspeliaca</i> , 411; polycerata, 411, var. <i>atlantica</i> , <i>Ball</i> , 411.		— <i>maritima</i>	687
<i>Trisetum flavescens</i> , 718; panicum, 718; pumilum, 718.		<i>Uromyces deformans</i> , <i>B. & Br. sp. n.</i> (Pl. II.)	52
<i>Tristan d'Acunha</i> , lichens of	222	<i>Urophyllosum Afzelii</i> , 265; callicarpoides, 265; hirtellum, 265; insulare, 265; micranthum, 265; paucinerve, 265; rubens, 265; viridiflorum, 265.	
<i>Triticum elongatum</i> , 731; hordeaceum, 730; junceum, 730; villosum, 730.		<i>Urospermum Dalechampii</i>	551
<i>Tritonia</i> , characters of.	78, 161	— <i>picroides</i>	551
— <i>aurca</i> , <i>Pappe, MSS.</i> , 163; Bolusii, 163; crocata, 162, var. <i>aurantiaca</i> , 162, var. <i>purpurea</i> , 162, var. <i>sanguinea</i> , 162; deusta, 162; hyalina, 163; miniata, 162; scillaris, 163; squalida, 163; undulata, 163.		<i>Urtica dioica</i> , 665, var. <i>procera</i> , 665; membranacea, 665; pilulifera, 664; urens, 664.	
<i>Trypethelium eruentum</i>	217	Uses of a commercial cane, note on	1
<i>Tunica compressa</i>	355	<i>Usnea ceratina</i> , 224; dasypoga, 228; microcarpa, 224; trullifera, <i>Nyl. sp. n.</i>	224
— <i>prolifera</i>	355	<i>Ustilago carbo</i>	52
<i>Turbinaria ornata</i>	6	<i>Vaillantia hispida</i>	275
<i>Typha angustifolia</i>	684	— <i>muralis</i>	483
<i>Ulex africanus</i> , 401; breticus, 401, var. <i>congestus</i> , <i>Ball</i> , 401; Boivini, 401, var. <i>megalarites</i> , <i>Ball</i> , 402.		<i>Valeriana tuberosa</i>	490
<i>Ulmus campestris</i>	666	<i>Valerianella Auricula</i> , 491; carinata, 492; discoides, 491; microcarpa, 492.	
		<i>Valonia agagropila</i>	7
		<i>Vangueria abyssinica</i> , 269; canthioides, 269; concolor, 269; edulis, 9, 269; euonymoides, 269; infausta, 262; membranacea, 269; pauciflora, 269; senegalensis, 269; tetraphylla, 269; umbellulata, 269; velutina, 269.	
		<i>Vangueria</i> , tribe of <i>Rubiaceæ</i>	268
		<i>Vaucheria</i>	6
		<i>Vegetable Poisons</i> , on the nature and mode of use of	55
		<i>Velezia rigida</i>	354
		<i>Verbascum calycinum</i> , 582; <i>Cossonianum</i> , <i>Ball</i> , 583; <i>Hænscleri</i> , 584; <i>Hookerianum</i> , <i>Ball</i> , 584; <i>Lychnitis</i> , 583; <i>sinuatum</i> , 583, 740.	
		<i>Verbena officinalis</i>	607
		— <i>supina</i>	607
		<i>Veronica agrestis</i> , 601; <i>Anagallis</i> , 599; <i>arvensis</i> , 600; <i>Beccabunga</i> , 599; <i>cuneifolia</i> , var. <i>atlantica</i> , <i>Ball</i> , 599; <i>Cymbalaria</i> , 601; <i>didyma</i> , 601; <i>hedrifolia</i> , 601; <i>Tournefortii</i> , 600; <i>triphylls</i> , 600.	
		<i>Verrucaria æthiobola</i> , 221; <i>albescens</i> , var. <i>subochracea</i> , 227; <i>cinchonæ</i> , 220; <i>congestula</i> , <i>Strn. sp.</i>	

	Page		Page
<i>n.</i> , 221; fumosaria, <i>Leight. n. sp.</i> , 2; neottizans, <i>Leight. n. sp.</i> , 2; paramera, <i>Nyl. sp. n.</i> , 227; per-tenuis, <i>Leight. n. sp.</i> , 2; rude-rella, <i>Nyl. sp. n.</i> , 217.		159; rosca, 158; spicata, 159; strictiflora, 158.	
<i>Viburnum Tinus</i>	481	Weihea, section of Geissorhiza, de-fined	95
<i>Vicia altissima</i> , 437; atropurpu-rea, 438; biflora, 437; Durandi, 438; elegans, 437; erviformis, 438; glauca, 437; Lens, 438; lutea, 439, var. hirtca, 439, var. muricata, 439, var. nitida, <i>Ball</i> , 439; onobrychoides, 436, var.?, 436; pubescens, 438; sativa, 438, var. abyssinica, 438, var. angustifolia, 438, var. cordata, 439.		Whangee, note on the uses of . . .	1
<i>Vicusseuxia</i> , characters of . . .	77, 132	Witsenia, characters of . . .	74, 108
— aristata, 133; fugax, 133; glaucopis, 133; mutila, 133; pa-ronia, 133; spiralis, 133; tenuis, 133; triuspis, 133; tripetaloi-des, 133; unguicularis, 133; vil-losa, 133.		— maura	108
<i>Vinca media</i>	565	<i>Woodsia mollis</i>	34
<i>Viola arboreseens</i> , 349; odorata, 349; Tezensis, <i>Ball</i> , 350.		— obtusa	34
<i>Virecta angustifolia</i> , 264; multi-flora, 264; procumbens, 264; setigera, 264.		<i>Woodwardia radicans</i>	28
<i>Vitex Agnus Castus</i>	607	— japonica	28
<i>Vitis cirrhosa</i> , 180; macropus, 180; vinifera, 392.		<i>Xanthium strumarium</i>	503
<i>Vittaria elongata</i> , 83, 205; falcata, 33; lineata, 33, 205; stipitata, 37.		—, var. antiquorum	503
<i>Volutarella erupinoides</i> , 525; leu-cantha, 526; Lippii, 525; mu-ricata, 526.		<i>Xeranthemum modestum</i>	518
<i>Watsonia</i> , characters of . . .	78, 157	<i>Xerotus tener</i> , 52; Bertierii, 53.	
— aletroides, 157; angusta, 157; brevifolia, 158; cylindrica, 157; densiflora, 158; humilis, 158; Meriana, 158, var. coccinea, 158, var. dubia, 158, var. iridifolia, 158, var. rosco-alba, 158; marginata, 158, var. minor, 158; plantaginea, 158, var. juncea, 159; punctata, 158, var. filifolia, 159, var. Zeyheri,		Xiphion, characters defined . . .	76, 122
		— Aitelhisoni, 124, var. chrysan-thum, 124; Aucheri, 124; cau-casicum, 124; Danfordiæ, 124; filifolium, 123, 675, var. interme-dium, 675, var. latifolium, 123; foetidissimum, 676; Fontanesii, 123; Histrio, 123; Junceum, 123; latifolium, 122; palæsti-num, 124; persicum, 123; pla-nifolium, 123; Pseudacorus, 676; reticulatum, 123, var. Kre-lagei, 123; Sisyrinchium, 675, 740; Stocksii, 124; tingitanum, 123, 675; vulgare, 122, var. lu-sitanicum, 122, var. serotina, 122, var. sordidum, 122, var. spectabilis, 122.	
		Xiphionideæ, tribe of Irids, charac-ters of	76
		<i>Xylaria hypoxylon</i>	48
		— piliceps, <i>B.</i>	44
		<i>Zanthoxylon</i>	9
		— paniculatum, <i>Balf. fil. sp. nov.</i>	11, 12
		<i>Zizyphus Lotus</i>	391, 739
		<i>Zonaria nigrescens</i>	6
		<i>Zostera marina</i>	683
		<i>Zygnema tetraspermum</i> , <i>Reinsch, n. sp.</i> (Pl. VI.)	243
		<i>Zygoon graveolens</i>	267
		<i>Zygophyllum Fontanesii</i>	381

END OF THE SIXTEENTH VOLUME.



